**Adaptation and competition in deteriorating environments**

Romana Limberger[1,2,3*] and Gregor F. Fussmann[2]

1 Research Department for Limnology, University of Innsbruck, Mondsee, Austria

2 Department of Biology, McGill University, Montreal, Quebec, Canada

3 Department of Evolutionary Biology and Environmental Studies, University of Zurich,

Zurich, Switzerland

E-mail addresses:

romana.limberger@ieu.uzh.ch

gregor.fussmann@mcgill.ca

* Corresponding author:

Romana Limberger

Department of Evolutionary Biology and Environmental Studies

Winterthurerstrasse 190, 8057 Zurich, Switzerland

e-mail:romana.limberger@ieu.uzh.ch

**Abstract**

Evolution might rescue populations from extinction in changing environments. Using

experimental evolution with microalgae, we investigated if competition influences adaptation

to an abiotic stressor, and vice versa, if adaptation to abiotic change influences competition. In

a first set of experiments, we propagated monocultures of five species with and without

increasing salt stress for ~180 generations. When assayed in monoculture, two of the five

species showed signatures of adaptation, that is, lines with a history of salt stress had higher

population growth rates at high salt than lines without prior exposure to salt. When assayed in

mixtures of species, however, only one of these two species had increased population size at

high salt, indicating that competition can alter how adaptation to abiotic change influences

population dynamics. In a second experiment, we cultivated two species in monocultures and

in pairs, with and without increasing salt. While we found no effect of competition on

adaptation to salt, our experiment revealed that evolutionary responses to salt can influence

competition. Specifically, one of the two species had reduced competitive ability in the no-salt

environment after long-term exposure to salt stress. Collectively, our results highlight the

complex interplay of adaptation to abiotic change and competitive interactions.

**Introduction**

Abiotic environmental change can render populations maladapted, leading to their decline, and potentially, extinction [1-3]. Population decline may be slowed or arrested by adaptation [4, 5], that is, by phenotypic change that improves fitness and has a genetic basis [6]. Both laboratory and field-based studies have demonstrated that organisms can adapt rapidly to a variety of abiotic drivers, including acidification [7], salt stress [8], pollution [9], nutrient limitation [10], and warming [11, 12]. Experiments have also demonstrated evolutionary responses to biotic drivers, such as competition [13-15] and predation [16]. However, only few experiments have explored the interactive effects of abiotic change and biotic interactions on adaptation [16-19]. Yet, species interactions might influence (i) whether populations adapt to abiotic change [16-18], and (ii) whether adaptation increases population size and persistence [20]. Moreover, within communities, the ecological consequences of adaptation to abiotic change could go beyond effects on population size, and include effects on species interactions and community dynamics [21]. This potentially complex interplay of biotic interactions and adaptation to abiotic change limits our understanding of whether evolution can contribute to the maintenance of diversity in changing environments.

Theory suggests several mechanisms whereby interspecific competition can influence adaptation to deteriorating environmental conditions. First, competition can constrain adaptation to environmental change by reducing population size, and thus the amount of standing genetic variation [22] and the supply of beneficial mutations [23]. Second, competition can reduce the time available for adaptation as species are outcompeted by competitors with broader environmental tolerance [24]. Third, competition can alter the strength and direction of selection [25]; depending on whether competition and environmental change impose selection on a trait in the same or in the opposite direction, competition can speed up or hinder adaptation to environmental change [25]. Finally, negative genetic

correlations between traits that mediate competition and stress tolerance could constrain

adaptation in environments with both biotic and abiotic selective agents [26]. Only few

experiments have analysed evolutionary responses to a factorial manipulation of abiotic

change and biotic interactions to test these theoretical predictions [16-19]. Moreover, most of

these empirical tests investigated biotic interactions other than interspecific competition [16,

17], or environmental amelioration rather than deterioration [17, 18, but see 19]. In our study,

however, we focused on interspecific competition, a key biotic interaction, and its interplay

with adaptation to deteriorating conditions, that is, to environmental changes that have a

negative effect on fitness (e.g. acidification, pollution).

Competition could not only alter how species adapt to abiotic change, but could also

modulate the effect of adaptation on population dynamics. When competition alters the fitness

landscape generated by abiotic change, adaptation of a focal species in isolation may not

translate into increased population size in a community context. Accordingly, an experiment

that manipulated $CO_2$ and diversity of a plant community found adaptation to elevated $CO_2$

only when plants were assayed in the same community context in which selection had

occurred [18]. Similarly, an experiment with a marine alga revealed that adaptation to $CO_2$

enrichment in isolation was not associated with increased population size within communities

[20]. The scenario of adaptation in isolation followed by competition might seem artificial,

but it is a plausible scenario in a spatial context. Species could adapt to changes in their local

environment before immigration of better-adapted competitors from the regional species pool

[27, 28]. Such changes in competitive interactions could prevent that adaptation of a focal

species to abiotic change translates into increased population size.

When investigating the ecological consequences of adaptation to an abiotic stressor, the

focus is often on effects on population size and persistence, but within communities,

adaptation of a focal species to the local environment may affect interactions with other

species [29-31]. Evolution can influence community dynamics directly, by altering traits that

underlie species interactions, and indirectly, by influencing population dynamics [32, 33]. In a

scenario with an abiotic stressor as the selective agent, adaptation could influence species

interactions if traits that confer adaptation to the stressor are correlated with traits that affect

biotic interactions. Some traits may even simultaneously mediate stress tolerance and species

interactions. For example, in *E. coli* reduced membrane permeability increases resistance to

antibiotics but decreases the ability to take up nutrients, such that adaptation to antibiotics can

result in lower competitive ability [34]. In addition to such trait-mediated effects, adaptation

of a focal species to an abiotic stressor might influence biotic interactions by increasing its

population size. Despite this range of possibilities of how adaptation to an abiotic stressor

could affect biotic interactions, its ecological consequences have rarely been considered in a

community context.

Here we used experimental evolution with freshwater algae to investigate if adaptation

to abiotic change influences competitive interactions, and, vice versa, if competition

influences adaptation to deteriorating environmental conditions. We used salt as a stressor

because it has a negative effect on the fitness of freshwater algae, and because it can easily be

manipulated without confounding changes in other abiotic factors. We did not intend to

mimic a specific environmental change scenario observed in natural systems; rather we

wanted to investigate conceptual questions about the interplay of competition and adaptation

to an abiotic stressor. We conducted two sets of experiments (Fig. 1): in the first experiment

we investigated if adaptation to salt in monoculture translated into increased population size

(i.e. cell number) within salt-stressed mixtures of species, to test the hypothesis that

competition can alter the effect of adaptation on population dynamics. In the second

experiment we tested the hypotheses that interspecific competition constrains adaptation to

salt stress, and that a history of salt stress leads to increased competitive ability at high salt.

## Material and methods

*Model organisms*

We tested our hypotheses using six species of microalgae from three different taxonomic
groups: the cyanobacteria *Synechococcus leopoliensis* (Canadian Phycological Culture Centre
(CPCC) 102) and *Anabaena variabilis* (CPCC 105), the diatoms *Navicula pelliculosa* (CPCC
552) and *Nitzschia palea* (CPCC 160), and the chlorophytes *Pseudokirchneriella subcapitata*
(CPCC 37) and *Scenedesmus acutus* (CPCC 10). We chose these six species because they
were available as axenic (i.e. bacteria-free) cultures and because we knew from previous
experiments that they grow well under the same experimental conditions [35, 36]. The algae
were cultivated in Bold's basal medium [37], which was modified by adding silicate (0.58 g
$L^{-1}$ of $Na_2SiO_3$) and vitamins [38] to allow growth of diatoms. In all experiments, the algae
were cultivated at 25 °C with light continuously provided at 100 $\mu E\ m^{-2}\ s^{-1}$. Cultures were
axenic at the start of the experiment and propagated under sterile conditions. At the end of the
experiment, we plated a subset of the cultures (18 cultures of *Anabaena* and *Scenedesmus*) on
agar and observed no bacterial contaminations. We did not make cultures isogenic at the start
of the experiment; the observed evolutionary responses could thus have resulted either from
*de novo* mutations or from sorting of standing genetic variation.

In a preliminary experiment, we measured salt tolerance of the six species (Fig. S1),
and found a low limit of salt tolerance in *Nitzschia* and *Pseudokirchneriella* (~6 g NaCl $L^{-1}$),
intermediate salt tolerance in *Anabaena*, *Scenedesmus* and *Navicula* (limit: ~14 g NaCl $L^{-1}$),
and high salt tolerance in *Synechococcus* (limit: > 20 g NaCl $L^{-1}$).

*Adaptation to salt in monoculture*

In a serial transfer experiment, we propagated monocultures of the six species with and
without increasing concentrations of NaCl (Fig. 1, Fig. S2). The algae were cultured in 125-

ml glass flasks filled with 50 ml of Bold's medium and continuously shaken at 250 rpm.

Every 3.5 days, we transferred 1 ml of culture to 50 ml of fresh medium for a total of 37

transfers (~ 185 generations; one transfer corresponding to five generations). For each of the

six species, three replicate control lines were cultivated without addition of salt and three

replicate selection lines were cultivated with increasing salt. We increased the salt

concentration in the selection lines by 0.25 g $L^{-1}$ every transfer until reaching 4 g NaCl $L^{-1}$ at

transfer 16 (Fig. S2). We maintained this salt concentration for 10 transfers and then

continued the increase in salt until reaching the final concentration of 6 g NaCl $L^{-1}$ at transfer

34, which we maintained for three more transfers. This salt concentration had a distinctly

negative effect on the growth rates of species with lowest salt tolerance but still allowed

persistence of all selection lines. A further increase in salt concentration would have probably

resulted in extinction of species with low tolerance and would thus have reduced the number

of species in the mixture assay.

The goal of the salt treatment was to simulate an environmental change scenario as it

would be experienced by the members of a community; we thus exposed all species to the

same change in environmental conditions rather than exposing each species to its respective

limit of salt tolerance. Because the species varied in salt tolerance (Fig. S1), they differed in

how strongly they were stressed by the salt treatment. Similarly, because the species varied in

growth rates and population sizes, they differed in the potential to adapt to salt. Such species-

specific differences are a property of any community and we thus made no attempt to reduce

these differences (e.g. by transferring equal cell numbers rather than equal volume).

After 37 transfers (~ 185 generations), we measured population growth rate of each

line in the two final salt environments (0 and 6 g NaCl $L^{-1}$) to test if the species had adapted to

salt stress. Before starting the growth assay, we cultivated all lines at identical environmental

conditions for three transfer cycles, that is two cycles in the ancestral environment (0 g NaCl

$L^{-1}$) and one cycle in the assay environment (0 and 6 g NaCl $L^{-1}$), to reduce non-evolutionary

effects and pre-condition the lines to the assay conditions. We then inoculated flasks

containing 50 ml medium of either 0 or 6 g NaCl $L^{-1}$ with 1 ml of acclimated culture and

measured absorbance through time. Depending on the time of individual cultures to reach

carrying capacity, we measured absorbance over 2.5 to 7.5 weeks with 23 to 34

measurements. At each measurement, we removed 200 µl of culture from each flask into 96-

well plates and measured absorbance at 660 nm on an optical plate reader (Synergy-HT,

BioTek, Winooski, VT). As done in other studies [8, 35], we calculated population growth

rates (see below) from raw absorbance values, i.e., we did not use standard curves to

transform absorbance to cell density. We thus did not take into account that changes in cell

size in response to salt might have influenced absorbance.

*Mixture assay*

To test if a history of salt stress can influence population size of species within salt-stressed

mixtures of species, we assembled mixtures either of control or of selection lines, transplanted

the mixtures into the two final salt environments of the selection experiment, and quantified

species abundances over four transfers. Prior to the mixture assay, the lines had been

cultivated for 37 transfers with or without increasing salt, followed by two transfers in the

ancestral environment (see above). We assembled mixtures with equal biovolume of the six

species. To this end, we determined species abundances in the monocultures by counting

Lugol-fixed samples with an inverted microscope, and then converted abundances to

biovolume using cellular biovolume estimates based on measurements of 20 individuals per

species (Table S1) [39]. We assembled three replicate mixtures of control lines and three

replicate mixtures of selection lines. From each of the six mixtures, we inoculated two flasks,

one with a salt concentration of 0 and one of 6 g NaCl $L^{-1}$. Every four days, we transferred 1

ml of the mixtures into 50 ml of fresh medium, with the flasks continuously shaken at 250

rpm. Over the course of four transfers, we sampled the mixtures at each transfer, fixed the

samples with Lugol's solution, and counted species abundances with an inverted microscope.

We used different approaches to quantify population growth rate in the monoculture

assay and population size in the  mixture assay (absorbance and abundance counts,

respectively) because high-throughput measurement of absorbance with a plate reader

allowed us to collect a dense time-series of absorbance data, while microscopical counts were

necessary to differentiate species in the mixture assay.

*Adaptation in monoculture vs. adaptation in pairs*

To test if competition can influence adaptation to salt stress, we propagated monocultures and

pairs of *Anabaena* and *Scenedesmus* with and without increasing salt for 46 transfers (~ 230

generations), then re-isolated the two species and quantified their competitive ability with and

without salt (Fig. 1). We chose the combination of *Anabaena* and *Scenedesmus* because in

other species combinations the inferior competitor was outcompeted quickly both with and

without increasing salt. In our experimental setup (i.e. a serial transfer experiment with

nutrient-replete conditions), competition is mediated by the difference in growth rates, with

the species with faster growth rate expected to outcompete the species with slower growth

rate. The monocultures and pairs of the two species were part of the selection experiment

described above and experienced the same changes in salt concentration except that cultures

spent nine more transfers at the final salt concentration (Fig. S2). To quantify how strongly

competition affected the abundances of the two species during the selection experiment, we

sampled the pairs and monocultures of *Anabaena* and *Scenedesmus* ten times over the course

of the selection experiment and microscopically determined the abundances of the two species

in Lugol-fixed samples.

After 46 transfers in the selection experiment, we re-isolated *Anabaena* and

*Scenedesmus* by plating the cultures on agar and isolating individual colonies. Monocultures

were treated the same way as the pairs. Individual colonies were resuspended in liquid Bold's

medium on well plates, and microscopically checked for contamination after a week of

growth. We then merged eight clean isolates per sample to one culture. The two isolation

steps were conducted in medium without addition of NaCl. We then pre-conditioned the

cultures at the two assay environments (0 and 6 g NaCl L$^{-1}$) for two transfer cycles. Hence,

before starting the growth and competition assays, the cultures were cultivated under identical

environmental conditions for four cycles (two isolation cycles without salt, two transfer cycles

in the assay environment) to reduce non-evolutionary effects.

The pre-conditioned cultures were used to start competition assays to test if competition

history and salt history influenced competitive ability. To this end, we assembled pairs of

*Anabaena* and *Scenedesmus* in the two final salt environments and determined changes in

biomass of the two species through time. We conducted two competition assays that differed

in how we assembled the pairs (Fig. 1). First, we assembled pairs of co-evolved *Anabaena*

and *Scenedesmus*, i.e. lines that had the same competition and salt history. However, because

pairs differed in both competitors in this assay (Fig. 1), differences among treatments could

not unequivocally be attributed to one of the two species. We thus conducted a second

competition assay where we competed each line of *Anabaena* with the same reference line of

*Scenedesmus*, and each line of *Scenedesmus* with the same reference line of *Anabaena*. As

reference line, we randomly selected one of the three replicate control lines (i.e. competition

history: monoculture, salt history: no salt). The second competition assay was started three

weeks after the first competition assay; cultures were stored meanwhile as monocultures at

their respective salt selection environment and then acclimated for one cycle to the salt assay

environment.

In the competition assays, we measured chlorophyll a concentrations of the two

competitors as proxy for biomass using a FluoroProbe (bbe Moldaenke, Kiel-Kronshagen,

Germany), a fluorometer which can discriminate among algal groups [40]. Before the start of

the assay, we calibrated the FluoroProbe with cultures of *Anabaena* and *Scenedesmus* with

known chlorophyll a concentrations. We started the competition assay with equal chlorophyll

a concentrations (2 µg) of both species in flasks filled with 50 mL of Bold's medium. We

transferred the cultures every 3.5 days for ten transfers in the first competition assay and for

four transfers in the second competition assay and measured chlorophyll a concentrations of

both species at each transfer.

*Data analysis*

All analyses were computed with R version 3.6.2 [41]. We estimated population

growth rates (absorbance day$^{-1}$) in monoculture assays by fitting logistic growth curves to

absorbance vs. time curves using the R package *grofit* [42]. Growth curves could not be fit

when absorbance did not increase; in this case (the three control lines of *Nitzschia* at 6 g NaCl

L$^{-1}$) growth rate was set to zero. Some of the growth curves of *Anabaena* were difficult to fit

because this species sometimes grew in clumps at high salt, which led to unreliable

absorbance measurements. We thus do not present monoculture growth rates of *Anabaena*.

To investigate treatment effects on growth rates, we computed two-way ANOVAs for

each of the five species using assay environment and salt history as explanatory variables. It

would have been preferential to include 'line' as a random effect because each of the six lines

(three control lines, three selection lines) was assayed in two environments (with and without

salt, respectively). However, computation of linear mixed models with 'line' as a random

effect often resulted in singular fits because of zero or almost zero variation among lines. We

thus dropped the random effect and computed ANOVAs instead. In case of a significant

interaction, we additionally computed two separate ANOVAs for each assay environment

with salt history as the explanatory variable. We used the function *lm* to compute ANOVAs

and the function *Anova* in the *car* package to calculate P-values.

We used the same approach (i.e. ANOVA) to test treatment effects on abundances of

species assayed within mixtures, using abundance averaged over the four transfers of the

mixture assay as response variable, and assay environment and salt history as explanatory

variables. Abundances were ln-transformed prior to analyses, except for abundances of

*Pseudokirchneriella* which remained untransformed due to better fit with model assumptions.

To test whether competition affected adaptation of *Anabaena* and *Scenedesmus* to salt, we

computed three-way ANOVAs for each of the two species and investigated the effects of

competition history, salt history and assay environment on relative chlorophyll a (arcsine

square-root transformed) in the competition assay. In all our analyses, we did not use model

selection to identify important predictor variables, but included all factors manipulated in the

experiment as explanatory variables in the model, that is, model structure followed the

experimental design.

**Results**

*Adaptation of monocultures*

After 185 generations of growth with and without increasing salt, respectively, we detected

adaptation to salt in two of five species, *Nitzschia* and *Pseudokirchneriella* (Fig. 2, Table S2),

that is, lines with a history of salt stress had higher population growth rates at 6 g NaCl L$^{-1}$

than lines without a history of salt stress (ANOVA: *Nitzschia*: $F_{1,4}= 16.19$, $P= 0.016$;

*Pseudokirchneriella*: $F_{1,4}= 77.70$, $P= 0.001$). In the no-salt assay environment, however, salt

history had no effect on the growth rates of these two species (ANOVA: *Nitzschia*: $F_{1,4}= 0.64$,

$P= 0.467$; *Pseudokirchneriella*: $F_{1,4}= 2.78$, $P= 0.171$). Among the remaining three species,

*Scenedesmus* and *Navicula* had reduced growth rates in the high-salt assay environment

irrespective of salt history, while *Synechococcus* was unaffected by both salt history and assay

environment (Fig. 2, Table S2).

*Mixture assay*

When assaying mixtures of either control or selection lines, we found that the effect of salt

history varied among species and depended on assay environment (Fig. 3 a-c, Table S3). Most

species declined in abundance over the four transfers of the assay, except for *Synechococcus,*

which increased (Fig. S3). Salt history affected the abundance of three species: *Nitzschia*,

*Pseudokirchneriella*, *Scenedesmus* (Fig. 3 a-c, Table S3). In *Nitzschia*, lines with a history of

salt stress had lower average abundance than control lines when assayed without salt (58%

lower abundance; $F_{1,4}= 25.97$, P= 0.007). At high salt, however, the effect of salt history on

average abundance of *Nitzschia* was not significant ($F_{1,4}= 4.54$, P= 0.10) and control and

selection lines declined similarly fast (Fig. S3a).  In *Pseudokirchneriella*, selection lines had

lower average abundance than control lines in the no-salt assay environment (24% lower

abundance; $F_{1,4}= 24.11$, P = 0.008), but 2.5 times higher abundance than control lines in the

high-salt assay environment ($F_{1,4}= 14.90$, P = 0.018). However, these treatment effects on

*Pseudokirchneriella* did not persist until the end of the mixture assay: within four transfers,

*Pseudokirchneriella* went extinct or declined to low abundance irrespective of salt history and

assay environment (Fig. S3b). In *Scenedesmus*, a history of salt stress had no effect on

average abundance in the no-salt assay environment ($F_{1,4}= 1.73$, P = 0.258), but a positive

effect at high salt (1.7 times higher abundance; $F_{1,4}= 11.86$, P = 0.026). Among the remaining

three species, abundance of *Anabaena* was negatively affected by salt irrespective of salt

history (Fig. S3c, Table S3), while *Navicula* and *Synechococcus* were unaffected by both salt

history and assay environment (Fig. 3d, e, Table S3).

*Adaptation in monoculture vs. adaptation in pairs*

Interspecific competition strongly influenced species abundances during the selection experiment, but did not lead to any evolutionary responses. In the selection experiment, *Anabaena* reduced the abundance of *Scenedesmus* as long as salt concentration was low (Fig. S4b). However, at the final salt concentration of 6 g NaCl $L^{-1}$ the competitive hierarchy reversed and *Anabaena* was negatively affected by *Scenedesmus* (Fig. S4a). This change in competitive hierarchy in response to salt stress was also apparent in the competition assay: *Anabaena* dominated without salt (Fig. 4a), while *Scenedesmus* dominated at high salt (Fig. 4b), irrespective of competition history.

The two competition assays revealed that lines of *Anabaena* with a history of salt stress had reduced competitive ability in the no-salt assay environment. When assayed with co-evolved competitors, high-salt lines of *Anabaena* had lower relative chlorophyll a than no-salt lines in the no-salt assay environment, and consequently high-salt lines of *Scenedesmus* had higher relative chlorophyll a than no-salt lines (salt history × assay: $F_{1,16} = 13.47$, P = 0.002; Table S4, Fig. S5). Because both competitors differed among treatment combinations, this experiment did not allow detecting if the observed pattern was due to altered competitive ability of *Anabaena* or of *Scenedesmus*. However, when assayed in competition with a reference line, only *Anabaena* was affected by salt history, again with lower relative chlorophyll a of high-salt than no-salt lines in the no-salt assay environment (salt history × assay: $F_{1,16} = 5.78$, P = 0.029, Table S5, Fig. 4a). As a consequence, in the no-salt assay environment the biomass (i.e. chlorophyll a concentration) of the competing reference line varied by a factor of 1.9 depending on the evolutionary history of *Anabaena* (Fig. S6a).

**Discussion**

With our experiments we addressed a range of scenarios that can arise from the interplay of

competition and adaptation to an abiotic stressor. In the first experiment, we found that

competition can alter how adaptation to an abiotic stressor influences population dynamics

(Fig. 2, 3): effects of salt history on population growth rate in monoculture did not necessarily

lead to increased population size within communities. Conversely, some evolutionary effects,

specifically, reduced population size of selection lines without salt, appeared only within

mixtures of species, not in monoculture. In the second experiment, we found no effect of

competition on adaptation to salt stress (Fig. 4). However, salt history influenced competition:

after selection at increasing salt, one of two species had reduced competitive ability without

salt (Fig. 4a). Collectively, our results highlight that ecological and evolutionary processes

interactively shape communities in changing environments, but that these eco-evolutionary

effects can be diverse and species-specific.

Species varied in their evolutionary response to salt stress. The two species with the

lowest limit of salt tolerance adapted to salt (Fig. 2a, b), while species that were less

negatively affected by salt showed no sign of adaptation (Fig. 2 c-e). This finding is in line

with the prediction that species with narrow tolerance of a stressor should evolve more than

species with broader tolerance because they experience stronger selection pressure [43].

Species-specific differences in genetic variance and covariance could be additional reasons

for variation among species in the rate of evolution [43, 44]. Accordingly, an experiment that

investigated thermal evolution in three phytoplankton species found largest fitness

improvements in the species with highest population density and the least complex genome

[45]. Given that only two species adapted in our experiment, our sample size is not large

enough to evaluate if species-specific differences other than ecological tolerance influenced

the evolutionary response to salt. Such variation among species in the ability to adapt to

environmental change could add further complexity to adaptation in a community context.

Our experiments suggest that competition can modulate how adaptation to abiotic

change influences population dynamics. When assayed at high salt in monoculture, two

species (*Nitzschia* and *Pseudokirchneriella*) showed signatures of adaptation, that is, higher

population growth rate of selection lines than control lines (Fig. 2a, b). When assayed at high

salt within mixtures, however, only one of these two species (*Pseudokirchneriella*) had higher

population size of selection than control lines (Fig. 3b), and this positive effect of salt history

was only transient (Fig. S3b). In contrast, control and selection lines of *Nitzschia* had similar

population size and declined equally fast within salt-stressed mixtures (Fig. 3a, Fig. S3a).

Similarly, adaptation of a marine alga to acidification was not associated with increased

population size in acidified communities [20]. Such a pattern can emerge when competition

changes the fitness landscape [18]; thereby adaptation of a species to abiotic change improves

its fitness only in the same community context in which it has evolved [18]. In our mixtures,

low competitive ability of the two species that adapted to salt might have also contributed to

this pattern. *Nitzschia* and *Pseudokirchneriella* were the weakest competitors in the

community and declined to extinction or to low abundance irrespective of salt history and salt

environment (Fig. S3). It remains to be determined if effects of adaptation on population

dynamics are less contingent on community context in stronger competitors. In addition to

low competitive ability, low starting abundance could have contributed to the fast extinction

of *Nitzschia*: this species was the largest in our community (Table S1) and was thus

inoculated with the lowest abundance among the six species. Taken together, the positive

effect of adaptation to abiotic change on population size may be reduced when community

context changes; such a scenario could result from immigration of novel competitors in

response to environmental change [27, 28, 46].

We also found evolutionary effects that appeared only within mixtures. In *Nitzschia* and

*Pseudokirchneriella,* adaptation to salt was associated with costs (i.e. reduced population size)

in the no-salt environment when these species were assayed within mixtures (Fig. 3a, b), but

not when assayed in isolation (Fig. 2a, b). A possible mechanism underlying such a pattern is

a trade-off between traits that mediate stress tolerance and competition [34]. We did not

measure traits and can thus only speculate about potential mechanisms. Tolerance of high salt

concentrations is conferred by the ability to efficiently export ions ($Na^+$ and $Cl^-$) and

accumulate compatible solutes, i.e. organic molecules that maintain osmotic balance without

interfering with metabolism [47]. High energetic costs for ion export and for synthesis of

compatible solutes [48] may result in reduced competitive ability. However, when adaptation

to an abiotic stressor occurs within communities, rather than in isolation as in our experiment,

reduced competitive ability might not evolve. Our experiment thus simulates a scenario in

which environmental change is followed by a change in community context, which could

result from immigration of novel competitors or from changes in local community

composition due to species-specific differences in environmental tolerance. To elucidate if

adaptation to an abiotic stressor is associated with changes in traits that mediate biotic

interactions, a trait-based approach would be an interesting avenue for future experiments.

Competition had no effect on adaptation to salt stress (Fig. 4). We had predicted that

competition during the selection experiment would constrain adaptation to salt because of

reduced population size and/or increased genetic constraints [23, 26]. Alternatively,

competition could facilitate adaptation to abiotic stress by increasing the strength of selection

[25]. The two species that we used to test this question, *Anabaena* and *Scenedesmus*, adapted

to salt neither in monoculture nor in pairs. Similarly, an experiment that manipulated $CO_2$

enrichment and competition in a selection experiment with two algae species found neither

adaptation to $CO_2$ nor to competition [19]. In our experiment, low selection pressure could

explain why the monocultures did not adapt to salt, given that the two species had not reached

their limit of salt tolerance in the selection experiment. However, the combination of salt and

competition in the selection experiment drove the inferior competitor to the brink of

extinction (Fig. S4), suggesting that the inferior competitor experienced strong selection

pressure to evolve faster growth rates. Yet, we did not find any effect of competition history

on adaptation. One possible explanation is that the competitive hierarchy in flasks with

increasing salt changed shortly before the end of the selection experiment when the final salt

concentration had been reached, from dominance of *Anabaena* to dominance of *Scenedesmus*.

Maybe relaxed selection pressure on *Scenedesmus* towards the end of the experiment and only

a short phase of strong selection on *Anabaena* explain why competition had no effect on the

evolutionary response of these two species.

425            Evolutionary change in response to salt influenced competitive ability in one of two

species (Fig. 4). We had predicted that a history of salt stress would lead to increased

competitive ability at high salt. In contrast, however, strains of *Anabaena* with a history of

salt stress had reduced competitive ability in the no-salt assay environment. Such a pattern is

an indication of conditionally deleterious mutations [49], i.e. mutations that are neutral in the

local environment and deleterious in an alternative environment. Because we have neither

molecular nor trait data, we cannot elucidate potential mechanisms behind the observed mal-

adaptation in the ancestral environment. The evolutionary response of *Anabaena* was

independent of competition history and of whether fitness was assayed in competition with a

co-evolved or a naïve competitor (Fig. 4, Fig. S5). The ecological effect of this evolutionary

response to salt is also illustrated by the response of the competing reference line: in the no-

salt assay, the chlorophyll a concentration attributable to the reference line of *Scenedesmus*

was 1.9 times higher when competing with selection lines of *Anabaena* than when competing

with control lines (Fig. S6a). Overall, our results suggest that evolutionary responses to

abiotic change can influence biotic interactions, such that populations with different histories

of abiotic stress might vary in their effects on communities and ecosystems.

The emerging field of eco-evolutionary dynamics investigates the effects of evolution

on populations, communities, and ecosystems [6]. Research on adaptation to abiotic change

usually explores the effect of evolution on population dynamics [4, 7, 11], while studies that

investigate the effect of evolution on species interactions often focus on biotic selective agents

[50-52]. Here we showed that evolutionary responses to an abiotic selective agent can alter

biotic interactions (Fig. 4a, Fig. S6a). Moreover, we found that competition can modulate the

effect of adaptation on population dynamics (Fig. 2, 3), suggesting that adaptation to abiotic

change may not enhance performance of a species when community context changes [18] e.g.

because of immigrating competitors [28]. We have observed a range of complexities arising

from adaptation in a multi-species context despite using a fairly simple model system. Higher

diversity and trophic structure of natural systems will pose an even greater challenge to

understanding eco-evolutionary dynamics in changing environments [33, 53]. Environmental

fluctuation is another complexity of natural systems that we were not able to address in our

experiment. Temporary environmental amelioration could promote adaptation to abiotic

change by increasing population size and thus the supply of beneficial mutations [11], but

could also reduce the potential for adaptation by relaxing selection pressure [54]. Taking these

complexities into account will be crucial in future work to further increase our understanding

of adaptation to abiotic change in a community context.

**Acknowledgements**

We thank Etienne Low-Décarie and Graham Bell for input on the design and analysis of

the experiment, and Yinci Yan for help with the implementation of the experiment.

**Funding**

This work was supported by an Erwin-Schrödinger scholarship of the Austrian Science

Fund (FWF), grant no. J3265-B16 to R.L., and an NSERC Discovery Grant to G.F.F.

**References**

1.    Ceballos G., Ehrlich P.R. 2002 Mammal population losses and the extinction crisis.

*Science* **296**, 904-907. (doi:10.1126/science.1069349).

2.    Both C., Bouwhuis S., Lessells C.M., Visser M.E. 2006 Climate change and

population declines in a long-distance migratory bird. *Nature* **441**, 81-83.

(doi:10.1038/nature04539).

3.    Stuart S.N., Chanson J.S., Cox N.A., Young B.E., Rodrigues A.S.L., Fischman D.L.,

Waller R.W. 2004 Status and trends of amphibian declines and extinctions worldwide.

*Science* **306**, 1783-1786. (doi:10.1126/science.1103538).

4.    Bell G., Gonzalez A. 2009 Evolutionary rescue can prevent extinction following

environmental change. *Ecol Lett* **12**, 942–948. (doi:10.1111/j.1461-0248.2009.01350.x).

5.    Gomulkiewicz R., Holt R.D. 1995 When does evolution by natural selection prevent

extinction? *Evolution* **49**, 201-207. (doi:10.2307/2410305).

6.    Hendry A.P. 2016 *Eco-evolutionary dynamics*. Princeton and Oxford, Princeton

University Press; 416 p.

7.    Lohbeck K.T., Riebesell U., Reusch T.B.H. 2012 Adaptive evolution of a key

phytoplankton species to ocean acidification. *Nat Geosci* **5**, 346-351. (doi:10.1038/ngeo1441).

8.    Lachapelle J., Bell G. 2012 Evolutionary rescue of sexual and asexual populations in a

deteriorating environment. *Evolution* **66**, 3508-3518. (doi:10.1111/j.1558-

5646.2012.01697.x).

9.      Reid N.M., Proestou D.A., Clark B.W., Warren W.C., Colbourne J.K., Shaw J.R.,

Karchner S.I., Hahn M.E., Nacci D., Oleksiak M.F., et al. 2016 The genomic landscape of

rapid repeated evolutionary adaptation to toxic pollution in wild fish. *Science* **354**, 1305-1308.

(doi:10.1126/science.aah4993).

10.     Collins S., de Meaux J. 2009 Adaptation to different rates of environmental change in

*Chlamydomonas*. *Evolution* **63**, 2952-2965. (doi:10.1111/j.1558-5646.2009.00770.x).

11.     Schaum C.E., Buckling A., Smirnoff N., Studholme D.J., Yvon-Durocher G. 2018

Environmental fluctuations accelerate molecular evolution of thermal tolerance in a marine

diatom. *Nat Commun* **9**, 1719. (doi:10.1038/s41467-018-03906-5).

12.     Schaum C.E., Barton S., Bestion E., Buckling A., Garcia-Carreras B., Lopez P., Lowe

C., Pawar S., Smirnoff N., Trimmer M., et al. 2017 Adaptation of phytoplankton to a decade

of experimental warming linked to increased photosynthesis. *Nat Eco Evo* **1**, 0094.

(doi:10.1038/s41559-017-0094).

13.     Lawrence D., Fiegna F., Behrends V., Bundy J.G., Phillimore A.B., Bell T.,

Barraclough T.G. 2012 Species interactions alter evolutionary responses to a novel

environment. *PLoS Biology* **10**, e1001330. (doi:10.1371/journal.pbio.1001330).

14.     Zuppinger-Dingley D., Schmid B., Petermann J.S., Yadav V., De Deyn G.B., Flynn

D.F.B. 2014 Selection for niche differentiation in plant communities increases biodiversity

effects. *Nature* **515**, 108-111. (doi:10.1038/nature13869).

15.     Castledine M., Padfield D., Buckling A. 2020 Experimental (co)evolution in a multi-

species microbial community results in local maladaptation. *Ecol Lett* **23**, 1673-1681.

(doi:10.1111/ele.13599).

16.     Hiltunen T., Cairns J., Frickel J., Jalasvuori M., Laakso J., Kaitala V., Künzel S.,

Karakoc E., Becks L. 2018 Dual-stressor selection alters eco-evolutionary dynamics in

experimental communities. *Nat Eco Evo* **2**, 1974-1981. (doi:10.1038/s41559-018-0701-5).

17.   Collins S. 2011 Competition limits adaptation and productivity in a photosynthetic

alga at elevated $CO_2$. *Proc R Soc B* **278**, 247-255. (doi:10.1098/rspb.2010.1173).

18.   Kleynhans E.J., Otto S.P., Reich P.B., Vellend M. 2016 Adaptation to elevated $CO_2$ in

different biodiversity contexts. *Nat Commun* **7**, 12358. (doi:10.1038/ncomms12358).

19.   Listmann L., Hattich G.S.I., Matthiessen B., Reusch T.B.H. 2020 Eco-evolutionary

interaction in competing phytoplankton: nutrient driven genotype sorting likely explains

dominance shift and species responses to $CO_2$. *Front Mar Sci* **7**.

(doi:10.3389/fmars.2020.00634).

20.   Bach L.T., Lohbeck K.T., Reusch T.B.H., Riebesell U. 2018 Rapid evolution of highly

variable competitive abilities in a key phytoplankton species. *Nat Eco Evo* **2**, 611-613.

(doi:10.1038/s41559-018-0474-x).

21.   Tatters A.O., Schnetzer A., Fu F., Lie A.Y.A., Caron D.A., Hutchins D.A. 2013 Short-

versus long-term responses to changing $CO_2$ in a coastal dinoflagellate bloom: implications

for interspecific competitive interactions and community structure. *Evolution* **67**, 1879-1891.

(doi:10.1111/evo.12029).

22.   Bocedi G., Atkins K.E., Liao J.S., Henry R.C., Travis J.M.J., Hellmann J.J. 2013

Effects of local adaptation and interspecific competition on species' responses to climate

change. *Ann N Y Acad Sci* **1297**, 83-97. (doi:10.1111/nyas.12211).

23.   Johansson J. 2008 Evolutionary responses to environmental changes: How does

competition affect adaptation? *Evolution* **62**, 421-435. (doi:10.1111/j.1558-

5646.2007.00301.x).

24.   de Mazancourt C., Johnson E., Barraclough T.G. 2008 Biodiversity inhibits species'

evolutionary responses to changing environments. *Ecol Lett* **11**, 380–388.

(doi:10.1111/j.1461-0248.2008.01152.x).

25.     Osmond M.M., de Mazancourt C. 2013 How competition affects evolutionary rescue.

*Phil Trans R Soc B* **368**, 20120085. (doi:10.1098/rstb.2012.0085).

26.     Lau J.A., Terhorst C.P. 2015 Causes and consequences of failed adaptation to

biological invasions: the role of ecological constraints. *Mol Ecol* **24**, 1987-1998.

(doi:10.1111/mec.13084).

27.     Urban M.C., De Meester L., Vellend M., Stoks R., Vanoverbeke J. 2012 A crucial step

toward realism: responses to climate change from an evolving metacommunity perspective.

*Evol Appl* **5**, 154-167. (doi:10.1111/j.1752-4571.2011.00208.x).

28.     De Meester L., Vanoverbeke J., Kilsdonk L.J., Urban M.C. 2016 Evolving

perspectives on monopolization and priority effects. *Trends Ecol Evol* **31**, 136-146.

(doi:10.1016/j.tree.2015.12.009).

29.     Gomez P., Paterson S., De Meester L., Liu X., Lenzi L., Sharma M.D., McElroy K.,

Buckling A. 2016 Local adaptation of a bacterium is as important as its presence in

structuring a natural microbial community. *Nat Commun* **7**, 12453.

(doi:10.1038/ncomms12453).

30.     Pantel J.H., Duvivier C., De Meester L. 2015 Rapid local adaptation mediates

zooplankton community assembly in experimental mesocosms. *Ecol Lett* **18**, 992-1000.

(doi:10.1111/ele.12480).

31.     terHorst C.P., Lennon J.T., Lau J.A. 2014 The relative importance of rapid evolution

for plant-microbe interactions depends on ecological context. *Proc R Soc B* **281**, 20141615.

(doi:10.1098/rspb.2014.0028).

32.     Matthews B., Narwani A., Hausch S., Nonaka E., Peter H., Yamamichi M., Sullam

K.E., Bird K.C., Thomas M.K., Hanley T.C., et al. 2011 Toward an integration of

evolutionary biology and ecosystem science. *Ecol Lett* **14**, 690-701. (doi:10.1111/j.1461-

0248.2011.01627.x).

33.    Hendry A.P. 2019 A critique for eco-evolutionary dynamics. *Funct Ecol* **33**, 84-94.

(doi:10.1111/1365-2435.13244).

34.    Phan K., Ferenci T. 2013 A design-constraint trade-off underpins the diversity in

ecologically important traits in species *Escherichia coli*. *Isme J* **7**, 2034-2043.

(doi:10.1038/ismej.2013.82).

35.    Low-Décarie E., Jewell M.D., Fussmann G.F., Bell G. 2013 Long-term culture at

elevated atmospheric CO2 fails to evoke specific adaptation in seven freshwater

phytoplankton species. *Proc R Soc B* **280**. (doi:10.1098/rspb.2012.2598).

36.    Limberger R., Low-Décarie E., Fussmann G.F. 2014 Final thermal conditions override

the effects of temperature history and dispersal in experimental communities. *Proc R Soc B*

**281**, 20141540. (doi:10.1098/rspb.2014.1540).

37.    Bold H. 1949 The morphology of *Chlamydomonas chlamydogama*, sp. nov. *Bull*

*Torrey Bot Club* **76**, 101-108. (doi:10.2307/2482218 ).

38.    Low-Décarie E., Fussmann G.F., Bell G. 2011 The effect of elevated $CO_2$ on growth

and competition in experimental phytoplankton communities. *Glob Change Biol* **17**, 2525–

2535. (doi:10.1111/j.1365-2486.2011.02402.x ).

39.    Hillebrand H., Durselen C.D., Kirschtel D., Pollingher U., Zohary T. 1999 Biovolume

calculation for pelagic and benthic microalgae. *J Phycol* **35**, 403-424. (doi:10.1046/j.1529-

8817.1999.3520403.x).

40.    Catherine A., Escoffier N., Belhocine A., Nasri A.B., Hamlaoui S., Yepremian C.,

Bernard C., Troussellier M. 2012 On the use of the FluoroProbe®, a phytoplankton

quantification method based on fluorescence excitation spectra for large-scale surveys of

lakes and reservoirs. *Water Res* **46**, 1771-1784. (doi:10.1016/j.watres.2011.12.056).

41.    R Core Team. 2015 R: A language and environment for statistical computing. R

foundation for statistical computing.  (Vienna, Austria. URL http://www.R-project.org/.

42.    Kahm M., Hasenbrink G., Lichtenberg-Frate H., Ludwig J., Kschischo M. 2010 grofit:

fitting biological growth curves with R. *Journal of Statistical Software* **33**, 1-21.

43.    Lande R. 1979 Quantitative genetic analysis of multivariate evolution, applied to

brain:body size allometry. *Evolution* **33**, 402-416. (doi:10.2307/2407630).

44.    Barraclough T.G. 2015 How do species interactions affect evolutionary dynamics

across whole communities? *Annu Rev Ecol Evol Syst* **46**, 25-48. (doi:10.1146/annurev-

ecolsys-112414-054030).

45.    Barton S., Jenkins J., Buckling A., Schaum C.E., Smirnoff N., Raven J.A., Yvon-

Durocher G. 2020 Evolutionary temperature compensation of carbon fixation in marine

phytoplankton. *Ecol Lett* **23**, 722-733. (doi:10.1111/ele.13469).

46.    Alexander J.M., Diez J.M., Levine J.M. 2015 Novel competitors shape species'

responses to climate change. *Nature* **525**, 515-518. (doi:10.1038/nature14952).

47.    Hagemann M. 2011 Molecular biology of cyanobacterial salt acclimation. *Fems

Microbiology Reviews* **35**, 87-123. (doi:10.1111/j.1574-6976.2010.00234.x).

48.    Oren A. 1999 Bioenergetic aspects of halophilism. *Microbiol Mol Biol R* **63**, 334-+.

(doi:10.1128/mmbr.63.2.334-348.1999).

49.    Mee J.A., Yeaman S. 2019 Unpacking conditional neutrality: genomic signatures of

selection on conditionally beneficial and conditionally deleterious mutations. *Am Nat* **194**,

529-540. (doi:10.1086/702314).

50.    Yoshida T., Jones L.E., Ellner S.P., Fussmann G.F., Hairston N.G.J. 2003 Rapid

evolution drives ecological dynamics in a predator-prey system. *Nature* **424**, 303–306.

(doi:10.1038/nature01767).

51.    Frickel J., Sieber M., Becks L. 2016 Eco-evolutionary dynamics in a coevolving host-

virus system. *Ecol Lett* **19**, 450-459. (doi:10.1111/ele.12580).

52.     Hart S.P., Turcotte M.M., Levine J.M. 2019 Effects of rapid evolution on species

coexistence. **116**, 2112-2117. (doi:10.1073/pnas.1816298116).

53.     De Meester L., Brans K.I., Govaert L., Souffreau C., Mukherjee S., Vanvelk H.,

Korzeniowski K., Kilsdonk L., Decaestecker E., Stoks R., et al. 2019 Analysing eco-

evolutionary dynamics - the challenging complexity of the real world. *Funct Ecol* **33**, 43-59.

(doi:10.1111/1365-2435.13261).

54.     Hao Y.-Q., Brockhurst M.A., Petchey O.L., Zhang Q.-G. 2015 Evolutionary rescue

can be impeded by temporary environmental amelioration. *Ecol Lett* **18**, 892-898.

(doi:10.1111/ele.12465).

**Figure legends**

**Figure 1:** Experimental design. In Experiment 1, we propagated monocultures of six species of microalgae with or without increasing salt for ~ 185 generations, and then quantified growth rate in monoculture and population size in mixtures of species with and without salt. White boxes are flasks without salt, grey boxes are flasks with salt. In Experiment 2, we propagated two species in monoculture and in pairs, with and without salt. After ~ 230 generations we re-isolated the species on agar plates and quantified the chlorophyll-a concentration (as proxy for biomass) of all lines when competing either with a competitor of the same salt and competition history (co-evolved competitor) or with a reference line (naïve competitor; salt history: no salt, competition history: monoculture). Bold species abbreviations denote lines cultivated in pairs during the selection experiment.

**Figure 2:** Effects of salt history and assay environment on the monoculture growth rates of five species of microalgae (sorted by salt tolerance). Control lines (blue) have a salt history of 37 transfers at 0 g NaCl L$^{-1}$, selection lines (red) have a salt history of 37 transfers at increasing salt stress with a final concentration of 6 g NaCl L$^{-1}$. Salt concentrations in the two assay environments were the same as in the final selection environments, i.e. 0 and 6 g NaCl L$^{-1}$, respectively. Inserts give the P-values for assay environment (A), salt history (H), and their interaction (AxH). Bold font denotes P-values < 0.05. For F-statistics see Table S2. Values are means ± SE, n = 3.

**Figure 3:** Effects of salt history and assay environment on species abundances in the mixture assay. Mixtures of six species were assembled either from control lines (blue) or from selection lines (red). Species abundances were averaged over four transfers in assay environments with and without salt, respectively. Inserts give the P-values for assay

environment (A), salt history (H), and their interaction (AxH). For F-statistics see Table S3.

Abundances were ln-transformed prior to analyses except for *Pseudokirchneriella*. For

consistency with Fig. 2, we do not show *Anabaena*, which was also part of the communities.

For time-resolved data of all six species, see supplementary figure S4. Values are means ± SE

(SE too small to be visible), n = 3.

**Figure 4:** Effects of competition history, salt history, and assay environment on competitive

abilities of *Anabaena* and *Scenedesmus*. Relative chlorophyll a of (a) *Anabaena* in

competition with a reference line of *Scenedesmus,* and (b) of *Scenedesmus* in competition

with a reference line of *Anabaena*. Values are means ± SE, n = 3.

**Experiment 1: Does competition alter the effect of adaptation on population dynamics?**

i) Selection experiment

0 g NaCl L$^{-1}$           0 g NaCl L$^{-1}$     ii) Monoculture assay     iii) Mixture assay

37 transfers

0 g NaCl L$^{-1}$           6 g NaCl L$^{-1}$

**Experiment 2: Does competition affect adaptation to salt? Does salt history affect competition?**

i) Selection experiment      ii) Re-isolation   iii) Competition assay     iv) Competition assay

0 g NaCl L$^{-1}$        0 g NaCl L$^{-1}$              (co-evolved competitor)    (naive competitor)

46 transfers

0 g NaCl L$^{-1}$        6 g NaCl L$^{-1}$