Adaptation and evolutionary rescue in a community context

Romana Limberger[1,2*] and Gregor F. Fussmann[2]

1 Research Department for Limnology, University of Innsbruck, Mondsee, Austria

2 Department of Biology, McGill University, Montreal, Quebec, Canada

* Corresponding author:

Romana Limberger

romana.limberger@uibk.ac.at

Abstract

Evolution might mitigate negative effects of environmental change on diversity by rescuing

populations from extinction. However, biotic interactions could affect adaptation to abiotic

change, and conversely, adaptation to an abiotic stressor could influence biotic interactions.

We used experimental evolution of microalgae to investigate reciprocal effects of competition

and adaptation to salt stress. We found that evolutionary history influenced competitive

interactions and community properties. In high-salt environments, communities assembled

from lines with a history of salt stress declined more slowly in diversity than communities

assembled from lines without prior exposure to salt. We also found that a history of salt stress

can lead to reduced competitive ability in the ancestral environment. While competition had

no effect on adaptation to salt, we found that competition can hinder evolutionary rescue by

preventing that adaptation translates into population persistence. Collectively, our results

highlight the importance of considering evolutionary rescue in a community context.

Introduction

Abiotic environmental change can render populations maladapted, leading to their decline,

and ultimately, extinction (Ceballos & Ehrlich 2002; Stuart *et al.* 2004; Both *et al.* 2006).

Adaptive evolution may rescue populations from extinction if genotypes rise in frequency that

have positive growth rates under the new environmental conditions (evolutionary rescue;

Gomulkiewicz & Holt 1995; Bell & Gonzalez 2009). Laboratory experiments have

demonstrated that microorganisms can adapt rapidly to a variety of abiotic stressors, including

acidification (Lohbeck et al. 2012), salt stress (Lachapelle & Bell 2012), resource limitation

(Bell 2013), and warming (Schaum *et al.* 2018). Experimental evolution has also helped to

reveal the demographic, genetic, and extrinsic factors that influence the probability of

evolutionary rescue (Carlson *et al.* 2014). For example, evolutionary rescue is facilitated by

larger population size (Bell & Gonzalez 2009), higher genetic variation (Lachapelle & Bell

2012), and a slower rate of environmental change (Lindsey *et al.* 2013). However, most

empirical studies on evolutionary rescue have considered single species in isolation. Yet,

interactions with other species might influence (i) whether populations adapt to abiotic

stressors, and (ii) whether adaptation translates into population persistence (Fig. 1a).

Moreover, within communities, the ecological consequences of evolutionary rescue could go

beyond population persistence and include effects on species interactions and community

dynamics (Fig. 1b). This potentially complex interplay of biotic interactions and adaptation to

abiotic stressors limits our understanding of whether evolution can contribute to the

maintenance of diversity in changing environments.

Theory suggests several mechanisms whereby interspecific competition can influence

adaptation to abiotic change. First, competition can constrain adaptation to environmental

change by reducing population size, and thus the amount of standing genetic variation

(Bocedi *et al.* 2013) and the supply of beneficial mutations (Johansson 2008). Second,

competition can reduce the time available for adaptation as species are outcompeted by better

adapted competitors (de Mazancourt *et al.* 2008). Third, competition can alter the strength and

direction of selection (Osmond & de Mazancourt 2013); depending on whether competition

and environmental change impose selection on a trait in the same or in the opposite direction,

competition can speed up or hinder adaptation to environmental change (Osmond & de

Mazancourt 2013). Finally, negative genetic correlations between traits that mediate

competition and stress tolerance could constrain adaptation in environments with both biotic

and abiotic selective agents (Lau & Terhorst 2015). Empirical tests of these theoretical

predictions are rare and focused on biotic interactions other than interspecific competition

(Collins 2011; Hiltunen *et al.* 2018) or on environmental amelioration rather than

deterioration (Kleynhans *et al.* 2016).

Competition could not only alter how species adapt to abiotic change, but could also

modulate the effect of adaptation on population dynamics. When competition alters the

selection environment generated by abiotic change, adaptation of a focal species in isolation

may not translate into increased population size in a community context. Accordingly, an

experiment that manipulated CO_2 and diversity of a plant community showed that

interspecific competition altered the selection environment created by CO_2 (Kleynhans *et al.*

2016): adaptation to elevated CO_2 was only observed when the plants were assayed in the

same community context in which selection had occurred. Similarly, an experiment with a

marine alga revealed that adaptation to acidification in isolation was not associated with

increased population size within communities (Bach *et al.* 2018). In the context of

evolutionary rescue, competition could prevent that adaptation translates into population

persistence and thus preclude that evolution mitigates negative effects of environmental

change on diversity.

The field of evolutionary rescue primarily considers the effect of evolution on population

dynamics, but within communities, adaptation of a focal species to the local environment may

affect interactions with other species (terHorst *et al.* 2014; Pantel *et al.* 2015; Gomez *et al.*

2016). Evolution can influence community dynamics directly, by altering traits that underlie

species interactions, and indirectly, by influencing population dynamics (Matthews *et al.*

2011; Hendry 2019). In a scenario with an abiotic stressor as the selective agent (Fig. 1b),

adaptation could influence species interactions if traits that confer adaptation to the stressor

are correlated with traits that affect biotic interactions. Some traits may even simultaneously

mediate stress tolerance and species interactions. For example, in *E. coli* reduced membrane

permeability increases resistance to antibiotics but decreases the ability to take up nutrients,

such that adaptation to antibiotics results in lower competitive ability (Phan & Ferenci 2013).

In addition to such trait-mediated effects, adaptation of a focal species to an abiotic stressor

might influence biotic interactions by increasing its population size. Despite this range of

possibilities of how adaptation to abiotic change could affect species interactions, the

ecological consequences of evolutionary rescue have been rarely considered in a community

context.

A further complexity of adaptation in a multi-species context is that species might vary in

their ability to adapt to environmental change, which could hamper maintenance of diversity.

The rate of evolution depends on selection pressure, genetic variance, and covariance (Lande

1979; Agrawal & Stinchcombe 2009). Variation among species in ecological tolerance, life-

history traits, and genomic architecture could thus result in variation in evolvability

(Barraclough 2015). Species with narrow tolerance of the abiotic stressor will have a higher

risk of extinction, but will also experience stronger selection pressure and might thus evolve

more than species with broader tolerance (Osmond & de Mazancourt 2013). Species also vary

widely in population density (Damuth 1987), mutation rate (Drake *et al.* 1998; Baer *et al.*

2007), recombination rate (Stapley *et al.* 2017), and genome complexity (Lynch *et al.* 2011),

all of which influence the rate of evolution. Such variation among species in evolutionary

potential could lead to changes in the composition and interaction structure of communities in

deteriorating environments.

Here we investigated if adaptation to abiotic change influences competitive interactions

and community properties, and, vice versa, if competition influences adaptation to

deteriorating environmental conditions. In a laboratory experiment, we propagated

monocultures of six species of microalgae for ~ 180 generations with and without increasing

salt stress. We then quantified monoculture growth rates of all lines in a reciprocal transplant

assay to test the hypothesis that species vary in their evolutionary response to salt stress

(Hypothesis 1). We then assembled communities, either from lines with a history of salt stress

or from lines without prior exposure to salt, and investigated if evolutionary history affected

community dynamics in environments with and without salt, respectively. We predicted that

adaptation to salt in isolation would translate into longer persistence of species within salt-

stressed communities and consequently into maintenance of diversity (Hypothesis 2).

Alternatively, competitive interactions could prevent that adaptation of monocultures

translates into persistence within communities. In a second set of experiments, we propagated

two algae species for ~ 200 generations as monocultures and as pairs, respectively, in

environments with and without increasing salt. We then re-isolated the two species, and

measured monoculture growth rates and competitive abilities in reciprocal transplant assays.

We predicted that interspecific competition during the selection experiment would constrain

adaptation to salt stress (Hypothesis 3). As in the first set of experiments, we expected that a

history of salt stress would lead to increased competitive ability at high salt (Hypothesis 4).

Our results highlight the complexities that arise from adaptation to abiotic change in a

community context. We found that adaptation to abiotic stress influenced competitive

interactions and community-level properties, but we observed no effect of competition on

adaptation to salt stress.

Material and methods

Model organisms

We tested our hypotheses using six species of microalgae from three different taxonomic

groups: the cyanobacteria *Synechococcus leopoliensis* (Canadian Phycological Culture Centre

(CPCC) 102) and *Anabaena variabilis* (CPCC 105), the diatoms *Navicula pelliculosa* (CPCC

552) and *Nitzschia palea* (CPCC 160), and the chlorophytes *Pseudokirchneriella subcapitata*

(CPCC 37) and *Scenedesmus acutus* (CPCC 10). The algae were cultivated in Bold's basal

medium (Bold 1949), which was modified by adding silicate (0.58 g L^{-1} of Na_2SiO_3) and

vitamins (Low-Décarie *et al.* 2011) to allow growth of diatoms. In all experiments, the algae

were cultivated at 25 °C with light continuously provided at 100 μE m^{-2} s^{-1}. Cultures were

propagated under sterile conditions. We did not make cultures isogenic at the start of the

experiment; the observed evolutionary responses could thus have resulted either from *de novo*

mutations or from sorting of standing genetic variation.

Salt tolerance of ancestral lines

We determined the limit of salt tolerance of the six species by measuring their growth along a

salt gradient implemented on 48-well plates. The length of the gradient was based on results

of a preliminary experiment, and was 0 to 10 g NaCl L^{-1} in 1 g increments for *Nitzschia* and

Pseudokirchneriella, and 0 to 20 g NaCl L^{-1} in 2 g increments for the remaining species. After

six days of acclimation at the respective salt concentration, we inoculated fresh plates with

acclimated cultures (20 μl culture in 1 ml medium), sealed the plates with sterile air-

permeable membranes (VWR), and cultivated them on shakers (350 rpm). Fifteen times over

the course of ten days, we measured absorbance at 660 nm on an optical plate reader

(Synergy-HT, BioTek, Winooski, VT). We estimated the limit of salt tolerance for each

species as the salt concentration where we observed no increase in absorbance. Each salt ×

species combination was replicated twice.

Adaptation to salt in monoculture

In a serial transfer experiment, we propagated monocultures of the six species with and

without increasing concentrations of NaCl (Fig. S1, S2). The algae were cultured in 125-ml

glass flasks filled with 50 ml of Bold's medium and continuously shaken at 250 rpm. Every

3.5 days, we transferred 1 ml of culture to 50 ml of fresh medium for a total of 37 transfers (~

180 generations). For each of the six species, three replicate control lines were cultivated

without addition of salt and three replicate selection lines were cultivated with increasing salt.

We increased the salt concentration in the selection lines by 0.25 g L^{-1} every transfer until

reaching 4 g NaCl L^{-1} at transfer 16 (Fig. S2). We maintained this salt concentration for 10

transfers and then continued the increase in salt until reaching 6 g NaCl L^{-1} at transfer 34. We

chose a final salt concentration of 6 g NaCl L^{-1} because this was the limit of salt tolerance for

the species with the lowest salt tolerance among the six species.

To test if the six species of algae differed in their evolutionary response to salt stress

(Hypothesis 1), we measured the growth rate of each line in the two final salt environments (0

and 6 g NaCl L^{-1}). Before starting the growth assay, we removed non-evolutionary effects by

cultivating all lines at identical environmental conditions for three transfer cycles, that is two

cycles in the ancestral environment (0 g NaCl L^{-1}) and one cycle in the assay environment (0

and 6 g NaCl L^{-1}). We then inoculated flasks containing 50 ml medium of either 0 or 6 g NaCl

L^{-1} with 1 ml of acclimated culture and measured absorbance through time. Depending on the

time of individual cultures to reach carrying capacity, we measured absorbance over 2.5 to 7.5

weeks with 23 to 34 measurements. Absorbance was measured twice a day during the first 5

days, once a day during the intermediate phase of the experiment, and once every other day

towards the end of the experiment. At each measurement, we removed 200 µl of culture from

each flask and measured absorbance in 96-well plates on an optical plate reader.

Community assay

To test if evolutionary history can influence community dynamics (Hypothesis 2), we

assembled communities either of control or of selection lines, transplanted the communities

into the two final salt environments of the selection experiment, and quantified species

abundances and diversity of the community over four transfers. Prior to the community assay,

the lines had been cultivated for 37 transfers with or without increasing salt, followed by two

transfers in the ancestral environment (see above). We assembled communities with equal

biovolume of the six species. To this end, we determined species abundances in the

monocultures by counting Lugol-fixed samples with an inverted microscope, and then

converted abundances to biovolume using cellular biovolume estimates based on

measurements of 20 individuals per species (Hillebrand *et al.* 1999). We assembled three

replicate communities of control lines and three replicate communities of selection lines.

From each of the six communities, we inoculated two flasks, one with a salt concentration of

0 and one of 6 g NaCl L^{-1}. Every four days, we transferred 1 ml of the community into 50 ml

of fresh medium, with the flasks continuously shaken at 250 rpm. Over the course of four

transfers, we sampled the communities at each transfer, fixed the samples with Lugol's

solution, and counted species abundances with an inverted microscope.

Adaptation in monoculture vs. adaptation in pairs

To test if competition can influence adaptation to salt stress (Hypothesis 3), we propagated monocultures and pairs of *Anabaena* and *Scenedesmus* with and without increasing salt for 46 transfers, then re-isolated the two species and quantified their fitness with and without salt (Fig. S1b). The monocultures and pairs of the two species were part of the selection experiment described above and experienced the same changes in salt concentration (Fig. S2). To quantify how strongly competition affected the abundances of the two species during the selection experiment, we sampled the pairs and monocultures of *Anabaena* and *Scenedesmus* ten times over the course of the selection experiment and microscopically determined the abundances of the two species in Lugol-fixed samples. We chose the combination of *Anabaena* and *Scenedesmus* because in other species combinations the inferior competitor was outcompeted quickly both with and without increasing salt.

After 46 transfers in the selection experiment, we re-isolated *Anabaena* and *Scenedesmus* by plating the cultures on agar and isolating individual colonies. Monocultures were treated the same way as the pairs. Individual colonies were resuspended in liquid Bold's medium on well plates, and microscopically checked for contamination after a week of growth. We then merged eight clean isolates per sample to one culture. The two isolation steps were conducted in medium without addition of NaCl. We then acclimated the cultures to the two assay environments (0 and 6 g NaCl L^{-1}) for two transfer cycles. Hence, before starting the growth and competition assays, the cultures were cultivated under identical environmental conditions for four cycles (two isolation cycles without salt, two transfer cycles in the assay environment) to remove any non-evolutionary effects.

The acclimated cultures were used to start growth and competition assays. Growth rates of all lines were determined in monoculture assays in the two final salt environments by measuring absorbance through time as described above. To test if salt history and community

history influenced competitive interactions (Hypothesis 4), we assembled pairs of *Anabaena*

and *Scenedesmus* in the two final salt environments and determined changes in biomass of the

two species through time. We conducted two competition assays that differed in how we

assembled the pairs. First, we assembled pairs of co-evolved *Anabaena* and *Scenedesmus*, i.e.

lines that had the same community and salt history. However, because both competitors

differed between flasks in this assay, differences among treatments could not unequivocally

be attributed to one of the two species. We thus conducted a second competition assay where

we competed each line of *Anabaena* with the same reference line of *Scenedesmus*, and each

line of *Scenedesmus* with the same reference line of *Anabaena*. As reference line, we

randomly selected one of the three replicate control lines (i.e. community history:

monoculture, salt history: no salt). The second competition assay was started three weeks after

the first competition assay; cultures were stored meanwhile as monocultures at their

respective salt selection environment and then acclimated for one cycle to the salt assay

environment.

In the competition assays, we measured chlorophyll a concentrations of the two

competitors as proxy for biomass. To quantify chlorophyll a concentrations of the two

species, we used a FluoroProbe (bbe Moldaenke, Kiel-Kronshagen, Germany), a fluorometer

which can discriminate among algal groups (Catherine *et al.* 2012). Before the start of the

assay, we calibrated the FluoroProbe using cultures of *Anabaena* and *Scenedesmus* with

known chlorophyll a concentrations. We started the competition assay with equal chlorophyll

a concentrations (2 µg) of both species in flasks filled with 50 mL of Bold's medium. We

transferred the cultures every 3.5 days for ten transfers in the first competition assay and for

four transfers in the second competition assay and measured chlorophyll a concentrations of

both species at each transfer.

Data analysis

We estimated growth rates in monoculture assays by fitting linear models to the exponential

part of the absorbance vs. time curves, using ln-transformed absorbance data as response

variable. We computed two-way ANOVAs to investigate effects of salt history and assay

environment on growth rates of the six species. We analyzed the time series data of the

community assay with linear mixed models, with ln-transformed abundance as response

variable, transfer, salt history and assay environment as fixed effects, and flask as random

effect to account for repeated sampling of communities. We used the same approach to

investigate treatment effects on species richness. To test whether competition affected

adaptation of *Anabaena* and *Scenedesmus* to increasing salt, we computed three-way

ANOVAs and investigated the effects of community history, salt history and assay

environment on monoculture growth rates (growth assay) and relative chlorophyll a

(competition assay). All analyses were computed with R version 3.4.1 (R Core Team 2015);

the package *lme4* was used to calculate linear mixed effect models.

267
Results

Salt tolerance of ancestral lines

Among the six species, *Nitzschia* and *Pseudokirchneriella* had the lowest limit of salt

tolerance (5 and 6 g NaCl L^{-1}, respectively), *Anabaena*, *Scenedesmus* and *Navicula* had

intermediate limits of salt tolerance (12-16 g NaCl L^{-1}), and *Synechococcus* had the highest

limit of salt tolerance (18-20 g NaCl L^{-1}).

Adaptation of monocultures

After 37 transfers with and without increasing salt, respectively, we detected adaptation to salt

in two of six species, *Nitzschia* and *Pseudokirchneriella* (Fig. 2). In both species, the growth

response to the assay environment depended on salt history: selection lines had higher growth

rates at 6 g NaCl L^{-1} than control lines, but did not differ from control lines in the no-salt assay

environment (Fig. 2 c, d). The effect of salt history was particularly strong in *Nitzschia*; the

control lines were unable to grow at the final salt concentration, while all three replicate

selection lines had positive growth rates. Among the remaining four species, *Scenedesmus*

and *Synechococcus* were affected neither by salt history nor assay environment, while

Navicula and *Anabaena* had reduced growth rates at high salt, irrespective of salt history (Fig.

2).

Community assay

When assaying communities of either control or selection lines, we found that the effect of

salt history varied among species and assay environment (Table 1, Fig. 3). The abundances of

most species declined over the four transfers except for *Synechococcus,* which increased.

Assay environment had no effect on the abundances of *Navicula* and *Synechococcus*, while

the other four species declined faster when the community was assayed at high salt than

without salt. Salt history affected three of the six species (*Nitzschia*, *Pseudokirchneriella*, and

Scenedesmus), but the effect depended on the assay environment (Table 1). The strongest

effect of salt history was observed for *Pseudokirchneriella* (Fig. 3): in the high-salt assay

environment, selection lines declined more slowly in abundance than control lines, resulting

in extinction being delayed by one transfer. Similarly, selection lines of *Scenedesmus* declined

more slowly at high salt than control lines, but the effect was of small magnitude (Fig. 3). In

contrast, selection lines of *Nitzschia* declined slightly faster than control lines when assayed in

the no-salt environment.

The effect of evolutionary history on species abundances translated into an effect on the

diversity of the community (Table S1, Fig. 4). Species richness declined with time (transfer: P

< 0.001), and this decline was faster when communities were assayed with than without salt

(assay: P < 0.001). However, communities assembled from selection lines declined more

slowly in species richness than communities assembled from control lines (salt history ×

assay; P = 0.047), because of slower extinction of *Pseudokirchneriella*.

Adaptation in monoculture vs. adaptation in pairs

Interspecific competition strongly influenced species abundances during the selection

experiment, but did not lead to any evolutionary responses. In the selection experiment,

Anabaena reduced the abundance of *Scenedesmus* as long as salt concentration was low (Fig.

S3). However, at the final salt concentration of 6 g NaCl L^{-1} the competitive hierarchy

reversed and *Anabaena* was negatively affected by *Scenedesmus* (Fig. S3). When assayed as

monocultures after the selection experiment, neither of the two species showed evolutionary

responses (Table S2, Fig. 5a, b). Salt had a negative effect on the growth rate of both species

(P < 0.001), irrespective of salt history and community history.

The two competition assays revealed that lines of *Anabaena* with a history of salt

stress had reduced competitive ability in the ancestral environment. When assayed with co-

evolved competitors, high-salt lines of *Anabaena* had lower relative chlorophyll a than no-salt

lines in the no-salt assay environment, and consequently high-salt lines of *Scenedesmus* had

higher relative chlorophyll a than no-salt lines (salt history × assay: P = 0.002; Table S3, Fig.

S4). Because both competitors differed among treatment combinations, this experiment did

not allow detecting if the observed pattern was due to altered competitive ability of *Anabaena*

or of *Scenedesmus*. However, when assayed in competition with a reference line, only

Anabaena was affected by salt history, again with lower relative chlorophyll a of high-salt

than no-salt lines in the no-salt assay environment (salt history × assay: P = 0.029, Table S4,

Fig. 5c, d).

Discussion

We found that adaptation to an abiotic stressor can influence competitive interactions and

community-level properties, but we observed no effect of competition on adaptation to salt.

Communities assembled from lines with a selection history of salt stress maintained diversity

longer under salt stress than communities assembled from control lines. The positive effect of

salt history on diversity resulted from longer persistence of selection lines of

Pseudokirchneriella in salt-stressed communities compared with control lines. However, we

also found that adaptation of monocultures to salt stress did not necessarily translate into

longer persistence within communities: *Nitzschia* adapted to salt in isolation, but selection and

control lines went extinct equally fast within salt-stressed communities. This result indicates

that competition can override the effect of evolution on population dynamics. Finally, the

second set of experiments revealed that a history of salt stress can lead to reduced competitive

ability in benign conditions. When assayed in competition, selection lines of *Anabaena*

performed worse than control lines in the no-salt assay environment. Taken together, we

found different scenarios of how adaptation to an abiotic stressor can play out in a community

context, ranging from a positive effect of adaptation on performance within stressed

communities, over no effect, to a negative effect on performance under benign conditions.

Species varied in their evolutionary response to salt stress. The two species with the

lowest limit of salt tolerance adapted to salt, while species that were less negatively affected

by salt showed no sign of adaptation (Fig. 2). This finding is in line with the prediction that

species with narrow tolerance of a stressor should evolve more than species with broader

tolerance (Osmond & de Mazancourt 2013) because they experience stronger selection

pressure. Species-specific differences in genetic variance and covariance could be additional

reasons for variation among species in the rate of evolution (Lande 1979; Barraclough 2015).

However, given that only two species adapted, our sample size is not large enough to evaluate

if species-specific differences other than ecological tolerance influenced the evolutionary

response to salt. To elucidate such effects, future experiments could expose a set of species to

their respective limit of tolerance and then investigate if differences in life-history traits (e.g.

population size) influence the rate of adaptation. In our experiment, however, we simulated a

scenario where all species experienced the same amount of stress.

Evolution transiently mitigated the negative effect of salt stress on diversity, but did not

maintain diversity at unstressed levels. In high-salt environments, diversity declined more

slowly when communities were assembled from lines with a history of salt stress (Fig. 4),

because selection lines of *Pseudokirchneriella* persisted for one transfer longer than control

lines (Fig. 3). In *Nitzschia*, however, adaptation to salt in isolation was not associated with

increased fitness in salt-stressed communities. Consequently, even in communities of

selection lines, diversity declined faster in the high-salt than the no-salt assay environment

(Fig. 4). Our results thus suggest that competitive interactions can hinder evolutionary rescue

by preventing that adaptation translates into persistence of the population. The scenario of

adaptation in isolation followed by competition with pre-adapted species might seem

artificial, but it is a plausible scenario in a spatial context. Species may adapt to changes in

their local environment before immigration of better-adapted species from the regional

species pool (Urban *et al.* 2012). Depending on how well the resident species manage to adapt

to the new environmental conditions, they may coexist with or even dominate over pre-

adapted, immigrating species (De Meester *et al.* 2016). In our experiment, however, the

positive effect of salt history on persistence of *Pseudokirchneriella*, and thus on the diversity

of the community, was only transient. By the end of the experiment, this species had gone

extinct at high salt irrespective of salt history. However, *Pseudokirchneriella* also declined to

low or zero abundance in the no-salt assay environment, indicating that this species was

generally a weak competitor, irrespective of salt concentration. The effect of adaptation on

diversity and community dynamics might be more pronounced and persistent for species with

high interaction strengths and strong ecosystem effects, such as superior competitors or

efficient predators (Pantel *et al.* 2015; Matthews *et al.* 2016).

Competition had no effect on adaptation to salt stress, neither in terms of growth rate nor

in terms of competitive ability (Fig. 5). We had predicted that competition during the

selection experiment would constrain adaptation to salt because of reduced population size

and/or increased genetic constraints (Johansson 2008; Lau & Terhorst 2015). Alternatively,

competition could facilitate adaptation to abiotic stress by increasing the strength of selection

(Osmond & de Mazancourt 2013). The two species that we used to test this question,

Anabaena and *Scenedesmus*, adapted to salt neither in monoculture nor in pairs. Given that

these species had not reached their limit of salt tolerance in the selection experiment, low

selection pressure could explain why the monocultures did not adapt to salt. However, the

combination of salt and competition in the selection experiment drove the inferior competitor

to the brink of extinction (Fig. S3), suggesting that the inferior competitor experienced strong

selection pressure to evolve faster growth rates. Yet, we did not find any effect of community

history on adaptation. One possible explanation is that the competitive hierarchy in flasks

with increasing salt changed shortly before the end of the selection experiment when the final

salt concentration had been reached, from dominance of *Anabaena* to dominance of

Scenedesmus. Maybe relaxed selection pressure on *Scenedesmus* towards the end of the

experiment and only a short phase of strong selection on *Anabaena* explain why competition

had no effect on the evolutionary response of these two species.

Evolutionary change in response to salt influenced competitive interactions (Fig. 5c).

We had predicted that a history of salt stress would lead to increased competitive ability at

high salt. In contrast, however, strains of *Anabaena* with a history of salt stress had reduced

competitive ability in the no-salt assay environment. Such a pattern is an indication of

conditionally deleterious mutations (Mee & Yeaman 2019), i.e. mutations that are neutral in

the local environment and deleterious in an alternative environment. Because we have neither

molecular nor trait data, we cannot elucidate potential mechanisms behind the observed mal-

adaptation in the ancestral environment. The evolutionary response of *Anabaena* was

independent of community history and of whether fitness was assayed in competition with a

co-evolved or a naïve competitor (Fig. 5, Fig. S4). Importantly, however, we observed

reduced fitness of selection lines of *Anabaena* only when assayed in competition, not in the

monoculture growth assay. It thus seems that the evolutionary response of *Anabaena* to salt

was associated with changes in traits that influenced competitive interactions. Similar

evidence comes from a functional genomics study on *Daphnia*, which suggested that

evolution of salt tolerance could entail fitness costs in environments with predators (Latta *et*

al. 2012). Overall, our results suggest that evolutionary responses to abiotic change can

influence biotic interactions, such that populations with different histories of abiotic stress

might vary in their effects on communities and ecosystems.

The emerging field of eco-evolutionary dynamics investigates the effects of evolution

on populations, communities, and ecosystems (Hendry 2016). Evolutionary rescue, as one

component of this field, has its focus on the effect of evolution on population dynamics,

specifically on population persistence. Here we showed that the ecological consequences of

evolutionary rescue can include effects on species interactions and the diversity of

communities. Moreover, we found that competition can override the effect of evolution on

population dynamics, illustrating that community context can modulate the ecological

consequences of evolutionary change. We have observed a range of complexities arising from

a multi-species context despite using a fairly simple model system. Higher diversity and

trophic structure of natural systems will pose an even greater challenge to understanding eco-

evolutionary dynamics (De Meester *et al.* 2019; Hendry 2019). Field studies have

demonstrated evolutionary rescue of populations embedded in diverse communities (Reid *et*

al. 2016; Oziolor *et al.* 2019), but it remains to be determined if stress tolerant genotypes have

the same ecosystem effects as ancestral genotypes. If adaptation to an abiotic stressor is

associated with changes in traits that mediate biotic interactions, evolution may maintain

populations but fail to maintain community and ecosystem structure. Collectively, our results

suggest that considering community context is of fundamental importance if we want to

evaluate the capacity of evolution to maintain diversity and species interactions in changing

environments.

Acknowledgements

We thank Etienne Low-Décarie and Graham Bell for input on the design and analysis of

the experiment, and Yinci Yan for help with the implementation of the experiment. This work

was supported by an Erwin-Schrödinger scholarship of the Austrian Science Fund (FWF),

grant no. J3265-B16 to R.L., and an NSERC Discovery Grant to G.F.F.

References

Agrawal, A.F. & Stinchcombe, J.R. (2009). How much do genetic covariances alter the rate of

adaptation? *Proc. R. Soc. B*, 276, 1183-1191.

Bach, L.T., Lohbeck, K.T., Reusch, T.B.H. & Riebesell, U. (2018). Rapid evolution of highly

variable competitive abilities in a key phytoplankton species. *Nat. Eco. Evo.*, 2, 611-

613.

Baer, C.F., Miyamoto, M.M. & Denver, D.R. (2007). Mutation rate variation in multicellular

eukaryotes: causes and consequences. *Nat. Rev. Genet.*, 8, 619-631.

Barraclough, T.G. (2015). How do species interactions affect evolutionary dynamics across

whole communities? *Annu. Rev. Ecol. Evol. Syst.*, 46, 25-48.

Bell, G. (2013). Evolutionary rescue of a green alga kept in the dark. *Biol. Lett.*, 9, 20120823.

Bell, G. & Gonzalez, A. (2009). Evolutionary rescue can prevent extinction following

environmental change. *Ecol. Lett.*, 12, 942–948.

Bocedi, G., Atkins, K.E., Liao, J.S., Henry, R.C., Travis, J.M.J. & Hellmann, J.J. (2013).

Effects of local adaptation and interspecific competition on species' responses to

climate change. *Ann. N. Y. Acad. Sci.*, 1297, 83-97.

Bold, H. (1949). The morphology of *Chlamydomonas chlamydogama*, sp. nov. *Bull. Torrey*

Bot. Club, 76, 101-108.

Both, C., Bouwhuis, S., Lessells, C.M. & Visser, M.E. (2006). Climate change and population

declines in a long-distance migratory bird. *Nature*, 441, 81-83.

Carlson, S.M., Cunningham, C.J. & Westley, P.A.H. (2014). Evolutionary rescue in a

changing world. *Trends Ecol. Evol.*, 29, 521-530.

Catherine, A., Escoffier, N., Belhocine, A., Nasri, A.B., Hamlaoui, S., Yepremian, C. *et al.*

(2012). On the use of the FluoroProbe®, a phytoplankton quantification method based

on fluorescence excitation spectra for large-scale surveys of lakes and reservoirs.

Water Res., 46, 1771-1784.

Ceballos, G. & Ehrlich, P.R. (2002). Mammal population losses and the extinction crisis.

Science, 296, 904-907.

Collins, S. (2011). Competition limits adaptation and productivity in a photosynthetic alga at

elevated CO_2. *Proc. R. Soc. B*, 278, 247-255.

Damuth, J. (1987). Interspecific allometry of population density in mammals and other

animals: the independence of body mass and population energy-use. *Biol. J. Linn.*

Soc., 31, 193-246.

de Mazancourt, C., Johnson, E. & Barraclough, T.G. (2008). Biodiversity inhibits species'

evolutionary responses to changing environments. *Ecol. Lett.*, 11, 380–388.

De Meester, L., Brans, K.I., Govaert, L., Souffreau, C., Mukherjee, S., Vanvelk, H. *et al.*

(2019). Analysing eco-evolutionary dynamics - the challenging complexity of the real

world. *Funct. Ecol.*, 33, 43-59.

De Meester, L., Vanoverbeke, J., Kilsdonk, L.J. & Urban, M.C. (2016). Evolving perspectives

on monopolization and priority effects. *Trends Ecol. Evol.*, 31, 136-146.

Drake, J.W., Charlesworth, B., Charlesworth, D. & Crow, J.F. (1998). Rates of spontaneous

mutation. *Genetics*, 148, 1667-1686.

Gomez, P., Paterson, S., De Meester, L., Liu, X., Lenzi, L., Sharma, M.D. *et al.* (2016). Local

adaptation of a bacterium is as important as its presence in structuring a natural

microbial community. *Nat. Commun.*, 7, 12453.

Gomulkiewicz, R. & Holt, R.D. (1995). When does evolution by natural selection prevent

extinction? *Evolution*, 49, 201-207.

Hendry, A.P. (2016). *Eco-evolutionary dynamics*. Princeton University Press, Princeton and

Oxford.

Hendry, A.P. (2019). A critique for eco-evolutionary dynamics. *Funct. Ecol.*, 33, 84-94.

Hillebrand, H., Durselen, C.D., Kirschtel, D., Pollingher, U. & Zohary, T. (1999). Biovolume

calculation for pelagic and benthic microalgae. *J. Phycol.*, 35, 403-424.

Hiltunen, T., Cairns, J., Frickel, J., Jalasvuori, M., Laakso, J., Kaitala, V. *et al.* (2018). Dual-

stressor selection alters eco-evolutionary dynamics in experimental communities. *Nat.*

Eco. Evo., 2, 1974-1981.

Johansson, J. (2008). Evolutionary responses to environmental changes: How does

competition affect adaptation? *Evolution*, 62, 421-435.

Kleynhans, E.J., Otto, S.P., Reich, P.B. & Vellend, M. (2016). Adaptation to elevated CO_2 in

different biodiversity contexts. *Nat. Commun.*, 7, 12358.

Lachapelle, J. & Bell, G. (2012). Evolutionary rescue of sexual and asexual populations in a

deteriorating environment. *Evolution*, 66, 3508-3518.

Lande, R. (1979). Quantitative genetic analysis of multivariate evolution, applied to

brain:body size allometry. *Evolution*, 33, 402-416.

Latta, L.C., Weider, L.J., Colbourne, J.K. & Pfrender, M.E. (2012). The evolution of salinity

tolerance in *Daphnia*: a functional genomics approach. *Ecol. Lett.*, 15, 794-802.

Lau, J.A. & Terhorst, C.P. (2015). Causes and consequences of failed adaptation to biological

invasions: the role of ecological constraints. *Mol. Ecol.*, 24, 1987-1998.

Lindsey, H.A., Gallie, J., Taylor, S. & Kerr, B. (2013). Evolutionary rescue from extinction is

contingent on a lower rate of environmental change. *Nature*, 494, 463-467.

Lohbeck, K.T., Riebesell, U. & Reusch, T.B.H. (2012). Adaptive evolution of a key

phytoplankton species to ocean acidification. *Nat. Geosci.*, 5, 346-351.

Low-Décarie, E., Fussmann, G.F. & Bell, G. (2011). The effect of elevated CO_2 on growth

and competition in experimental phytoplankton communities. *Glob. Change Biol.*, 17,

2525–2535.

Lynch, M., Bobay, L.M., Catania, F., Gout, J.F. & Rho, M. (2011). The repatterning of

eukaryotic genomes by random genetic drift. *Annu Rev Genomics Hum Genet*, 12,

347-366.

Matthews, B., Aebischer, T., Sullam, K.E., Lundsgaard-Hansen, B. & Seehausen, O. (2016).

Experimental evidence of an eco-evolutionary feedback during adaptive divergence.

Current Biology, 26, 483-489.

Matthews, B., Narwani, A., Hausch, S., Nonaka, E., Peter, H., Yamamichi, M. *et al.* (2011).

Toward an integration of evolutionary biology and ecosystem science. *Ecol. Lett.*, 14,

690-701.

Mee, J.A. & Yeaman, S. (2019). Unpacking conditional neutrality: genomic signatures of

selection on conditionally beneficial and conditionally deleterious mutations. *Am.*

Nat., 194, in press.

Osmond, M.M. & de Mazancourt, C. (2013). How competition affects evolutionary rescue.

Phil. Trans. R. Soc. B, 368, 20120085.

Oziolor, E.M., Reid, N.M., Yair, S., Lee, K.M., VerPloeg, S.G., Bruns, P.C. *et al.* (2019).

Adaptive introgression enables evolutionary rescue from extreme environmental

pollution. *Science*, 364, 455-457.

Pantel, J.H., Duvivier, C. & De Meester, L. (2015). Rapid local adaptation mediates

zooplankton community assembly in experimental mesocosms. *Ecol. Lett.*, 18, 992-

1000.

Phan, K. & Ferenci, T. (2013). A design-constraint trade-off underpins the diversity in

ecologically important traits in species *Escherichia coli*. *Isme J.*, 7, 2034-2043.

R Core Team (2015). R: A language and environment for statistical computing. R foundation

for statistical computing. Vienna, Austria. URL http://www.R-project.org/.

Reid, N.M., Proestou, D.A., Clark, B.W., Warren, W.C., Colbourne, J.K., Shaw, J.R. *et al.*

(2016). The genomic landscape of rapid repeated evolutionary adaptation to toxic

pollution in wild fish. *Science*, 354, 1305-1308.

Schaum, C.E., Buckling, A., Smirnoff, N., Studholme, D.J. & Yvon-Durocher, G. (2018).

Environmental fluctuations accelerate molecular evolution of thermal tolerance in a

marine diatom. *Nat. Commun.*, 9, 1719.

Stapley, J., Feulner, P.G.D., Johnston, S.E., Santure, A.W. & Smadja, C.M. (2017). Variation

in recombination frequency and distribution across eukaryotes: patterns and processes.

Phil. Trans. R. Soc. B, 372, 20160455.

Stuart, S.N., Chanson, J.S., Cox, N.A., Young, B.E., Rodrigues, A.S.L., Fischman, D.L. *et al.*

(2004). Status and trends of amphibian declines and extinctions worldwide. *Science*,

306, 1783-1786.

terHorst, C.P., Lennon, J.T. & Lau, J.A. (2014). The relative importance of rapid evolution for

plant-microbe interactions depends on ecological context. *Proc. R. Soc. B*, 281,

20141615.

Urban, M.C., De Meester, L., Vellend, M., Stoks, R. & Vanoverbeke, J. (2012). A crucial step

toward realism: responses to climate change from an evolving metacommunity

perspective. *Evol. Appl.*, 5, 154-167.

Table 1: Community assay. Results (i.e. P-values) of linear mixed models testing for effects

of transfer (T), assay environment (A), and salt history (H) on the abundances of the six

species within communities assembled either from control or selection lines. Transfer was

treated as a continuous variable. Abundances of species were ln-transformed prior to analysis,

$P < 0.05$ in bold.

Species	Transfer	Assay	History	$A \times H$	$T \times A$	$T \times H$	$T \times A \times H$
Anabaena	**< 0.001**	**< 0.001**	0.807	0.538	**0.048**	0.770	0.822
Navicula	**< 0.001**	0.836	0.947	0.749	**0.010**	0.128	0.664
Nitzschia	**< 0.001**	**< 0.001**	0.285	**0.031**	**0.004**	0.151	0.648
Pseudokirchneriella	**< 0.001**	**< 0.001**	0.100	**0.005**	0.502	0.450	0.843
Scenedesmus	**< 0.001**	**< 0.001**	0.682	**0.017**	**0.005**	0.901	0.189
Synechococcus	**< 0.001**	0.289	0.781	0.545	0.328	0.611	0.633

(a) Effects of competition on evolutionary rescue

Abiotic stressor   Competition

N

Stress traits ← Interaction traits

Competition

$N_{Focal\ species}$

(b) Effects of evolutionary rescue on competition

Abiotic stressor

Stress traits = Interaction traits - - - ▸ $N_{Competitor}$

Interaction traits - - - ▸ $N_{Competitor}$

$N_{Focal\ species}$ - - - ▸ $N_{Competitor}$

Figure 1: Conceptual figure of how evolutionary rescue from an abiotic stressor (a) is affected by competition, and (b) affects competition. Evolutionary effects are denoted with solid lines, ecological effects with dashed lines. Stress traits indicate traits that mediate the response to a stressor, interaction traits indicate traits that mediate competitive interactions. Evolutionary rescue from a stressor occurs when the focal species responds with the evolution of stress traits such that population size (N) increases and the population persists. (a) Competition can influence adaptation to an abiotic stressor by reducing population size and thus genetic variation of the focal species, by imposing selection on stress traits, or by imposing selection on interaction traits that are genetically correlated with stress traits. Competition could also influence evolutionary rescue by altering the effect of adaptation on population size. (b) Evolutionary rescue can affect competition when stress traits simultaneously act as interaction traits, when stress traits are correlated with interaction traits, or when increased population size of the focal species influences the competitor.

Figure 2: Effects of salt history and assay environment on the monoculture growth rates of six species of microalgae. Control lines (blue) have a salt history of 37 transfers at 0 g NaCl L^{-1}, selection lines (red) have a salt history of 37 transfers at increasing salt stress with a final concentration of 6 g NaCl L^{-1}. Salt concentrations in the two assay environments were the same as in the final selection environments, i.e. 0 and 6 g NaCl L^{-1}, respectively. Inserts give the P-values for assay environment (A), salt history (H), and their interaction (AxH). Bold font denotes P-values < 0.05. Values are means ± SE, n = 3.

Figure 3: Effects of salt history and assay environment on species abundances in the community assay. Communities of six species were assembled either from control lines (blue) or from selection lines (red). Species abundances were quantified over four transfers in assay environments without salt (solid circles, solid lines) and with salt (open triangles, dotted lines), respectively. Values are means ± SE, n = 3.

Figure 4: Effects of salt history and assay environment on species richness in the community assay. Communities of six species were assembled either from control lines (blue) or from selection lines (red). Species richness was quantified over four transfers in assay environments without salt (solid circles, solid lines) and with salt (open triangles, dotted lines), respectively. Values are means ± SE, n = 3.

Figure 5: Effects of community history, salt history, and assay environment on the growth rates and competitive abilities of Anabaena and Scenedesmus. Monoculture growth rate of (a) Anabaena and (b) Scenedesmus, relative chlorophyll a of (c) Anabaena in competition with a reference line of Scenedesmus and (d) of Scenedesmus in competition with a reference line of Anabaena. Inserts indicate significant effects ($P < 0.05$) of community history (C), salt history (H), assay environment (A), or their interactions. Values are means ± SE, n = 3.