Local human movement patterns and land use impact exposure to zoonotic malaria in Malaysian

Borneo

Kimberly M Fornace*[1], Neal Alexander[2], Tommy R Abidin[3], Paddy M Brock[4], Tock H Chua[3], Indra

Vythilingam[5], Heather M. Ferguson[4], Benny O. Manin[3], Meng L. Wong[5], Sui Hann Ng[3], Jon Cox[1], Chris

J Drakeley[1]

1. Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine,

London, UK

2. Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical

Medicine, London, UK

3. Department of Pathobiology and Medical Diagnostics, Faculty of Medicine and Health

Sciences, Universiti Malaysia Sabah, Kota Kinabalu, Malaysia

4. Institute of Biodiversity, Animal Health and Comparative Medicine, College of Medical,

Veterinary and Life Sciences, University of Glasgow, UK

5. Parasitology Department, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia

* Corresponding author:

Kimberly M Fornace

Kimberly.Fornace@lshtm.ac.uk

London School of Hygiene and Tropical Medicine, Keppel Street, London, WC1E 7HT, UK

+44 (0) 20 7927 2010

Abstract: (150 max)

Human movement into insect vector and wildlife reservoir habitats determines zoonotic disease

risks; however, few data are available to quantify the impact of land use on pathogen transmission.

Here, we utilise GPS tracking devices and novel applications of ecological methods to develop fine-

scale models of human space use relative to land cover to assess exposure to the zoonotic malaria

Plasmodium knowlesi in Malaysian Borneo. Combining data with spatially explicit models of

mosquito biting rates, we demonstrate the role of individual heterogeneities in local space use in

disease exposure. At a community level, our data indicate that areas close to both secondary forest

and houses have the highest probability of human *P. knowlesi* exposure, providing quantitative

evidence for the importance of ecotones. Despite higher biting rates in forests, incorporating human

movement space use into exposure estimates illustrates the importance of intensified interactions

between pathogens, insect vectors and people around habitat edges.

Keywords (min 3):

Disease ecology, spatial epidemiology, *Plasmodium knowlesi*, malaria, human movement, land use

Introduction:

Environmental change and human encroachment into wildlife habitats are key drivers in the emergence and transmission of zoonotic diseases (1, 2). Individual movements into different habitats influence exposure to disease vectors and animal reservoirs, determining risk and propagation of vector-borne diseases (3-5). Increased contact between these populations is theorised to drive increases of the zoonotic malaria *Plasmodium knowlesi* in Malaysian Borneo, now the main cause of human malaria within this region. *P. knowlesi* is carried by long- and pig-tailed macaques (*Macaca fascicularis* and *M. nemestrina*) and transmitted by the *Anopheles leucospryphus* mosquito group, both populations highly sensitive to land cover and land use change (6). Although higher spatial overlap between people, macaques and mosquito vectors likely drives transmission, the impact of human movement and land use in determining individual infection risks is poorly understood (7).

The emergence of the zoonotic malaria *Plasmodium knowlesi* has been positively associated with both forest cover and historical deforestation (8, 9). However, out of necessity, statistical approaches to assess environmental risk factors for *P. knowlesi* and other infectious diseases typically evaluate relationships between disease metrics and local land cover surrounding houses or villages. While an individual may spend most of their time within the vicinity of their residence, this area does not necessarily represent where they are most likely to be exposed to a disease. This is supported by varying associations between *P. knowlesi* occurrence and landscape variables at different distances from households, ranging from 100m to 5km, likely partially due to human movement into different surrounding habitats (8, 10). Although land cover variables describing physical terrestrial surfaces are frequently incorporated into disease models, land use is rarely quantified. Land use is commonly defined as "the arrangements, activities, and inputs that people undertake in certain land cover types" (11). Places with the similar types of land cover may be used

very differently, with the activities and frequencies with which people visit these places determining

the spatial distribution of disease (1).

Mathematical modelling studies have revealed the importance of spatial variation in contact rates

due to the movement of individuals through heterogeneous environments with varying transmission

intensity (12). A multi-species transmission model of *P. knowlesi* highlighted the role of mixing

patterns between populations in different ecological settings in determining the basic reproductive

rate and subsequent modelling studies illustrate the sensitivity of this disease system to population

densities of both people and wildlife hosts (7, 13). However, although mechanistic models have been

extended to explore the potential importance of these heterogeneities in disease dynamics, there

are inherent constraints on model complexity and most models make simplistic assumptions about

the habitat uses of different populations.

Empirical data on human population movement is increasingly available, allowing assessment of the

impact of mobility on infectious disease dispersion and risks (5). On larger spatial scales, mobile

phone data has revealed the role of human migration in the transmission of infectious diseases such

as malaria, dengue and rubella (14-16).Although this data can provide insights into long range

movements, spatial resolution of this data is limited, particularly in areas with poor or no mobile

coverage, such as forested areas (17). Alternatively, the advent of low-cost GPS tracking devices

allows quantification of fine-scale movements, demonstrating marked heterogeneity in individual

movement and risk behaviours (3, 18). Combining these data with detailed data on land cover and

vector dynamics can provide new insights into how landscapes affect *P. knowlesi* transmission.

Previous studies of *P. knowlesi* have relied on questionnaire surveys, identifying self-reported travel

to nearby plantations and forest areas as a risk factor for *P. knowlesi* and other malaria infections

(e.g. (19-21)). However, the resultant spatial range and frequency of these movements remain

unknown and the definition of different habitat types is entirely subjective. Further, little is known

about differences in local movement patterns in different demographic groups. While infections in

male adults have been linked to forest and plantation work, it is unknown whether infections

reported in women and young children are likely to arise from exposure to similar environments

(22). The main mosquito vector in this area, *An. balabacensis*, is primarily exophagic and has been

identified in farm, forest and village areas near houses (23, 24). Macaque populations are reported

in close proximity to human settlements and molecular and modelling studies suggest transmission

remains primarily zoonotic in this area (7, 25, 26). A case control study detected higher abundances

of *An. balabacensis* near *P. knowlesi* case housesholds, suggesting the possibility of peri-domestic

transmission (24). Understanding the importance of these habitats is essential to effectively target

intervention strategies and predict impacts of future environmental changes.

Key questions remain about where individuals are likely to be exposed to *P. knowlesi* and how

landscape determines risk. Functional ecology approaches allow the distribution of different

populations to be modelled based on biological resources and relate transmission to landscape and

environmental factors (27). Within wildlife ecology, numerous methods have been developed to

estimate utilisation distributions (UDs), the probability of an individual or species being within a

specific location during the sampling period (28). Although these methods traditionally rely on kernel

density smoothing, kernel density estimates may not actually reflect time individuals spend in a

specific location if there is substantial missing data or irregular time intervals. Alternatively, biased

random bridges (BRBs) improve on these methods by estimating the UD as a time-ordered series of

points, taking advantage of the autocorrelated nature of GPS tracks to bias movement predictions

towards subsequent locations in a time series (29). This allows for interpolation of missing values

and adjustment for spatial error to estimate UDs representing both the intensity (mean residence

time per visit) and frequency of individual visits to specific locations. By integrating these estimates

of individual space use with detailed spatial and environmental data in a Bayesian framework, fine-

scale patterns of human land use can be predicted and overlaid with spatiotemporal models of

mosquito distribution. This allows exploration how landscape composition, as well as configuration

and connectivity between habitats, impacts human exposure to *P. knowlesi* and other vector-borne

and zoonotic diseases.

Focusing on one aspect of land use, human movement and time spent within different land cover

types, we explored the role of heterogeneity in local space use on disease exposure. Rolling cross-

sectional GPS tracking surveys were conducted in two study areas with on-going *P. knowlesi*

transmission in Northern Sabah, Malaysia (Matunggong and Limbuak (30)). We aimed to

characterise local movement patterns and identify individuals and locations associated with

increased *P. knowlesi* exposure risks by: 1. analysing individual movement patterns and developing

predictive maps of human space use relative to spatial and environmental factors, 2. modelling

biting rates of the main vector *An. balabacensis*, and 3. assessing exposure risks for *P. knowlesi*

based on predicted mosquito and human densities (Figure 1) Integrating these three approaches

allowed a uniquely spatially explicit examination of disease risk.

Figure 1. Analysis methods used to estimate individual and community-level exposure to *P. knowlesi*

sporozoite positive *An. balabacencis* bites

Methods

Study site

This study was conducted in two rural communities in Northern Sabah, Malaysia: Matunggong,

Kudat (6°47N, 116°48E, population: 1260) and Limbuak, Pulau Banggi (7°09N, 117°05E, population:

1009) (Figure 2). These areas were the focus for integrated entomology, primatology and social

science studies for risk factors for *P. knowlesi* (https://www.lshtm.ac.uk/research/centres-projects-

[groups/monkeybar](#)), with clinical cases and submicroscopic infections reported from both sites and

P. knowlesi sero-prevalence estimated as 6.8% and 11.7% in Matunggong and Limbuak respectively

(30).

Demographic data and GPS locations of primary residences were collected for all individuals residing

in these areas (30). Potential spatial and environmental covariates for these sites were assembled

from ground-based and remote-sensing data sources (Supplementary Table 1). The enhanced

vegetation index (EVI) was used to capture temporal changes in vegetation levels; this index

captures photosynthetic activity and has higher sensitivity in high biomass areas compared to the

normalised difference vegetation index (NDVI) frequently used. Due to the high cloud cover within

this area, EVI at a high spatial resolution could not be obtained for all time periods. Instead, EVI data

at a lower spatial but higher temporal resolution was used and monthly averages were calculated

from all available cloud-free data and resampled to 30m per pixel (31).

Figure 2. A. Location of study sites and tracked houses (households with one or more individual GPS

tracked) and survey houses (households with only questionnaire data collected and used for

prediction) in B. Matunggong, Kudat and C. Limbuak, Banggi; description of land cover classification

and survey methodology in (30)

GPS tracking survey

A minimum of 50 participants per site were targeted in a rolling cross-sectional survey (32). During

pre-defined two-week intervals, randomly selected participants from comprehensive lists of eligible

community members were asked to carry a QStarz BT-QT13000XT GPS tracking device (QStarz,

Taipei, Taiwan) programmed to record coordinates continuously at one-minute intervals for at least

14 days regardless of individual movement. Individuals were excluded if they were not primarily

residing in the study area, under 8 years old or did not consent. Trained fieldworkers visited the

participant every two days to confirm the device was functioning, replace batteries and administer

questionnaires on locations visited and GPS use. Fieldworkers recorded whether the device was

working and if the individual was observed carrying the GPS device to assess compliance. Individuals

were excluded from analysis if insufficient GPS data were collected (less than 33% of sampling

period) or individuals were observed not using the device for two or more visits.

Human space use

Biased random bridges (BRBs) were used to calculate individual utilisation distributions (UDs), the

probability of an individual being in a location in space within the sampled time period (29). Within

this study, large proportions of GPS fixes were missed due to technical issues with batteries and GPS

tracking; BRBs were used to interpolate between known locations and adjust for missing data, using

the time series GPS data to provide a more accurate estimate of space use. UDs were calculated

separately for each individual for all movement and night-time only movements (6pm – 6am).

To fit BRBs, we estimated the maximum threshold between points before they were considered

uncorrelated (T_{max}) as 3 hours based on typical reported activity times. The minimum distance

between relocations (L_{min}), the distance below which an individual is considered stationary, was set

at 10m to account for GPS recording error based on static tests. Finally, the minimum smoothing

parameter (h_{min}), the minimum standard deviation in relocation uncertainty, was set as 30m to

account for the resolution of habitat data and capture the range of locations an individual could

occupy while being recorded at the same place (28, 29). Estimates of the core utilisation area (home

range) were based on the 99[th] percentile, representing the area with a 99% cumulative probability

distribution of use by the sampled individual.

To assess relationships between space use and environmental factors and develop predictive maps

of community space use, we fit resource utilisation functions, regression models in which the UDs

are used as the response variable, improving on models using raw GPS count points as the response

when there is location uncertainty and missing data (33). The probability density function (UD) per

individual was rasterised to 30m^2 grid cells and environmental and spatial covariates extracted for

each grid cell. Potential environmental covariates included distance to the individual's own house,

distance to closest house, distance to roads, land use class (forest, agriculture, cleared or water),

distance to forest edge, elevation and slope (Supplementary table 1). Resource utilisation was

modelled as a Bayesian semi-continuous (hurdle) model with two functionally independent

components, a Bernoulli distribution for probability of individual i visiting a specific grid cell j (ω_{ij})

and a gamma distribution for the UD in grid cells visited (y_{ij}) (34, 35). For each individual, we defined

absences to be all grid cells with a UD less than 0.00001, indicating a very low probability the

individual visited this grid cell during the study period. We included all presences (grid cells with a

UD > 0.00001) and randomly subsampled equal numbers of absences (grid cells not visited) for each

individual as including equal numbers of presences and absences can improve predictive abilities of

species distribution models (36). The utilisation distribution for grid cells visited is defined as:

$$y_{ij} = \left\{ \begin{array}{c} \text{Gamma}\left(\dfrac{\mu_{ij}^2}{\sigma^2}, \dfrac{\sigma^2}{\mu_{ij}}\right) \text{with probability } 1 - \phi_{ij} \\ 0 \text{ with probability } \phi_{ij} \end{array} \right\}$$

216   Where the mean of y_{ij} is given by:

$$\mu_{ij} = \text{E}\big(y_{ij}|X_{ij}^T\big) = \big(1 - \omega_{ij}\big)v_{ij}$$

217   The full model was specified as:

$$\omega_{ij} \sim \text{Bernoulli}\big(\phi_{ij}\big)$$

220   With the linear predictor for the Bernoulli model specified as:

$$\text{logit}\big(\phi_{ij}\big) = \beta_0 + X_{ij}^T\beta_i + \gamma_j$$

223 Where β_0 represents the intercept, $X_{ij}^T \beta_i$ represents a vector of covariate effects and γ_j represents

224 the additive terms of random effects for individual. For the Gamma component, σ^2 is the variance

225 and the linear predictor v_{ij} is specified as:

$$\log(v_{ij}) = \alpha_0 + X_{ij}^T \alpha_i + \varphi_j$$

With α_0 representing the intercept, $X_{ij}^T \alpha_i$ representing a vector of coordinates and φ_j representing

the random effects. Weakly informative normal priors specified as Normal (0,1/0.01) were used for

all intercepts and coefficients. Bayesian inference was implemented using integrated nested Laplace

approximation (INLA) (37). This approach uses a deterministic algorithm for Bayesian inference,

increasing computational efficiency relative to Markov chain Monte Carlo and other simulation-

based approaches (34). We did not explicitly include spatial autocorrelation as several distance-

based covariates were included (e.g. distance from own house) (33). Predictive models used data for

all individuals aged 8 or over residing in these communities (Table 1) and models were limited to

land areas within 5km of households included in the study site. Separate models were fit for each

site.

Exposure to infected vectors

To estimate vector biting rates, we assembled data from 328 nights of human landing catches (HLCs)

conducted with 5km of the Matunggong study site while GPS tracking was on-going, including:

monthly longitudinal surveillance (23), investigations surrounding households of cases and controls

(24), and environmentally stratified outdoor catches (38) (Supplementary table 2). We limited this

data to counts of *An. balabacensis*, the primary *knowlesi* vector which comprises over 95% of

Anopheles caught in this region. As one experiment only collected mosquitoes for 6 hours, we fit a

linear model of all available data vs totals after 6 hour catches to estimate the total numbers of *An.*

balabacensis which would have been caught over 12 hours for these data (R^2= 0.85). Plausible

environmental covariates were assembled, including land use type, slope, aspect, elevation,

topographic wetness index, EVI, population density and average monthly temperature and rainfall.

To select variables for inclusion, Pearson correlation analysis was used to assess multicollinearity

between selected environmental variables. As topographic slope and TWI had a strong negative

correlation, only TWI was included in the analysis. The autocorrelation function (ACF) and partial

autocorrelation function (PACF) were used to explore correlation between time lags.

A Bayesian hierarchical spatiotemporal model was implemented using counts of *An. balabacensis*

bites as the outcome, denoted as m_{it}; j = 1…n; t = 1…n; where j indexes location and t indexes month.

The log number of person-nights per catch was included as an offset to adjust for numbers of

catchers conducting HLCs during different experiments. As the data were over dispersed, a negative

binomial distribution was used to model m_{it}. The linear predictor was specified as:

$$\log(\mu_{jt}) = \log(N_{jt}) + Z_0 + D_{jt}^T Z + w_j + e_t$$

263   Where N_{ijt} represents the number of person-nights for each HLC catch, Z_0 represents the intercept,

264    $D_{jt}^T Z$ represents a vector of covariates, w_j is the spatial effect and e_t is the temporal effect. The

265   temporal effect e_t was included as a fixed effect, random effect or temporally structured random

266   walk model of order 1 in candidate models (39). The spatial effect w_j was modelled as a Matern

267   covariance function between locations s_j and s_k:

$$W \sim Multivariate\ Normal\ (0, \Sigma)$$

$$\Sigma_{hk} = Cov(\xi(s_h), \xi(s_k)) = Cov(\xi_h, \xi_k) = \frac{\sigma^2}{\Gamma(\lambda)2^{\lambda-1}} (\kappa||s_h - s_k||)^\lambda K_\lambda(\kappa||s_h - s_k||)$$

Where $||s_h - s_k||$ denotes the Euclidean distance between locations s_h and s_k, $\xi(s)$ is the latent

Gaussian field accounting for spatial correlation, σ^2 is the spatial process variance and K_λ is a

modified Bessel function of the second kind and order $\lambda > 0$. κ is a scaling parameter related to r, the

distance at which spatial correlation becomes negligible, by $r = \sqrt{8\lambda}/\kappa$. A stochastic partial

differential equations (SPDE) approach was used, representing the spatial process by Gaussian

Markov random fields (GMRF) by partitioning the study area into non-intersecting triangles (40). This

approach represents the covariance matrix Σ by the inverse of the precision matrix Q of the GMRF

(34, 40). Prior distributions were specified on fixed effects and hyperparameters. A vague normal

prior distribution was used for the intercept. Weakly informative priors were used for fixed effects

specified as *N(1,1/0.01)*. Priors for spatial hyperparameters were specified as range $r \sim N(10, 1/0.01)$

and standard deviation $\sigma \sim N(0.1, 1/0.01)$ as described by Lindgren and Rue (39).

As these vectors are rarely reported indoors (24) and HLCs were primarily conducted outside, we

excluded areas within houses for calculations of exposure risks. The proportion of infectious

mosquitoes, c, was parameterised using a beta distribution for *P. knowlesi* sporozoite rates within

this site; with only 4 out of 1524 collected mosquitoes positive, it was not possible to look at

variations of infection rates by time and space. Spatially explicit exposure risks were calculated as

derived quantity from human resource utilisation, mosquito biting rate models and probability of *P.*

knowlesi sporozoite positivity. Individual exposure risk was explored using a simple exposure

assessment model where the number of infected bites received by an individual is the sum of bites

by infected vector across all locations visited, with the number of infectious bites received by

individual i in month t as:

$$r_{it} = c \sum_{j=1}^{J} y_{ij} m_{jt}$$

292   Where j indexes the grid cells visited, y_{ij} is the utilisation distribution, m_{jt} is the number of bites per

293   individual in that cell and month, and c is the proportion of infectious mosquitoes (4). To evaluate

294   places associated with exposure for the entire community, we calculated the number of infectious

295   bites per grid cell each month as:

$$r_{jt} = c \sum_{i=1}^{I} Y_{ij} m_{jt}$$

Where Y_{ij} is the predicted utilisation distribution for all individuals within the community per grid cell

j. All analyses were conducted in R version 3.5, with Bayesian models implemented using Integrated

Nested Laplace Approximation (INLA) (37). Model fit was assessed using deviance information

criteria (DIC) and area under the receiver operating curve (AUC), root mean square error (RMSE) or

conditional predictive ordinate (CPO) (41).

Ethics approval

This study was approved by the Medical Research Sub-Committee of the Malaysian Ministry of

Health and the Research Ethics Committee of the London School of Hygiene and Tropical Medicine.

Written informed consent was obtained from all participants or parents or guardians and assent

obtained from children under 18.

Results:

Between February 2014 and May 2016, 285 consenting people participated in the GPS tracking study

with 243 included in the final analysis including 109 in Limbuak and 134 in Matunggong (Table 1).

The most commonly reported occupation was farm or plantation work (n=73), primarily conducted

within the immediate vicinity of the house. A total of 3,424,913 GPS points were collected,

representing 6,319,885 person-minutes of sampling time. Median sampling duration was 16.27 days

(IQR 13.72 – 19.97), with points recorded for a median of 59.1% (IQR: 46.9% - 71.1%) of the sampling

duration. Maximum distances travelled ranged from no travel outside the house to 116km, with a

median distance travelled of 1.8km. Utilisation distributions (UDs), the probability of an individual

being in a location in space within a given time (Figure 3), varied by gender and occupation .

Individuals at the more rural Limbuak site covered larger distances (Table 2), with the largest

distances covered by individuals reporting primary occupations of fishing (n=5) and office work

(n=9). Although substantial differences were reported in all movements (24 hour sampling) between

seasons, no seasonal differences were observed in human movements during peak *Anopheles* biting

times (6pm-6am).

Table 1. Baseline characteristics of study site communities and sampled populations

	Matunggong		Limbuak	
	Sampled	Community*	Sampled	Community*
N	134	958	109	633
Gender				
Male, % (n)	51.5% (69)	46.1% (442)	47.7% (52)	46.1% (292)
Women, % (n)	48.5% (65)	53.9% (516)	52.3% (57)	53.9% (341)
Age in years, median (IQR)	31 (17 − 53)	32.5 (8 − 51)	29 (15 − 46)	30 (15 − 47)
Main occupation, % (n)				
Farming	29.9% (40)	28.6% (274)	7.3% (8)	10.2% (65)
Plantation work	10.4% (14)	8.6% (82)	10.1% (11)	7.6% (48)
Student	26.1% (35)	27.7% (265)	26.6% (29)	21.0% (133)
Other	6.7% (9)	9.1% (87)	15.6% (17)	14.4% (91)
No employment/ housewife	26.9% (36)	26.1% (250)	40.4% (44)	46.8% (296)

326   * Community includes all individuals eligible for these surveys (residents ages 8 and over)

328   **Figure 3.** Human movement relative to habitat. A. Example of GPS tracks from a 22-year-old male

329   plantation worker in Matunggong over aerial imagery, B. Individual utilisation distribution calculated

330   from GPS tracks

332   **Table 2.** Home range estimates by demographic group and site

	Area of 99% UD for all movement (hectares) Median (IQR)	Area of 99% UD from 6pm – 6am (hectares) Median (IQR)
Demographic group		
Men	32.09 (7.07, 148.93)	4.50 (2.79, 19.53)
Women	74.25 (12.24, 320.74)	6.08 (2.79, 24.17)
Children (under 15)	26.01 (6.39, 151.94)	3.83 (2.79, 8.73)
Occupation		
Farming	29.34 (8.15, 324.38)	6.75 (2.79, 19.80)
Plantation work	49.14 (9.72, 201.33)	4.59 (2.79, 27.72)
Fishing	442.49 (40.07, 1189.00)	227.16 (4.05, 465.14)
Office work	96.80 (63.61, 256.75)	13.63 (2.88, 20.14)
Other	19.98 (6.30, 26.82)	2.97 (2.61, 18.27)
No employment/ housewife	43.38 (11.97, 157.59)	3.60 (2.79, 19.12)
Site		
Limbuak	99.99 (24.57, 387.54)	7.74 (2.88, 58.05)
Matunggong	12.02 (3.94, 85.55)	2.97 (2.70, 11.77)
Season		
Dry (February – July)	28.62 (5.45, 252.45)	4.19 (2.79, 19.60)
Wet (August – January)	54.90 (17.23, 160.99)	4.64 (2.79, 19.35)

For both study areas, we developed models of community space use during peak mosquito biting

hours (6pm – 6am), in the form of resource utilisation functions, predictions of time- and space-

specific UDs on the basis of spatial and environmental variables (28). Between 6pm – 6am, human

space use (UDs) was mostly predictable and negatively correlated with distance from the individual's

house, other houses, roads and slope. The AUC for presence/ absence models was 0.936 for

Matunggong and 0.938 for Limbuak and RMSE for the overall model was 0.0073 and 0.0043 for

Matunggong and Limbuak respectively. While individuals were more likely to use areas further away

from forests in the Matunggong site, human space use was positively correlated with proximity to

forests in the Limbuak site (Table 3). Despite marked differences between different demographic

groups and seasons observed during 24 hour movements, these factors did not improve the

predictive power of the model for movements between 6pm and 6am.

Table 3. Estimated coefficients for fixed effects of resource utilisation functions (6pm – 6am)

	Matunggong			Limbuak		
	Mean	SD	95% CI	Mean	SD	95%CI
Probability of presence/ absence						
Intercept	3.383	0.839	3.218, 3.547	3.571	0.104	3.368, 3.775
Distance from own house (km)	-0.954	0.006	-0.966, -0.942	-0.543	0.003	-0.548, -0.539
Distance from forest (km)	5.997	0.177	-5.650, 6.344	-1.845	0.050	-1.944, -1.746
Distance from road (km)	-5.552	0.057	-5.663, -5.441	-3.656	0.019	-3.694, -3.618
Distance from houses (km)	-0.504	0.030	-0.563, -0.444	0.176	0.007	0.162, 0.189
Elevation (100 MSL)	-0.710	0.025	-0.759, -0.662	-1.268	0.037	-1.340, -1.197
Slope (degrees)	-0.0244	0.002	-0.028, -0.021	-0.009	0.001	-0.012, -0.006
Utilisation distributions for locations present						
Intercept	-6.846	0.866	-8.549, -5.147	-5.676	1.017	-7.673, -3.681
Distance from own house (km)	-0.583	0.004	-0.590, -0.576	-0.308	0.002	-0.311, -0.305
Distance from forest (km)	12.012	0.199	11.621, 12.403	-1.771	0.049	-1.868, -1.675
Distance from road (km)	-0.833	0.054	-0.939, -0.728	-1.532	0.011	-1.554, -1.511
Distance from houses (km)	-0.819	0.023	-0.864, -0.773	-0.239	0.006	-0.249, -0.228
Elevation (100 MSL)	0.664	0.027	0.610, 0.718	-0.297	0.003	-0.303, -0.297
Slope (degrees)	-0.021	0.002	-0.024, -0.018	-0.034	0.001	-0.036, -0.031

Between August 2013 and December 2015, 4814 *An. balabacensis* were caught from 328 sampling

nights in 155 unique locations. The median biting rate was 2.1 bites per night per person, ranging

from 0 – 28 bites per person per night (Figure 4). Despite monthly variation, no significant time lags

or seasonal patterns in mosquito biting rates were identified (Table 4). Although no associations

were identified between land classification and vector density in this site, models identified positive

relationships with enhanced vegetation indices (EVI) and negative associations with distance to

forest and human population density (Table 5). Of 1524 mosquitoes tested for *Plasmodium*

sporozoites, the median sporozoite rate was 0.24% (95% CI: 0.09 – 0.58%).

Table 4. Model selection statistics for mosquito biting rates

Model		DIC*	Marginal Likelihood	Model complexity*	RMSE*	Mean log-score (CPO)
M1	No spatial or temporal effect	2367.03	-1196.61	4.12	4.99	3.61
M2	Spatial effect only	2292.97	-1175.47	40.03	4.42	4.16
M3	Spatial effect + month as	2282.88	-1173.68	43.99	4.24	3.90

	fixed effect					
M4	Spatial effect + month as random effect	2222.89	-1155.91	50.28	4.05	3.61
M5	Spatial effect + month as random walk	2225.43	-1167.79	47.55	4.09	3.63

Table 5. Posterior rate ratio estimates and 95% Bayesian credible interval (BCI) for model 4 of
mosquito biting rates

Covariate	95% BCI Rate Ratio		
	Mean	2.5%	97.5%
Population density	0.963	0.916	1.004
EVI	3.185	1.185	8.532
Distance to forest (100m)	0.926	0.871	0.976
Spatial range (km)	3.120	0.514	6.926

364
365

Figure 4. Mosquito biting rates. A. *An. balabacensis* biting rate per person-night from data collected
in Matunggong, B. Predicted mean *An. balabacensis* biting rates per month from spatiotemporal
models, C. Predicted number of bites for all individuals residing in Matunggong by distance from
secondary forest, and by D. Distance from households

For individuals included in the GPS tracking study in Matunggong, where both human movement
and entomology data was available, we calculated exposure risks as a derived quantity from
utilisation distributions and mosquito biting rate models. Exposure varied markedly between
individuals, with an overall 150-fold difference in predicted mean probabilities of infected bites per
night (range: 0.00005-0.0078) (Table 6). No clear differences were observed between genders, age
groups or occupations of individuals sampled and there was no association between risk and
distance travelled.

380 **Table 6.** Probabilities of infected bites per night for sampled individuals in Matunggong by

381 demographic characteristics

	Predicted infectious bites per night (median (IQR))
Demographic group	
Men	0.00157 (0.000804, 0.00289)
Women	0.00219 (0.000864, 0.00307)
Children (under 15)	0.00131 (0.000812, 0.00330)
Occupation	
Farming	0.00180 (0.00101, 0.00362)
Plantation work	0.00216 (0.000680, 0.00278)
Student	0.00143 (0.000915, 0.00304)
Other	0.00225 (0.000852, 0.00302)
No employment/ housewife	0.00142 (0.000297, 0.00263)

Using the resource utilisation function with demographic and spatial data for all individuals in

Matunggong, we predicted community-wide space use and estimated exposure to infected

mosquitoes (Figure 5). The predicted number of person nights per grid cell for the entire community

ranged from 0 to 12.79 (median: 0.01, IQR: 0.0004 – 0.99), with the mean probability of a

community member exposed to an infected bite per grid cell of 0.00082 (IQR: 0.00001, 0.00050).

Although over 43% of the study site is forest and relatively high biting rates were predicted in forests

during the study period (mean: 1.94, range: 0.04 – 12.59), this habitat was rarely used by people in

the evenings, with less than 8% of predicted person-nights in forests. Models only based on

mosquito biting rates and not including human space use predicted 42% of infectious bites occurred

in forested areas and only 8.6% of bites occurring within 100m of houses (Figure 5C). In contrast,

when space use patterns are included, over 91% of predicted infected bites were predicted within

500m of houses (Figure 5D). Highest exposure risks were consistently found near forest edges and in

close proximity to households, despite spatial and temporal heterogeneity and model uncertainty

(Figure 4).

Figure 5. A. Land use in Matunggong site, B. Predicted number of person- nights for entire

community per grid cell, C. Predicted mosquito biting rates, D. Predicted infected bites per grid cell

Discussion

This study highlights the importance of human space use in different land cover types in determining

exposure to zoonotic and vector-borne diseases such as *P. knowlesi*. Although *P. knowlesi* has

previously been associated with forest exposure (e.g. (19)) and higher biting rates have been

reported in forest interiors (23), this novel approach incorporating both mosquito and human space

use data provides a new perspective on peri-domestic transmission, with more than 90% of

infectious bites predicted in areas surrounding households at forest edges. This study additionally

demonstrates the utility of ecological methods to understand human movement and identify

geographical areas associated with higher contact with disease vectors.

Within these communities, local movement patterns during peak vector times were largely

predictable and could be explained by spatial and environmental factors. However, despite this

finding, there was substantial variation in predicted exposure between individuals as a result of

heterogeneity in habitats used. No significant differences in exposure were predicted between men

and women, with individuals with high exposure risks identified across occupational and age groups.

Although this finding differs from clinical reports, a comprehensive survey within this community

identified equal proportions of men and women exposed to *P. knowlesi* as evidenced by specific

antibody responses and data on asymptomatic infections suggests higher numbers of non-clinical

infections in women (30, 42). While infrequent events or long-range movements (such as hunting

trips) may contribute to these differences in clinical cases and may not have been captured within

this two-week sampling period within the study site, this analysis highlights the importance of

routine movements into local environments in shaping exposure risks.

This improved understanding of how local human land use is related to exposure risk has important

implications for surveillance and control programmes. Malaria control programmes often rely on

interventions within the house, such as insecticide treated bednets and indoor residual spraying;

however, movements outside during peak biting times illustrate the importance of also targeting

outdoor transmission. The identification of areas where exposure is likely to occur can further be

used to refine interventions; for example, although insecticide treated hammocks have been

proposed for deep forest environments, larval source management may be more appropriate to

target environments in close proximity to houses. Although initial *P. knowlesi* cases were primarily

identified in adult men living and working in forests (20), this study illustrates the potential

importance of peri-domestic habitats in transmission and provides quantitative insight on mixing

between people and infected mosquitoes in forest fringe areas. As Malaysia moves towards malaria

elimination, surveillance systems are incorporating novel focal investigation methods, including

monitoring changes in local land use and populations at risk (43). In additional to routine vector

surveillance, this study highlights the need to incorporate measures of human space when defining

risk zones.

Even with the large and highly detailed movement dataset analysed, this study was limited by the

availability of mosquito data; as human landing catch data were assembled from other studies, there

was not uniform spatial and temporal coverage of the study site increasing uncertainty. The limited

mosquito data availability precluded development of mosquito biting rate models for Limbuak and

other outlying islands. An additional limitation to estimating mosquito biting rates was the difficulty

obtaining spatially and temporally resolute remote sensing data for predictors due to high cloud

cover (44). As few positive mosquitoes were identified, uniform estimates of sporozoite rates based

on available data were used across the Matunggong site; if further data was available, these models

could be refined to incorporate estimates of human and macaque density, mosquito biting

preferences in different habitats and infection levels in all hosts (45). Additionally, as this study was

designed to quantitatively estimate time spent in different landscapes, further studies could explore

other aspects of land use, such as the purposes of travel, activities undertaken or practices used to

modify or management land cover.

Despite these limitations, this is the first large-scale study to utilise GPS tracking data and ecological

methods to create fine-scale maps of exposure risk. This study highlights the importance of

incorporating heterogenous patterns of human space use into disease models, as the majority of

human exposure may occur in areas with lower vector biting rates but greater probabilities of

human use. Further, results quantitatively illustrate the importance of forest edges and local habitat

in *P. knowlesi* transmission and can inform understanding of other zoonotic and vector-borne

diseases.

Acknowledgements

We would like to thank the MONKEYBAR project team and the participants in Sabah, Malaysia for

their help with these studies. We acknowledge the Medical Research Council, Natural Environmental

Research Council, Economic and Social Research Council and Biotechnology and Biosciences

Research Council for funding received for this project through the Environmental and Social Ecology

of Human Infectious Diseases Initiative, grant no. G1100796. This work was additionally supported

by the National Socio-Environmental Synthesis Center (SESYNC).

References:

1.   Lambin EF, Tran A, Vanwambeke SO, Linard C, Soti V. Pathogenic landscapes: interactions between land, people, disease vectors, and their animal hosts. Int J Health Geogr. 2010;9:54.
2.   Patz JA, Daszak P, Tabor GM, Aguirre AA, Pearl M, Epstein J, et al. Unhealthy landscapes: Policy recommendations on land use change and infectious disease emergence. Environ Health Perspect. 2004;112(10):1092-8.
3.   Stoddard ST, Forshey BM, Morrison AC, Paz-Soldan VA, Vazquez-Prokopec GM, Astete H, et al. House-to-house human movement drives dengue virus transmission. Proc Natl Acad Sci U S A. 2013;110(3):994-9.
4.   Stoddard ST, Morrison AC, Vazquez-Prokopec GM, Paz Soldan V, Kochel TJ, Kitron U, et al. The role of human movement in the transmission of vector-borne pathogens. PLoS Negl Trop Dis. 2009;3(7):e481.
5.   Pindolia DK, Garcia AJ, Wesolowski A, Smith DL, Buckee CO, Noor AM, et al. Human movement data for malaria control and elimination strategic planning. Malar J. 2012;11(205).
6.   Moyes CL, Shearer FM, Huang Z, Wiebe A, Gibson HS, Nijman V, et al. Predicting the geographical distributions of the macaque hosts and mosquito vectors of Plasmodium knowlesi malaria in forested and non-forested areas. Parasit Vectors. 2016;9(1):242.
7.   Imai N, White MT, Ghani AC, Drakeley CJ. Transmission and control of Plasmodium knowlesi: a mathematical modelling study. PLoS Negl Trop Dis. 2014;8(7):e2978.
8.   Fornace KM, Abidin TR, Alexander N, Brock P, Grigg MJ, Murphy A, et al. Association between Landscape Factors and Spatial Patterns of Plasmodium knowlesi Infections in Sabah, Malaysia. Emerg Infect Dis. 2016;22(2):201-8.
9.   Shearer FM, Huang Z, Weiss DJ, Wiebe A, Gibson HS, Battle KE, et al. Estimating Geographical Variation in the Risk of Zoonotic Plasmodium knowlesi Infection in Countries Eliminating Malaria. PLoS Negl Trop Dis. 2016;10(8):e0004915.
10.   Brock PM, Fornace KM, Grigg MJ, Anstey NM, William T, Cox J, et al. Predictive analysis across spatial scales links zoonotic malaria to deforestation. Proceedings of the Royal Society B. 2019;286(1894).
11.   IPCC. Land use, land-use change, and forestry. Watson RT, Noble IR, Bolin B, Ravindranath NH, Verado DJ, Dokken DJ, editors. UK: Cambridge University Press; 2000.
12.   Acevedo MA, Prosper O, Lopiano K, Ruktanonchai N, Caughlin TT, Martcheva M, et al. Spatial heterogeneity, host movement and mosquito-borne disease transmission. PLoS One. 2015;10(6):e0127552.
13.   Yakob L, Lloyd AL, Kao RR, Ferguson HM, Brock PM, Drakeley C, et al. Plasmodium knowlesi invasion following spread by infected mosquitoes, macaques and humans. Parasitology. 2018;145(1):101-10.
14.   Chang HH, Wesolowski A, Sinha I, Jacob CG, Mahmud A, Uddin D, et al. Mapping imported malaria in Bangladesh using parasite genetic and human mobility data. Elife. 2019;8.
15.   Wesolowski A, Qureshi T, Boni MF, Sundsoy PR, Johansson MA, Rasheed SB, et al. Impact of human mobility on the emergence of dengue epidemics in Pakistan. Proc Natl Acad Sci U S A. 2015;112(38):11887-92.
16.   Wesolowski A, Metcalf CJ, Eagle N, Kombich J, Grenfell BT, Bjornstad ON, et al. Quantifying seasonal population fluxes driving rubella transmission dynamics using mobile phone data. Proc Natl Acad Sci U S A. 2015;112(35):11114-9.
17.   Wesolowski A, Buckee CO, Engo-Monsen K, Metcalf CJE. Connecting Mobility to Infectious Diseases: The Promise and Limits of Mobile Phone Data. J Infect Dis. 2016;214(suppl_4):S414-S20.
18.   Vazquez-Prokopec GM, Bisanzio D, Stoddard ST, Paz-Soldan V, Morrison AC, Elder JP, et al. Using GPS technology to quantify human mobility, dynamic contacts and infectious disease dynamics in a resource-poor urban environment. PLoS One. 2013;8(4):e58802.

19. Grigg MJ, Cox J, William T, Jelip J, Fornace KM, Brock PM, et al. Individual-level factors
associated with the risk of acquiring human Plasmodium knowlesi malaria in Malaysia: a case control
study. Lancet Planetary Health. 2017;1:e97-104.
20. Singh B, Kim Sung L, Matusop A, Radhakrishnan A, Shamsul SS, Cox-Singh J, et al. A large
focus of naturally acquired Plasmodium knowlesi infections in human beings. Lancet.
2004;363(9414):1017-24.
21. Yasuoka J, Levins R. Impact of deforestation and agricultural development on anopheline
ecology and malaria epidemiology. Am J Trop Med Hyg. 2007;76(3):450-60.
22. Barber BE, William T, Dhararaj P, Anderios F, Grigg MJ, Yeo TW, et al. Epidemiology of
Plasmodium knowlesi malaria in north-east Sabah, Malaysia: family clusters and wide age
distribution. Malar J. 2012;11:401.
23. Wong ML, Chua TH, Leong CS, Khaw LT, Fornace K, Wan-Sulaiman WY, et al. Seasonal and
Spatial Dynamics of the Primary Vector of Plasmodium knowlesi within a Major Transmission Focus
in Sabah, Malaysia. PLoS Negl Trop Dis. 2015;9(10):e0004135.
24. Manin BO, Ferguson HM, Vythilingam I, Fornace K, William T, Torr SJ, et al. Investigating the
Contribution of Peri-domestic Transmission to Risk of Zoonotic Malaria Infection in Humans. PLoS
Negl Trop Dis. 2016;10(10):e0005064.
25. Lee KS, Divis PC, Zakaria SK, Matusop A, Julin RA, Conway DJ, et al. Plasmodium knowlesi:
reservoir hosts and tracking the emergence in humans and macaques. PLoS Pathog.
2011;7(4):e1002015.
26. Chua TH, Manin BO, Daim S, Vythilingam I, Drakeley C. Phylogenetic analysis of simian
Plasmodium spp. infecting Anopheles balabacensis Baisas in Sabah, Malaysia. PLoS Negl Trop Dis.
2017;11(10):e0005991.
27. Hartemink N, Vanwambeke SO, Purse BV, Gilbert M, Van Dyck H. Towards a resource-based
habitat approach for spatial modelling of vector-borne disease risks. Biol Rev Camb Philos Soc.
2015;90(4):1151-62.
28. Papworth SK, Bunnefeld N, Slocombe K, Milner-Gulland EJ. Movement ecology of human
resource users: using net squared displacement, biased random bridges and resource utilization
functions to quantify hunter and gatherer behaviour. Methods in Ecology and Evolution.
2012;3(3):584-94.
29. Benhamou S. Dynamic approach to space and habitat use based on biased random bridges.
PLoS One. 2011;6(1):e14592.
30. Fornace KM, Herman LS, Abidin TR, Chua TH, Daim S, Lorenzo PJ, et al. Exposure and
infection to Plasmodium knowlesi in case study communities in Northern Sabah, Malaysia and
Palawan, The Philippines. PLoS Negl Trop Dis. 2018;12(6):e0006432.
31. DAAC NL. MODIS/ Terra Vegetation Indices 16-Day L3 Global 250m Grid SIN V006. Sioux
Falls, South Dakota: USGS Earth Resources Observation and Science (EROS) Center.
32. Johnston R, Brady HE. The rolling cross-section design. Electoral Studies. 2002;21(2):283-95.
33. Hooten MB, Hanks EM, Johnson DS, Alldredge MW. Reconciling resource utilization and
resource selection functions. J Anim Ecol. 2013;82(6):1146-54.
34. Blangiardo M, Cameletti M. Spatial and Spatio-temporal Bayesian Models with R-INLA.
Chichester, United Kingdom: John Wiley & Sons, Ltd; 2015.
35. Sadykova D, Scott BE, De Dominicis M, Wakelin SL, Sadykov A, Wolf J. Bayesian joint models
with INLA exploring marine mobile predator-prey and competitor species habitat overlap. Ecol Evol.
2017;7:5212-26.
36. Barbet-Massin M, Jiguet F, Albert CH, Thuiller W. Selecting pseudo-absences for species
distribution models: how, where and how many? Methods in Ecology and Evolution. 2012;3:327-38.
37. Rue H, Martino S, Chopin N. Approximate Bayesian inference for latent Gaussian models by
using integrated nested Laplace approximations. Journal of the Royal Statistical Society.
2009;71(Part 2):319-92.

38.   Ng SH, Homathevi R, Chua TH. Mosquitoes of Kudat: species composition and their medical
importance (Diptera: Culicidae). Serangga. 2016;21(2):149-62.
39.   Lindgren F, Rue H. Bayesian Spatial Modelling with R-INLA. Journal of Statistical Software.
2015;63(19).
40.   Lindgren F, Rue H, Lindstrom J. An explicit link between Gaussian fields and Gaussian Markov
random fields: the stochastic partial differential equation approach. Statistical Methodology.
2011;73(4):423-98.
41.   Held L, Schrodle B, Rue H. Posterior and Cross-validatory Predictive Checks: A Comparison of
MCMC and INLA. In: Kneib T, Tutz G, editors. Statistical Modelling and Regression Structures:
Physica-Verlag HD; 2010.
42.   Fornace KM, Nuin NA, Betson M, Grigg MJ, William T, Anstey NM, et al. Asymptomatic and
Submicroscopic Carriage of Plasmodium knowlesi Malaria in Household and Community Members of
Clinical Cases in Sabah, Malaysia. J Infect Dis. 2015.
43.   Sektor Penyakit Bawaan Vektor Bahagian Kawalan Penyakit Kementarian Kesihatan
Malaysia. Garis Panduan Pencegahan Malaria Re-Introduction di Malaysia. Putra Jaya: Kementerian
Kesihatan Malaysia; 2016.   Contract No.: KKM/TM/620/2016.
44.   Weiss DJ, Mappin B, Dalrymple U, Bhatt S, Cameron E, Hay SI, et al. Re-examining
environmental correlates of Plasmodium falciparum malaria endemicity: a data-intensive variable
selection approach. Malar J. 2015;14:68.
45.   Yakob L, Bonsall MB, Yan G. Modelling knowlesi malaria transmission in humans: vector
preference and host competence. Malar J. 2010;9:329.

Human Movement

Human GPS tracks collected

- Two week sampling of randomly selected individuals (over age 8)

Biased random bridges (individual space use)

- Estimate size of area used per individual for all movements and 6pm – 6am
- Compare between individuals

Resource utilization functions (community space use)

- Identify spatial and environmental characteristics associated with increased human space use
- Develop predictive models of human space use during peak mosquito biting times

Mosquito Ecology

Human landing catches

- Numbers of *An. balabacensis*
- Identification of *P. knowlesi* sporozoite positive mosquitoes

Spatiotemporal model of *An. balabacensis* biting rates

- Identify spatial and environmental factors associated with increased mosquito density
- Check residual spatial and temporal correlation
- Develop predictive model of biting rates

Probability of human space use during mosquito biting hours

Predicted *An. balabacensis* biting rate

Probability of infection with *P. knowlesi*

**Exposure to *P. knowlesi*
Number of infectious bites on humans**

| ● Tracked houses | ■ Forest | □ Cleared |
| ● Survey houses | ■ Agriculture | □ Water |

Meters	
0 10 20 40 60 80	

○ Human movement

Secondary forest

Houses

Forest

Agriculture

Clearing

Water

0   500   1,000   2,000   Meters

		Meters
0	500	1,000	2,000

● Houses

Total Person Nights

High : 12.7924

Low : 0

Meters

0   500   1,000   2,000

● Houses

Mean biting rate

High : 5.1001

Low : 0.91939

Meters

0   500   1,000   2,000

● Houses

Mean Infected Bites

High : 0.0869811

Low : 0