1   **Title: Augmenting propulsion demands during split-belt walking increases**
2   **locomotor adaptation in the asymmetric motor system**
3

4   **Authors:** Carly J. Sombric, Ph.D.[1], Gelsy Torres-Oviedo, Ph.D.[1]
5   [1]Department of Bioengineering, University of Pittsburgh, Pittsburgh, PA, United States
6
Corresponding Author:
Dr. Gelsy Torres-Oviedo
Department of Bioengineering
4420 Bayard Street
Suite 110, Pitt
Pittsburgh, PA 15213
Univerity of Pittsburgh
gelsyto@pitt.edu

Keywords: gait, locomotion, kinematics, kinetics, stroke, motor learning,
neurorehabilitation, hemiparesis

Abstract

Background:

We previously found that increasing propulsion demands during split-belt walking (i.e.,

legs moving at different speeds) facilitates locomotor adaptation. There is a clinical

interest to determine if this is also the case in stroke survivors.

Objective:

We investigated the effect of propulsion forces on locomotor adaptation during and after

split-belt walking in the asymmetric motor system post-stroke.

Methods:

To test this, 12 chronic stroke subjects experienced a split-belt protocol in a flat and

incline session so as to contrast the effects of two different propulsion demands. Step

length asymmetry and propulsion forces were used to compare the motor behavior

between the two sessions because these are clinically relevant measures that are altered

by split-belt walking.

Results:

The incline session resulted in more adaptation (i.e., less limping) during late split-belt

walking and larger after-effects following split-belt walking. In both testing sessions,

stroke subjects adapted to regain speed and slope-specific leg orientations similarly to

younger adults. These leg orientations achieved during split-belt walking were predictive

of the post-adaptation behavior.

Conclusion:

These results indicated that the asymmetric motor system post-stroke can adapt to meet

leg-specific kinetic demands. This promising finding suggests that augmenting

propulsion demands during split-belt walking could favor symmetric walking in stroke

survivors, perhaps making split-belt interventions a more effective gait rehabilitation

strategy.

Introduction

Brain lesions, such as stroke, may result in asymmetric gait (i.e., limp). Post-

stroke disability is largely due to such gait impairments, which may be why improving

walking is the most common goal of stroke survivors (Jørgensen et al., 1995). It is of

clinical interest to reduce post-stroke gait asymmetry because it can lead to comorbidities

affecting mobility such as musculoskeletal injuries (Jørgensen et al., 2000) and joint pain

(Patterson et al., 2012). Promising studies show that split-belt walking, in which the legs

move at different speeds, could correct gait asymmetries post-stroke (Reisman et al.,

2007, 2009). However, it is not effective in all individuals (Reisman et al., 2013). It has

been suggested that each subject's baseline asymmetries are a factor limiting their ability

to adjust their gait (Malone and Bastian, 2014), raising the question of whether locomotor

adaptation could be increased in this clinical population.

63  Our previous work indicates that locomotor adaptation in young, healthy subjects

64 increases by augmenting propulsion demands during split-belt walking. More

65 specifically, we found that baseline kinetic demands were predictive of step lengths at

66 steady state and after-effects, such that greater propulsion demands led to more

67 adaptation and larger after-effects in every individual (Sombric et al., 2019). It is unclear

68 if the same could be observed post-stroke given their known propulsion deficits

69 (Balasubramanian et al., 2007; Bowden et al., 2006). However, it has been shown that

70 stroke survivors can augment their propulsion forces when required by the task (Awad et

71 al., 2014; Hsiao et al., 2015, 2016; Kesar et al., 2011; Reisman et al., 2013). Thus, we

72 tested whether locomotor adaptation in stroke survivors could be augmented by

73 increasing propulsion demands with inclined split-belt walking.

74  We hypothesized that increasing propulsion demands would lead to more

75 adaptation and after-effects following split-belt walking in stroke survivors. To this end

76 stroke subjects experienced a split-belt adaptation protocol both in a flat and incline

77 environment that had different propulsion demands (Lay et al., 2006, 2007). We

78 expected that stroke subjects' gait adaptation and recalibration would be augmented by

79 incline split-belt walking relative to flat split-belt walking. We also anticipated that that

80 the adaptation and recalibration would be achieved through similar unilateral changes to

81 one step length during adaptation and the other step length following adaptation. These

82 changes in step length were expected to be achieved by recovering speed and slope-

83 specific baseline leg orientations. These anticipated findings would suggest that therapies

84 increasing propulsion demands during walking would be a good strategy for improving

85 post-stroke gait.

Methods

We investigated the effect of augmenting propulsion demands during split-belt

walking on gait adaptation under distinct slopes (i.e., flat and incline), which naturally

increase propulsion forces (Lay et al., 2006, 2007). To this end, we evaluated the

adaptation and after-effects of 12 stroke patients (8 male and 4 female, 61.1 +/- 10.6

years of age) in the chronic phase of recovery (>6 months post-stroke) during separate

flat and incline testing sessions. Stroke subjects were eligible if they (1) had only

unilateral and supratentorial lesions (i.e., without brainstem or cerebellar lesion) as

confirmed by MRI, (2) were able to walk without assistance for 5 minutes at a self-

selected pace, (3) were free of orthopedic injury or pain that would interfere with testing,

(4) had no other neurological condition other than stroke, (5) had no severe cognitive

impairments defined by a mini mental state exam score below 24, (6) could perform

moderate intensity exercise, and (7) did not take medications that altered cognitive

function. Written and informed consent was obtained from all participants prior to

participation. The University of Pittsburgh Institutional Review Board approved the used

experimental protocol, which conformed to the standards set by the Declaration of

Helsinki except for registration in the database.

Table 1. Clinical characteristics of stroke survivors

ID	Age	Gender	Affected Side	Lesion Location	Fugl-Meyer Score	Mid Speed (m/s)	Adapt Strides (flat/ incline)	Post Strides (flat/ incline)	Incline Session Slope (°)
P1	43	Female	R	Left MCA and basal ganglia	33	1.13	907/609	605/303	8.5°
P2	64	Female	R	Left MCA and ACA, temporal lobe, basal ganglia	26	0.81	867/301	642/300	5°
P3	64	Female	R	Left MCA, frontal, parietal lobe and basal ganglia	29	0.60	617/368	307/10	5°
P4	58	Female	R	Left medial, frontal and parietal area's	21	0.45	901/406	625/10	5°
P5	66	Male	R	Left MCA, frontal, temporal and parietal lobes	30	0.77	606/452	599/302	5°
P6	60	Female	R	Left frontal	26	0.9	907/597	600/300	5°
P7	77	Male	R	Thalamus	30	0.35	589/605	598/302	5°
P8	59	Male	R	Left MCA	32	0.7	905/608	600/306	8.5°
P9	52	Male	R	Left MCA	32	0.96	903/602	603/302	5°
P10	66	Male	L	Right frontal superior, parietal and posterior area's	29	0.76	908/519	602/299	8.5°
P11	75	Male	R	Left periventricular, temporal and basal ganglia	32	0.94	913/497	552/306	5°
P12	49	Male	R	Frontotemporal parietal	33	0.71	931/450	303/300	5°

2.1 General Paradigm

All subjects experienced a split-belt protocol while either walking flat or incline throughout two separate experimental sessions (Figure 1A). The flat session was always performed first. The protocol was tailored (i.e., slope, duration, and speed) so that each subject could complete both testing sessions at the same walking speed. The subject-specific walking speed on the treadmill was determined by subtracting 0.35 m/s from each subject's overground walking speed during a Six-Minute Walking Test (Rikli and Jones, 1998). We selected this procedure since it leads to treadmill walking speeds that participants of similar age ranges to our population can sustain during long durations of the split-belt walking condition (Iturralde and Torres-Oviedo, 2019). Treadmill walking speed, labeled as Mid speed, for each participant is presented in Table 1. The speeds experienced during split-belt walking were selected based on subject's mid walking speed. The slow speed was defined as 66.6% of the medium speed, and the fast speed as 133.3% of the same. In this way, the average belt-speed during split-belt walking matched that of baseline and washout, and the belt-speed ratio during split-belt walking was 2:1. We selected an inclination of either 5° or 8.5° based on the level of the subject-specific motor impairments to ensure that all participants could complete the incline session.

Experimental protocols for both sessions consisted of three epochs (i.e., Baseline, Adaptation, and Post-Adaptation). These epochs were used to assess subjects' baseline walking characteristics and subjects' ability to adjust and recalibrate their gait for each

session-specific slope. Baseline: Subjects first experienced a baseline epoch, lasting at

least 50 strides, was used to characterize their baseline gait at the specific inclination used

throughout each session. Subjects walked with both belts moving at the same Mid speed

(Table 1). A baseline epoch with the belts moving at the slow walking speed (i.e., 66.6%

of the Mid speed) was also measured during the flat session. However, this epoch was

removed in the incline session to ensure that all subjects could complete the entire

protocol. Adaptation: Next, the Adaptation epoch was used to assess subjects' ability to

adjust their locomotor pattern in response to a split-belt perturbation. During this epoch,

the non-paretic leg walked twice as fast as the paretic leg. The paretic leg was confirmed

with MRI. The speeds for the fast and slow belts and the duration of the Adaptation

epoch for each subject are shown in Table 1. Post-Adaptation: Finally, The Post-

Adaptation epoch was used to assess the after-effects when the split-belt condition was

removed. Both bets moved at the same Mid speed as in the Baseline epoch. We counted

the number of strides in real-time to regulate the duration for each epoch, where a stride

was defined as the period between two consecutive heel-strikes (i.e., foot landings) of the

same leg. All participants took resting breaks as requested. Also, all subjects wore a

safety harness attached to a sliding rail in the ceiling to prevent falls. In addition, there

was a handrail in front of the treadmill for balance support, but individuals were

encouraged to hold on to it only if needed.

2.2 Data Collection

153   Kinematic and kinetic data were used to characterize subjects' ability to adapt

their gait during Adaptation, and retain the learned motor pattern during Post-

Adaptation. <u>Kinematic Data</u>: Kinematic data were collected with a passive motion

analysis system at 100 Hz (Vicon Motion Systems, Oxford, UK). Subjects' behavior was

characterized with passive reflective markers placed symmetrically on the ankles (i.e.,

lateral malleolus) and the hips (i.e., greater trochanter) and asymmetrically on the shanks

and thighs (to differentiate the legs). The origin of the kinematic data was rotated with

the treadmill in the incline conditions such that the z-axis ('vertical' in the flat condition)

was always orthogonal to the surface of the treadmill (Figure 1B). Gaps in raw kinematic

data were filled with a quintic spline interpolation (Woltring; Vicon Nexus Software,

Oxford Uk). <u>Kinetic Data</u>: Kinetic data were collected with an instrumented split-belt

treadmill at 1,000 Hz (Bertec, Columbus, OH). Force plates were zeroed prior to each

testing session so that each force plate's weight did not affect the kinetic measurements.

In addition, the reference frame was rotated at the inclination of each specific experiment

such that the anterior-posterior forces were aligned with the surface on which the subjects

walked. A heel-strike was identified in real-time when the raw normal force under each

foot reached a threshold of 30 N. This threshold was chosen to ensure accurate counting

of strides at all slopes. On the other hand, we used a threshold of 10 N on median filtered

data (with a 5 ms window) to detect the timing of heel strikes more precisely for data

processing.

2.3 Data Analysis

2.3.1 Kinematic Data Analysis

Kinematic behavior was characterized with step length asymmetry, which exhibits robust adaptation in split-belt paradigms (e.g., Reisman et al., 2005) and is of clinical interest. It is calculated as the difference in step length between the two legs on consecutive steps. Step length (SL) is defined as the distance in millimeters between the ankle markers at heel strike. Therefore, equal step lengths result in zero step length asymmetry. A positive step length asymmetry indicates that the non-paretic leg's step length was longer than the paretic leg's step length. Step length asymmetry was normalized by stride length, which is the sum of two consecutive step lengths, resulting in a unitless parameter that is robust to inter-subject differences in step size.

Each step length was also decomposed into anterior and posterior foot distances relative to the hip position (Figure 1B) as in previous work (Finley et al., 2015). This was done to quantify the leading and trailing legs' positions relative to the body when taking a step because inclination is known to affect these measures (Dewolf et al., 2017; Leroux et al., 2002). The leading leg's position ('α') was computed as the distance in millimeters between the leading leg's ankle and the hip at heel strike; similarly, the trailing leg's position ('X') was computed as the distance in millimeters between the trailing leg's ankle and the hip at heel strike. The hip position, which is a proxy for the body's position, was estimated as the mean instantaneous position across hip markers. By convention

positive α values indicate that the foot landed in front of the hips, whereas negative X

values indicate that the trailing leg was behind the hips. Note that the magnitudes of α

and X summed to the leading leg's step length. As indicated in Figure 1B, α and X were

computed aligned to the treadmill's surface in all sloped conditions.

2.3.2 Kinetic Data Analysis

Kinetic data were used to characterize the adaptation of ground reaction

forces. We focused our analysis on the propulsion component of the anterior-posterior

ground reaction forces for three reasons: 1) propulsion forces are augmented by incline

walking (e.g., Lay et al., 2006, 2007), 2) these are associated with augmented kinematic

adaptation during split-belt walking (Sombric et al., 2019), and 3) they are associated

with hemiparetic gait pathologies (Balasubramanian et al., 2007; Bowden et al.,

2006). The anterior-posterior ground reaction forces (AP forces) were first low-pass

filtered with a cutoff frequency of 20 Hz. Then, they were normalized by each subject's

body weight to account for inter-subject differences. Similar to our previous work, we

computed peak propulsion forces (Sombric et al., 2019) as the maximum AP force ($P_{Paretic}$

and $P_{Non-Paretic}$) excluding the initial positive AP forces following heel strike. Note that we

did not remove slope-specific biases due to gravity because we focused on analyzing

changes in propulsion forces between epochs of interest.

219
Figure 1: Experimental Paradigm and Kinetic and Kinematic
Analysis| (A) Paradigm used for both the flat and incline sessions to assess locomotor
adaptation during and after split-belt walking. Subjects walked flat for the entire flat
session, and incline (either 5° or 8.5°) for the entire incline session. The walking speeds,
duration of epochs, resting breaks and inclination were based on each subject's ability.
(B) The decomposition of step length into leading (α) and trailing (X) leg positions with
respect to the body is illustrated for each sloped condition. This decomposition was done
because it is known that inclination affects these aspects of step length differently
(Dewolf et al., 2017, 2018; Leroux et al., 2002). Also note that when taking a step, the
step length will depend on the position of the leading and trailing leg, which are
generating a braking and propulsion force, respectively. (C) We used the peak propulsion
force for each step to compute outcome measures of interest, such as the ΔAdapt
measure. This measure was computed to quantify increments or reductions in magnitude
within the adaptation epoch of each specific parameter. Note that increases in magnitude
were defined as positive changes, whereas decreases in magnitude were defined as
negative changes.
236

2.3.3 Kinetic and Kinematic Outcome Measures

Outcome measures were used to characterize kinematic and kinetic changes during the Adaptation and Post-Adaptation epochs relative to Baseline or within the Adaptation epoch. Outcome measures of interest were Baseline, Late Adaptation, After-Effects, ΔAdapt, and ΔPost. **Baseline** was defined as the average of the last 40 strides of the Baseline epoch for all parameters. This outcome measure characterized subjects' baseline gait characteristics at each sloped environment and was used as a reference for Late Adaptation and After-Effects. **Late Adaptation** was defined as the difference between the average of the last 40 strides of the Adaptation epoch and Baseline for all parameters. This outcome measure indicated the steady state behavior reached at the end of the Adaptation epoch. **After-Effects** were defined as the difference between the average of the first 5 strides of Post-Adaptation and Baseline values (e.g., Post-Adaptation - Baseline). Positive After-Effect values indicated increments in magnitude of a specific parameter during Post-Adaptation relative to Baseline, and vice versa for negative values. We also characterized the behavioral changes within Adaptation and Post-Adaptation with ΔAdapt and ΔPost, respectively. **ΔAdapt** was computed as the difference between Late Adaptation and Early Adaptation values (i.e., average of the first 5 strides during the Adaptation epoch). **ΔPost** was computed as the difference between Baseline and early Post-Adaptation (e.g., Baseline - Post-Adaptation). Baseline was used instead of late Post-Adaptation because the duration of the Post-Adaptation epoch was not sufficiently long enough in all individuals to extinguish split-belt After-Effects. Thus, Baseline behavior was used a proxy for the late Post-Adaptation behavior. ΔAdapt and

ΔPost were calculated such that an increase in the magnitude of a parameter during either

Adaptation or Post-Adaptation resulted in positive values and a reduction of a parameter

during these epochs resulted in negative values. Figure 1C illustrates an example, to

illustrate ΔAdapt for propulsion forces.

2.4 Statistical Analysis

A significance level of $\alpha=0.05$ was used for all statistical tests. All statistical

analyses were performed either with Stata (StataCorp LP, College Station, TX) or with

MATLAB (The MathWorks, Inc., Natick, Massachusetts, United States).

2. 4. 1. Group analyses

Session averages were compared to determine the effect of slope on each of our

outcome measures using kinetic (e.g., $P_{paretic}$) and kinematic parameters (e.g., step length

asymmetry). We considered that slope might influence gait (Baseline), the extent of

movement update during Adaptation or Post-Adaptation (ΔAdapt and ΔPost), the final

adapted stated reached (Late Adaptation), and/or After-Effects. Thus, the influence of

slope was assessed for each of these outcome measures with paired t-tests. A post-hoc,

one sample t-test was utilized to determine if ΔPost was significantly different from zero

in the flat session.

During baseline walking, it was also of interest to identify differences between the

paretic and non-paretic legs in addition to determining the effect of slope on outcome

measures. Therefore, we performed ANOVAs with individual subjects as a random factor

to account for the paired nature of the data set and slope and leg as fixed, repeated

factors. These ANOVAs were performed on the peak propulsion values and the trailing

leg's position because our study was focused on the propulsion phase of the gait cycle,

which is associated to these two parameters.

The changes of both step lengths during the Adaptation and Post-Adaptation

epochs for the flat and incline sessions was also of interest. Therefore, we performed an

ANOVA with individual subjects as a random factor to account for the paired nature of

the data, and the fixed factors are slope, leg, and epoch. Slope and leg were considered

repeated factors in the analysis. Epoch is not repeated and is treated as a between-subject

factor given that these epochs are not directly associated (Sombric et al., 2019).

2. 4. 2. Regression analyses

We tested the association between leg positions ('α' and 'X') during speed-

specific Baseline and Late Adaptation to determine if Late Adaptation values could be

predicted from Baseline values in stroke survivors, as observed in young unimpaired

subjects (Sombric et al., 2019). We specifically tested the model $|y| = a*|z|$, where y is the

predicted leg position during Late Adaptation and z is the leg position recorded during

Baseline. We also tested the ipsilateral association between the leading leg's position

during Late Adaptation and Post-Adaptation and the contralateral association between the

trailing leg's position during these two epochs in stroke survivors, since these relations

were also observed in young individuals (Sombric et al., 2019). Thus, we tested the

model $|y| = a*|z|$, where y is each leg's position during Early Post-Adaptation and z is

either the ipsilateral 'α' position recorded during Late Adaptation or the contralateral 'X'

position recorded during Late Adaptation. An absolute model was utilized so that the

data would not bias the results of the regression to be linear by having a cluster of

positive (α) and negative (X) data points.

Results

Adaptation and recalibration of step length asymmetry are augmented when

walking incline

Step length asymmetry adaptation and recalibration were augmented by incline

walking. Figure 2A illustrates the evolution of step length asymmetry throughout the flat

and incline sessions. Figure 2B shows a wide range of baseline step length asymmetries

across individuals (colored lines) for each slope condition. On average, these baseline

biases were not different between flat and incline walking (p=0.30). During Adaptation,

participants exhibited similar changes in step length asymmetry from early to late

adaptation (Figure 2D, p=0.75), but they were more symmetric in the incline than the flat

session in Late Adaptation (Figure 2C, p=0.004). Furthermore, the incline session had

larger magnitudes of After-Effects during early Post-Adaptation relative to the flat

 session (Figure 2E, p=0.008). Thus, incline walking augmented the symmetry in step

 lengths during Late Adaptation and the magnitude of After-Effects.

 Figure 2: Step Length Asymmetry Adaptation and Recalibration| (A) Stride-by-
stride time course of step length asymmetry during Baseline, Adaptation, and Post-
Adaptation for each session are shown. Note that each subject's baseline bias has been
removed. Each data point represents the average of 5 consecutive strides and shaded
regions indicate the standard error for each session. For display purposes only, we
include in the time courses stride values that were computed with a minimum of 10
subjects. The black arrow indicates a discontinuity in the data caused by many subjects
taking a resting break at the same time. **(B-E)** The height of the bars indicates group
average step length asymmetry ± standard errors. Individual subjects are represented with
colored dots connected with lines. **(B)** Baseline: Baseline step length asymmetry is not
influenced by slope. **(C)** Late Adaptation: Note that each session plateaued at different
step length asymmetry values during the Adaptation epoch such that subjects reached
more symmetric step lengths in the incline session than the flat session **(D)** ΔAdapt:
Participants changed their gait by similar amounts during the Adaptation epoch in both
sessions. **(E)** After-effects: Subjects had larger After-Effects during early Post-
Adaptation in the incline session than the flat session, which is consistent with the Late
Adaptation differences across sessions.
349

Both step lengths contribute to step length asymmetry adaptation and after-effects during incline walking in the asymmetric motor system

Stroke subjects adjusted both step lengths during split-belt walking. Stroke subjects modulate both their slow (paretic) and fast (non-paretic) step lengths during Adaptation and have After-Effects during Post-Adaptation (Figure 3A). The change of each step length during the Adaptation and Post-Adaptation epochs are quantified in Figure 3B. There was a significant effect of epoch (p_{epoch}=0.001) and interaction between leg and epoch ($p_{leg\#epoch}$<0.001) indicating that the step length with the paretic leg is reduced during Adaptation, but increased during Post-Adaptation and vice versa for the non-paretic leg. Overall, slope did not alter step length changes (p_{slope}=0.16, $p_{slope\#leg}$=0.18, $p_{slope\#epoch}$=0.17), except for the paretic leg's de-adaptation quantified by ΔPost ($p_{leg\#epoch\#slope}$=0.016). More specifically, the paretic step lengths did not exhibit de-adaptation in the flat session (i.e., ΔPost is not different from zero, p=0.38), whereas step lengths for both legs had significant de-adaptation in the incline session (i.e., non-zero ΔPost, p<0.001). Stroke subjects use both their paretic and non-paretic leg to counteract the split-belt perturbation and both legs are recalibrated following incline adaptation.

368
Figure 3: Step length Adaptation and After-Effects.| (A) Stride-by-stride time courses
of step lengths when either the non-paretic leg (top panel, fast leg during Adaptation) or
the paretic leg is leading (bottom panel, slow leg during Adaptation) are shown during
Baseline, Adaptation, and Post-Adaptation. Each data point represents the average of 5
consecutive strides and shaded regions indicate the standard error for each group. For
display purposes only, we include stride values during Post-Adaptation that were
computed with a minimum of 10 subjects. **(B)** The effect of slope on each leg's change
during Adaptation (ΔAdapt) and Post-Adaptation (ΔPost) is illustrated. Note that both the
paretic and non-paretic leg adapted similarly. While the non-paretic leg has recalibrated
(ΔPost≠0) following both the flat and incline session, the paretic leg is only recalibrated
following incline Adaptation.
380

Slope and speed-specific walking demands determine the distinct step length

asymmetries across inclination conditions

Speed and slope-specific leg orientations mediated the distinct step length

asymmetries selected during Late Adaptation and early Post-Adaptation. Figure 4A

illustrates a top-down view of the baseline leg orientations that contribute to each step

length relative to the hips. The stroke subjects' leg positions are diverse across subjects

(colored lines, $p_{Individual}$=0.002), but the trailing leg position, X, is increased in both legs

during incline walking (p_{Slope}=0.042, p_{Leg}=0.22, $p_{Slope\#Leg}$=0.76). The schematic in figure

4B illustrates how young subjects are able to recover speed-specific leg orientations

(Sombric et al 2019). Figure 4C indicates that this is also true for stroke individuals

based on the paretic leg's leg orientation during slow baseline walking and late

adaptation (solid cyan line; $|y|=a*|x|$; 95% confidence interval of a=[0.92, 1.13], R^2=0.76,

p<0.001). We also show as a reference, the relation between (recorded) Baseline and

(predicted) Late Adaptation leg orientation values for both legs and inclinations in young

unimpaired individuals (magenta dashed line; ($|y|=a*|x|$; 95% confidence interval of

a=[0.91, 0.96], R^2=0.89, p<0.001). Note the similarity between the intact and lesioned

behavior (cyan vs. magenta lines). The leg orientations selected during Late Adaptation

influence Post-Adaptation behavior as is illustrated in Figure 4D. Specifically, the

leading leg's orientations were similar before and after removal of the split-belt

perturbation (i.e., Late Adaptation $\alpha_{Paretic}$=Post-Adaptation $\alpha_{Paretic}$ and vice versa)

whereas the trailing legs' orientations were swapped between the legs (i.e. Late

Adaptation $X_{Paretic}$ = Post-Adaptation $X_{Non-Paretic}$ and vice versa). This is supported by the

significant relationship between Late Adaptation and Post-Adaptation leg orientations

observed when individual subjects' values for each leg and both sloped sessions are

regressed (Figure 4E; solid cyan line; $|y|=a*|x|$; 95% confidence interval of a=[0.94,

1.02], R^2=0.79, p<0.001). We also show as a reference, the relation between (recorded)

Late Adaptation and (predicted) Post-Adaptation leg orientation values for both legs and

both sloped conditions in young unimpaired individuals (magenta dashed line; ($|y|=a*|x|$;

95% confidence interval of a=[0.95, 1.03], R^2=0.78, p<0.001). Note the similarity

between the intact and lesioned behavior (cyan vs. magenta lines). Similar to the intact

motor system, the lesioned motor system is able to recover speed and slope-specific leg

orientations during Late Adaptation, which propagate to Post-Adaptation in a predictable

way.

416
Figure 4: Leg orientation Adaptation and After-Effects. | (A) Leg orientations are
depicted for individual subjects (as indicated with different colors) in both the flat and
incline conditions. Note that subjects orient their legs about their bodies differently and
that subjects alter their leg orientation based on slope. Thick vertical black lines indicated
that there is a significant effect of leg (i.e., paretic or non-paretic) and slope (i.e., flat or
incline) on trailing leg positions. **(B)** Schematic of the slow and fast (predicted) baseline
behavior for the paretic and non-paretic leg orientations, respectively. The speed-specific
leg orientations were regained during Late Adaptation. **(C)** The similarity between leg
orientations across the speed-specific Baseline and Late Adaptation epochs is illustrated
by the significant regression (solid cyan line; $|y| = a*|x|$, 95% Confidence interval for a =
[0.92, 1.13]). Recall that a slow Baseline was only collected in the flat session, thus only
the slow Baseline and Late Adaptation for the paretic leg (which walked slow during
Adaptation) are shown. Note that the regression line closely overlaps with the idealized
situation in which baseline and late adaptation values are identical (dashed gray line;
slope of one, i.e., $y = x$) and the behavior of young, healthy adults (sombric et al. 2019,

dashed magenta line). **(D)** Schematic of the leg orientations during early Post-Adaptation. The forward leg positions are ipsilaterally and the trailing leg positions are contralaterally maintained from split-to-tied walking. **(E)** The ipsilateral and contralateral similarity between α and X, respectively, across the Late Adaptation and early Post-Adaptation epochs is quantified with a significant correlation (solid cyan line; $|y| = a*|x|$, 95% Confidence interval for a = [0.94, 1.02]). The idealized situation in which Late Adaptation and early Post-Adaptation values are identical (dashed gray line; slope of one, i.e., $y = x$) and the behavior of young, healthy adults (Sombric et al. 2019, dashed magenta line) are presented as a reference.

442
Figure 5: Propulsion force Adaptation and After-Effects.| (A) Stride-by-stride time
courses of propulsion forces of the non-paretic (top panel) and paretic leg (bottom panel)
are shown during self-selected Baseline, Adaptation, and Post-Adaptation. Each data
point represents the average of 5 consecutive strides and shaded regions indicate the
standard error for each group. For display purposes only, we include stride values during

Post-Adaptation that were computed with a minimum of 10 subjects. **(B-E)** We display
group average values for propulsion force outcome measures ± standard errors.
Individual subjects are represented with colored dots connected with lines. **(B)** Baseline:
Thick horizontal black lines indicated that there is a significant effect of leg (i.e., paretic
or non-paretic) and slope (i.e., flat or incline) on propulsion forces. On average, stroke
subjects generate larger propulsion forces with their non-paretic leg, and they generate
larger propulsion forces with both legs when walking incline. However, some individual
stroke subjects generate larger propulsion forces with their paretic than their non-paretic
leg. **(C)** ΔAdapt: Propulsion forces were similarly modulated during the Adaptation
epoch for both sloped conditions. **(D)** Late Adaptation: Stroke subjects were closer to
their baseline propulsion forces in the incline than the flat sessions. Moreover, baseline
propulsion forces in the incline session were larger than the flat session (Figure 2C).
Taken together, these results suggest that stroke subjects are forced to propel more during
incline split-belt walking with both legs compared to flat split-belt walking. **(E)** After-
Effects: Even though both sloped sessions did not change the extent of propulsion force
adaptation (ΔAdapt), slope influenced the After-Effects for the non-paretic leg, but not
the paretic leg.
465

Larger after-effects of propulsion forces split-belt incline walking

Sloped walking influenced the extent of recalibration of the non-paretic

propulsion forces. It can be seen in Figure 5A that propulsion forces were altered during

the Adaptation epochs. These data are plotted relative to baseline propulsion forces,

which were larger in the incline condition and the non-paretic leg for both sloped

conditions (Figure 5B: $p_{Individual}$=0.007, p_{Slope}<0.0001, p_{Leg}=0.040, $p_{Slope\#Leg}$=0.43). Note

that subjects approached better the larger baseline propulsion values in the incline than

flat session for both legs. In the case of the paretic side, this indicates that subjects were

generating larger propulsion forces in the incline compared to flat during Late Adaptation

(Figure 5C). Even though the Late Adaptation behavior was different across sessions

(Figure 5C; non-paretic propulsion: p=0.032, paretic propulsion: p=0.015), the changes in

propulsion forces from early to late Adaptation were similar across sloped conditions

(Figure 5D; non-paretic propulsion: p=0.92, paretic propulsion: p=0.33). While paretic

propulsion After-Effects are similar in either sloped conditions (Figure 5E, p=0.43), the

non-paretic After-Effects are larger in magnitude following incline adaptation (p=0.015).

Note that the paretic propulsion forces change the most during Adaptation (Figure 5C),

whereas the non-paretic propulsion forces are the ones exhibiting after-effects during

Post-Adaptation (Figure 5E). In summary, incline walking demands greater propulsion

forces in general, which lead to larger paretic propulsion during split-belt walking and

after-effects that reduce the non-paretic propulsion.

Discussion

We investigated the influence of propulsion demands of walking on locomotor

adaptation and recalibration in the asymmetric motor system post-stroke. We find that

subjects adapt more during incline than flat split-belt walking. We also find that leg

orientations during Adaptation are predictive of those Post-Adaptation, leading to greater

step length asymmetry after-effects in the incline than flat sessions. Lastly, the larger

after-effects in step length asymmetry result from shorter paretic step lengths and lower

non-paretic propulsion forces during Post-Adaptation compared to Baseline walking in

the incline session. In summary, the ability to control leg orientation to meet speed and

force demands during split-belt walking is maintained post-stroke, which can be

exploited for designing effective gait rehabilitation interventions in this clinical

population.

Post-stroke gait adapts more in response to larger propulsion demands

We find that stroke subjects behave similarly to young, healthy adults in their

response to sloped split-belt walking. Specifically, stroke subjects are able to augment

their propulsion forces and adjust their leg orientations in response to incline split-belt

walking as observed in young, healthy adults (Sombric et al., 2019). This is consistent

with previous literature indicating that post-stroke patients at the chronic stage can

modulate their gait in response to task demands (Awad et al., 2014; Hsiao et al., 2015,

2016; Kesar et al., 2011, 2014; Reisman et al., 2013). Additionally, we observe that

stroke survivors recover speed and slope-specific leg orientations in the paretic leg for the

flat session. We speculate that the same would have been observed in both slope

conditions and both legs, as observed in young individuals (Sombric et al., 2019). We

think that this is a reasonable expectation given that stroke survivors exhibit similar

control of leg orientations to young adults during Late Adaptation and early Post-

Adaptation for both legs and sloped conditions. Thus, our results provide further evidence

that steady state in the split-belt walking task can be predicted from baseline walking.

It has been previously suggested that baseline gait asymmetries determine the

patients motor behavior at steady state split-belt walking (Malone and Bastian, 2014;

Reisman et al., 2007). Our results provide new insights into the influence of baseline

walking on adaptation. Specifically, we find that stroke survivors recover their baseline

asymmetry in the incline, but not in the flat condition. Thus, it is not baseline gait

asymmetry, but kinetic demands that seem to govern patients' motor patterns. More

specifically, our results suggest that that post-stroke survivors aim to recover the baseline

leg orientations for the specific kinetic demands for each leg in the split condition, as

observed in young adults (Sombric et al., 2019). Further, the forward leg orientation in

the split condition might be adjusted to harness energy from the treadmill (Sánchez et al.,

2019). However, the trailing leg orientation does not match the behavior predicted from

an objective function solely based on minimizing work (Supplementary Figure 1). Thus,

there might be some other factors such as stability (Buurke et al., 2018) or metabolic

energy (Gordon et al., 2009) regulating leg orientation in walking. In summary, the forces

generated to propel one's body forward constitute an important control variable

regulating the adaptation of movements in the intact and asymmetric motor systems.

Bilateral adaptation in stroke survivors contrasts unilateral adaptation in young

adults

Stroke subjects recruit both legs in order to adapt their gait, whereas young adults

primarily adapt one leg. Notably, our results show that stroke survivors adjusted the step

lengths taken with the paretic and non-paretic legs walking on the slow and fast belts,

respectively. In contrast, we have previously observed that young individuals mostly

adjusted the step length of the leg walking on the fast belt (Sombric et al., 2019). This

could be because stroke survivors may require more repetitions in the altered

environment to recover their baseline leg orientation with their paretic leg, whereas intact

subjects can do so immediately after the split condition is introduced. Alternatively, it

could be that the larger neural coupling post-stroke (Kloter et al., 2011) enhances

bilateral adaptation. Regarding post-adaptation, after-effects are only observed in the

paretic leg in the incline condition. More specifically, paretic step lengths become longer

than in baseline walking, which is beneficial for stroke survivors regularly taking short

paretic step lengths (Balasubramanian et al., 2007). On the other hand, after-effects are

observed in the non-paretic leg, regardless of the sloped condition. This is atypical since

the non-paretic leg walked fast in the split condition and young adults only exhibit after-

effects in the leg that walked slow (Sombric et al., 2019). This atypical behavior consists

of shortening step lengths compared to baseline walking and might be a strategy to

recover balance (e.g., Eng et al., 1994), which is challenged upon removal of the split

condition (Buurke et al., 2018; Iturralde and Torres-Oviedo, 2019). In summary, stroke

subjects adapt both legs during split-belt walking, but the paretic step lengths only change

after the incline split condition.

Neurorehabilitation through reinforcement of a corrective pattern during

adaptation, rather than short-lived after-effects post-adaptation.

The long-term therapeutic effect of locomotor adaptation with split-belt treadmills

may be due to walking with the motor demands of the split-belt task, rather than the

After-Effects. Split-belt walking has been shown to reduce step length asymmetries

(Lewek et al., 2018; Reisman et al., 2013). However, it remains an open question what

aspects of the split-belt walking underlie these long term changes. Aftereffects could lead

to motor improvements (Bastian, 2008). Namely, some patients exhibit reduced gait

asymmetry immediately after split-belt walking (Choi et al., 2009; Reisman et al., 2007).

However, these after-effects are short lived and decrease as individuals experience

multiple days of practicing the split-belt condition (Larish et al., 1988; Leech et al., 2018;

Sombric et al., 2017). It is known that regular treadmill walking cannot modify gait

asymmetries post-stroke (Kautz et al., 2005; Den Otter et al., 2006; Silver et al., 2000),

suggesting that the specific motor demands of the split-belt task might be important for

neurorehabilitation. For example, we observe that the split condition forces post-stroke

individuals to take longer paretic step lengths and generate greater paretic propulsion.

Perhaps practice of these gait features through multiple exposure to the split situation

might lead to long term changes in gait symmetry. It is also possible that the strenuous

nature of split-belt walking increases neural plasticity, as shown with other high-intensity

exercises (Andrews et al., 2019). Thus, incline split-belt walking may be beneficial not

only for inducing greater paretic propulsion, but also because it is more demanding than

level walking (Johnson et al., 2002). Lastly, it is also possible that the initial disruption of

step length asymmetry, which is experienced multiple times during split-belt training, is

fundamental for individuals to start exploring new patterns that could converge to more

metabolically efficient gait than their baseline walking pattern (Sánchez et al., 2019;

Selinger et al., 2015). In summary, the long term benefit of split-belt walking may

originate from practicing motor patterns specific to split-belt walking, rather than

reinforcement of those observed during post-adaptation.

Clinical implications

Split-belt walking has been shown to induce long term changes that could

improve the mobility of stroke survivors (Betschart et al., 2018; Lewek et al., 2018;

Reisman et al., 2013). However, there is little understanding for what leg should walk on

the slow or the fast belt during the split condition (Finley et al., 2015; Malone and

Bastian, 2014; Reisman et al., 2007, 2009). For instance, one may consider that the

paretic leg should walk on the slow belt to force stroke survivors to use it more, as a form

of "constrained use therapy" (e.g., Kwakkel et al., 2015). On the other hand, our results

here and in a previous study (Sombric et al., 2019) indicate that placing the paretic leg on

the fast leg would force subjects to augment their paretic propulsion and lengthen their

paretic steps during split-belt walking, which would be beneficial if these were the

patterns that one would like to reinforce. Future studies are needed to determine if stroke

survivors could actually augment their paretic propulsion forces during split-belt walking,

as observed in the fast leg of young adults (Sombric et al., 2019). This remains an open

question given that we observed limited changes in paretic propulsion post-adaptation

compared to those reported in controls (Sombric et al., 2019). In sum, our study provides

greater understanding of the motor demands associated to the split-belt task, which could

be harnessed for gait neurorehabilitation.

The ability to augment locomotor adaptation and recalibration in the lesioned

motor system with incline split-belt training is promising, but future studies are needed to

determine if this will be a more effective intervention than flat split-belt walking. Notably

not all stroke survivors re-learn to walk symmetrically following several training weeks

of flat split-belt walking (Betschart et al., 2018; Lewek et al., 2018; Reisman et al., 2013).

Thus, it is clinically relevant to explore alternative strategies to augment adaptation in

stroke survivors other than increasing the speed difference (Yokoyama et al., 2018) since

not all patients can walk with large speed differences. This work supports that the extent

of adaptation in the split condition can be augmented by increasing the propulsion

demands required by the task, which in turn would possibly make split-belt walking more

effective to more patients. Previous research, however, indicates that overground walking

post-stroke is most improved following decline, rather than incline, interventions due to

greater similarity in motor patterns between the flat and decline conditions (Carda et al.,

2013). Thus, the augmented adaptation in the incline environment may not transfer to

overground walking. Therefore, the efficacy of our protocol as an intervention will

depend on future work assessing its generalization to level walking.

Competing interests:

The authors have no conflicts of interest to report.

Author Contributions

All experiments were performed in the Sensorimotor Learning Laboratory. G.T. and C.S. were

involved with the conception and design of the work. C.S. collected and analyzed the data. C.S

and G.T. interpreted the results. C. S. drafted the manuscript, which was carefully revised by all

authors. The final version of the manuscript has been approved by all the authors who agree to

be accountable for all aspects of the work in ensuring that questions related to the accuracy or

integrity of any part of the work are appropriately investigated and resolved. All authors qualify

for authorship and all those who qualify for authorship are listed.

Funding

639   C.S. is funded by a fellowship from the National Science Foundation (NSF-GRFP); CRDF 4.30204;

640   NSF1535036; AHA 15SDG25710041

Acknowledgements

642   The authors acknowledge the valuable input from Pablo Iturralde and Digna de Kam.

Supplementary Material

644
Supplementary Figure 1: Leg Position Adaptation and After-Effects in the
Asymmetric Motor System| (A) Stride-by-stride time courses of leg positions (α and X)
for the non-paretic and paretic leg are shown during self-selected Baseline, Adaptation,
and Post-Adaptation. Each data point represents the average of 5 consecutive strides and

shaded regions indicate the standard error for each group. The beginning and Late
Adaptation group average behavior are shown for the Adaptation epoch. For display
purposes only, we include stride values during Post-Adaptation that were computed with
a minimum of 10 subjects. **(B)** Schematic of the self-selected Baseline, early Adaptation,
and late Adaptation behavior for the paretic and non-paretic leg orientations, respectively.
Note that there is a general forward movement of the leg position of the non-paretic leg,
but the paretic leg increases both the leading and trailing positions.

Bibliography

Andrews, S. C., Curtin, D., Hawi, Z., Wongtrakun, J., Stout, J. C., and Coxon, J. P.

(2019). Intensity Matters: High-intensity Interval Exercise Enhances Motor Cortex

Plasticity More Than Moderate Exercise Sophie. *Cereb. Cortex*, 1–12.

doi:10.1093/cercor/bhz015.

Awad, L. N., Reisman, D. S., Kesar, T. M., and Binder-Macleod, S. A. (2014). Targeting

paretic propulsion to improve poststroke walking function: A preliminary study.

Arch. Phys. Med. Rehabil. 95, 840–848. doi:10.1016/j.apmr.2013.12.012.

Balasubramanian, C. K., Bowden, M. G., Neptune, R. R., and Kautz, S. A. (2007).

Relationship Between Step Length Asymmetry and Walking Performance in

Subjects With Chronic Hemiparesis. *Arch. Phys. Med. Rehabil.* 88, 43–49.

doi:10.1016/j.apmr.2006.10.004.

Bastian, A. A. J. (2008). Understanding sensorimotor adaptation and learning for

rehabilitation. *Curr. Opin. Neurol.* 21, 628–633.

doi:10.1097/WCO.0b013e328315a293.Understanding.

Betschart, M., McFadyen, B. J., and Nadeau, S. (2018). Repeated split-belt treadmill

walking improved gait ability in individuals with chronic stroke: A pilot study.

Physiother. Theory Pract. 34, 81–90.

Bowden, M. G., Balasubramanian, C. K., Neptune, R. R., and Kautz, S. a. (2006).

Anterior-Posterior Ground Reaction Forces as a Measure of Paretic Leg

Contribution in Hemiparetic Walking. *Stroke* 37, 872–876.

doi:10.1161/01.STR.0000204063.75779.8d.

Buurke, T. J. W., Lamoth, C. J. C., Vervoort, D., van der Woude, L. H. V., and den Otter,

R. (2018). Adaptive control of dynamic balance in human gait on a split-belt

treadmill. *J. Exp. Biol.* 221, jeb174896. doi:10.1242/jeb.174896.

Carda, S., Invernizzi, M., Baricich, A., Cognolato, G., and Cisari, C. (2013). Does

altering inclination alter effectiveness of treadmill training for gait impairment after

stroke? A randomized controlled trial. *Clin. Rehabil.* 27, 932–938.

doi:10.1177/0269215513485592.

Choi, J. T., Vining, E. P. G., Reisman, D. S., and Bastian, A. J. (2009). Walking

flexibility after hemispherectomy: split-belt treadmill adaptation and feedback

control. *Brain* 132, 722–33. doi:10.1093/brain/awn333.

Dewolf, A. H., Ivanenko, Y. P., Lacquaniti, F., and Willems, P. A. (2017). Pendular

energy transduction within the step during human walking on slopes at different

speeds. *PLoS One* 12, 1–25. doi:10.1371/journal.pone.0186963.

Dewolf, A. H., Ivanenko, Y., Zelik, K. E., Lacquaniti, F., and Willems, P. A. (2018).

Kinematic patterns while walking on a slope at different speeds. *J. Appl. Physiol.*

125, 642–653. doi:10.1152/japplphysiol.01020.2017.

Eng, J. J., Winter, D. A., and Patla, A. E. (1994). Strategies for recovery from a trip in

early and late swing during human walking. *Exp. Brain Res.* 102, 339–349.

Finley, J. M., Long, A., Bastian, A. J., and Torres-Oviedo, G. (2015). Spatial and

Temporal Control Contribute to Step Length Asymmetry During Split-Belt

Adaptation and Hemiparetic Gait. *Neurorehabil Neural Repair* 29, 786–795.

doi:1545968314567149.

Gordon, K. E., Ferris, D. P., and Kuo, A. D. (2009). Metabolic and Mechanical Energy

Costs of Reducing Vertical Center of Mass Movement During Gait. *Arch. Phys.*

Med. Rehabil. 90, 136–144. doi:10.1016/j.apmr.2008.07.014.

Hsiao, H., Knarr, B. A., Higginson, J. S., and Binder-macleod, S. A. (2015). Mechanisms

to increase propulsive force for individuals poststroke. *J. Neuroeng. Rehabil.* 12, 1–

8. doi:10.1186/s12984-015-0030-8.

Hsiao, H. Y., Zabielski, T. M., Palmer, J. A., Higginson, J. S., and Binder-Macleod, S. A.

(2016). Evaluation of measurements of propulsion used to reflect changes in

walking speed in individuals poststroke. *J. Biomech.* 49, 4107–4112.

doi:10.1016/j.jbiomech.2016.10.003.

Iturralde, P. A., and Torres-Oviedo, G. (2019). Corrective Muscle Activity Reveals

Subject-Specific Sensorimotor Recalibration. *Eneuro* 6, ENEURO.0358-18.2019.

doi:10.1523/eneuro.0358-18.2019.

Johnson, A. T., Benhur Benjamin, M., and Silverman, N. (2002). Oxygen consumption,

heat production, and muscular efficiency during uphill and downhill walking. *Appl.*

Ergon. 33, 485–491. doi:10.1016/S0003-6870(02)00031-5.

Jørgensen, H. S., Nakayama, H., Raaschou, H. O., and Olsen, T. S. (1995). Recovery of

walking function in stroke patients: the Copenhagen Stroke Study. *Arch. Phys. Med.*

Rehabil. 76, 27–32. Available at: http://www.ncbi.nlm.nih.gov/pubmed/7811170.

Jørgensen, L., Crabtree, N. J., Reeve, J., and Jacobsen, B. K. (2000). Ambulatory level

and asymmetrical weight bearing after stroke affects bone loss in the upper and

lower part of the femoral neck differently: Bone adaptation after decreased

mechanical loading. *Bone* 27, 701–707. doi:10.1016/S8756-3282(00)00374-4.

Kautz, S. A., Duncan, P. W., Perera, S., Neptune, R. R., and Studenski, S. A. (2005).

Coordination of hemiparetic locomotion after stroke rehabilitation. *Neurorehabil.*

Neural Repair 19, 250–258. doi:10.1177/1545968305279279.

Kesar, T. M., Reisman, D. S., Perumal, R., Jancosko, A. M., Higginson, J. S., Rudolph,

K. S., et al. (2011). Combined effects of fast treadmill walking and functional

electrical stimulation on post-stroke gait. *Gait Posture* 33, 309–313.

doi:10.1016/j.gaitpost.2010.11.019.

Kesar, T. M., Sauer, M. J., Binder-Macleod, S. A., and Reisman, D. S. (2014). Motor

learning during poststroke gait rehabilitation: A case study. *J. Neurol. Phys. Ther.*

38, 183–189. doi:10.1097/NPT.0000000000000047.

Kloter, E., Wirz, M., and Dietz, V. (2011). Locomotion in stroke subjects: Interactions

between unaffected and affected sides. *Brain* 134, 721–731.

doi:10.1093/brain/awq370.

Kwakkel, G., Veerbeek, J. M., van Wegen, E. E. H., and Wolf, S. L. (2015). Constraint-

induced movement therapy after stroke. *Lancet Neurol.* 14, 224–234.

doi:10.1016/S1474-4422(14)70160-7.

Larish, D. D., Martin, P. E., and Mungiole, M. (1988). Characteristic Patterns of Gait in

the Healthy Old. *Ann. N. Y. Acad. Sci.* 515, 18–32. doi:10.1111/j.1749-

6632.1988.tb32960.x.

Lay, A. N., Hass, C. J., and Gregor, R. J. (2006). The effects of sloped surfaces on

locomotion: A kinematic and kinetic analysis. *J. Biomech.* 39, 1621–1628.

doi:10.1016/j.jbiomech.2005.05.005.

Lay, A. N., Hass, C. J., Nichols, T. R., and Gregor, R. J. (2007). The effects of sloped

surfaces on locomotion: An electromyographic analysis. *J. Biomech.* 40, 1276–1285.

doi:10.1016/j.jbiomech.2006.05.023.

Leech, K. A., Roemmich, R. T., Day, K. A., and Bastian, A. J. (2018). Movement and

perception recalibrate differently across multiple days of locomotor learning. *J.*

Neurophysiol. 114, 608–623. doi:10.1152/jn.00628.2014.

Leroux, A., Fung, J., and Barbeau, H. (2002). Postural adaptation to walking on inclined

surfaces: I. Normal strategies. *Gait Posture* 15, 64–74.

Lewek, M. D., Braun, C. H., and Wutzke, C. (2018). The role of movement errors in

modifying spatiotemporal gait asymmetry post stroke: a randomized controlled trial.

Clin. Rehabil. 32, 161–172. doi:10.1177/0269215517723056.

Malone, L. A., and Bastian, A. J. (2014). Spatial and Temporal Asymmetries in Gait

Predict Split-Belt Adaptation Behavior in Stroke. *Neurorehabil. Neural Repair* 28,

230–240. doi:10.1177/1545968313505912.

Den Otter, A. R., Geurts, A. C. H., Mulder, T. H., and Duysens, J. (2006). Gait recovery

is not associated with changes in the temporal patterning of muscle activity during

treadmill walking in patients with post-stroke hemiparesis. *Clin. Neurophysiol.* 117,

4–15.

Patterson, K. K., Nadkarni, N. K., Black, S. E., and McIlroy, W. E. (2012). Gait

symmetry and velocity differ in their relationship to age. *Gait Posture* 35, 590–594.

766    doi:10.1016/j.gaitpost.2011.11.030.

767 Reisman, D. S., Block, H. J., Bastian, A. J., Darcy, S., Block, H. J., and Bastian, A. J.

768    (2005). Interlimb coordination during locomotion: what can be adapted and stored?

769    *J. Neurophysiol.* 94, 2403–15. doi:10.1152/jn.00089.2005.

770 Reisman, D. S., McLean, H., Keller, J., Danks, K. A., and Bastian, A. J. (2013). Repeated

771    split-belt treadmill training improves poststroke step length asymmetry.

772    *Neurorehabil. Neural Repair* 27, 460–468. doi:10.1177/1545968312474118.

773 Reisman, D. S., Wityk, R., Silver, K., and Bastian, A. J. (2007). Locomotor adaptation on

774    a split-belt treadmill can improve walking symmetry post-stroke. *Brain* 130, 1861–

775    72. doi:10.1093/brain/awm035.

776 Reisman, D. S., Wityk, R., Silver, K., and Bastian, A. J. (2009). Split-belt treadmill

777    adaptation transfers to overground walking in persons poststroke. *Neurorehabil.*

778    *Neural Repair* 23, 735–744. doi:10.1177/1545968309332880.

779 Rikli, R. E., and Jones, C. J. (1998). The reliability and validity of a 6-minute walk test as

780    a measure of physical endurance in older adults. *J. Aging Phys. Act.* 6, 363–375.

781 Sánchez, N., Simha, S. N., Donelan, J. M., and Finley, J. M. (2019). Taking advantage of

782    external mechanical work to reduce metabolic cost: the mechanics and energetics of

783    split-belt treadmill walking. *J. Physiol.* 15, JP277725. doi:10.1113/JP277725.

784 Selinger, J. C., Shawn, M., Connor, O., Wong, J. D., Maxwell, J., Selinger, J. C., et al.

785    (2015). Humans Can Continuously Optimize Energetic Cost during Walking Report

786    Humans Can Continuously Optimize Energetic Cost during Walking. *Curr. Biol.* 25,

787    2452–2456. doi:10.1016/j.cub.2015.08.016.

788 Silver, K., Macko, R., Forrester, L., Goldberg, A., and Smith, G. (2000). Effects of

aerobic treadmill training on gait velocity, cadence, and gait symmetry in chronic

hemiparetic stroke: a preliminary report. *Neurorehabil. Neural Repair* 14, 65–71.

Sombric, C. J., Calvert, J. S., and Torres-Oviedo, G. (2019). Large propulsion demands

increase locomotor learning at the expense of step length symmetry. in *Frontiers in*

Physiology doi:10.3389/fphys.2019.00060.

Sombric, C. J., Harker, H. M., Sparto, P. J., and Torres-oviedo, G. (2017). Explicit

Action Switching Interferes with the Context-Specificity of Motor Memories in

Older Adults. *Front. Aging Neurosci.* 9, 40.

Yokoyama, H., Sato, K., Ogawa, T., Yamamoto, S.-I., Nakazawa, K., and Kawashima, N.

(2018). Characteristics of the gait adaptation process due to split-belt treadmill

walking under a wide range of right-left speed ratios in humans. *PLoS One* 13,

e0194875. doi:10.1371/journal.pone.0194875.