Progression from feature-specific brain activity to hippocampal

spatial binding during episodic encoding

Rose A. Cooper *[1] & Maureen Ritchey [1]

[1] Department of Psychology, Boston College

* Correspondence concerning this article should be addressed to Dr. Rose Cooper, Boston

College Department of Psychology, 140 Commonwealth Ave, Chestnut Hill, MA 02467. Email:

rose.cooper@bc.edu. Phone: +1 617 552 3805.

Abbreviated title: Evolving encoding-related brain activity

Pages: 30

Figures: 4

Abstract: 225

Introduction: 649

Discussion: 1497

ACKNOWLEDGEMENTS

This work was supported by NIH R00MH103401 grant (M.R.). We thank Max Bluestone,

Rosalie Samide, and Emily Iannazzi for their assistance with data collection. This research was

carried out at the Harvard Center for Brain Science, involving the use of instrumentation

supported by the NIH Shared Instrumentation Grant Program; grant number S10OD020039.

ABSTRACT

The hallmark of episodic memory is recollecting multiple perceptual details tied to a specific spatial-temporal context. To remember an event, it is therefore necessary to integrate such details into a coherent representation during initial encoding. Here we tested how the brain encodes and binds multiple, distinct kinds of features in parallel, and how this process evolves over time during the event itself. We analyzed data from 27 subjects who learned a series of objects uniquely associated with a color, a panoramic scene location, and an emotional sound while fMRI data were collected. By modeling brain activity at event onset and event offset, we were able to test how the neural signatures of individual features as well as the integrated event changed over the course of encoding. We observed a striking dissociation between early and late encoding processes: left inferior frontal and sensory signals at event onset tracked the amount of detail subsequently recalled and were dissociable based on distinct remembered features. In contrast, memory-related brain activity shifted to the left hippocampus at event offset, which was particularly sensitive to binding item color and sound associations with spatial information. These results provide evidence of simultaneous feature-specific neural responses at the start of episodic encoding that predict later remembering and show that the hippocampus acts to integrate these features into a spatially-coherent experience at an event transition.

SIGNIFICANCE STATEMENT

Understanding and remembering complex experiences is crucial for many socio-

cognitive abilities, including being able to navigate our environment, predict the future, and

share experiences with others. Probing the neural mechanisms by which features become

bound into meaningful episodes is a vital part of understanding how we view and reconstruct the

rich detail of our environment. Our findings show a functional dissociation between initial

encoding processes, engaging lateral frontal and sensory regions to prioritize event features,

and later encoding processes that recruit hippocampus to embed these features within a

contextual framework. Identifying the mechanisms of complex event encoding is an important

step toward understanding why many neurological and psychiatric disorders exhibit differences

in how the world is perceived and remembered.

INTRODUCTION

Our ongoing perceptual experience of the world includes a stream of disparate, multimodal features unfolding in parallel. Memory-related increases in brain activity during encoding are often found in visuo-perceptual brain regions (Spaniol et al., 2009; Kim, 2011), emphasizing that stronger, more precise representations of perceptual information support memory formation. For instance, comparing encoding of different kinds of information, such as words and pictures, reveals category-selective patterns of sensory activity that predict subsequent recollection of these individual features (Gottlieb et al., 2010; Duarte et al., 2011; Park and Rugg, 2011). Yet natural experience not only involves representing different perceptual features but, crucially, it requires us to encode them simultaneously. Moreover, we later remember this information as a unified event. In this study, we investigated how perceptual features are uniquely represented during encoding and the neural operations that bind them together.

Studies addressing these issues have largely examined memory for simple paired associations studied as short "events". This research involves presenting subjects with pairs of features within the same trial, such as a location, person, emotion, or color. By predicting subsequent memory separately for each component feature (Uncapher et al., 2006; Staresina and Davachi, 2008; Gottlieb et al., 2012; Ritchey et al., 2018) or contrasting events with overlapping or non-overlapping feature types (Horner et al., 2015), these studies take a step toward understanding how the brain represents and encodes distinct kinds of information in parallel. Moreover, some regions, most notably the hippocampus, show a preference for binding pairs of features during encoding rather than promoting memory for either detail alone (Uncapher et al., 2006; Horner et al., 2015). However, restricting encoding trials to bimodal associations leaves it unclear how distinct feature signals of more complex experiences are simultaneously distinguished, prioritized, and integrated as the event unfolds over time.

The hippocampus has long been considered to be crucial for binding elements of our

experience contextual information (Davachi, 2006; Diana et al., 2007; Eichenbaum et al., 2007;

Ranganath, 2010), though less is known about how exactly the hippocampus organizes

multimodal information. Its involvement might be tied to total memory content, including details

and their associations, during event encoding (Addis and McAndrews, 2006; Staresina and

Davachi, 2008; Park and Rugg, 2011), such that activity tracks the amount of information

subsequently recalled (Qin et al., 2011) regardless of its content or relational structure.

Alternatively, hippocampus could be specifically recruited to organize our environment in a

structured way, perhaps around a spatial framework (Horner et al., 2015; Deuker et al., 2016;

Zeidman and Maguire, 2016). Another outstanding issue is the timing of a binding process: does

it proceed throughout the event or occur once all relevant information has been gathered?

Interestingly, recent evidence using video stimuli suggests that the hippocampus may be

particularly useful for encoding at the end of an event (Ben-Yakov et al., 2014; Ben-Yakov and

Henson, 2018). However, research cannot determine if this later signal serves a binding

function until specific qualities of the resulting memory are measured. Finally, it is important to

distinguish the role of the hippocampus in multimodal event encoding from that of other regions

across the brain that also reliably show encoding effects related to organization and integration,

such as left inferior frontal gyrus (Ranganath et al., 2004; Addis and McAndrews, 2006;

Staresina and Davachi, 2006; Park and Rugg, 2011).

Here, we presented participants with multimodal features constituting a single event. In

line with previous work (Horner and Burgess, 2013; Joensen et al., 2019), we recently showed

that successful recall, but not the precision, of event associations encoded in our task - an

object with a color, scene location, and sound - exhibits a dependent, integrated structure in

memory (Cooper and Ritchey, 2019). Therefore, this paradigm allows us to cleanly test memory

for each event dimension, understand how these features are segregated and prioritized at

encoding, and assess how they are bound into a unified representation across the brain.

MATERIALS AND METHODS

Portions of this dataset have been previously reported (Cooper and Ritchey, 2019). Whereas the previous paper was focused on functional connectivity analyses of the retrieval phase, the current paper reports on univariate activation analyses of the encoding phase. Methods for MRI data collection, the task procedure, and behavioral analyses have been previously detailed in (Cooper and Ritchey, 2019), and so are summarized here.

Participants

27 participants took part in the current experiment (16 females, 11 males). All participants were 18-35 years of age (mean = 21.7 years, SD = 3.58) and did not have a history of any psychiatric or neurological disorders. Seven additional subjects took part but were excluded from data analyses: two participants did not complete the experiment, one due to anxiety and the other due to excessive movement in the MRI scanner, four additional participants had chance-level performance on the memory task, and one subject was excluded after data quality checks revealed 3/6 encoding functional runs exceeded our motion criteria. Informed consent was obtained from all participants prior to the experiment and participants were reimbursed for their time. Procedures were approved by the Boston College Institutional Review Board.

Experimental Design

Paradigm

Participants were presented with a series of 144 unique object "events" in an MRI scanner, 24 per scan run. Each object was presented in a color from a continuous CIEL*A*B color spectrum, in a scene location within one of 6 panoramic environments, and in conjunction with one of 12 sounds - 6 that were emotionally negative and 6 that were neutral. All sounds contained natural, easily recognizable content and were 6 seconds in duration, corresponding to

the time each event was displayed during encoding. Events were separated by a 1 second

fixation. Participants were instructed to integrate the object and its associated features into a

meaningful event, but no response was required (Figure 1A). Allocations of features to objects

as well as the presentation order of events were randomized within each subject. This memory

encoding phase is the focus of all fMRI analyses presented here; see Cooper and Ritchey

(2019) for results from the retrieval phase. After encoding 24 events, subjects were tested on

their memory for the features associated with each object, presented in grayscale as memory

cues. On each retrieval trial, participants attempted to remember all of the features in their mind

for 4s, following which they were prompted to remember if the object was paired with a negative

or neutral sound (2s), and to reconstruct the object's color and spatial location (6s each). For

the sound feature, participants reported their confidence in their decision ('maybe' or 'sure'),

whereas, for the visual features, participants changed the object's color and moved it around the

panorama to recreate its visual appearance as precisely as possible in 360-degree space.

fMRI Data Acquisition

MRI scanning was performed using a 3-T Siemens Prisma MRI scanner at the Harvard

Center for Brain Science, with a 32-channel head coil. Structural MRI images were obtained

using a T1-weighted (T1w) multiecho MPRAGE protocol (field of view = 256 mm, 1 mm isotropic

voxels, 176 sagittal slices with interleaved acquisition, TR = 2530 ms, TE = 1.69/ 3.55/ 5.41/

7.27 ms, flip angle = 7º, phase encoding: anterior-posterior, parallel imaging = GRAPPA,

acceleration factor = 2). Functional images were acquired using a whole brain multiband echo-

planar (EPI) sequence (field of view = 208 mm, 2 mm isotropic voxels, 69 slices with interleaved

acquisition, TR = 1500 ms, TE = 28 ms, flip angle = 75º, phase encoding: anterior-posterior,

parallel imaging = GRAPPA, acceleration factor = 2), for a total of 466 TRs encompassing

encoding and retrieval trials per scan run. Fieldmap scans were acquired to correct the EPI

images for signal distortion (TR = 314 ms, TE = 4.45/ 6.91 ms, flip angle = 55º).

Statistical Analysis

Behavioral Data Processing

As previously reported (Cooper and Ritchey, 2019), participants' responses for the object color and scene location questions were measured in terms of error -- the difference between the target (encoded) feature value and the reported feature value in 360-degree space. Responses to the sound question were considered in terms of accuracy and confidence. Each encoding event was characterized in terms of its pattern of memory success (correct vs. incorrect subsequent retrieval) for each of these features. We used a more conservative method for identifying "correct" memory retrieval than in our previous analysis of this data (Cooper and Ritchey, 2019) to ensure that we only included trials associated with confident subsequent recollection, and to more evenly balance the distribution of trials across different remembered feature combinations. Specifically, the sound feature was considered correctly recalled if the participant successfully identified the associated sound as negative or neutral *and* reported this memory with high confidence. For color and scene location features, we first fitted a mixture model (Zhang and Luck, 2008; Bays et al., 2009) to the aggregate group error data, per feature. The mixture model includes a uniform distribution to estimate the proportion of responses that reflect guessing, as well as a circular Gaussian (von Mises) distribution to estimate the proportion of responses that reflect successful remembering. We defined "correct" color and scene memory by the probability that an error had at least a 75% chance of belonging to the von Mises distribution and not the uniform distribution, resulting in a threshold of +/- 42 degrees for color and +/- 24 degrees for scene memory. For each encoding event, a composite measure of memory detail was calculated as the number of features subsequently recalled (0-3). Of note, this is a slightly different composite measure of memory than we have used previously (Cooper and Ritchey, 2019), where we accounted for variations in confidence and precision within "successful" recall. However, we have already restricted our measure of successful memory for each feature to a confident and precise judgment, and, in our previous behavioral analyses, it

was successful recall but not the precision of recall that showed a dependent structure in

memory.

fMRI Data Preprocessing

All data preprocessing was performed using FMRIPrep v1.0.3 (Esteban et al., 2018) with

the default processing steps. Each T1w volume was corrected for intensity non-uniformity and

skull-stripped. Spatial normalization to the ICBM 152 Nonlinear Asymmetrical template version

2009c was performed through nonlinear registration, using brain-extracted versions of both the

T1w volume and template. All analyses reported here use images in MNI space. Brain tissue

segmentation of cerebrospinal fluid (CSF), white-matter (WM) and gray-matter (GM) was

performed on the brain-extracted T1w image. Functional data was slice time corrected, motion

corrected, and corrected for field distortion. This was followed by co-registration to the

corresponding T1w using boundary-based registration with 9 degrees of freedom. Physiological

noise regressors were extracted applying aCompCor. Framewise displacement was also

calculated for each functional run. For further details of the pipeline, including the software

packages utilized by FMRIPrep for each preprocessing step, please refer to the online

documentation: https://fmriprep.readthedocs.io/en/1.0.3/index.html. After preprocessing, the first

116 TRs were selected from each scan run, capturing all encoding trials and extending 6s

beyond completion of the final trial. Functional encoding runs were checked for motion and were

excluded from data analyses if more than 20% of TRs exceeded a framewise displacement of

0.3 mm. Participants were retained for analyses if at least 4/6 runs passed criteria. One subject

was excluded as result of 3 runs having high motion, but no other subjects had runs removed

because of movement. Four participants successfully completed only 5 out of the 6 scan runs, 3

as a result of exiting the scanner early and 1 due to a technical problem with the sound system

during the first run.

Whole Brain Parcellation

To test which regions across the whole brain were sensitive to subsequent memory effects, we divided the cortex into 200 functionally homogeneous regions using a recent cortical parcellation (Schaefer et al., 2018; https://github.com/ThomasYeoLab/CBIG). Four medial temporal areas were added to this whole-brain atlas (left and right hemisphere as separate regions). The hippocampus was obtained from a probabilistic medial temporal lobe atlas (Ritchey et al., 2015; https://neurovault.org/collections/3731/) and amygdala was added from the Harvard-Oxford subcortical atlas, consistent with our prior methods (Cooper and Ritchey, 2019). We used a parcellation rather than voxel-based approach to i) utilize what we know about anatomical and functional similarities across the brain to maximise statistical power and ii) allow us to draw conclusions about the function of individual brain regions. The results of all whole brain analyses were displayed using BrainNet Viewer (Xia et al., 2013), where, for visualisation purposes, every voxel within a region was allocated the same t statistic calculated from an analysis of that region's mean activity.

Single Trial Analyses

All analyses were run using single trial estimates of activity during encoding. Single trial betas were computed using MATLAB SPM12, implementing the least squares-separate (LS-S) method (Mumford et al., 2012) where a separate general linear model (GLM) was constructed for each encoding trial. In each single trial GLM, the first regressor corresponds to the trial of interest and a second regressor codes for all other encoding trials in that functional run. Functional data were denoised by including framewise displacement, 6 aCompCor principal components, and 6 movements parameters as nuisance regressors. Spike regressors were also included to exclude data points with high motion (> 0.6mm). Two types of single trial model were constructed: 1) a delta function at stimulus *onset* convolved with the canonical HRF, and 2) a delta function at stimulus *offset* convolved with the canonical HRF. Mean single trial onset

and offset beta series were extracted by averaging across voxels within each of the 204 regions

per subject. Analyses of these mean single trial betas were conducted using RStudio (RStudio

Team, 2016).

Before running any analyses, data were first cleaned by z-scoring the single trial beta

values within each subject and region and removing any trials that exceed a z value of |4|. A

total of 103 (2.72%) and 113 (2.98%) out of 3792 trials were removed for analyses of event

onsets and offsets, respectively. The relationship between each region's encoding beta series

and subsequent memory was modeled using linear mixed effects analysis (*lme4*; Bates et al.,

2015). In all models, subject was included as a random effect with a variable intercept and

variable slopes for all fixed effects. This 'maximal' approach guards against an inflated Type 1

error rate (Barr et al., 2013) but can commonly result in model convergence failures with

increasing complexity of the random effect variance-covariance structure (Matuschek et al.,

2017). To address this problem, random effect correlations were constrained to zero in all

models (Barr et al., 2013; Matuschek et al., 2017) using the formula *lmer(Y ~ X_1 + X_2 + (1 + X_1 +*

X_2 || Subject)). To avoid singular fits for more complex models, the random effects structure of

models with more than two fixed effects were additionally simplified using the *lmerTest* package

(Kuznetsova et al., 2017) 'step' function, which iteratively removes random slopes that do not

improve the model fit. Each fixed effect was tested against zero using *lmerTest*, which

implements t-tests with Satterthwaite's approximation method, estimating the effective degrees

of freedom for each effect. All p-values reported were FDR-corrected for multiple comparisons

across all 204 brain regions within each type of mixed effects model.

Code Accessibility

Single trial encoding data for all subjects and brain regions as well as R scripts to run the

analyses described here have been made freely available through GitHub:

RESULTS

Neural correlates of subsequent episodic detail are sensitive to distinct memory features

262   We first sought to test where across the brain activity at the onset of a multi-feature

263 event was sensitive to the amount of detail later recalled (Figure 1B). Activity of each brain

264 region was predicted from an objective measure of memory detail coding each trial according to

265 whether 0, 1, 2, or 3 features were subsequently recalled.

Fig.1. Task design and neural correlates of subsequent memory detail at event onset. A) During

encoding, participants learned a series of events presented for 6s, each uniquely associating an

object to a color, sound, and scene location. Each event was coded by its level of detail,

specified by the number of features (0-3) that were subsequently successfully recollected. B)

Brain regions whose activity at event onset is significantly ($p < .05$ FDR corrected) predicted by

the number of features subsequently recalled. All voxels within a functionally homogeneous

region, defined based on a whole-brain parcellation, are color-coded by the region's t-statistic.

Subsequent memory-related increases in activity at onset were found in a number of

sensory regions, including bilateral occipital and ventral visual regions (ts > 2.94, ps < .047), as

well as bilateral parahippocampal cortex (ts > 3.07, ps < .038), and right retrosplenial cortex (t =

2.96, p = .042), amygdala (t = 3.29, p = .032), and superior temporal cortex (t = 4.54, p = .007).

Notably, left inferior frontal gyrus onset activity also tracked memory detail (ts > 3.40, ps < .024),

with this region being frequently associated with successful associative encoding. In contrast,

activity of default network regions, including bilateral medial prefrontal cortex (mPFC) (ts < -

3.35, ps < .027) and posterior cingulate (ts < -3.14, ps < .037), showed a reduction in activity at

event onset that tracked increasing memory detail. Therefore, left lateral frontal and sensory

regions appear to prioritize event features early on in encoding to support a detailed memory

representation.

We next wanted to determine which of these regions were associated with subsequent

memory in a feature-specific way-- that is, to what extent does encoding a multi-feature event

involve the parallel activation of feature-specific patterns? To this end, mixed effects models

were used to identify patterns of event-onset activity that were *uniquely* predicted by

subsequent memory for color, sound, or scene perspective, with memory coded according to

whether the feature was subsequently correct or incorrect (Figure 2). Whereas deactivation of

default regions was generally important for remembering any feature, regardless of its modality,

more pronounced feature-specific effects emerged in terms of positive subsequent memory

correlates. Successful encoding of the object's color was associated with enhanced activity of

right ventral visual cortical areas (ts > 3.25, ps < .030), whereas subsequent memory for the

sound was positively related to activity in bilateral superior temporal (auditory) cortex (ts > 4.16,

ps < .014). Neural correlates of successful scene memory were more widespread, suggesting a

dominance of spatial information during episodic encoding. Positive changes in activity with

subsequent scene recall were observed in right lateral frontal cortex (ts = 3.49, p = .018),

bilateral dorsal medial parietal and occipital cortex (ts > 2.98, ps < .049), as well as bilateral

retrosplenial and parahippocampal cortex (*t*s > 3.49, *p*s < .032), frequently reported to be

sensitive to specific representations of spatial locations and perspectives (Epstein, 2008; Robin

et al., 2018; Berens et al., 2019). Even though spatial information appears to be prioritized

across the brain, we were able to detect unique patterns of brain activity at the onset of an event

that track different kinds of episodic features encoded simultaneously.

Fig.2. Regions whose activity at event onset is uniquely predicted by subsequent color, sound,

or scene memory while these features are encoded in parallel. Task-positive effects are labeled,

which show mostly distinct patterns across the three features. Effects shown are FDR-corrected

for multiple comparisons at *p* < .05, and all voxels within a region are color-coded by the

region's t-statistic.

Hippocampal offset signals bind episodic features within a spatial framework

The previous analysis showed that there are partially distinct patterns of encoding

activity that support memory for individual features in parallel. Yet a fundamental aspect of

episodic memory is the binding of multimodal features into a coherent event representation.

How does this emerge over the course of encoding? Most models of episodic memory assume

that the hippocampus supports this binding process. Interestingly, we found that hippocampal

onset activity was not predicted by subsequent memory detail ($ts < 0.60$, $ps > .77$), or memory

for any individual feature ($ts < 0.71$, $ps > .68$), despite being frequently reported in this context

(Ranganath et al., 2004; Addis and McAndrews, 2006; Ritchey et al., 2018), specifically for

encoding multiple episodic features (Horner et al., 2015). Yet this onset-locked analysis does

not capture the full temporal dynamics of episodic encoding. We reasoned that feature-specific

brain regions might be important for prioritizing features early on whereas other regions, such as

hippocampus, might be engaged later in encoding to integrate such features into a coherent

memory representation. Thus, we tested whether the pattern of subsequent memory effects

would shift from event onset to event offset, with the hippocampus playing a binding role that is

especially apparent at offset (Figure 3A).

First, activity of each brain region at event *offset* was predicted from a continuous

measure of memory detail. Overall, we found that fewer regions exhibited sensitivity to the

number of features subsequently recalled at the end of an event (Figure 3B), although the

sensitivity of some default regions to memory detail was sustained over encoding, with an

overlapping subset of mPFC regions showing decreased activity with greater subsequent

memory detail at both onset ($ts < - 3.35$, $ps < .027$) and offset ($ts < -2.78$, $ps < .038$). A transition

of subsequent memory effects was particularly the case for left inferior frontal and sensory

regions that tracked later memory detail and feature-specific recall early on during event

learning -- these regions did not sustain their sensitivity to memory over the full course of

encoding. Importantly, enhanced activity at event offset in left hippocampus was predictive of

the number of event details subsequently remembered (t = 3.25, p = .034), reflecting the

emergence of a hippocampal memory effect over the course of the encoding trial. Of note, the

same pattern was also present in right hippocampus, but it did not reach significance after

correction for multiple comparisons across the brain (t = 2.23, p = .13). Positive correlates of

memory detail at event offset were also seen in left inferior parietal cortex (t = 3.29, p = .017)

and middle temporal gyrus (t = 3.58, p = .018). This temporal transformation in the neural

correlates of memory detail during encoding is unlikely to be influenced by the proximity of each

event offset to the onset of the next trial. Specifically, a mixed effects model predicting

subsequent memory detail on the next trial from subsequent memory detail on the current trial,

accounting for nested random effects of subject and task run, revealed no statistical

dependency between encoding success on successive trials (t = 1.16, p = .26).

Fig.3. Whole brain correlates of subsequent memory detail at event offset and the change

memory-related activity in 5 ROIs between event onset and offset. A) Single trial estimates were

constructed by convolving either the event offset (dark gray), compared to event onset (light

gray), with the canonical HRF (TR=1.5s). B) Brain regions whose offset activity is significantly (*p*

< .05 FDR corrected) predicted by the number of details subsequently recalled (0-3). All voxels

within a region are color-coded by the region's t-statistic. C-D) Visualization of the change in

memory-related activity between onset and offset, for subsequent memory detail (C) in left

hippocampus (L HIPP) and left inferior frontal gyrus (L IFG), and for subsequent color, sound,

and scene memory (D) in bilateral ventral temporal cortex (VTC), auditory cortex (AUD), and

retrosplenial/parahippocampal cortex (RSC/PHC), respectively. Bars = fixed effect beta

estimates of memory-modulated activity, error bars = SEM. ** $p <$ .005, * $p <$ .05.

To complement the previous onset-related analyses, we next tested the unique influence

of memory for each feature - color, sound, and scene - on offset activity across the brain.

Whereas there were no significant increases in activity with subsequent color or sound memory,

some regions did, however, track subsequent scene memory. These effects were not present in

retrosplenial/parahippocampal cortex and occipital regions as previously seen; rather, activity of

left hippocampus (t = 3.91, p = .006), bilateral inferior frontal gyrus ($ts>$ 3.16, $ps <$ .034), and left

temporoparietal junction ($ts>$ 3.17, $ps <$ .034) tracked scene encoding at the end of the event.

To visualize the transformation of memory-related activity between event onset and offset, we

focused on 5 regions (Figure 3 C-D) -- left hippocampus (L HIPP), left inferior frontal gyrus (L

IFG), showing sensitivity to total memory detail at event offset and onset, respectively, as well

as ventral temporal cortex (VTC), auditory cortex (AUD), and retrosplenial/parahippocampal

cortex (RSC/PHC), which tracked color, sound, and scene encoding, respectively, at event

onset. To confirm, sensitivity to memory detail was significantly greater at event offset than

onset in L HIPP (F(1,52.9) = 4.95, p = .030), whereas the opposite was true in L IFG (F(1,61.3)

= 6.63, p = .012). AUD and RSC/PHC also showed significantly greater sensitivity to later sound

(F(1,50.5) = 24.30, $p <$ .001) and scene memory (F(1,60.3) = 12.18, $p <$ .001), respectively, at

event onset than offset, but this pattern was not significant for VTC and color encoding

(F(1,59.4) = 1.02, p = .32). Therefore, lateral frontal and sensory memory-related onset signals

transition to a hippocampal offset signal that is sensitive to later scene recollection and the

amount of information recalled.

The observed pattern of hippocampal activity raises two possibilities about its memory-

related function: 1) it may promote memory for the spatial context at an event transition

regardless of other episodic features, or 2) it could reflect a binding signal, integrating event

features within a spatial framework after the individual features have been processed. To

distinguish these explanations, we predicted the mean activity of left hippocampus with all

combinations of features recalled (each feature remembered alone, each possible pair of

features remembered together but the third forgotten, or all three features remembered), and

tested if hippocampal offset activity is greater in situations where space was bound to either

color or sound features, or both, than when space was recalled in the absence of other

information (Figure 4). Indeed, hippocampal activity was only increased at offset for trials where

scene information was successfully remembered in conjunction with color, sound, or both

features (ts > 3.06, ps < .003) and not when scene was the only feature subsequently recalled (t

= 1.29, p = .20; scene-alone vs. scene-binding fixed effects: $F_{(1,2553.9)}$ = 3.51, p = .061, two-

tailed). Hence, although hippocampal activity was uniquely predicted by subsequent memory for

space, it seems particularly sensitive to binding other event features within a spatial framework

at the end of a studied event.

Fig.4. The pattern of L HIPP activity based on all possible feature conjunctions in memory,

relative to an implicit baseline of no features recalled. Hippocampal activity at event onset (left)

was not sensitive to memory for any combination of features. In contrast, hippocampal activity at

offset (right) was increased when spatial information was successfully bound with other episodic

features in memory. Co = Color, Sc = Scene, So = Sound. Bars = fixed effect beta estimates of

activity in each condition, error bars = SEM. * $p < .05$, ~ $p < .1$.

DISCUSSION

This study tested the neural correlates of distinct features encoded in parallel within a

complex event, and how those features are integrated into a coherent representation. First, we

found that left IFG and visuo-perceptual regions were sensitive to the amount of detail encoded

at the onset of each event. Whereas IFG selectively tracked the number of associations

recalled, we uncovered unique variance in sensory brain activity predicted by subsequent

memory for color, sound, and scene features. In contrast, modeling activity at event offset

revealed a shift in the positive neural correlates of successful encoding. Specifically, activity in

left hippocampus at the end of each event was predicted by the amount of detail later recalled

as well as memory for spatial information. Probing specific feature combinations revealed that

these hippocampal effects were driven by binding color and sound information with an

associated scene.

Our findings of feature-specific encoding effects demonstrate that multimodal details of

complex episodes can be decomposed in the brain, even when encoded simultaneously. The

specific pattern of our results converges with univariate and multivariate decoding studies

previously using these types of features, including color (Uncapher et al., 2006; Favila et al.,

2018), sounds (Gottlieb et al., 2012), and scenes (Park and Chun, 2009; Morgan et al., 2011;

Staresina et al., 2011) to study perception and encoding. Moreover, by modeling the unique

variance of each feature, we found that scenes showed the most prominent subsequent

memory effects. This suggests that spatial information may be prioritized during encoding and

serve as the dominant framework for integrating other features within memory (Robin et al.,

2016; Robin, 2018). In fact, complementary work using multivariate pattern analysis of multi-

feature mental imagery also found more widespread neural sensitivity, primarily in posterior

medial cortex, to distinct spatial environments compared to different people or objects (Robin et

al., 2018). We additionally showed that these visuo-perceptual neural correlates were only

present early on in the encoding trial, which is consistent with relatively earlier subsequent

memory effects in ventral visual areas shown using intracranial EEG (Long and Kahana, 2015).

Therefore, while the strength of sensory signals predicts feature-specific encoding, the

importance of this activity for memory generally decreases over the trial. In contrast, we

observed that deactivation of default network regions, particularly medial prefrontal cortex,

correlated with subsequent memory at both the onset and offset of multi-feature events,

suggesting that reallocation of task resources to the external stimulus (Daselaar et al., 2004) is

consistently important for memory encoding.

In addition to feature-specific encoding effects showing an early, transient profile, left

IFG also showed an early relationship to total memory detail that diminished over time. Notably,

its onset activity was not sensitive to memory for any feature alone. This pattern supports the

proposal that IFG processes goal-relevant information and initiates an organizational framework

(Blumenfeld and Ranganath, 2007; Blumenfeld et al., 2011) to support integration of perceptual

features. In our paradigm, participants were asked to generate a story linking the associations

together, and so early IFG activity may help to generate meaningful semantic associations

necessary to support feature binding (Gabrieli et al., 1998). Interestingly, and in contrast to prior

literature, we did not find a similar relationship between hippocampal onset activity and later

encoding success. Instead, we observed a shift whereby left hippocampal activity at event *offset*

significantly predicted the number of features recalled. Collectively, these findings suggest that

both hippocampus and left IFG contribute to the integration of event components (Staresina and

Davachi, 2006; Zeithamova and Preston, 2010) but reveal that their memory signals are

temporally dissociable. As such, while left IFG may control organizational strategies,

hippocampus may act upon this organizational structure to bind an item with its context.

Previous research supports this distinction, showing that IFG generates associations whereas

hippocampus is sensitive to an existing relational structure (Addis and McAndrews, 2006), with

the former tracking subsequent memory confidence and the latter the number of details

remembered (Qin et al., 2011). Another interesting contrast suggests that IFG may actively

control encoding processes while hippocampal signals are driven by inherent memorability of a

scene's perceptual features (Bainbridge et al., 2017). Furthermore, intracranial EEG work has

demonstrated that memory-related activity in IFG precedes that in hippocampus (Long and

Kahana, 2015), thus providing convincing evidence for their distinct functional and temporal

roles in integration during memory encoding.

The absence of hippocampal correlates of subsequent memory at the onset of an event

is particularly interesting in light of inconsistency in the literature. Although hippocampal

encoding effects have often been reported (Spaniol et al., 2009; Kim, 2011), many individual

studies have not found them (Sommer et al., 2005; Haskins et al., 2008; Cooper et al., 2017;

Tibon et al., 2019). Although variability in hippocampal encoding effects might be attributed to

differences in memory strength and trial selection (Henson, 2005), the current results suggest

an additional possible explanation-- namely, that such inconsistencies might be related to the

duration of encoding event used in fMRI analyses. Most event-related fMRI tasks use short

trials, often 2-3 seconds, and analyze activity locked to an event onset, meaning that memory

correlates at different times will be virtually indistinguishable. Meanwhile, studies using longer,

naturalistic events have consistently shown increased hippocampal activity at the end of an

event (Ben-Yakov et al., 2014; Ben-Yakov and Henson, 2018), which correlates with later

memory reinstatement both neurally and behaviorally (DuBrow and Davachi, 2016; Baldassano

et al., 2017). Here, like our whole brain results, this offset response is mostly selective to the

hippocampus. The hippocampal offset signal is thought to reflect a binding operation based on

event discontinuities, linking together all the features that just co-occurred within the same

spatial-temporal context before transitioning to a new environment (Staresina and Davachi,

2009). Evidence that earlier visuo-perceptual activity predicts a hippocampal event boundary

response within continuous experience (Baldassano et al., 2017) provides support for this

proposal and complements the temporal shift from sensory memory correlates to hippocampal

memory effects observed here.

To our knowledge, this is the first study to explicitly test the hypothesis that hippocampal

activity at event onset reflects binding per se; previous studies have not separately tested

memory for distinct kinds of episodic features. Our paradigm allowed us to tease apart which

features and conjunctions influenced this signal most, where left hippocampal activity was

enhanced selectively on trials were spatial information was successfully remembered *and*

integrated with color or sound features, or both. This neural finding corroborates previous

explanations of the hippocampal event offset effect (Cohen et al., 2015; Ben-Yakov and

Henson, 2018) as well as theoretical accounts that hippocampus binds information within a

contextual framework (Davachi, 2006; Ranganath, 2010; Eichenbaum, 2017). Moreover, it

aligns with our previous behavioral analysis of this dataset showing that dependencies in recall

are centered around memory for the object's spatial context, such that color and sound features

were more likely to be bound to the object's scene location than each other (Cooper and

Ritchey, 2019). Evidence of an offset-type hippocampal encoding signal is also present in other

event-related tasks: When learning overlapping associations across trials, hippocampal activity

predicts subsequent memory after being exposed to component event associations (Horner et

al., 2015) and binds previously studied associations based on shared context (Zeithamova and

Preston, 2010; Zeithamova et al., 2012), suggesting a prominent function is the integration of

previously encoded event features.

When interpreting the results of this study, it is important to consider some limitations of

the current experimental approach. First, although our events were longer than those in most

event-related encoding studies, 6 seconds is still relatively short and as such, we were unable to

map out the full time course of memory-related signals over a longer period. This partially

reflects the fact that, even though participants were encouraged to generate a story to encode

the events, the features were static rather than evolving over time. Although this design enabled

us to measure memory for distinct event features, future studies would benefit from using

temporally dynamic videos while controlling event features that can be objectively tested later

on. Second, our findings of spatial primacy, in terms of sensory signals and later hippocampal

binding, might not necessarily demonstrate that space is always the dominant contextual

framework. Rather, the object's background scene may have been particularly salient in our

task, meaning that attention and binding was centered around this feature. Future work could

therefore manipulate both the content of an event and the relative salience of those features to

explore how these factors modulate onset and offset binding effects in IFG and hippocampus.

In summary, we demonstrated that feature-specific neural representations are

dissociable and support encoding early in an event. Integration of these features into a coherent

episode as an event unfolds is likely supported by complementary lateral frontal and

hippocampal signals, with early left IFG operations providing a meaningful structure, and later

hippocampal operations integrating these associations within a unified spatial-temporal

framework. This multifaceted process thus promotes subsequent episodic retrieval, helping us

to mentally reconstruct the rich detail of our environment and predict the future.

REFERENCES

Addis DR, McAndrews MP (2006) Prefrontal and hippocampal contributions to the generation

and binding of semantic associations during successful encoding. Neuroimage 33:1194–

1206.

Bainbridge WA, Dilks DD, Oliva A (2017) Memorability: A stimulus-driven perceptual neural

signature distinctive from memory. Neuroimage 149:141–152.

Baldassano C, Chen J, Zadbood A, Pillow JW, Hasson U, Norman KA (2017) Discovering Event

Structure in Continuous Narrative Perception and Memory. Neuron 95:709–721.e5.

Barr DJ, Levy R, Scheepers C, Tily HJ (2013) Random effects structure for confirmatory

hypothesis testing: Keep it maximal. J Mem Lang 68 Available at:

http://dx.doi.org/10.1016/j.jml.2012.11.001.

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4.

Journal of Statistical Software, Articles 67:1–48.

Bays PM, Catalao RFG, Husain M (2009) The precision of visual working memory is set by

allocation of a shared resource. J Vis 9:7.1–11.

Ben-Yakov A, Henson R (2018) The hippocampal film-editor: sensitivity and specificity to event

boundaries in continuous experience. J Neurosci Available at:

http://dx.doi.org/10.1523/JNEUROSCI.0524-18.2018.

Ben-Yakov A, Rubinson M, Dudai Y (2014) Shifting gears in hippocampus: temporal

dissociation between familiarity and novelty signatures in a single event. J Neurosci

34:12973–12981.

Berens SC, Joensen BH, Horner AJ (2019) Tracking the emergence of location-based spatial

representations. bioRxiv:547976 Available at:

https://www.biorxiv.org/content/10.1101/547976v1.

Blumenfeld RS, Parks CM, Yonelinas AP, Ranganath C (2011) Putting the pieces together: the

role of dorsolateral prefrontal cortex in relational memory encoding. J Cogn Neurosci

23:257–265.

Blumenfeld RS, Ranganath C (2007) Prefrontal cortex and long-term memory encoding: an

integrative review of findings from neuropsychology and neuroimaging. Neuroscientist

13:280–291.

Cohen N, Pell L, Edelson MG, Ben-Yakov A, Pine A, Dudai Y (2015) Peri-encoding predictors of

memory encoding and consolidation. Neurosci Biobehav Rev 50:128–142.

Cooper RA, Richter FR, Bays PM, Plaisted-Grant KC, Baron-Cohen S, Simons JS (2017)

Reduced Hippocampal Functional Connectivity During Episodic Memory Retrieval in

Autism. Cereb Cortex 27:888–902.

Cooper RA, Ritchey M (2019) Cortico-hippocampal network connections support the

multidimensional quality of episodic memory. Elife 8 Available at:

http://dx.doi.org/10.7554/eLife.45591.

Daselaar SM, Prince SE, Cabeza R (2004) When less means more: deactivations during

encoding that predict subsequent memory. Neuroimage 23:921–927.

Davachi L (2006) Item, context and relational episodic encoding in humans. Curr Opin Neurobiol

16:693–700.

Deuker L, Bellmund JL, Navarro Schröder T, Doeller CF (2016) An event map of memory space

in the hippocampus. Elife 5 Available at: http://dx.doi.org/10.7554/eLife.16534.

Diana RA, Yonelinas AP, Ranganath C (2007) Imaging recollection and familiarity in the medial

temporal lobe: a three-component model. Trends Cogn Sci 11:379–386.

Duarte A, Henson RN, Graham KS (2011) Stimulus content and the neural correlates of source

memory. Brain Res 1373:110–123.

DuBrow S, Davachi L (2016) Temporal binding within and across events. Neurobiol Learn Mem

134 Pt A:107–114.

Eichenbaum H (2017) Memory: Organization and Control. Annu Rev Psychol 68:19–45.

Eichenbaum H, Yonelinas AP, Ranganath C (2007) The medial temporal lobe and recognition

memory. Annu Rev Neurosci 30:123–152.

Epstein RA (2008) Parahippocampal and retrosplenial contributions to human spatial

navigation. Trends Cogn Sci 12:388–396.

Esteban O, Markiewicz CJ, Blair RW, Moodie CA, Ilkay Isik A, Erramuzpe A, Kent JD,

Goncalves M, DuPre E, Snyder M, Oya H, Ghosh SS, Wright J, Durnez J, Poldrack RA,

Gorgolewski KJ (2018) fMRIPrep: a robust preprocessing pipeline for functional MRI. Nat

Methods:1.

Favila SE, Samide R, Sweigart SC, Kuhl BA (2018) Parietal representations of stimulus features

are amplified during memory retrieval and flexibly aligned with top-down goals. J Neurosci

Available at: http://dx.doi.org/10.1523/JNEUROSCI.0564-18.2018.

Gabrieli JD, Poldrack RA, Desmond JE (1998) The role of left prefrontal cortex in language and

memory. Proc Natl Acad Sci U S A 95:906–913.

Gottlieb LJ, Uncapher MR, Rugg MD (2010) Dissociation of the neural correlates of visual and

auditory contextual encoding. Neuropsychologia 48:137–144.

Gottlieb LJ, Wong J, de Chastelaine M, Rugg MD (2012) Neural correlates of the encoding of

multimodal contextual features. Learn Mem 19:605–614.

Haskins AL, Yonelinas AP, Quamme JR, Ranganath C (2008) Perirhinal cortex supports

encoding and familiarity-based recognition of novel associations. Neuron 59:554–560.

Henson R (2005) A mini-review of fMRI studies of human medial temporal lobe activity

associated with recognition memory. Q J Exp Psychol B 58:340–360.

Horner AJ, Bisby JA, Bush D, Lin W-J, Burgess N (2015) Evidence for holistic episodic

recollection via hippocampal pattern completion. Nat Commun 6:7462.

Horner AJ, Burgess N (2013) The associative structure of memory for multi-element events. J

Exp Psychol Gen 142:1370–1383.

Joensen BH, Gaskell MG, Horner AJ (2019) United we fall: All-or-none forgetting of complex

episodic events. J Exp Psychol Gen Available at: http://dx.doi.org/10.1037/xge0000648.

Kim H (2011) Neural activity that predicts subsequent memory and forgetting: a meta-analysis

of 74 fMRI studies. Neuroimage 54:2446–2461.

Kuznetsova A, Brockhoff P, Christensen R (2017) lmerTest Package: Tests in Linear Mixed

Effects Models. Journal of Statistical Software, Articles 82:1–26.

Long NM, Kahana MJ (2015) Successful memory formation is driven by contextual encoding in

the core memory network. Neuroimage 119:332–337.

Matuschek H, Kliegl R, Vasishth S, Baayen H, Bates D (2017) Balancing Type I error and power

609  in linear mixed models. J Mem Lang 94:305–315.

610 Morgan LK, Macevoy SP, Aguirre GK, Epstein RA (2011) Distances between real-world

611  locations are represented in the human hippocampus. J Neurosci 31:1238–1245.

612 Mumford JA, Turner BO, Ashby FG, Poldrack RA (2012) Deconvolving BOLD activation in

613  event-related designs for multivoxel pattern classification analyses. Neuroimage 59:2636–

614  2643.

615 Park H, Rugg MD (2011) Neural correlates of encoding within- and across-domain inter-item

616  associations. J Cogn Neurosci 23:2533–2543.

617 Park S, Chun MM (2009) Different roles of the parahippocampal place area (PPA) and

618  retrosplenial cortex (RSC) in panoramic scene perception. Neuroimage 47:1747–1756.

619 Qin S, van Marle HJF, Hermans EJ, Fernández G (2011) Subjective sense of memory strength

620  and the objective amount of information accurately remembered are related to distinct

621  neural correlates at encoding. J Neurosci 31:8920–8927.

622 Ranganath C (2010) Binding Items and Contexts: The Cognitive Neuroscience of Episodic

623  Memory. Curr Dir Psychol Sci 19:131–137.

624 Ranganath C, Yonelinas AP, Cohen MX, Dy CJ, Tom SM, D'Esposito M (2004) Dissociable

625  correlates of recollection and familiarity within the medial temporal lobes. Neuropsychologia

626  42:2–13.

627 Ritchey M, Montchal ME, Yonelinas AP, Ranganath C (2015) Delay-dependent contributions of

628  medial temporal lobe regions to episodic memory retrieval. Elife 4 Available at:

629  http://dx.doi.org/10.7554/eLife.05025.

630 Ritchey M, Wang S-F, Yonelinas AP, Ranganath C (2018) Dissociable medial temporal

631  pathways for encoding emotional item and context information. Neuropsychologia Available

632  at: http://www.sciencedirect.com/science/article/pii/S0028393218307826.

633 Robin J (2018) Spatial scaffold effects in event memory and imagination. Wiley Interdiscip Rev

634  Cogn Sci 9:e1462.

Robin J, Buchsbaum BR, Moscovitch M (2018) The Primacy of Spatial Context in the Neural

Representation of Events. J Neurosci 38:2755–2765.

Robin J, Wynn J, Moscovitch M (2016) The spatial scaffold: The effects of spatial context on

memory for events. J Exp Psychol Learn Mem Cogn 42:308–315.

RStudio Team (2016) RStudio: Integrated Development for R. RStudio, Inc., Boston, MA URL

http://www.rstudio.com/

Schaefer A, Kong R, Gordon EM, Laumann TO, Zuo X-N, Holmes AJ, Eickhoff SB, Yeo BTT

(2018) Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional

Connectivity MRI. Cereb Cortex 28:3095–3114.

Sommer T, Rose M, Weiller C, Büchel C (2005) Contributions of occipital, parietal and

parahippocampal cortex to encoding of object-location associations. Neuropsychologia

43:732–743.

Spaniol J, Davidson PSR, Kim ASN, Han H, Moscovitch M, Grady CL (2009) Event-related fMRI

studies of episodic encoding and retrieval: meta-analyses using activation likelihood

estimation. Neuropsychologia 47:1765–1779.

Staresina BP, Davachi L (2006) Differential encoding mechanisms for subsequent associative

recognition and free recall. J Neurosci 26:9162–9172.

Staresina BP, Davachi L (2008) Selective and shared contributions of the hippocampus and

perirhinal cortex to episodic item and associative encoding. J Cogn Neurosci 20:1478–

1489.

Staresina BP, Davachi L (2009) Mind the gap: binding experiences across space and time in the

human hippocampus. Neuron 63:267–276.

Staresina BP, Duncan KD, Davachi L (2011) Perirhinal and parahippocampal cortices

differentially contribute to later recollection of object- and scene-related event details. J

Neurosci 31:8739–8747.

Tibon R, Fuhrmann D, Levy DA, Simons JS, Henson R (2019) Multimodal integration and

vividness in the angular gyrus during episodic encoding and retrieval. J Neurosci Available

at: http://dx.doi.org/10.1523/JNEUROSCI.2102-18.2018.

Uncapher MR, Otten LJ, Rugg MD (2006) Episodic encoding is more than the sum of its parts:

an fMRI investigation of multifeatural contextual encoding. Neuron 52:547–556.

Xia M, Wang J, He Y (2013) BrainNet Viewer: a network visualization tool for human brain

connectomics. PLoS One 8:e68910.

Zeidman P, Maguire EA (2016) Anterior hippocampus: the anatomy of perception, imagination

and episodic memory. Nat Rev Neurosci 17:173–182.

Zeithamova D, Dominick AL, Preston AR (2012) Hippocampal and ventral medial prefrontal

activation during retrieval-mediated learning supports novel inference. Neuron 75:168–179.

Zeithamova D, Preston AR (2010) Flexible memories: differential roles for medial temporal lobe

and prefrontal cortex in cross-episode binding. J Neurosci 30:14676–14684.

Zhang W, Luck SJ (2008) Discrete fixed-resolution representations in visual working memory.

Nature 453:233–235.