Full Title: Quantitative trait loci (QTL) underlying phenotypic variation in bioethanol-

related processes in *Neurospora crassa*.

Short Title: Bioethanol QTL

Joshua C. Waters[1,2], Deval Jhaveri[2], Justin C. Biffinger[3], and Kwangwon Lee[1,2,*]

[1] Department of Biology, Rutgers, The State University of New Jersey - Camden, 315

Penn Street, Camden, NJ, United States of America

[2] Center for Computational & Integrative Biology, Rutgers, The State University of New

Jersey - Camden, 200 Federal Street, Camden, NJ, United States of America

[3] Chemistry Department, University of Dayton, Dayton OH., 300 College Park, Dayton

OH, United States of America

[*] Corresponding author

E-mail: kwagnwon.lee@rutgers.edu

Keywords: Fungal breeding, bioethanol, QTL, natural variation

Abstract

Bioethanol production from lignocellulosic biomass has received increasing attention over the past decade. Many attempts have been made to reduce the cost of bioethanol production by combining the separate steps of the process into a single-step process known as consolidated bioprocessing. This requires identification of organisms that can efficiently decompose lignocellulose to simple sugars and ferment the pentose and hexose sugars liberated to ethanol. There have been many attempts in engineering laboratory strains by adding new genes or modifying genes to expand the capacity of an industrial microorganism. There has been less attention in improving bioethanol-related processes utilizing natural variation existing in the natural ecotypes. In this study, we sought to identify genomic loci contributing to variation in saccharification of cellulose and fermentation of glucose in the fermenting cellulolytic fungus *Neurospora crassa* through quantitative trait loci (QTL) analysis. We identified one major QTL contributing to fermentation of glucose and multiple putative QTL's underlying saccharification. Understanding the natural variation of the major QTL gene would provide new insights in developing industrial microbes for bioethanol production.

Introduction

Fermentation of cellulosic biomass by microorganisms is a complex process requiring coordinated regulation of multiple pathways, including production and secretion of at least three distinct classes of catabolic enzymes (cellulase, hemicellulase, and lignin peroxidase enzymes), transport of substrates into the cell, followed by catabolism and fermentation of the resultant monomeric C5 and C6 sugars. Identification of organisms efficient in all requisite processes has been a major limitation toward realizing the vision of the consolidated bioprocessing (CBP) of biomass to ethanol for fuels (1–4). While many researchers attempt to engineer existing strains by introducing the genes for pathways lacking in the given organisms, success has been limited due to the increased physiological burdens imposed by genetic manipulation. There is also a growing concern that genetic manipulations that are effective in laboratory strains may not be translatable to more robust strains used in industry (3, 5). There are, however, organisms that already possess the genetic elements needed to perform all of the requisite processes, such as the model filamentous fungus *Neurospora crassa* [9].

These types of native microorganisms present excellent potential sources of useful industrial enzymes and products. Breeding has proven to be a reliable method for manipulating genomes to select for desired traits, as evidenced by the successes of plant breeding (6). Selective mating has long been used in plant breeding to improve traits ranging from fruit size and yield to appearance or flavor (7, 8). Recently, selective breeding has also been used in fungi in attempts to select for increased saccharification and fermentation potential (9).

The advent of computational tools such as genome-wide association studies

(GWAS) and quantitative trait loci (QTL) analysis have had drastic impacts on how plant

geneticists design new lineages with improved traits and qualities (10). GWAS and QTL

analysis are computational tools for investigating the genetic elements underlying

quantitative traits. Unlike qualitative traits such as color, which are inherited discretely

and exhibit discontinuous variation, quantitative traits exhibit continuous variation and

cannot be discretely grouped (10). GWAS and QTL use polymorphic genetic markers to

predict the extent to which a given genetic locus contributes to observed variation in

quantitative traits, such as plant height, and researchers have been able to use genetic

methods to identify combinations of desired alleles in individuals that are then used for

breeding, known as marker assisted selection (MAS).

QTL analysis and MAS have revolutionized plant genetics and should be equally

useful for improvement of quantitative traits in fungi, such as saccharification and

fermentation of cellulose. While a wealth of information is available on the effects of

gene deletions or overexpression of cellulolytic enzymes or fermentation pathways, there

have been few studies to characterize the variation resulting from allelic variation in

populations, most of which focus solely on fermentation (11–18, 18–21). While gene

deletions and overexpression studies can implicate genetic elements in these processes,

these manipulations increase cellular burden due to unknown pleiotropic effects, and

multiple manipulations can result in compounding deficits (3, 5, 22, 23).

The inherent complexity of lignocellulose metabolism allows for abundant

sources of variation that will affect CBP performance. Understanding how allelic

variation attenuates performance will lead to a more robust understanding of cellulose

metabolism and identify superior combinations of alleles for trait enhancement for CBP.

To this end, we chose to perform QTL analysis of bioethanol-related processes,

specifically saccharification and fermentation, on a laboratory generated population of

the model filamentous fungus *Neurospora crassa*. 111 offspring and two parental strains

were genotyped by sequencing, evaluated for their ability to decompose cellulose and

ferment glucose, and subjected to QTL analysis. A major QTL was identified for

fermentation and multiple putative QTL's were identified for saccharification of

cellulose.

Materials and methods

Strains

Parental strains FGSC2223 (Mat a, collected in Iowa, USA) and FGSC4825 (Mat A,

collected in Ivory Coast, Africa) from the Fungal Genetics Stock Center (FGSC) (24),

and their cross progeny (designated as the N6 population) were grown from long-term

stock prior to each experiment on Vogel's minimal media slants (1X Vogel's Salts, 2%

Sucrose, 1.5% Agar, pH 5.8). Spore suspensions in High Glucose Liquid Media (HGLM)

(1X Vogel's salts, 2% glucose, 0.5% L-Arginine, pH 5.8) were used to generate mycelial

mats in petri plates, which were used to generate replicate mycelial pads for each

experiment using a bore punch.

Enzyme activity assay

A modified FPA assay was performed using 96-well plates as described by Camassola

and Dillon, in which secreted protein extracts were taken from wells in 6-well plates

containing 1% CMC broth (1X Vogel's salts, 1% CMC) 4 days after inoculation with 10

gauge mycelial pads (25). Before inoculating the mycelia pads, they were rinsed in sterile

water three times to remove media. Culture broth containing secreted enzyme was

filtered and centrifuged at 13.2k rpm to remove any fungal cells or debris. 50μL of

supernatant was added to 100μL of 50mM sodium acetate buffer pH 5.6 in a 96-well

deep-well plate, which was then equilibrated to 50°C for 5 min in a hot-water bath. A

5mg strip of Whatman Grade No. 1 filter paper was submerged in the solution, and the

plate was incubated at 50°C for 60 min. After 60 min, 300μL of 3,5-dinitrosalicylic acid

(DNS) Reagent was added to stop the enzymatic reaction and visualize glucose

equivalents. The plate was incubated at 100°C for 10 min to develop color, then

transferred to an ice bath to stop color development. 100μL of Enzyme/DNS mixture was

transferred to a clear-bottom assay plate, diluted with 200μL of deionized H_2O, and

absorbance was measured at 545 nm. The concentration of reducing equivalents released

was determined with standard curves generated with glucose standards of known

concentration. All samples were performed in biological triplicates. The activity could

not be quantified in conventional Filter Paper Units because of the low concentration of

secreted enzymes. Therefore, these results were presented in the form of the

concentration of reducing equivalents released in each aliquot.

Fermentations and ethanol quantification

To characterize fermentation of glucose among the lab generated N6 first filial (F1)

generation and the parental strains (FGSC 2223 and FGSC4825), fermentation was

carried out in a 96-well format in deep-well plates. Biological replicates for each strain

were collected from mycelial mats grown from spore suspension in high glucose liquid

media (HGLM) with a 6 gauge punch and inoculated into 750µl of HGLM (2% glucose),

sealed with aluminum Thermowell^{TM} seals, and allowed to ferment for 7 days in 12:12

Light/Dark conditions at 25°C. All samples were performed in biological quadruplicate.

After fermentation, 600 µL of media was recovered and cell debris was removed by

sequential centrifugation at 13.2k rpm for 5 min. The recovered supernatant was

aliquoted into HPLC vials for ethanol analysis by HPLC. HPLC quantitation was

performed using a Varian ProStar HPLC with a Varian ProStar Autosampler and a

Varian 356-LC Refractive Index Detector. An isocratic elution was used with an Agilent

Hi-Plex H$^+$ (300 mm x 7.7 mm ion-exchange column) with 5mM H$_2$SO$_4$ at a flow rate of

0.7 ml/min at 60°C. The concentration of ethanol present was determined from a standard

curve based on ethanol standards with a known concentration.

Qualitative Trait Loci (QTL) Analysis

To investigate the underlying sources of variation in saccharification and fermentation, a

qualitative trait loci (QTL) analysis was performed using R to identify candidate genes

contributing to the observed variation. N6 population has been used for QTL analysis

using simple sequence repeat markers. For the current study, we generated a single

nucleotide polymorphism (SNP) library using the Genotyping By Sequencing (GBS)

approach (26). GBS data for the current study are available at (SRA accession:

PRJNA594422.). Briefly, 50,000 SNP markers were filtered to 146 evenly distributed

informative markers using Python and later Excel. First, all markers that were not

polymorphic between parents were removed, followed by any markers in which one of

the two parents was missing data. The resulting 4900 markers were formatted for Excel

for further filtering. Chi-square tests were performed for each marker and those markers

with unequal segregation among progeny (>20% disparity) were removed, followed by

markers with >10% missing data among progeny. The filtered genotype data was

combined with phenotype data from FPA and glucose fermentation assays and formatted

for R-QTL. The formatted data was imported into R-QTL as and the create map function

was used to generate a linkage map from a physical map based on recombination

frequencies. Markers were then hand curated to generate a linkage map for all 7

chromosomes (Linkage groups) of *Neurospora crassa* with approximately 22 evenly

spaced markers per chromosome.

After generating a linkage map for the N6 population, QTL scans were performed

for each phenotype. Composite interval mapping (CIM) was used to scan for single

QTL's using the maximum likelihood (em), Haley-Knott regression (hk), and multiple

imputation (imp) methods in the R-QTL package, allowing for up to 4 covariate markers

within a 20-marker window. Likelihood odds ratio (LOD) thresholds were generated for

95% and 90% confidence intervals for each method based on 1000 permutations of the

data. Putative QTL were designated as major QTL if they explained greater than 10% of

the observed variation (27). Peak marker positions for each chromosome (based on LOD

score) were used to identify linkage blocks between flanking markers to identify

chromosomal regions containing the suspected QTL, according to their physical

positions. These chromosomal regions were searched in the Fungi Database

(FungiDB.org) to identify all genes within the region to find candidate genes contributing

to the observed variation.

Results

To test if there exist a natural variation of the bioethanol-related processes in one

mapping population N6, we performed a QTL analyses in N6. Significant variation was

observed among the N6 population for both saccharification of cellulose and

fermentation of glucose (Fig 1). Both traits demonstrated transgressive segregation, with

21 or 58 of 111 offspring outperforming the more proficient parent and 66 or 33 of 111

underperforming the lesser parent in the FPA assay or fermentation of glucose,

respectively (t-test, p = 0.00232, 0.0340).

Figure 1. Natural variation in cellulolytic activity and fermentation among the N6

population. Significant variation was observed for saccharification of cellulose (left axis)

and fermentation of glucose (right axis) among the N6 population (ANOVA p=3.99×10^{-33}

and p=8.75×10^{-17} respectively). Each data point represents the mean of 3 biological

replicates. Red bars represent mean with SEM. Red filled triangle represents FGSC4825.

Blue filled triangle represents FGSC2223.

Due to the low level of recombination from only 1 meiotic event in a single cross,

generating a map with evenly spaced markers required filtering a large number of

polymorphic markers, reducing the power of QTL analysis. A linkage map based on

recombination frequencies was generated from a physical map of SNP markers with

approximately 22 evenly spaced markers on each of *N. crassa's* 7 chromosomes (Fig 2).

Figure 2. Linkage map of SNP markers for the N6 population. A linkage map created

in R-QTL from physical marker positions, with approximately 22 evenly distributed

markers per chromosome. Locations and distance are measured in centimorgans (cM).

A major QTL region contributing to variation in fermentation was identified on

Linkage Group (LG) I (Nc-M878 = CM002236: 7449359: 361 Genes - 160 Annotated,

142 Hypothetical, 59 Unspecified) within a 90% confidence interval (CI) (p-value=0.075)

(Fig 3). Additionally, minor QTL's were observed for fermentation on LG IV (Nc-

M2879 = CM002239: 4344007: 344 Genes - 158 Annotated, 134 Hypothetical, 52

Unspecified), LG VI (Nc-M4092 = CM002241: 1218567: 62 Genes - 28 Annotated, 32

Hypothetical, 2 Unspecified), and LG VII (Nc-M4866 = CM002236: 1218567: 77 Genes

- 30 Annotated, 37 Hypothetical, 10 Unspecified), however, none of these were above the

LOD threshold for 90% CI. Marker Nc-M878 on LG I lies within DUF1212 domain

membrane protein (NCU00717). Markers Nc-M2879 on LG IV and Nc-M4092 on LG VI

lie within hypothetical proteins (NCU07332 and NCU05646 respectively), while marker

Nc-M4866 on LG VII lies within an intergenic region.

Figure 3. QTL analysis of Fermentation of Glucose to Ethanol by the N6 population.

Composite Interval Mapping single-QTL scan for fermentation of glucose using linear

regression (EM Algorithm). The y-axis is a measure of the logarithm of the odds ratio

(LOD). Tick marks along the x-axis delineate relative marker positions, while alternating

grey bands denote different chromosomes Blue line represents 90% confidence interval.

Using the fitqtl function, it was observed that the 4 QTL markers account for

~26% of the observed variation in fermentation of ethanol. The QTL on LG I accounts

for 13% of the observed variation, while QTL on LG IV, VI and VII account for 4, 7, and

6% of the variation, respectively. Interestingly, marker segregation analysis revealed that

the parental genotypes contributing to greater ethanol production varied between

markers, with parental genotype A accounting for higher production at marker Nc-M878

on LG I and genotype B accounting for higher production at marker Nc-M4092 on LG VI

(Fig 4A and B).

Figure 4. Phenotype Segregation at QTL Markers for Fermentation. A) Phenotype
segregation at the QTL marker on LG I according to genotype. B) Phenotype segregation
at the QTL marker on LG VI according to genotype. The error bars represent the error of
the mean, n = 4 biological replicates. The data points in red are missing data at the given
marker, however based on imputation it is inferred that they would be that genotype at
the given marker and they are therefore plotted with that group.
235

Of particular interest in the linkage block in LG I, indicated by QTL analysis for

fermentation, was a cluster of annotated glycolytic enzymes, sugar transporters, enzymes

involved in alternate fates of pyruvate, including lactate dehydrogenase enzymes, and

components of energy production machinery (Table 1). Alternatively, the only annotated

gene of interest from the QTL on LG VI was another sugar transporter, *sut-7*.

Table 1. Candidate QTL genes for ethanol fermentation.

Gene ID	Gene Name	Description	Linkage Group	Location
NCU00629	emp-3	6-phosphofructokinase	I	CM002236: 7,764,287 - 7,768,767 (-)
NCU00575	gki-2	glucokinase-2	I	CM002236: 7,953,683 - 7,958,506 (-)
NCU00742	glp-1	glycerol-3-phosphate dehydrogenase	I	CM002236: 7,360,766 - 7,362,918 (+)
NCU00720	tca-17	L-lactate dehydrogenase	I	CM002236: 7,427,034 - 7,428,944 (+)
NCU00775	tca-6	isocitrate dehydrogenase subunit 1	I	CM002236: 7,214,344 - 7,217,183 (-)
NCU00780	ldh-4	lactate dehydrogenase-4	I	CM002236: 7,202,837 - 7,204,284 (+)
NCU00606	null	short-chain alcohol dehydrogenase	I	CM002236: 7,858,582 - 7,859,522 (-)
NCU00801	cdt-1	MFS lactose permease, Sugar (and other) transporter	I	CM002236: 7,121,829 - 7,125,021 (-)
NCU00821	sut-15	sugar transporter-15	I	CM002236: 7,056,000 - 7,058,684 (-)
NCU00809	sut-26	sugar transporter-26	I	CM002236: 7,088,949 - 7,091,552 (+)
NCU05627	sut-7	sugar transporter-7	VI	CM002241: 1,275,879 - 1,278,130 (-)

Similarly, multiple QTL were observed for saccharification on LG II (Nc-M1321

= CM002240: 1924972: 184 Genes - 84 Annotated, 64 Hypothetical, 36 Unspecified),

LG IV (Nc-M2984 = CM002239: 4860803: 184 Genes - 74 Annotated, 72 Hypothetical,

38 Unspecified), and LG VI (Nc-M4097 = CM002241: 1342332: 144 Genes – 44

Annotated, 54 Hypothetical, 16 Unspecified), although none were above the LOD

threshold for 90% CI (Fig 5). Marker Nc-M1321 on LG II lies within an intergenic

region, while markers Nc-M2984 on LG IV lies within a hypothetical protein

(NCU07004) and marker Nc-M4097 on LG VI lies within phosphatidylinositol 3-kinase

tor2 (NCU05608). Combined, the putative QTL account for 19.5% of the total variation

observed in saccharification of cellulose. The major QTL on LG II accounts for 11% of

the observed variation, while minor QTL on LG IV and VI account for 4.6% and 5.7%,

respectively.

Figure 5. QTL analysis of Saccharification of Cellulose by the N6 population.

Composite Interval Mapping single-QTL scan for FPA assay using linear regression (EM

Algorithm). The y-axis is a measure of the logarithm of the odds ratio (LOD). Tick marks

along the x-axis delineate relative marker positions, while alternating grey bands denote

different chromosomes.

Marker segregation analysis was performed to confirm the estimation of variance

analysis; which showed that the QTL in Chr2 contributed 11% of the variance (ANOVA

F=14.647, p = 0.0002), while the one in Chr4 only contributed 4.6% variance (ANOVA

F=6.078, p = 0.015) (Fig 6A and B). From this finding, we chose to investigate the

potential for epistatic interactions among the QTL markers for saccharification of

cellulose using R/QTL's Scantwo function. Scantwo analysis revealed that the QTL

marker on LG II (Nc-M1321) exhibited strong interactions with markers across all

chromosomes, with the strongest interactions with the adjacent marker on LG II and the

putative QTL marker on LG IV (Nc-M2984) (Fig 7A). Marker interaction plots

demonstrated that the genotype B specific increases in FPA activity only held for the

interactions between QTL markers on LG II and LG IV, but not with the interactions

between adjacent markers on LG II (Fig 7B and C).

Figure 6. Phenotype Segregation at QTL Markers for Saccharification. A)

Phenotype segregation at the QTL marker on LG II according to genotype. B) Phenotype

segregation at the QTL marker on LG IV according to genotype. The error bars represent

the error of the mean, n = 3 biological replicates. The data points in red are missing data

at the given marker, however based on imputation it is inferred that they would be that

genotype at the given marker and they are therefore plotted with that group.

Figure 7. Saccharification QTL Marker Interactions. (A) Scantwo QTL analysis for

epistatic marker interactions. Heatmap legend represents logarithm of the odds ratio

(LOD) scores for marker interactions. (B) Average phenotypes according to genotype at

sequential markers on LG II [Marker Nc-1321 is QTL marker on LG II]. (C) Average

phenotypes according to genotype at QTL markers on LG II and LG IV. Colors represent

genotype at marker indicated in legend, while letters on x-axis indicate genotype at

marker indicated on x-axis title. Error bars represent standard deviations.

Among those genes within the QTL regions for saccharification there were some

glucanase and xylanase enzymes, and transcription factors, which may have a role in

sensing or degrading cellulose. However, two major regulatory genes of interest were

also identified, *vib-1* and *xlr-1*, on LG II and LG IV respectively (Table 2).

Table 2. Candidate QTL genes involved in saccharification of cellulose.

Gene ID	Gene Name	Description	Linkage Group	Location
NCU03725	vib-1	VIB-1, variant 1	II	CM002240: 2,215,819 - 2,219,221 (+)
NCU06971	xlr-1	transcriptional activator xlnR	IV	CM002239: 4,706,112 - 4,709,560 (-)
NCU01050	gh61-4	endoglucanase II	II	CM002240: 2,769,232 - 2,770,004 (+)
NCU03641	gh3-1	Beta-glucosidase 2, variant 1	II	CM002240: 1,920,114 - 1,923,804 (+)
NCU01059	gh47-3	glycosyl hydrolase family 47 protein	II	CM002240: 2,790,429 - 2,792,645 (-)
NCU01050	gh61-4	endoglucanase II	II	CM002240: 2,769,232 - 2,770,004 (+)
NCU03657	gh63-1	mannosyl-oligosaccharide glucosidase, variant 1	II	CM002240: 1,992,478 - 1,995,662 (+)
NCU01080	gh64-1	glucanase B	II	CM002240: 2,848,665 - 2,850,629 (-)
NCU06961	gh28-2	Exopolygalacturonase	IV	CM002239: 4,669,933 - 4,671,737 (+)
NCU07326	gh43-6	hypothetical protein	IV	CM002239: 4,359,254 - 4,361,686 (-)
NCU07005	gh76-9	glycosyl hydrolase family 76-9	IV	CM002239: 4,865,028 - 4,867,282 (-)
NCU07269	gh92-2	alpha-1,2-mannosidase	IV	CM002239: 4,569,439 - 4,572,146 (-)

Discussion

In current study, we have identified that there exists a substantial variation in bioethanol-

related traits in the mapping population, N6. We also identified a few major candidate

genes with the potential to contribute to the observed variation, including the clustered

glucose metabolism genes on LG I for fermentation, and transcriptional regulators *vib-1*

and *xlr-1* on LG II and LG IV, respectively, for saccharification. The clustering of

glucose metabolism genes on LG I makes identification of candidate genes for further

analysis more complicated, requiring gene expression analysis for identification of those

genes within the region that are differentially regulated and contributing to the observed

variation in ethanol fermentation potential. While it is possible that an increased rate of

glycolysis is responsible for increased fermentative capacity, the presence of lactate

dehydrogenase enzymes within the QTL regions could present an alternative answer,

with decreased lactic acid fermentation resulting in increased rates of ethanol

fermentation.

The presence of major metabolic regulators within QTL regions for

saccharification provide more promising QTL candidate genes. The first major gene of

interest for saccharification of cellulose is the transcription factor, vegetative

incompatibility blocked-1 (*vib-1*), located on LG II, which is involved in sensing carbon

starvation and signaling for induction of the cellulolytic response (28). The second major

gene of interest for saccharification of cellulose indicated is the xylose degradation

regulator (*xlr-1*) located on LG IV, as it is involved in induction of the hemicellulose

response, and regulation of cellodextrin transporters and cellulose degradation regulator

clr-1, along with *vib-1*, identified in the putative QTL on LG II (12, 29).

Although *xlr-1* has not been shown to directly bind to promoters for cellulase

enzymes, it has been implicated in cellulase production. Transcriptional analyses

demonstrated that *xlr-1* knockouts have reduced hemicellulase and cellulase production

when grown in xylan inducing conditions (12, 17). ChIP-seq experiments further

revealed that although *xlr-1* does not directly bind to cellulase promoters it does bind to

the promoter of transcription factor *clr-1*, a major regulator of cellulase expression, and

can be found bound to targets in conjunction with *clr-1* and/or *clr-2* (12). Furthermore,

xlr-1 is involved in the regulation of *vib-1*, which in turn further regulates expression of

clr-1 (12). Similarly, Cai et al., demonstrated that *xlr-1*, not *clr-1/2*, is required for

expression of cellodextrin transporter-2 (*cdt-2*), which is involved in sensing cellulose in

the environment as well as transmembrane transport (30). Although cdt-2 was mildly

reduced in the clr-1 knockout, cdt-2 expression was completely abolished with the

deletion of xlr-1, suggesting that that xlr-1 is the main positive regulator of cdt-2 in N.

crassa (27). *Cdt-2* has been implicated as a major cellodextrin importer for facilitating

induction of the cellulase response, as cellulase production and growth on cellulose are

significantly reduced in *cdt-2* knockouts (30, 31). Overexpression of *cdt-2* has also been

shown to increase cellulase and hemicellulase production under cellulose and xylan

conditions (30). These data could suggest pleiotropic or epistatic roles for *xlr-1* as a key

component for cellulase production through its role in regulating of *vib-1* and *cdt-2* for

carbon sensing and cellulase induction, *clr-1* for inducing the cellulolytic response via

clr-2, and the co-regulation of genes targeted by *clr-1/2*. However, further experiments

would be needed to address this.

Among other filamentous fungi, *xlr-1* orthologues (*xlnR/xyr-1*) have been shown

to directly regulate cellulase expression alongside hemicellulase, although this role has

been lost in *Neurospora* (12, 15, 17, 18, 32). However, a vestige of this role may be

evident in *xlr-1*'s indirect role in the response to cellulose through its regulation of

critical genes involved in sensing cellulose in the environment, especially through *vib-1*,

clr-1, *cdt-2* and major facilitator superfamily (MFS) sugar transporters. While RNA-seq

and ChIP-seq experiments have previously been employed to decipher the role of *xlr-1* in

N. crassa, these studies have used knockout and overexpression strains that share the

same genetic scaffold as the wild type reference. Similar studies using diverse natural

strains may demonstrate that differential expression of *xlr-1* leads to differences in

cellulolytic potential or illustrate potentially disparate roles arising from allelic variation

at the *xlr-1* locus or its targets.

Allelic variation in lignocellulose degradation regulators and their target genes

may provide unique insights into the lignocellulose response of *N. crassa*. Transcriptomic

profiling and expression analysis may reveal differential interactions between *xlr-1* and

its' targets, resulting in differential enzyme induction and degradation of cellulosic

substrates. Furthermore, expression QTL (eQTL) analysis would be important for

discerning the pleiotropic effects of *xlr-1* from its epistatic effects. Understanding the

implications of allelic effects on lignocellulose metabolism should allude to elite

genotypes for further enhancement and possible industrial applications in bioethanol

production.

Acknowledgements

369   The authors thank Drs. Sunil Shende and Charot Rodeget for curating Python scripts,

370   critical reading and insightful comments.

Author Contributions

Conceptualization: Kwangwon Lee and Joshua C. Waters

Project Administration: Kwangwon Lee

Funding Acquisition: Kwangwon Lee

Methodology: Joshua C. Waters and Kwangwon Lee

Supervision: Joshua C. Waters and Kwangwon Lee

Investigation: Joshua C. Waters, Deval Jhaveri, Justin C. Biffinger

Validation: Joshua C. Waters, Deval Jhaveri, Justin C. Biffinger, Kwangwon Lee

Data curation: Deval Jhaveri

Software: Joshua C. Waters

Formal Analysis: Joshua C. Waters

Visualization: Joshua C. Waters

Writing-original draft: Joshua C. Waters and Kwangwon Lee

Writing-review and editing: Joshua C. Waters and Kwangwon Lee

Conflict of Interest Disclosure: The authors declare no conflict of interest.

References

1.   M. Dashtban, H. Schraft, W. Qin, Fungal Bioconversion of Lignocellulosic Residues;
Opportunities & Perspectives. *Int. J. Biol. Sci.* **5**, 578–595 (2009).

2.   I. Dogaris, D. Mamma, D. Kekos, Biotechnological production of ethanol from renewable
resources by Neurospora crassa: an alternative to conventional yeast fermentations? *Appl.*
Microbiol. Biotechnol. **97**, 1457–1473 (2013).

3.   L. Favaro, *et al.*, Designing industrial yeasts for the consolidated bioprocessing of starchy
biomass to ethanol. *Bioengineered* **4**, 97–102 (2013).

4.   H. Mattila, J. Kuuskeri, T. Lundell, Single-step, single-organism bioethanol production and
bioconversion of lignocellulose waste materials by phlebioid fungal species. *Bioresour.*
Technol. **225**, 254–261 (2017).

5.   G.-C. Zhang, T. L. Turner, Y.-S. Jin, Enhanced xylose fermentation by engineered yeast
expressing NADH oxidase through high cell density inoculums. *J. Ind. Microbiol. Biotechnol.*
44, 387–395 (2017).

6.   A. R. Hallauer, M. J. Carena, J. B. M. Filho, *Quantitative Genetics in Maize Breeding*, 3rd Ed.
(Springer-Verlag, 2010) (June 27, 2019).

7.   H. Muranty, *et al.*, Accuracy and responses of genomic selection on key traits in apple
breeding. *Hortic. Res.* **2**, 15060 (2015).

8.   X. Qiu, *et al.*, Examining two sets of introgression lines reveals background-independent
and stably expressed QTL that improve grain appearance quality in rice (Oryza sativa L.).
Theor. Appl. Genet., 1–17 (2017).

9.   J. C. Waters, A. Nixon, M. Dwyer, J. C. Biffinger, K. Lee, Developing elite Neurospora crassa
strains for cellulosic ethanol production using fungal breeding. *J. Ind. Microbiol.*
Biotechnol., 1–8 (2017).

10.   J. Kumar, *et al.*, Quantitative trait loci from identification to exploitation for crop
improvement. *Plant Cell Rep.*, 1–27 (2017).

11.   J. P. Benz, *et al.*, A comparative systems analysis of polysaccharide-elicited responses in
Neurospora crassa reveals carbon source-specific cellular adaptations. *Mol. Microbiol.* **91**,
275–299 (2014).

12.   J. P. Craig, S. T. Coradetti, T. L. Starr, N. L. Glass, Direct Target Network of the Neurospora
crassa Plant Cell Wall Deconstruction Regulators CLR-1, CLR-2, and XLR-1. *mBio* **6**, e01452-
15 (2015).

13.   A. Hildebrand, E. Szewczyk, H. Lin, T. Kasuga, Z. Fan, Engineering Neurospora crassa for
Improved Cellobiose and Cellobionate Production. *Appl. Environ. Microbiol.* **81**, 597–603
(2015).

14. M. C. Reilly, L. Qin, J. P. Craig, T. L. Starr, N. L. Glass, Deletion of homologs of the SREBP
pathway results in hyper-production of cellulases in Neurospora crassa and Trichoderma
reesei. *Biotechnol. Biofuels* **8**, 1–18 (2015).

15. A. R. Stricker, R. L. Mach, L. H. de Graaff, Regulation of transcription of cellulases- and
hemicellulases-encoding genes in Aspergillus niger and Hypocrea jecorina (Trichoderma
reesei). *Appl. Microbiol. Biotechnol.* **78**, 211 (2008).

16. J. Sun, N. L. Glass, Identification of the CRE-1 Cellulolytic Regulon in Neurospora crassa.
PLoS ONE **6**, e25654 (2011).

17. J. Sun, C. Tian, S. Diamond, N. L. Glass, Deciphering Transcriptional Regulatory Mechanisms
Associated with Hemicellulose Degradation in Neurospora crassa. *Eukaryot. Cell* **11**, 482–
493 (2012).

18. S. Tani, S. Kanamasa, J. Sumitani, M. Arai, T. Kawaguchi, XlnR-independent signaling
pathway regulates both cellulase and xylanase genes in response to cellobiose in
Aspergillus aculeatus. *Curr. Genet.* **58**, 93–104 (2012).

19. C. Tian, *et al.*, Systems analysis of plant cell wall degradation by the model filamentous
fungus Neurospora crassa. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 22157–22162 (2009).

20. Z. Wang, *et al.*, QTL analysis reveals genomic variants linked to high-temperature
fermentation performance in the industrial yeast. *Biotechnol. Biofuels* **12**, 59 (2019).

21. P. Marullo, *et al.*, Natural allelic variations of Saccharomyces cerevisiae impact stuck
fermentation due to the combined effect of ethanol and temperature; a QTL-mapping
study. *BMC Genomics* **20** (2019).

22. H. Sakuragi, K. Kuroda, M. Ueda, Molecular Breeding of Advanced Microorganisms for
Biofuel Production. *J. Biomed. Biotechnol.* **2011** (2011).

23. G. Wu, *et al.*, Metabolic Burden: Cornerstones in Synthetic Biology and Metabolic
Engineering Applications. *Trends Biotechnol.* **34**, 652–664 (2016).

24. K. McCluskey, A. Wiest, M. Plamann, The Fungal Genetics Stock Center: a repository for 50
years of fungal genetics research. *J. Biosci.* **35**, 119–126 (2010).

25. M. Camassola, A. J.P. Dillon, Cellulase Determination: Modifications to Make the Filter
Paper Assay Easy, Fast, Practical and Efficient. *J. Anal. Bioanal. Tech.* **01** (2012).

26. R. J. Elshire, *et al.*, A Robust, Simple Genotyping-by-Sequencing (GBS) Approach for High
Diversity Species. *PLoS ONE* **6** (2011).

27. Y. Hartman, D. A. P. Hooftman, M. Eric Schranz, P. H. van Tienderen, QTL analysis reveals
the genetic architecture of domestication traits in Crisphead lettuce. *Genet. Resour. Crop
Evol.* **60**, 1487–1500 (2013).

28.   Y. Xiong, J. Sun, N. L. Glass, VIB1, a Link between Glucose Signaling and Carbon Catabolite
Repression, Is Essential for Plant Cell Wall Degradation by Neurospora crassa. *PLOS Genet*
10, e1004500 (2014).

29.   T. Benocci, M. V. Aguilar-Pontes, M. Zhou, B. Seiboth, R. P. de Vries, Regulators of plant
biomass degradation in ascomycetous fungi. *Biotechnol. Biofuels* **10**, 152 (2017).

30.   P. Cai, *et al.*, Evidence of a Critical Role for Cellodextrin Transporte 2 (CDT-2) in Both
Cellulose and Hemicellulose Degradation and Utilization in Neurospora crassa. *PLoS ONE* **9**,
e89330 (2014).

31.   J. M. Galazka, *et al.*, Cellodextrin Transport in Yeast for Improved Biofuel Production.
Science **330**, 84–86 (2010).

32.   S. Tani, T. Kawaguchi, T. Kobayashi, Complex regulation of hydrolytic enzyme genes for
cellulosic biomass degradation in filamentous fungi. *Appl. Microbiol. Biotechnol.* **98**, 4829–
4837 (2014).

Supporting Information

S1 File. N6_QTL_Data. This .csv file contains genotype and phenotype information

used for QTL analysis using the R-qtl package in R.

S2 Text. N6 QTL script for analysis in R. This file contains R script used for QTL

analysis in R. The numbers in row two is a chromosome number and row 3 a physical

base-pair position.

S3 File. GenesByLocation_summary (Chr1-Ferm).xlsx. This file contains all of the

genes within the QTL region on chromosome 1 for fermentation.

S4_GenesByLocation_summary (Chr2-Sacch).xlsx. This file contains all of the genes

within the QTL region on chromosome 2 for saccharification.

S5_GenesByLocation_summary (Chr4-Ferm).xlsx. This file contains all of the genes

within the QTL region on chromosome 4 for fermentation.

S6_GenesByLocation_summary (Chr4-Sacch).xlsx. This file contains all of the genes

within the QTL region on chromosome 4 for saccharification.

S7_GenesByLocation_summary (Chr6-Ferm).xlsx. This file contains all of the genes

within the QTL region on chromosome 6 for fermentation.

S8_GenesByLocation_summary (Chr6-Sacch).xlsx. This file contains all of the genes

within the QTL region on chromosome 6 for saccharification.

S9_GenesByLocation_summary (Chr7-Ferm).xlsx. This file contains all of the genes

within the QTL region on chromosome 7 for fermentation.

Genetic map

A

B Marker Interaction

C Marker Interaction