Evaluation of 20 enset (*Ensete ventricosum*) landraces for response to *Xanthomonas vasicola*

pv. *musacearum* infection

Sadik Muzemil, Alemayehu Chala, Bezuayehu Tesfaye, David J. Studholme, Murray Grant,

Zerihun Yemataw, Temesgen Magule Olango

Corresponding author: Sadik Muzemil (mik.mub2@gmail.com)

Abstract

Bacterial wilt, caused by *Xanthomonas vasicola* pv. *musacearum* (Xvm), formerly *X. campestris*

pv. *musacearum*, is the most threatening and economically important disease of enset (*Ensete*

ventricosum), the multipurpose food security crop orphan to south and southwestern Ethiopia.

Xvm has also had a major impact on banana and plantain production in East Africa following its

detection in Uganda in 2001 and subsequent spread. Effective control of this disease currently

relies on integrated disease management (IDM) strategies including minimization of field

pathogen inoculum and deployment of wilt resistant enset landraces. Identifying landraces with

stable and durable Xvm resistance will greatly accelerate breeding of varieties that can be included

as a component of IDM. In this study, 20 enset landraces previously reported to exhibit lower

susceptibility to Xvm were grown in pots under open field conditions and inoculated with an

aggressive Xvm inoculum isolated from a disease hotspot area. Longitudinal and survival analyses

were applied to each landrace, based on disease units representing a combination of area-under-

disease progress stairs, disease index and apparent infection rate. Considerable variation was

observed among the 20 landraces; however, none exhibited full immunity to Xvm infection. Three

landraces, viz. Hae'la, Mazia and Lemat (HML), showed lowest susceptibility to Xvm as

evidenced by lower disease units and higher survival rates. Landraces Kuro, Gezewet, Bededet,

and Alagena showed similar levels of Xvm infection as did HML, but with lower survival rates.

By contrast, landrace Arkia showed the highest infection level and lowest survival rate, suggesting

a high degree of susceptibility to Xvm. This study identifies new material that can be used in future

breeding programmes to develop Xvm-resistant enset varieties.

Keywords: Enset landrace, bacterial wilt disease, resistance, susceptibility

Author Addresses

- Sadik Muzemil: Southern Agricultural Research Institute, Areka Agricultural Research
Center, Areka, Ethiopia,

- Alemayehu Chala: Hawassa University, College of Agriculture, Hawassa Ethiopia

- Bezuayehu Tesfaye: Hawassa University, College of Agriculture, Hawassa Ethiopia

- David J. Studholme: Biosciences, University of Exeter, Exeter, United Kingdom

- Murray Grant: School of Life Sciences, Gibbet Hill, University of Warwick, Coventry,
CV4 7AL, UK

Zerihun Yemataw: Southern Agricultural Research Institute, Hawassa Agricultural
Research Center, Hawassa, Ethiopia,

- Temesgen Magule Olango: Hawassa University, College of Agriculture, Hawassa Ethiopia

1. Introduction

Enset (*Ensete ventricosum* (Welw.) Cheesman) is a diploid (2n=18), herbaceous, perennial monocarpic crop belonging to the family *Musaceae*. Enset is often referred to as false banana, due to its phenotypic resemblance to banana (*Musa* species). The crop is cultivated exclusively in Ethiopia, where it has a considerable economic and social importance for millions (Brandt *et al.*, 1997; Borrel et al., 2019).

Enset is tolerant to prolonged drought and provides a year-round source of staple nutritious food. It is thus widely cultivated across south and southwestern Ethiopia contributing to improved food security for more than 20 million Ethiopians (Yesuf and Hunduma, 2012, Yemataw et al., 2017). For farmers, enset is more than a year-round staple food, as it provides multiple additional daily benefits yet requires little crop management husbandry. The multipurpose benefits are derived from different enset landraces that are particularly suited for feed, fiber, packaging, construction material as well as providing a medicinal role (Brandt et al., 1997; Nurfeta et al., 2008; Yemataw et al., 2017). Due to its long history of cultivation across diverse ethnic groups, enset has significant cultural and socio-economic value in Ethiopia (Shigeta, 1990; Olango et al., 2014, Borrell et al., 2019).

Enset production is threatened by bacterial wilt disease caused by *Xanthomonas vasicola* pv. *musacearum* (Xvm) (Ashagari, 1981, 1985; Archido and Tessera, 1993; Tessera et al., 2008; Yesuf and Hunduma, 2012; Nakato et al., 2018). The bacterial disease was first reported in Kaffa district of Ethiopia in 1960's initially on enset (Yirgou and Bradbury, 1968) and subsequently on banana (Yirgou and Bradbury, 1974). However, the first observations for the bacterial disease on enset in Ethiopia dates back to 1930s (Castellani, 1939). The causative agent was previously known as

Xanthomonas musacearum and *X. campestris* pv. *musacearum*, but a recent taxonomic study been

transferred it to the species *Xanthomonas vasicola* (Studholme et al. 2019 in press). Currently, the

disease is found distributed in all enset growing areas of southern and southwestern Ethiopia,

where it has a devastating impact on enset production (SARI-McKnight CCRP, 2013; Blomme et

al., 2017; and Nakato et al., 2018). Other diseases caused by fungi (Tessera and Quimio, 1993),

viruses (Tessera et al*.,* 2003) and nematodes (Bogale et al., 2004) also affect enset. Mammals and

pests such as porcupines, mole rats, wild pigs and insects such as mealybugs (Azerefegne et al.,

2009) also impact enset production (Handoro et al*.,* 2012).

Enset landraces in Ethiopia might offer enhanced resistance to Xvm, thus potentially offering a

huge genetic resource to farmers though no systematic patho-testing for responses to Xvm has yet

been implemented. Currently, farmers claim that certain enset landraces show relatively low level

of infection to Xvm and they often incorporate these landraces in their backyard landrace mixture

as one option for disease management (Ashagari, 1985; SARI-McKnight CCRP, 2013; Yemataw

et al., 2016). Nevertheless, identification of resistant landraces has been challenging. Literature

has reported enset landraces with lower but varying susceptibility to Xvm infection (Ashagari,

1985; Archaido and Tessera, 1993; Handoro and Welde-Michael, 2007; Welde-Michael et al.,

2008; Haile et al., 2014; Hunduma, 2015; Wolde *et al.,* 2016; Handoro and Said, 2016). However,

inconsistency in enset landraces' responses to Xvm has been observed (Ashagari, 1985; Handoro

and Welde-Michael, 2007; Welde-Michael *et al.,* 2008; Tadesse *et al.,* 2008), with variation in

virulence among batches of Xvm inoculum being the most probable cause. Notably, existing Xvm

resistance phenotyping data primarily comprise assessments of individual landraces.

In banana transgenic approaches have been demonstrated for resistance against Xvm (Namukwaya

et al., 2012) due to absence of genetic resistance in the cultivated banana germplasm pool. While

initial results look promising, this approach would require biosafety regulations for the adoption

and use of transgenic plants, and adoption of the approach to enset. Hence, the availability of

sources of resistance to Xvm in enset germplasm could serve as a breeding stock against Xvm in

enset as well as banana plant.

To date, no systematic screening has been undertaken on the same enset landraces that are reported

to have high tolerance to Xvm infection. Such a study would greatly assist breeding/selection

efforts to identify elite landraces exhibiting enhanced Xvm resistance. In this paper we have

undertaken field experiments to re-evaluate selected enset landraces previously reported to have

reduced susceptibility to Xvm. We undertook a detailed study of the infection reaction and

identified specific enset landraces with potential to contribute towards sustainable management of

the disease, using a common garden experiment at a single site to minimize environmental

variability and enable a detailed study of the infection reaction and identify specific enset landraces

with potential to contribute towards sustainable management of the disease.

2. Materials and Methods

2.1. Description of the study site

Evaluation of enset landraces for resistance to *Xanthomonas vasicola* pv. *musacearum* (Xvm) was carried out using an open-field potted experiment at Southern Agricultural Research Institute (SARI), Hawassa, Ethiopia. The site is located at $7^04'$ N and $38^031'$ E with an elevation of 1700 meters above sea level and having an annual rain fall of 1100 mm. The annual average, minimum, and maximum temperatures at Hawassa are 20.6^0C, 13.5^0C and 27.6^0C, respectively.

2.2. Plant material acquisition and multiplication

This study used 18 enset landraces previously reported to exhibit low level infection or having contrasting infection phenotypes following Xvm inoculation. Landrace 'Arkia', was also included for comparison because it was previously reported to support a high level of Xvm growth. Also included was landrace 'Bota Arkia', which has a similar name to landrace 'Arkia' but of different origin. This gave a total 20 landraces for the present study (Table 1).

To initiate the experiment, underground corms of 2-3 year old mother plants from each of the 20 landraces were macro-propagated to produce true-to-type suckers for the experiment. Mother corms of landrace Mazia were kindly provided by Areka Agricultural Research Center (AARC). The remaining 19 enset landraces were collected from enset farmers in six enset growing zones of southern Ethiopia viz. Dawro, Gedio, Gurage, Kambata, Sidama and Wolaita. Collection of the landraces from these areas was undertaken by first confirming that the landrace described was true for its reaction to Xvm, followed by a rapid appraisal on site to avoid the possiblilty of mis-identification (Figure 1).

121   Table 1. Enset landraces studied for their reaction to *Xanthomonas vasicola* pv. *musacearum*

122   (Xvm)and previously recorded reaction to Xvm.

No	Enset landraces	Collection zones*	Previously reported response**	References for landrace reaction to Xvm
1	Abatemerza	Kambata	Resistance /Tolerant	Handoro and Said (2016)
2	Ado	Sidama	Moderately Tolerant	Ashagari (1985), Welde-Michael et al. (2008)
3	Alagena	Wolaita	Resistance /Tolerant	Handoro and Said (2016)
4	Arkia	Wolaita	Susceptible	Handoro and Said (2016)
5	Bededet	Gurage	Moderately Tolerant	Wolde et al. (2016)
6	Bota Arkia	Dawro	NA	
7	Dirbo	Kambata	Resistance /Tolerant	Handoro and Said (2016)
8	Gefetano	Wolaita	Resistance /Tolerant	Handoro and Said (2016)
9	Genticha	Sidama	Moderately Tolerant	Ashagari (1985), Welde-Michael et al. (2008)
10	Gezewet	Gurage	Susceptible	Welde-Michael et al. (2008)
		Gurage	Resistant	Wolde et al. (2016)
11	Ginbuwa	Gurage	Tolerant	Handoro and Said (2016)
12	Godere	Wolaita	Resistance /Tolerant	Handoro and Said (2016)
13	Hae'la	Kambata	Highly Tolerant	Tessera et al. (2000), Gizachew et al. (2008)
14	Kuro	Dawro	Moderately Tolerant	Handoro and Said (2016)
15	Lemat	Gurage	Moderately Resistance /Tolerant	Welde-Michael et al. (2008), Handoro and Said (2016)
	Lemat	Gurage	Susceptible	Mekuria et al. (2016)
16	Mazia	AARC (Orginally Collected from Dawro)	Resistance /Tolerant	Handoro and Welde-Michael, (2007), Welde-Michael et al. (2008), Tessera et al. (2008), Tariku et al. (2015), Handoro and Said (2016)
17	Nechuwe	Gurage	Moderatel Tolerant	Welde-Michael et al. (2008),
		Gurage	Susceptible	Wolde et al. (2016)
18	Unjeme	Kambata	Resistance /Tolerant	Handoro and Said (2016)
19	Wachiso	Kambata	Resistance /Tolerant	Handoro and Said (2016)
20	Yesha	Dawro	Resistance /Tolerant	Handoro and Said (2016)

123   * Administrative zone in Southern Ethiopia where the respective enset landrace was collected from farmers
124   field. Original collection history of the landraces obtained from the articles listed.
125   ** The recorded Xvm response indicated was derived from the accompanying cited article(s).
126

Figure 1. Geographical locations of enset landraces collection and experimental sites.

Following collection, enset landraces were multiplied at Areka Agricultural Research Center

(AARC) using an enset macro-propagation method previously described (Yeshitila et al. 2009).

Briefly, the mother corm was dissected into two equal halves and the apical tissue removed from

the center of each half corm to allow secondary meristems to develop into suckers for planting.

These dissected corms are dried in the shade for 3 hours prior to planting. Each half corm was

planted in slanted orientation in holes of 1 m x 1m and covered with soil.

After 12 months, multiplied enset suckers were uprooted and transplanted into a 5 kg capacity

plastic pot filled with sun dried mixtures of soil: sand: manure at a ratio of 3:1:1 (Quimio, 1992)

and placed in an open field experimental site at SARI. Suckers were allowed to establish for two

months to get sufficient leaves (>3) to allow inoculation with Xvm. During this establishment

period, suckers in plastic pots were watered daily prior to inoculation with Xvm and for a

subsequent two months after Xvm inoculation. Watering thereafter was reduced to two times per

weeks for remaining experimental period.

2.3. Preparation of bacterial suspension for inoculation

Virulent Xvm was collected from a disease hot spot in the Hagereselam area, Sidama Zone,

southern Ethiopia. Xvm bacterial ooze from young leaves and /or pseudostem of diseased enset

plants was harvested into sterile distilled water and preserved at 4^0C until use. Two sets of bacterial

suspensions were prepared for hypersensitivity and pathogenicity tests. One comprised the

uncultured bacterial suspension. The other was derived from a day-old preserved field harvested

Xvm isolates streaked on YPSA (5 g yeast extract, 10 g peptone, 20 g sucrose and 15 g agar) – a

growth and isolation medium used for selecting pure Xvm colonies (Haile *et al.*, 2014). These

plates were incubated at 28°C for 24 h (Schaad and Stall, 1998). Single colonies with a yellow,

convex, mucoid morphology typical of Xvm were harvested and preserved in YPSA slants at 4^0C.

2.4. Hypersensitivity and pathogenicity tests

Both uncultured and cultured suspension were tested for hypersensitivity and pathogenicity on

two-month-old tobacco (*Nicotiana tabacum*) or the highly susceptible control enset landrace

'Arkia'. In hypersensitivity test conducted on *N. tabacum* (Bobosha, 2003) using both inoculum

types independently, 2 ml of a bacterial suspension containing ~1.0×10^8 colony forming units

(cfu) per mL (approximately $OD_{600} = 0.5$) was used for inoculation. A positive hypersensitive

response was scored if tissue exhibited yellow clearing chlorosis limited to around the point

injection.

For initial assessment of pathogenicity tests, 14-month-old (i.e. two months after transplanting)

disease-free enset suckers of the susceptible landrace 'Arkia' (Handoro and Welde-Michael, 2007)

were infected with 4 mL of bacterial suspension ($\sim 10^8$ cfu/mL at $OD_{600} = 0.5$) from uncultured and

cultured inocula.

2.5. Inoculation of Xvm to test landraces

As the uncultured bacterial suspension resulted in shorter incubation period and more severe

disease on both *N. tabacum* and enset landrace Arkia during the pathogenicity tests, field-

harvested, uncultured Xvm suspensions preserved at 4 ^{0}C were used as inoculum for landrace

evaluation. Suckers of 14-months-old enset landraces (two months after transplanting in potted

soil mix) were inoculated with a 4 mL aliquot of the bacterial suspension, adjusted to $\sim 10^8$ cfu/mL

as described above, by infiltration with a hypodermic sterile syringe into the youngest innermost

leaf petiole (Figure 2). A new sterile hypodermic syringe was used for inoculating each sucker of

every landrace. Control plants were infiltrated with the same volume of sterile distilled water. The

pot experiment was organized in four replicates of 20 landraces, each landrace containing 10 plants

and landraces within each replication were randomized. Thus, the entire experiment comprised a

total of 800 plants. This included a total of 30 individuals of the 20 enset landraces inoculated with

uncultured Xvm suspension and a further 10 individuals of the 20 plants comprising negative

controls inoculated with sterile distilled water.

Figure 2. Evaluation of enset landraces grown under potted soil mix (A and B) and inoculation of

landraces (C and D) with uncultured *Xanthomonas vasicola* pv. *musacearum* (Xvm) suspension

2.6. Measurements and data analysis

Data recording and quantifying disease progression

Data collection was initiated a week after inoculation and continued weekly for the first three

weeks, then at two-week intervals for 155 days thereafter. It was notable that in the first two weeks

following infection, some enset landraces showed unusual infection responses including twisting

of the leaf blade in some landraces, rolling or curling of the leaf tip and leaf edge in other enset

landraces. From the third week after Xvm infection onward, we observed typical wilting symptoms

such as severe drooping of the top 25% of leaf blade, collapsing of the leaf blade, wilting of both

inoculated and adjacent uninoculated leaves, chlorosis across the majority of leaves and, in certain

landraces, complete death. At each evaluation time, the number of leaves per plant that showed

these consistent wilting symptoms and total number of asymptomatic leaves per plant were

recorded. Data from the "inconsistent" first two-week period were excluded from analysis as it ws

not possible to associate these initial responses with a phytopathological response.

Four response variables were used to quantify enset disease phenotypes following Xvm infection:

disease index (DI), area under disease progress stairs (AUDPS), apparent infection rate (AIR) and

survival. DI is the percentage of symptomatic leaves per individual at each evaluation period

averaged across the 4 replicates (Schandry, 2017). DI helps to quantify disease symptoms over the

evaluation period using a scale of 0-4, with 1 and 4 corresponding to 25% and 100% of total wilted

leaves per plant respectively.

$$\textbf{Disease Index (DI)} = \frac{w}{t} x\,4$$

Where, w is the number of symptomatic leaves and t is the number of total leaves of a single plant.

AUDPS (Simko and Piepho, 2012) was used to estimate disease accumulation and progress over

time. AUDPS is proposed to provide better estimates of the disease by giving weight closer to the

optimal than that derived from AUDPC (area under disease progress curve) assessments (Madden

et al., 2007). AIR, corresponding to the speed at which an epidemic develops (Meena *et al.*, 2011),

was calculated as the slope of disease index development. Finally, survival analysis was applied

to study the fraction of survivors per time point among enset landraces challenged with Xvm.

Survival data was generated according to Schandry (2017) with a customized DI cut-off point of

2.23. This cut-off DI was used for comparison purpose and is the maximum DI value of landrace

Mazia that is frequently cited (Table 1) for its resistance/tolerance to Xvm.

$$\textbf{Area under disease Progress Curve (AUDPC)} = \sum_{i=1}^{n-1} \frac{y_i + y_{i+1}}{2} x\,(t_{i+1} - t_i)$$

$$\textbf{Area Under Disease Progress Staris (AUDPS)} = \text{AUDPC} + \left[\frac{y_1 + y_n}{2} x \frac{D}{n-1}\right]$$

Where, y_i is disease index at the i^{th} observation, t_i is time in days at the i^{th} observation, n is the

number of total observations, and D is duration from the first to the last observation ($D=t_n-t_1$)., n

is the number of total observations, and D is duration from the first to the last observation ($D=t_n-$

t_1).

Data analysis and visualization

Various functions from the following R packages were used for data manipulation and formatting:

broom (Robinson et al, 2015), *dplyr* (Wickham and Francois, 2015), *magrittr* (Bache and

Wickham, 2014), *modelr* (Wickham, 2016a), *stargazer* (Hlavac, 2015), *stringr* (Wickham, 2010),

tidyr (Wickham et al., 2017) and *tidyverse* (Wickham, 2016b). For most of the disease parameters

in this study, data were fitted to linear mixed model using functions in *lme4* package (Bates et al.,

2015) to properly account for differential disease development in enset landraces over time. The

linear model (lm) function from the base *stats* package was employed. For model comparison in

the disease units and survival analysis, Akaike information criterion (AIC) and Schwarz's

Bayesian information criterion (BIC) were used to identify the best model among the alternatives.

Models with the lower AIC and BIC value were used for analysis. Functions from *emmeans*

(Russell, 2019) and *lmerTest* (Kuznetsova *et al.*, 2016) were used to extract means from data fitted

to linear and linear mixed models respectively, applying the process "data manipulation for result

visualization". Graphic visualizations were done using *ggplot2* (Wickham, 2009).

Functions from the following packages were used for analysis: *MESS* (Ekstrøm, 2016), *survival*

(Therneau and Lumley, 2011), *survcomp* (Schröder et al., 2011), *rms* (Harrell Jr, 2016), coxme

(Therneau, 2015b), *lme4* (Bates et al., 2015), *lmerTest* (Kuznetsova *et al.*, 2016), *multcomp* (Hoth

on *et al.*, 2008), and *rcompanion* (Mangiafico, 2017). Packages *rmarkdown* (Allaire et al., 2015)

and *knitr* (Xie, 2014, 2015, 2016) were used to generate R Markdown (.rmd) files. Furthermore,

scripts used in data manipulation, analysis and visualization are available in the Rmarkdown

(**supplementary R_script and Enset_Xvm_data**) which contains a full description of data mana

gement and analysis.

Generalized linear hypothesis testing, adjusted for multiple comparisons using Tukey's method

(Hothorn et al., 2008), was used to assess statistically significant differences at *P*-value cut-off of

0.05. Statistical significance among treatments was determined using the "compact letter displays

(cld)" method in the graphic visualization; cld function from *multcomp* (Hothorn et al., 2008)

package. Treatments within the same "letter group" are not significantly different whereas

treatments with different letters display a significant difference.

3. Results

3.1. Hypersensitivity and pathogenicity tests

A typical hypersensitive response (HR) was observed on tobacco leaves from both of uncultured

and medium based Xvm suspension. However, uncultured Xvm suspension caused a severe HR

on tobacco within three days post inoculation whereas symptoms were delayed for up to eight days

when cultured Xvm was used (data not shown). The effect of inoculum type on symptom

appearance was more pronounced during pathogenicity testing on the susceptible landrace Arkia.

It took 21 days after inoculation (DAI) with uncultured Xvm suspension to cause wilting and

complete collapse of enset leaves of the susceptible landrace Arkia whereas the same disease level

was not attained in the cultured suspension inoculation until 30 to 45 DAI.

3.2. Symptom description

A range of symptoms was observed during the course of infection and subsequent disease

development on Xvm-challenged enset landraces. Necrosis around the point of inoculation and

surrounding tissues was observed 3 DAI in most landraces (Figure 3). At early stages of infection,

up to 15 DAI, landraces showed a varying range of symptoms. Included among these symptoms

were twisting and slight leaf curling, and drooping of the blade and tip of the inoculated leaf. The

leaf blade around the Xvm inoculated area often became deformed, twisted or curving inwards.

These symptoms were replaced by severe curling of the leaf edge, drooping and folding back of

leaf blade from 15 DAI, symptoms that were consistently observed in all landraces. Gradually,

drooping from the leaf apex and folding back or collapsing of leaves became the most prominent

symptoms as the disease developed. All tested enset landraces showed one or more these

symptoms.

Figure 3. Xvm infection symptoms during the early stage of disease development on enset landraces. Labels are placed near to Xvm infected leaves. A) Curling of inoculated leaf, B) Twisting of inoculated leaf, C) Drooping of leaf apex of infected and other leaves and D) Folding around point of inoculation.

On severely infected enset landraces such as Dirbo and Arkia (susceptible control), the symptoms further developed into yellowing of leaves starting from leaf apex, and then gradual collapsing and clear wilting of the inoculated leaf and spreading the symptoms to other leaves. Eventually the whole leaves wilt, leading to their death and subsequent rotting of the whole plant (Figure 3A). However, on landraces that showed mild infection symptoms, for example Lemat and the resistant/tolerant control, Mazia, the inoculated leaf collapses and then dries (reminiscent of a classical hypersensitive response) or the symptoms extended to just a few adjacent leaves and the whole plant remains green afterward (Figure 4B). During the study period, a classical HR or HR-like symptoms were not observed in all of tested enset landraces.

Figure 4. Comparison of Xvm disease symptoms on enset landraces. Development of Xvm disease

sysmptoms on the susceptible enset landrace Dirbo (A) contrasted to the resistant/tolerant landrace

Mazia (B).*DAI= Days after inoculation.

3.4. Area under disease progress stairs (AUDPS)

Plot of the area under disease progress stairs showed more or less distinct curves and disease build-

up for the tested 20 enset landraces (Figure 5A). Analysis of AUDPS also showed a spectrum of

significant differences ($P<0.05$) among landraces with landrace Arkia having the highest and

Hae'la the lowest AUDPS value (Figure 5B). In addition, the analysis revealed that landraces could

be grouped to fewer clusters based on their AUDPS values; those at lower tail, middle to upper

tail, the last landrace Arkia that showed the peak AUDPS value.

3.5. Disease Index and apparent infection rate

Disease severity expressed as unit of DI illustrated the progress of disease development of Xvm

infection in the 20 enset landraces. Apparent Infection Rate (AIR) calculates the rate of disease

index development as a function of slope at each disease evaluation period (See Material and

Methods). Analysis of variance of both the DI and AIR showed marked differences (P< 0.05)

between landraces (Table 2).

Table 2. Analysis of variance for area under disease progress stairs, disease index and apparent

infection rate of the twenty enset landraces infected with enset bacterial wilt disease.

	Degrees of freedom	Sum squares	Mean squares	F value
Area Under Disease Progress Stairs (AUDPS)				
Landrace	19	3,176,321	167,175	19.29 ***
Disease Index (DI)				
Landrace	19	573	30	37.90 ***
Apparent Infection Rate (AIR)				
Landrace	19	1.04×10^{-3}	5.5×10^{-4}	7.18 ***
Residuals	578	0.044	7.63×10^{-5}	

Figure 5. Area under the disease progression stairs (AUDPS) for 20 the landraces following

controlled inoculation with Xvm. (A) Disease progression curve of tested enset landraces across

blocks. The area under curve for each replicate in landrace represented by three shades of gray in

each landrace. (B) Means (thick horizontal lines) and 95% confidence intervals (CIs) (shaded

boxes) of the AUDPS values as estimated by a linear mixed effects model. Calculated areas for all

individuals are plotted, with symbols indicating different replicates. The compact letter display

above the landrace names indicates the results of multiple comparison tests using of means

AUDPS using Tukey's method at $P<0.05$. Enset landraces with similar letters are statistically not

different from each other and vice versa.

Mean DI values ranged from 0.65, 0.80, 0.81 in Hae'la, Mazia and Lemat respectively,

corresponding to 16.25%, 20.00%, 20.25% severity. This contrasted with DI values of 1.98

(49.50%), 2.05 (51.25%), and 2.59 (64.75%) in Yesha, Godere and Arkia, respectively, (Table 3).

As evidenced from the lowest AIR, disease development rate is the slowest in landrace Lemat, and

fastest in Genticha, Abatemerza, Nechuw, Gefetano and Godere. Interestingly, AIR did not show

a direct linear relationship with disease index suggesting that AIR varies among enset landraces

irrespective of disease severity of landraces as measured by DI.

3.6 Survival Analysis

A DI of ≥ 2.23 was used as a time-to-event cutoff point for survival analysis. This cut-off DI is

the peak of infection level of the resistant/tolerant control enset landrace "Mazia". The survival

data generated with the survfit function of the survival package against landraces stratified by

block, best fitted a Gaussian distribution that showed the minimum AIC and BIC values (Table 4)

among Weibull, Logistic, Lognormal distributions. The linear fit of survival showed non-

proportionality of hazard ratio and, alternatively, data fitted to a mixed effects Cox model for

subsequent comparison of estimates (**supplementary R_script**).

Enset landraces displaying mild and severe infection to Xvm only showed significant estimates of

hazard ratio (HR) and Gaussian distribution (Table 5). Lemat, Hae'la and Mazia had estimated

hazard ratios of 0.230, 0.272 and 0.294, respectively, indicative of a mild infection to Xvm. This

result indicates that for landraces Lemat, Hae'la and Mazia there is, respectively, an 87.0%, 86.8%,

80.6% decrease in risk of getting severely infected with Xvm higher than disease index (DI) value

of ≥ 2.23 which was taken as a cutoff point. The susceptible control landrace Arkia included in

this study showed an estimated hazard ration of 1.866 that suggests 86.6% higher risk of showing

sever infection higher than DI value ≥ 2.23 after challenged with Xvm.

A survival fit to Gaussian distribution revealed that a higher proportion of plants survive from

landraces that showed mild infection than those landraces that showed severe infection (see peak

disease index) during the course of disease development (Figure 6). For example, right after 95

days post inoculation 50% of landrace Arkia was estimated to have experienced little probability

of surviving whereas for Lemat, Hae'la and Mazia at this time point it was less than 7%.

Figure 6. Kaplan-Meier estimates of survival and fits produced by survival regression analysis for

disease index of the 12 enset landraces that fitted to Gaussian distribution. Landraces that showed

non-significant values for hazard ratio and Gaussian distribution were omitted in this display. 6a)

Kaplan-Meier survival estimates for individual landrace with different linetypes indicating

experimental replicates. 6b) Combined display of Kaplan-Meier survival estimates

for the 12 enset landraces. The event of interest was defined as a disease index of 2.23.

4. Discussion

The bacterial wilt disease caused by *Xanthomonas vasicola* pv. *musacearum* (formerly *X. campestris* pv. *musacearum*) is a major bottleneck to enset (*Ensete ventricosum*) cultivation in south and southwestern Ethiopia, where diverse enset landraces and wild types of the crop are found. Farmers traditionally adopted host plant resistance as a management option for this bacterial wilt disease by incorporating certain landraces perceived to be more resistant/tolerant to Xvm (Ashagari, 1985; Yemataw et al., 2017). Several studies have indicated the presence of enset landraces that show low level of susceptibility to Xvm (Ashagari, 1985; Archaido and Tessera, 1993; Handoro and Welde-Michael, 2007; Welde-Michael et al., 2008; Haile et al., 2014; Hunduma, 2015; Wolde *et al.,* 2016; Handoro and Said, 2016).

To date there has been no systematic comparison of enset landraces' responses to Xvm infection in a common environment with a consistent inoculum. This detailed and, by nature of enset propagation, time consuming study undertook a detailed assessment of disease development on enset landraces previously reported for their low susceptibility to Xvm with the objective of gaining a definitive insight into Xvm disease phenotypes on enset landraces with the future objective of identifying suitable landraces to study the underlying molecular components of Xvm resistance. Characterisation of early infection symptoms through to the percentage of symptomatic leaves at each evaluation stage were considered. Various parameters such as area under disease progress stairs, disease index, apparent infection rate and survival rate were incorporated into detailed statistical analyses.

A limitation of this study was the unavailability of a standard stock of virulent inoculum. Given the constraints of available infrastructure in the geographic region of this study, the only practical

possibility was to use freshly harvested inoculum from already infected plants. This approach was

validated though preliminary experiments demonstrating that culturing the bacteria resulted in a

significant attenuation of virulence (based on HR in tobacco and symptoms on Arkia); therefore,

uncultured inoculum (i.e. bacterial suspension derived directly from bacterial ooze of infected

plans) was used. Most previous enset-Xvm interaction studies have used Xvm isolates from

disease hot-spot areas in southern Ethiopia in the District of Hagere Selam where highly virulent

Xvm isolates were identified by Handoro and Welde-Michael (2007), based upon the observation

that the tolerance/resistant landrace Mazia showed comparatively higher-level of infection to Xvm

from Hager Selam districts than isolates from four other districts. However, comprehensive studies

on classical and molecular epidemiology on diverse Xvm strains from enset growing areas in

Ethiopia is required, and is currently underway, to identify virulent Xvm strains for future breeding

efforts. Comparison of results between different studies has been hampered by the inevitable

variability between batches of freshly collected inoculum. However, within our present study, we

used a single uniform batch of inoculum, simultaneously inoculating all plants, thus eliminating

such confounding batch effects.

The majority of vascular pathogens, including Xvm, manifest wilting symptom in their host after

infection. Yet, even in well developed model systems, knowledge of the underlying infection

mechanisms remain poorly resolved. Despite such challenges, it is important to understand the

temporal spatial development of symptoms from initial visible phenotypes to the actual wilting

and subsequent death. Furthermore, it is important to understand how different closely related

genotypes respond to the same Xvm challenge. In this study we show that during early stages of

Xvm infection on enset landraces, i.e. up to 15 days after inoculation, a range of transient

symptoms were observed, including twisting and leaf curling, and drooping of the leaf blade and

apex. Symptoms associated with wilting, clear folding or collapsing of leaf blade, severe drooping

from leaf apex and wilting tended to appear from the third week following Xvm infection. Thus,

for robustness of scoring, analyses were restricted to disease data gathered from the third week

onwards.

This study revealed HR-like symptoms in a limited number of Xvm challenged enset landraces,

reminiscent of that seen in resistance-gene mediated HR. This raises the possibility that disease

classical resistance genes are deployed in restricting Xvm ingress. Involvement of HR cannot be

excluded even in those landraces that do not show HR-like symptoms; evidence suggests that local

resistant gene mediated reactions can also manifest as cellular changes contributing to wilting

(Jakobek and Lindgren, 1993; Chasan, 1994; Pajerowska-Mukhtar and Dong, 2009). All 20 tested

enset landraces showed at least partial symptoms of Xvm infection at different times of assessment

while mock infected plants were asymptomatic during the course of evaluation. Destructive

bacterial multiplication was not assessed at each disease evaluation stage, but rather the focus was

to distinguish symptoms associated with disease and potential symptoms associated with an

immune response. However, it is important to note that resistant plants also display a low level of

infection symptoms as reported in several crops. Indeed, bacterial enumeration of most classical

gene-for-gene interactions demonstrate a low level of accumulation of the pathogen until effective

resistance sets in and this is probably reflected in the weak phenotypes recorded in these landraces.

Previous reports which claim a lack of complete immunity to Xvm infection in enset due to the

presence of some level of infection in tested landraces (Archaido and Tessera, 1993; Handoro and

Welde-Michael, 2007; Welde-Michael et al., 2008; Haile et al., 2014; Hunduma, 2015; Wolde *et*

al., 2016; Handoro and Said, 2016) probably highlight the challenge in deploying effective

resistance to a vascular pathogen compared to a more classical foliar infection.

Analysis of area under disease progress stairs and disease index discriminated those enset

landraces with lower and higher level of infection, identifying elite landraces for further studies.

Accordingly, landrace Hae'la, Mazia and Lemat showed markedly lower susceptibility to Xvm

where as Yesha, Godere and Arkia were significantly more susceptible. Apparent infection rates

indicated that disease development was slowest in Lemat and fastest in Yesha compared to the

other landraces.

Complementing these results, survival analysis provided insight into the inherent infection risk.

Inoculations leading to a cutoff disease index value of ≥ 2.23 corresponded to 55.75% disease

incidence. The maximum disease value recorded was from the resistant control landrace Mazia

and this was used as a reference in the survival analysis. The significant difference in proportional

hazard model and Kaplan-Meier estimates of survival time demonstrated the variation in gross

impact of Xvm in different enset landraces. Analysis of the hazard ratio suggested 2.4 to 3.9 fold

changes in the resistance/susceptibility of landrace to disease caused by Xvm. Survival analysis

suggested a higher proportion of landraces from Hae'la would be estimated to survive after Xvm

infection followed by Lemat and Mazia whereas most plants in landrace Areki would succumb to

Xvm infection, followed by landraces Yesha and Godere.

This study confirmed the lower level of susceptibility of previously reported enset landraces to

Xvm infection after subjecting the landraces to a longer infection period, undertaking in parallel a

detail account of symptomologies and rigorous analysis of disease data. It is important to mention

that there is a great deal of evidence of sharing vernacular of enset landraces within and in

neighboring enset growing districts in Ethiopia (Olango et al, 2015; Yemataw et al., 2014; 2016)

that might not be phenotypical or genetically identical. For example, Olango et al. (2015) revealed

by Simple Sequence Repeat based genetic diversity analysis that two enset landraces with identical

vernaculars 'Gena' from Wolaita and Sidama zones in Southern Ethiopia were genetically distinct.

Futhermore, genome-wide comparison of SNP data from the landrace Mazia sourced from Dawro

and Wolaita zones of Southern Ethiopia revealed different SNP profiles (Yemataw *et al.*, 2018).

Thus, care needs to be taken when identifying enset landraces for study as variation in the reaction

of landraces with identical or similar vernaculars might not be only due to the variation in virulence

of Xvm isolates or environmental factor but also due to of the presence possible homonym enset

landraces.

Most of the landraces that showed higher tolerance/resistance to Xvm infection viz. Hae'la, Mazia,

Lemat, Kuro, Bededet, and Gezewet originated from areas where landrace enset landrace diversity

is richer with a high number of unique landraces present (Yemataw et al., 2014; 2016).   These

areas are the Dawro (landrace Mazia and Kurro), Kembata (landrace Hea'la), and Gurage (landrace

Lemat, Bededet, and Gezewet) districts of southern Ethiopia. The recent disease pressure in these

districts is comparatively higher than other enset growing areas in southern Ethiopia except for

Kembata Tembaro district (SARI-McKnight CCRP, 2013; and personal observation by Sadik

Muzemil). Furthermore, these areas also reside along the belts of the initial discovery of Xvm by

Castellani (1939) in southern Ethiopia. Hence, we hypothesize that the co-existence of Xvm with

these enset might have contributed to evolution of landraces with lower susceptibility to Xvm

infection.

Enset landraces with lower susceptibility to Xvm identified in this study i.e landraces Hae'la,

Lemat and Mazia, along with highly susceptible landraces provide the foundation for further

testing in multi-stage and environment conditions in order to better understand the underpinning

resistance/tolerance components. We note that it is important that not only the source of landraces,

but also the pathogen inoculum need to be considered for future studies.

Acknowledgement

The authors would like to thank McKnight Foundation Collaborative Research Program and the

European Community Horizon 2020 grant Project ID 727624, "Microbial uptakes for sustainable

management of major banana pests and diseases (MUSA) for financial and technical support. We

thank Areka and Hawassa Agricultural Research Centers, Southern Agricultural Research Institute

and Ethiopian Agricultural Research Institute for hosting and provision of germplasm as well as

necessary services and facilities during field study. We are indebted to School of plant and

horticultural science of Hawassa University for hosting the study, undertaken as part of SM's MSc

research work.

Compliance with Ethical Standards'

The authors would like to undertake that all authors listed have contributed sufficiently to the

project to be included as authors. To the best of our knowledge, no conflict of interest, financial

or other, exists. We have included acknowledgements, conflicts of interest, and funding sources

after the discussion.

6. References

Allaire, J.J., Xie, Y., McPherson, J., Luraschi, J., Ushey, K., Atkins, A., Wickham, H., Cheng, J.,

Chang, W. and Iannone, R. (2019). *rmarkdown*: Dynamic Documents for R. R package

version 1.13. doi: https://rmarkdown.rstudio.com

Azerefegne F.; Addis, T.; Alemu, T.; Lemawork, S.; Tadesse, E.; Gemu, M.; Blomme, G.

(2009). An IPM guide for enset root mealybug *(Cataenococcus ensete) in enset production*.

(V. Johnson, Ed.). Bioversity International. doi:

https://www.bioversityinternational.org/index.php?id=244&tx_news_pi1%5Bnews%5D=12

09&tx_news_pi1%5Bcontroller%5D=News&tx_news_pi1%5Baction%5D=detail&cHash=

87494e2eaf2d7f63a3c149c12bd330b1

Anita, S.H.C., Taboge E., and Woldemichael, G. 1996. Enset Need Assessment Project. Phase I

Report. Awassa, Ethiopia.

Archido, A.J. and Tezera, M. 1993. Diseases of Enset. An overview on enset research and future

technological needs for enhancing its production and utilization. *In*: Tsedeke, A., Hiebisch,

C., Brandt, S.A. and Seifu Gebremariam (eds.) Enset Based Sustainable Agriculture in

Ethiopia. Proceedings from the International Workshop on enset held in Addis Ababa, 13-20

December, pp. 3-12. doi: http://hdl.handle.net/123456789/1047

Ashagari. D. 1981. Studies on Bacterial Wilt of Enset. Paper Presented at the 13[th] National Crop

Improvement Conference. Addis Ababa, Ethiopia. 23-25 March 1981. IAR, Ethiopia.

Ashagari. D. 1985. Studies on the Bacterial Wilt of Ensat (*Ensete ventricosum*) and Prospects for

its Control. Eth J Agric Sci. 7(1): 1-14. doi: http://agris.fao.org/agris-

search/search.do?recordID=ET2010000090

Bache, S.M., & Wickham, H. (2014). *magrittr*: A Forward-Pipe Operator for R. R package version

1.5. doi: https://CRAN.R-project.org/package=magrittr

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models

using lme4. *Journal of Statistical Software*. doi: 10.18637/jss.v067.i01

Befekadu, H., Girma, A., & Fikre, H. (2014). Physiological characteristics and pathogenicity of

Xanthomonas campestris pv. *musacearum* strains collected from enset and banana in

Southwest Ethiopia. *African Journal of Biotechnology*, *13*(24), 2425–2434.

doi:10.5897/AJB2014.13794

Blomme, G., Dita, M., Jacobsen, K. S., Pérez Vicente, L., Molina, A., Ocimati, W., et al. (2017).

Bacterial Diseases of Bananas and Enset: Current State of Knowledge and Integrated

Approaches Toward Sustainable Management. *Frontiers in Plant Science*, *8*(1290), 1–23.

doi:10.3389/fpls.2017.01290

Bogale, M., Speijer, P. R., Mekete, T., Tessera, M., & Gold, C. (2004). Survey of plant parasitic

nematodes and banana weevil on *Ensete ventricosum* in Ethiopia. *Nematologia Mediterranea*,

32, 223–227. doi: https://journals.flvc.org/nemamedi/article/view/86797

Borrell, J. S., Biswas, M. K., Goodwin, M., Blomme, G., Schwarzacher, T., Heslop-Harrison, J.

S. (Pat), et al. (2019). Enset in Ethiopia: a poorly characterized but resilient starch staple.

Annals of Botany, 123(5), 747–766. doi:10.1093/aob/mcy214

Brandt, S., Spring, A., Hiebsch, C., McCabe, J. T., Tabogie, E., Wolde-Michael, G., et al. (1997).

The "Tree against hunger." Enset based agricultural systems in Ethiopia. American

Association for Advancement of Science with Awassa Agricultural Research Center, Kyoto

University center for African Areas Studies and University of Florida. Directorate for

International programs 1200 New York Avenue, NW, Washington, DC 20005. 56pp doi :

http://hdl.handle.net/123456789/1284

Chasanm R. (1994). Disease resistance: Beyond the R genes, 6, 461–463. doi:10.1105/tpc.6.4.461

Ekstrøm, C.T. (2019). *MESS*: Miscellaneous Esoteric Statistical Scripts. R package version 0.5.5.

doi : https://CRAN.R-project.org/package=MESS

Haile, B., Adugna, G., & Handoro, F. (2014). Physiological characteristics and pathogenicity of

Xanthomonas campestris pv. *musacearum* strains collected from enset and banana in

Southwest   Ethiopia.   *African   Journal   of   Biotechnology*,   *13*(24),   2425–2434.

doi:10.5897/AJB2014.13794

Handoro, F., & Michael, G. W. (2007). Evaluation of enset clone meziya against enset bacterial

wilt. *8th African Crop Science Society Conference,* El-Minia, Egypt, 27-31 October 2007, 8,

887–890.   doi :   https://www.cabdirect.org/cabdirect/abstract/20103268232.   Accessed   12

August 2018

Handoro, F., & Said, A. (2016). Enset clones responses to bacterial wilt disease (X*anthomonas*

campestris pv. *musacearum*). *International Journal of Applied and Pure Science and*

Agriculture (IJAPSA) , *02*(09), 45–53. doi: https://ijapsa.com/published-papers/volume-

2/issue-9/enset-clones-responses-to-bacterial-wilt-disease-xanthomonas-campestris-pv-

musacearum.pdf. Accessed 13 August 2018

Harrell Jr, F. E. (2019). *rms*: Regression Modeling Strategies. R package version 5.1-3.1. doi:

https://CRAN.R-project.org/package=rms

Hlavac, M. (2018). *stargazer*: Well-Formatted Regression and Summary Statistics Tables. R

package version 5.2.1. doi: https://CRAN.R-project.org/package=stargazer

Hothorn, T., Bretz, F., & Westfall, P. (2008). Simultaneous Inference in General Parametric

Models. *Biometrical Journal*, *50*(3), 346–363. doi:10.1002/bimj.200810425

Hunduma, T. (2015). Evaluation of Enset Clones Resistance against Enset Bacterial Wilt Disease,

6(3). doi:10.4172/2157-7579.1000232

Jakobek, J. L., & Lindgren, P. B. (1993). Generalized Induction of Defense Responses in Bean Is

Not Correlated with the Induction of the Hypersensitive Reaction. *The Plant cell*. 5:49-56.

doi: https://doi.org/10.1105/tpc.5.1.49

Kuznetsova, A., Brockhoff, P., & Christensen, R. (2017). *lmerTest* Package: Tests in Linear Mixed

Effects Models. *Journal of Statistical Software 82:(13), 1-26*. doi: 10.18637/jss.v082.i13

Lenth, R. (2019). *emmeans*: Estimated Marginal Means, aka Least-Squares Means. R package

version 1.3.4. doi: https://CRAN.R-project.org/package=emmeans

Mangiafico, S. (2019). *rcompanion*: Functions to Support Extension Education Program

Evaluation. R package version 2.2.1. doi: https://CRAN.R-project.org/package=rcompanion

Meena, P. D., Chattopadhyay, C., Meena, S. S., & Kumar, A. (2011). Area under disease progress

curve and apparent infection rate of alternaria blight disease of indian mustard (*Brassica*

juncea) at different plant age. *Archives of Phytopathology and Plant Protection*, *44*(7), 684–

693. doi:10.1080/03235400903345281

Nakato, V., Mahuku, G., & Coutinho, T. (2018). *Xanthomonas campestri*s pv. *musacearum*: a

major constraint to banana, plantain and enset production in central and east Africa over the

past decade. Molecular Plant Pathology, 19(3), 525–536. doi:10.1111/mpp.12578

Nurfeta, A., Eik, L. O., Adugna Tolera, & Sundstøl, F. (2008). Chemical composition and in sacco

dry matter degradability of different morphological fractions of 10 enset (*Ensete ventricosum*)

varieties. *Animal Feed Science and Technology*, *146*(1–2), 55–73.

doi:10.1016/j.anifeedsci.2007.12.003

Ocimati, W., Bouwmeester, H., Groot, J. C. J., Tittonell, P., Brown, D., & Blomme, G. (2019).

The risk posed by Xanthomonas wilt disease of banana: Mapping of disease hotspots, fronts

and vulnerable landscapes. PLoS ONE. doi:10.1371/journal.pone.0213691

Olango, T. M., Tesfaye, B., Catellani, M., & Pè, M. E. (2014). Indigenous knowledge, use and on-

farm management of enset (*Ensete ventricosum* (Welw.) Cheesman) diversity in Wolaita,

Southern Ethiopia. *Journal of ethnobiology and ethnomedicine*, *10*(1), 41. doi:10.1186/1746-

4269-10-41

Olango, T. M., Tesfaye, B., Pagnotta, M. A., Pè, M. E., & Catellani, M. (2015). Development of

SSR markers and genetic diversity analysis in enset (*Ensete ventricosum* (Welw.) Cheesman),

an orphan food security crop from Southern Ethiopia. *BMC Genetics*, *16*(1), 98.

doi:10.1186/s12863-015-0250-8

Oli, M., Biratu, K. S., Hunduma, T., & Selvaraj, T. (2014). Assessment of Disease Intensity and

Evaluation of Enset Clones Against Bacterial Wilt (*Xanthomonas campestris* pv.

musacearum) in Tikur Inchini and Jibat Districts of West Shewa , Ethiopia. *International*

Journal of Research in Science, *1*(2), 83–97. doi: http://www.i-

scholar.in/index.php/sijrs/article/view/67547/58173

Pajerowska-Mukhtar, K., & Dong, X. (2009). A kiss of death - Proteasome-mediated membrane

fusion and programmed cell death in plant defense against bacterial infection. *Genes and*

Development, *23*(21), 2449–2454. doi:10.1101/gad.1861609

Robinson, D and Hayes, A. (2019). *broom*: Convert Statistical Analysis Objects into Tidy Tibbles.

R package version 0.5.2. doi: https://CRAN.R-project.org/package=broom

Schandry, N. (2017). A Practical Guide to Visualization and Statistical Analysis of *R.*

solanacearum Infection Data Using R. *Frontiers in Plant Science*, *8*, 1–14.

doi:10.3389/fpls.2017.00623

Shimwela, M. M., Ploetz, R. C., Beed, F. D., Jones, J. B., Blackburn, J. K., Mkulila, S. I., & van

Bruggen, A. H. C. (2016). Banana xanthomonas wilt continues to spread in Tanzania despite

an intensive symptomatic plant removal campaign: an impending socio-economic and

ecological disaster. *Food Security*, (8), 939–951. doi:10.1007/s12571-016-0609-3

Schröder, M. S., Culhane, A. C., Quackenbush, J., & Haibe-Kains, B. (2011). *survcomp*: An

R/Bioconductor package for performance assessment and comparison of survival models.

Bioinformatics. 27(22): 3206-3208. doi:10.1093/bioinformatics/btr511

Shigeta, M. (1990). Folk in-situ conservation of ensete (*Ensete ventricosum* (welw.) E.E.

Cheesman): Towards the interpretation of indigenous agricultural science of the Ari,

southwestern Ethiopia. *African Study Monographs*, *10*(3), 93–107. doi:

https://repository.kulib.kyoto-u.ac.jp/dspace/bitstream/2433/68056/1/ASM_10_93.pdf

Simko, I., & Piepho, H. (2012). The Area Under the Disease Progress Stairs : Calculation ,

Advantage , and Application. *Phytopathology*, *102*(4), 381–389. doi:10.1094/PHYTO-07-11-

0216

Studholme, D. J., Wicker, E., Abrare, S. M., Aspin, A., Bogdanove, A., Broders, K., et al. (2019).

Transfer of *Xanthomonas campestris* pv. arecae, and *Xanthomonas campestris* pv.

musacearum to *Xanthomonas vasicola* (Vauterin) as *Xanthomonas vasicola* pv. *arecae* comb.

nov., and *Xanthomonas vasicola* pv. *musacearum* comb. nov. and description of

Xanthomonas vasicola pv. *vasculorum* pv. nov. bioRxiv, 571166. doi:

https://doi.org/10.1101/571166

Therneau, T. (2015). *survival*: A Package for Survival Analysis in S. version 2.38, doi:

https://CRAN.R-project.org/package=survival

Therneau, T.M. (2018). *coxme*: Mixed Effects Cox Models. R package version 2.2-10. doi:

https://CRAN.R-project.org/package=coxme

Welde-Michael, G., Bobosha, K., Blomme, G., Addis, T., Mengesha, T., & Mekonnen, S. (2008).

Evaluation of enset clones against enset bacterial wilt. *African Crop Science Journal*, *16*(1),

89–95. doi: http://dx.doi.org/10.4314/acsj.v16i1.54348

Wickham, H. (2017). *tidyverse*: Easily Install and Load "Tidyverse" Packages. R package version

1.2.1. doi: https://CRAN.R-project.org/package=tidyverse

Wickham, H. (2019). *modelr*: Modelling Functions that Work with the Pipe. R package version

0.1.4. doi: https://CRAN.R-project.org/package=modelr

Wickham, H. (2019). *stringr*: Simple, Consistent Wrappers for Common String Operations. R

package version 1.4.0. doi: https://CRAN.R-project.org/package=stringr

Wickham, H. and Henry, L. (2019). *tidyr*: Easily Tidy Data with 'spread()' & 'gather()' Functions.

R package version 0.8.3. doi: https://CRAN.R-project.org/package=tidyr

Wickham, H., Chang, W., Henry, L., Pedersen, T. L., Takahash, K., Wilke, C., & Woo, C. (2018).

ggplot2: Create Elegant Data Visualisations Using the Grammar of Graphics. R package

version 3.1.0. doi:10.1093/bioinformatics/btr406

Wickham, H., François, R., Henry, L., & Müller, K. (2019). *dplyr*: A Grammar of Data

Manipulation. R package version 0.8.1. doi: https://CRAN.R-project.org/package=dplyr

Yemataw, Z., Muzemil, S., Ambachew, D., Tripathi, L., Tesfaye, K., Chala, A., et al. (2018).

Genome sequence data from 17 accessions of *Ensete ventricosum*, a staple food crop for

millions in Ethiopia. *Data in Brief*, *18*, 285–293. doi:10.1016/J.DIB.2018.03.026

Yemataw, Z., Mekonen, A., Chala, A., Tesfaye, K., Mekonen, K., Studholme, D. J., & Sharma,

K. (2017). Farmers' knowledge and perception of enset Xanthomonas wilt in southern

Ethiopia. *Agriculture and Food Security*, *6*(1). doi:10.1186/s40066-017-0146-0

Yemataw, Z., Tesfaye, K., Zeberga, A., & Blomme, G. (2016). Exploiting indigenous knowledge

of subsistence farmers' for the management and conservation of Enset (*Ensete ventricosum*

(Welw.) Cheesman) (musaceae family) diversity on-farm. *Journal of Ethnobiology and*

Ethnomedicine, *12*(1), 34. doi:10.1186/s13002-016-0109-8

Yemataw, Z., Mohamed, H., Diro, M., Addis, T., & Blomme, G. (2014). Ethnic-based diversity

and distribution of enset (*Ensete ventricosum*) clones in southern Ethiopia. *Journal of Ecology*

and the Natural Environment, *6*(7), 244–251. https://cgspace.cgiar.org/handle/10568/66049.

Accessed 12 August 2018

Yirgou, D., & Bradbury, J. (1974). A note on wilt of banana caused by Enset wilt organism,

Xanthomonas musacearum. East African Agricultural and Forestry Journal. 40:1, 111-114.

doi: 10.1080/00128325.1974.11662720

Yirgou, D., & Bradbury, J. F. (1968). Bacterial wilt of Enset (*Ensete ventricosum*) incited by

Xanthomonas musacearum sp.n. *Phytopathology*, *58*(1), 111–112. doi:

http://www.musalit.org/seeMore.php?id=10016

Xie, Y. (2019). *knitr*: A General-Purpose Package for Dynamic Report Generation in R. R package

version 1.23.

655