Flotillin mediated membrane fluidity controls peptidoglycan synthesis and MreB

movement.

Aleksandra Zielińska[#,1], Abigail Savietto[#,2,8], Anabela de Sousa Borges[1], Denis Martinez[3],

Melanie Berbon[3], Joël R. Roelofsen[1], Alwin M. Hartman[4,5,6], Rinse de Boer[7], Ida J. van der

Klei[6], Anna K. H. Hirsch[4,5,6], Birgit Habenstein[3], Marc Bramkamp[*,2,8], Dirk-Jan Scheffers[*,1]

[#]equal contribution; [*]corresponding author

[1]Molecular Microbiology, Groningen Biomolecular Sciences and Biotechnology Institute, University

of Groningen, the Netherlands; [2]Biozentrum, Ludwig-Maximilians-Universität München, Germany; [3]

Institute of Chemistry & Biology of Membranes & Nanoobjects (UMR5248 CBMN), IECB, CNRS,

Université Bordeaux, Institut Polytechnique Bordeaux, Pessac, France; [4]Department of Drug Design

and Optimization (DDOP), Helmholtz-Institute for Pharmaceutical Research Saarland (HIPS) -

Helmholtz Centre for Infection Research (HZI), Saarbrücken, Germany; [5]Department of Pharmacy,

Saarland University, Saarbrücken, Germany; [6]Stratingh Institute for Chemistry, University of

Groningen, The Netherlands; [7]Molecular Cell Biology, Groningen Biomolecular Sciences and

Biotechnology Institute, University of Groningen, the Netherlands. [8]Institute for General Microbiology,

Christian-Albrechts-University Kiel, Germany.

[*]correspondence: marc.bramkamp@lmu.de; d.j.scheffers@rug.nl

Abstract

The bacterial plasma membrane is an important cellular compartment. In recent years

it has become obvious that protein complexes and lipids are not uniformly distributed

within membranes. Current hypotheses suggest that flotillin proteins are required for the

formation of complexes of membrane proteins including cell-wall synthetic proteins. We

show here that bacterial flotillins are important factors for membrane fluidity

homeostasis. Loss of flotillins leads to a decrease in membrane fluidity that in turn leads

to alterations in MreB dynamics and, as a consequence, in peptidoglycan synthesis. These

alterations are reverted when membrane fluidity is restored by a chemical fluidizer. _In_

***vitro*, the addition of a flotillin increases membrane fluidity of liposomes. Our data**

support a model in which flotillins are required for direct control of membrane fluidity

rather than for the formation of protein complexes via direct protein-protein interactions.

Introduction

The shape of a bacterium is predominantly defined by the structure of its peptidoglycan.

Although there is a great variety in bacterial shapes, the overall chemistry of peptidoglycan is

very similar between bacteria and thus the shape of peptidoglycan is primarily determined by

the temporal and spatial regulation of peptidoglycan synthesis. In rod-shaped bacteria,

peptidoglycan synthesis is thought to be mediated by two protein assemblies, the elongasome

and the divisome, that synthesize peptidoglycan along the long axis and across the division

plane of the cell, respectively[1,2]. These complexes contain a set of proteins required for the

final steps of synthesis and translocation of the peptidoglycan precursor, LipidII, from the inner

to the outer leaflet of the cytoplasmic membrane, and proteins that incorporate LipidII into

peptidoglycan. These include SEDS (Shape, Elongation, Division and Sporulation) proteins

that can perform glycosyl transferase reactions[3-5], and Penicillin Binding Proteins (PBPs) that

are divided in class A PBPs (aPBPs) that catalyse both glycosyl transferase and transpeptidase

reactions, class B PBPs (bPBPs) that only catalyse transpeptidase reactions and low molecular

weight PBPs that modify peptidoglycan, as well as hydrolases[2,6].

Coordination of these complexes is linked to cytoskeletal elements, MreB (-like proteins) for

the elongasome and FtsZ for the divisome. In models, the cytoplasmic membrane is often

depicted as a passive environment in which these machineries are embedded. However, it is

becoming clear that the structure of the membrane plays a critical role in the coordination of

peptidoglycan synthesis[7]. Inward membrane curvature serves as a localization trigger for MreB

and the elongasome, and enhanced local synthesis at bulges straightens out the membrane

sufficient to convert spherical cells to a rod shape[8,9]. In *Bacillus subtilis,* the motion of MreB

along the membrane is associated with elongasome activity[10,11] and the velocity of MreB

patches is related to growth rate[12], indicating that MreB motion can be used as a marker for

elongasome activity. Interestingly, MreB localizes to and organizes regions of increased

membrane fluidity (RIF)[13], which in turn is linked to the presence of LipidII, which favours a

more fluid membrane and promotes local membrane disorder[14,15]. Inhibition of LipidII

synthesis by genetic or chemical means results in a dissolution of membrane structures

observed with the dye FM 4-64 and release of MreB from the membrane[10,11,16,17].

Next to RIFs, membrane regions of decreased fluidity have been identified in bacteria[7,18,19].

These so-called functional membrane microdomains (FMMs) are thought to be organized by

the bacterial flotillin proteins, are enriched in isoprenoid lipids [20,21], and can be found in so-

called Detergent Resistant Membrane (DRM) fractions of the membrane. Since the formulation

of the FMM hypothesis, FMMs have been linked to many processes, such as protein secretion,

biofilm formation, competence and cell morphology[22-25]. Cell morphology effects are linked

to cell wall synthesis, and analysis of the protein content of *Bacillus subtilis* DRMs identified

several PBPs, MreC and other proteins involved in cell wall metabolism as well as the two

flotillins, FloA and FloT[21,24,26]. FloA is constitutively expressed, whereas FloT is expressed

primarily during stationary growth, cell wall stress and sporulation [27-29]. Super resolution

microscopy showed that the flotillins and other proteins found in DRMs do not colocalize and

have different dynamics[30], so it is unlikely that FMMs are regions in the membrane that offer

a favourable environment in which these membrane proteins are continuously present and

active. Recently, the hypothesis has been put forward that FMMs/flotillins form a platform for

the formation of functional protein oligomers, as work in *Staphylococcus aureus* showed that

multimerization of Type 7 secretion systems and PBP2a depends on FMMs[19,20,31].

Here, we have analysed the role of flotillins in peptidoglycan synthesis in *B. subtilis.* Our

results show that, at high growth rates, flotillins control membrane fluidity in a manner that is

critical for peptidoglycan synthesis and MreB dynamics, but have no effect on PBP

oligomerization. This results in a new model for flotillin function in the physical organization

of membranes during fast growth in rich medium.

Results

Absence of flotillins shifts peptidoglycan synthesis to division septa

In previous studies, a double deletion of *floA/floT* was either reported to suffer severe shape

defects and perturbed membrane structure[25], or to not have strong shape defects but with a

change in the overall lipid ordering of the membrane[24]. We grew wild type and Δ*floAT* strains

and analysed exponentially growing cells. We did not observe striking shape defects, but did

see an increase in median cell length and distribution of cell lengths in the absence of flotillins

(**Figure 1A, G**). To look at effects on peptidoglycan synthesis, we labelled cells with HADA,

a fluorescent D-Alanine analogue that reports on sites of active peptidoglycan synthesis[32], and

with fluorescent vancomycin (Van-FL), which labels LipidII and peptidoglycan containing

pentapeptide side chains[33,34]. This revealed a significant accumulation of peptidoglycan

synthesis stains at division septa in the Δ*floAT* strain (**Figure 1A-C**). To look at membrane

structure, cells were labelled with FM4-64, Nile-Red and DiI-C12, which are lipid dyes that

accumulate in zones enriched in fluid lipids[13]. Again, the stains accumulated at the septa in the

Δ*floAT* strain, which also showed some accumulation of FM4-64 and DiI-C12 in patches,

suggesting that the more fluid regions of the membrane are coalescing into larger regions

(**Figure 1A, D-F**). The HADA, FM4-64 and Nile-Red measurements were repeated using a

wild type strain expressing endogenous GFP, allowing simultaneous imaging of both strains

on the same slide, and gave similar results, confirming that the observed signal increase is not

due to variation between microcopy experiments (**Figure 1 - figure supplement 1 A,B**). In

this mixed-strain experiment, Nile-Red labelling at the lateral membrane was the same between

wild type and Δ*floAT* strains, indicating that there is no difference in dye diffusion between the

strains (**Figure 1 - figure supplement 1D**). Inspection of the septa by electron microscopy

revealed that there was no difference between the thicknesses of the septa between the wild

type and Δ*floAT* strain, ruling out that the increase in signal was due to formation of thicker

septa (**Figure 1 - figure supplement 1C**). The shift of peptidoglycan synthesis to the division

site could hint at stress in the overall peptidoglycan synthesis route. This was confirmed by

growing cells at a sublethal concentration of fosfomycin, which limits synthesis of LipidII[35],

but that does not impact growth rate at the concentration used. This resulted in bulging cells

and some lysis, which was exacerbated in the Δ*floAT* strain (**Figure 1 - figure supplement

1F,G**). It should be noted that the peptidoglycan synthesis stress caused by fosfomycin is not

the same as the stress caused by the absence of flotillins, as the phenotypes of wild type cells

with sublethal fosfomycin are quite distinct from Δ*floAT* cells without fosfomycin.

We ruled out that the peptidoglycan synthesis stress was caused by a change in the folding or

complex formation by PBPs in the absence of flotillins, as there were no differences in the

overall PBP-profiles of Bocillin-FL labelled wild type or flotillin deletion strains (**Figure 1 -

figure supplement 2A**). PBP complex formation was analysed by analysis of Bocillin-labelled

membrane fractions using a combination of Native-PAGE and SDS-PAGE[36] and showed that

various PBPs can be found in a high-MW complex (notably PBPs 1, 2, 3 and 4), but that

complex formation is similar in the Δ*floAT* strain (**Figure 1 - figure supplement 2B**). Also,

none of the five functional GFP-PBPs examined changed their localization in the Δ*floAT* strain

(**Figure 1 - figure supplement 2C**). Overall, the data suggest that in the absence of flotillins,

peptidoglycan synthesis is affected and relatively increased at division septa, with a

concomitant accumulation of membrane dyes that are indicative of higher membrane fluidity.

Figure 1. Accumulation of peptidoglycan synthesis and membrane material at division

sites in a flotillin mutant. A. Morphology of the exponentially growing wild type (WT) and

Δ*floAT* strains labelled with HADA, fluorescent Vancomycin (Van-FL), FM 4-64, Nile Red,

and DiI-C12. Scale bar 5 μm. **B-F**. Peak intensity of HADA (B), Van-FL (C), Nile Red (D),

FM4-64 (E) and DiI-C12 (F) labelled division sites of the cells shown in (A). Cells from each

strain (n ≥ 100, except E, n = 60) were analysed using the ObjectJ macro tool PeakFinder. **G.**

Distribution of the cell length (n = 100) of the strains analyzed in (A).

All data analysed with Prism and depicted as box plots with whiskers indicating 5-95%

percentile. Significant differences are based on the two-tailed Mann-Whitney test (* p<0.05;

** p<0.01).

The absence of both flotillins and PBP1 causes a severe phenotype, linked to a loss of

membrane fluidity.

We reasoned that a non-lethal defect in septal peptidoglycan synthesis could reveal more about

the role of flotillins and constructed a flotillin mutant that lacks PBP1, a bifunctional glycosyl

transferase/transpeptidase that is required for efficient cell division[37]. Simultaneous deletion of

pbp1, *floA*, and *floT* resulted in strong filamentation and delocalization of peptidoglycan

synthesis as well as membrane dyes to patches **(Figure 2A, Figure 2 - figure supplement 1A)**.

Deletion of single flotillin genes and PBP1 had similar, albeit less severe effects **(Figure 2 -**

figure supplement 1B, C). To exclude the possibility that an alteration of peptidoglycan

modification resulted in the delocalization of HADA and Van-FL, we used D-Alanine-D-

Propargylglycine (D-Ala-D-Pra), a clickable dipeptide analogue which is exclusively

incorporated into peptidoglycan via LipidII[38]. D-Ala-D-Pra incorporation was delocalized in

the *Δpbp1ΔfloAT* strain, indicating that peptidoglycan synthesis itself is delocalized **(Figure 2**

- figure supplement 1D). So far, our experiments were done with fast growing cells and

Lysogeny Broth (LB) as the growth medium. Strikingly, none of the mutant strains had an

apparent phenotype when cultivated in Spizizen's minimal medium (SMM, **Figure 2B**), and

peptidoglycan synthesis and lipid dyes were no longer accumulating at division sites in the

ΔfloAT strain **(Figure 2 - figure supplement 2)**. SMM has a higher Mg^{2+} concentration, which

is known to rescue various cell shape mutations by inhibition of cell wall hydrolysis[39].

However, the increase in Mg^{2+} was not sufficient to explain the reversal of phenotype as cells

grown on LB supplemented with Mg^{2+} (6 mM, concentration in SMM, or 20 mM) still

displayed the elongated phenotype with delocalized peptidoglycan synthesis **(Figure 2 - figure**

supplement 3). This indicated that the phenotypes associated with the absence of flotillins are

growth-rate and/or nutrient related.

Figure 2. Cell morphology and cell wall synthesis localization is dependent on growth
conditions. Morphology of the *WT*, *ΔfloAT*, *Δpbp1*, and *Δpbp1ΔfloAT* strains grown in **(A)**
rich (LB), **(B)** minimal (SMM) medium, and in **(C)** rich medium with membrane fluidizing
conditions (0.1 % benzyl alcohol, LB BnOH). Cells were labelled with HADA, and aberrant
cell shape and peptidoglycan synthesis are indicated with arrowheads. Panels on the right
indicate corresponding cell length distributions (n ≥ 100). Distributions were analysed using
Dunn's multiple comparison tests after Kruskal–Wallis. Statistically significant cell length

distribution classes (*p < 0.001*) are represented as letters above each graph – in B and C there

were no significant differences. Whiskers represent 5-95% percentile, "+" represent the means

Scale bar: 4 μm.

Next, we determined lipid packing order in the different strains using the fluorescent dye

Laurdan, a reporter for flotillin-mediated lipid ordering[24]. LB-grown cells lacking flotillins

displayed an increased generalized polarization (GP)[24], indicative of an overall increase in

ordered lipid packing in the membrane, but the effect of flotillins on membrane ordering

completely disappeared when cells were grown on SMM **(Figure 3)**. The resolution obtained

with Laurdan does not allow the detection of local differences in fluidity between the lateral

membrane and the septa, but does report on overall lipid ordering. Overall, lipid order was

increased in cells grown on SMM compared to LB **(Figure 3)**, whereas the absence of PBP1

had no significant effect on membrane fluidity, also not when combined with flotillin deletions

(Figure 3). The changes in lipid ordering were not due to changes in the overall fatty acid

composition of the membranes - the ratios of C17/C15 side chains and *iso/anteiso* fatty acids,

which are indicative of fluidity[13], were identical for wild type and Δ*floAT* strains grown on LB,

and very similar for cells grown on SMM (**Figure 3 - figure supplement 1**).

Figure 3. Flotillins increase overall membrane fluidity at high growth rate. Changes in overall membrane fluidity were assessed by Laurdan microscopy in cells grown on LB (**A**), SMM (**B**) and LB+BnOH (**C**). Micrographs show colour-coded generalized polarization (GP) maps in which red indicates regions of decreased fluidity (scale bar: 4 μm). Correspondent theoretical GP measurements in the graphs vary from -1 (more fluid) to 1 (less fluid). Significant statistical differences according to Dunn's multiple comparison tests after Kruskal–Wallis are represented as letters above each graph in panel **A**. Datasets labeled as A are significantly different from datasets labeled as B; datasets with the same letter are not significantly different. No statistically significant difference was observed for the data in panels **B** and **C** (*p < 0.001;* n ≥ 150, two biological replicates).

Restoring membrane fluidity rescues normal peptidoglycan synthesis.

The GP values indicated that membranes are more ordered when cells are grown on minimal medium, and this suggests that the flotillin-associated increase in overall membrane fluidity is important for cell shape control at high growth rates. This was tested by growing the strains lacking flotillins and PBP1 on LB in the presence of benzyl alcohol, an extensively used membrane fluidizer that increases membrane hydration due to disordering of membrane

structure[40]. Notably, the addition of benzyl alcohol increased membrane fluidity to similar

extents in the wild type and the mutant strains (see Figure 3C), but did not affect the growth

rates of the strains (**Figure 2 – figure supplement 4**). The increase in membrane fluidity

restored normal cell length and normal peptidoglycan synthesis patterns to the *pbp1/floA/floT*

strain **(Figure 2C)**.

In *B. subtilis*, the rate of growth and of peptidoglycan synthesis is linked to the speed of MreB

movement – in minimal media, the speed of MreB patches is reduced compared to the speed

in rich media[12]. Analysis of the movement of a fully functional mRFPruby-MreB fusion[10] by

time lapse TIRF (Total Internal Reflection Fluorescence) microscopy, confirmed that MreB

patch mobility is higher in cells grown on LB than in cells grown on SMM, with MreB speeds

similar to those reported previously[12] **(Figure 4, supplemental movie 1, 2)**. Strikingly, in the

absence of flotillins, MreB patch mobility was notably decreased in cells grown on LB, while

in SMM grown cells MreB patch mobility was independent of the presence of flotillins (**Figure**

4, supplemental movie 3,4). Fluidizing the membrane with benzyl alcohol, which does not

alter the growth rate, almost completely restored MreB mobility in LB grown cells **(Figure 4,**

supplemental movie 5, 6). These results indicate that the MreB patch mobility is not only

controlled by growth rate but also by membrane fluidity. Thus, in fast growing cells with

decreased membrane fluidity there is a decrease in elongasome mediated peptidoglycan

synthesis, reflected by the reduction of MreB mobility. This fits with an observed increase in

peptidoglycan synthesis at the division site, which may act as a compensatory mechanism.

Figure 4. MreB speed is linked to membrane fluidity. (A) The MreB speed in different strain backgrounds and growth conditions was analysed by time-lapse TIRF microscopy. Scatter plot of the speed of patches obtained from individual tracks in 5 different cells are represented per fusion and condition. Average speeds are shown; error bars indicate the standard deviation. Significant statistical differences according to Dunn's multiple comparison tests after Kruskal–Wallis are represented ($p < 0.001$). **(B)** Representative kymographs showing fast and slow patches of mRFPruby-MreB in *B. subtilis* cells lacking endogenous *mreB* (WT) or *mreB* and *floAT* (Δ*floAT*). See supplementary movies S1-S6 for corresponding raw image series.

Flotillin increases fluidity of model membranes in vitro.

To assess whether the influence of flotillins on membrane fluidity is direct, we determined the membrane fluidity of model membranes with purified flotillin using solid-state NMR (ssNMR). [2]H ssNMR is a biophysical tool that assesses lipid mobility in native-like model membranes on the atomic level, by monitoring the carbon-deuterium order parameter of a deuterated lipid along the acyl chain (here POPC-d31)[41,42]. We purified *B. subtilis* FloT and tested the impact of FloT on the membrane, when reconstituted in POPC-d31 liposomes (schematically depicted in Figure 5B). FloT decreases the spectral width of the [2]H quadrupolar splitting, reflecting an increase in motion on the atomic scale (**Figure 5A**). The [2]H spectrum encodes the local order parameter S_{CD} of the carbon-deuterium in absence and in presence of FloT. Strikingly, FloT has an important impact on the order parameter along the entire acyl chain. It is remarkable that the protein significantly decreases the order parameter S_{CD}, reaching

even the inner carbon atoms of the acyl chain, indicating a different packing behavior and

increased membrane fluidity upon interaction with FloT (**Figure 5B**). The strong fluidizing

effect described for FloT is notably different from the effects other proteins have when

reconstituted into liposomes, such as plant remorins[42] or the membrane binding peptide of the

nonreceptor tyrosine kinase Src [43]. The anisotropic lineshape of the ^{31}P spectra indicates that

the membrane is in the lamellar phase as expected for POPC at the chosen temperature

(298K)[44]. Upon interactions with FloT the lamellar phase remains intact with formation of a

few smaller objects, indicating that the overall liposome structure is not affected and that its

phase is maintained (**Figure 5 figure supplement 1**).

Figure 5. Lipid ordering of FloT probed by ^{2}H solid-state NMR. **(A)** Wide-line ^{2}H spectra of

POPC-d31 liposomes with or without FloT at a lipid-to-protein molar ratio of 25:1 acquired at

298 K. **(B)** Effect of FloT on the C-^{2}H order parameters of the PC acyl chain. De-Pake-ing and

simulations were applied on the ^{2}H solid-state NMR spectra to determine accurately individual

quadrupolar splittings. Order parameters of POPC-d31 acyl chain were derived from

experimental quadrupolar splittings and plotted as a function of the labeled carbon position.

Insert: schematic depiction of a liposome with added FloT which attaches to the membrane via

a hairpin loop[45].

Discussion

Our data provide evidence that flotillins play a direct role in controlling membrane fluidity and

that membrane fluidity is critical for peptidoglycan synthesis at certain growth conditions. *In*

vitro, flotillins enhance the fluidity of a model membrane, and *in vivo*, the membranes of fast

growing flotillin-mutant cells are less fluid even though the fatty acid composition in these

cells is identical. Therefore, we propose that the effect of flotillins on membrane fluidity is

direct, through a change in the packing behaviour of the lipids resulting in an efficient

separation of states of liquid ordered and disordered lipid domains in the membrane bilayer[24].

We found that membrane fluidity is not solely a function of temperature, but also of growth

conditions. *In vivo*, flotillins may also recruit specific, more rigid lipids, such as hopanoids and

carotenoids[18,20,21] which have been found in association with FMMs, and whose synthesis

could be growth condition dependent. The predominantly physical role in membrane

organization for flotillins fits with our observation that adding a chemical fluidizer is sufficient

to restore MreB dynamics and cell shape to fast growing cells that lack flotillins. We propose

that in fast-growing cells on rich medium, flotillin-mediated control of membrane fluidity is

critical and sufficient to allow essential membrane bound processes, such as peptidoglycan

synthesis, to proceed normally.

A sufficiently fluid membrane is necessary for the efficient recruitment and movement of

MreB, and provides a more favorable environment for the peptidoglycan precursor LipidII[8,9,17].

It has recently been shown that modulation of either MreBCD or PBP1 levels is sufficient to

alter the shape of *B. subtilis* cells[46], underscoring the importance of both systems. In the

absence of flotillins, the activity of the MreBCD component is strongly reduced – as evidenced

by the reduction of MreB speed – and the overall rigidity of the membrane is increased. This

results in a less favourable environment for the peptidoglycan precursor LipidII, which prefers

more liquid, disordered membrane phases[14,15,47]. Our data indicate that the reduction in

elongasome activity, which does not impact the growth rate itself, is compensated by increased

peptidoglycan synthesis activity around division sites in flotillin mutants, which is sufficient

to keep the overall cell shape intact, although cells are elongated. The accumulation of lipid

dyes indicative of increased fluidity at division sites is in line with a recent study that showed

phases of different fluidity in *Streptococcus pneumoniae* membranes, with more fluid

membranes and LipidII localizing at midcell[47] where the membrane is most bent. Our findings

are also in agreement with the recent observation that *B. subtilis* cells elongate and lose

organization of MreB when membrane fluidity is decreased by altering the membrane fatty

acid composition[48] (H. Strahl, personal communication). It could very well be that the shift of

fluidity towards the septum is only relative as the overall fluidity of the membrane is decreased

in the absence of flotillins. This is yet to be determined, as the resolution of Laurdan imaging

does not allow conclusive statements about local fluidity changes at the septum. It may seem

paradoxical that cells elongate when elongasome activity is reduced, but one has to remember

that a large amount of the peptidoglycan synthesis contributing to elongation of bacterial cells

is actually taking place at midcell, before the ingrowth of the septum[49-51]. The increased

peptidoglycan synthesis at future division sites makes the activity of PBP1 critical and explains

why its deletion has such a dramatic effect in cells lacking flotillins. Restoring fluidity using a

chemical fluidizer allows the MreBCD component to again efficiently drive peptidoglycan

synthesis during elongation, which is sufficient to suppress the flotillin mutant phenotype. The

net effect of this is that cells lacking PBP1 and flotillins grown with benzyl alcohol behave as

cells that only lack PBP1, which is quite similar to wild type. At low growth rates, there is no

difference between wild type cells and cells lacking flotillins with respect to membrane fluidity,

and the speed of MreB is similar between the two cell types. Thus, the deletion of flotillins

does not exacerbate the phenotype of cells lacking PBP1. The reason for the change in

membrane fluidity between cells grown on rich or minimal medium is not yet clear – it does

not seem to be caused by a large shift in the fatty acid composition of the membranes. Various

factors could play a role, such as the synthesis of specific lipids (hopanoids, isoprenoids) on

either type of medium, but also protein crowding, which is higher in membranes of fast-

growing cells than in slow-growing cells[52]. It will be an important future challenge to establish

the cause for this difference. An overall rigidification of the membrane may also lead to

retardation of processes which require membrane modifications such as division and

sporulation, which is indeed observed in *B. subtilis* flotillin mutants[25,53].

One of the proposed roles for flotillin proteins is that they form a 'platform' that transiently

interacts with membrane proteins that need to oligomerize into functional complexes[19]. We

tested this hypothesis for *B. subtilis* PBPs by comparing their localization and oligomerization

in wild type and flotillin mutant strains. Although we were capable of detecting a high MW

complex containing various PBPs (notably PBP1, 2a, 2b, 3 and 4), the complex was not

dependent on the presence of flotillins. We also note that PBP1 was present in the complex, as

well as present in a large smear in the first dimension native gel, which would explain why

PBP1 was detected by mass spectrometry analysis of a native PAGE band containing FloA[27].

PBP5, on the other hand, was not part of the high MW complex, which fits with its role in

processing of the terminal D-Ala from stem-peptides that have not been cross-linked, which it

exercises over the entire surface of the cell[32]. Although we cannot exclude that flotillins may

affect PBPs that are not easily detected by Bocillin-FL, our results do not provide any evidence

for a role for flotillins in the oligomerization of PBPs in *B. subtilis*. This extends the finding of

the Graumann lab that found either transient or no colocalization between flotillins and other

proteins present in DRM fractions[30]. Although it is obvious that peptidoglycan synthesis is

altered in the absence of flotillins, our data strongly suggest that the basis for this alteration is

in the physical organisation of the membrane rather than inefficient formation of divisome or

elongasome complexes in the absence of flotillins, because flotillin mutants strains show no

synthetic phenotype on minimal medium, and the defects on rich medium can be reverted by

chemically fluidizing the membrane.

In conclusion, our data provide a new model for flotillin function in the physical organization

of membranes during fast growth. The observation that flotillins differentially affect the

membrane in different growth conditions also explains the diversity of phenotypes described

for flotillin mutants in the literature.

Methods

B. subtilis strains and growth conditions

All *B. subtilis* strains used in this study are derived from strain 168 and are listed in **Table 1**.

Construction of new strains was based on natural competence of *B. subtilis*[54]. Gene integration

or deletion was validated by colony PCR whereas the expression and localization of the

fluorescent fusions was additionally validated by microscopy. Cells were grown either in LB

Lennox (5 g/L yeast extract; 5 g/L NaCl; 10 g/L tryptone)[55] or Spizizen minimal medium

(SMM)[56] supplemented with 1% glucose, at 37°C and 200 rpm, unless indicated otherwise.

Induction of the P_{xyl} promoter was triggered by addition of 0.2 - 0.5 % xylose. Cell cultures

were supplemented with spectinomycin (50 µg/ml), tetracycline (10 µg/ml), chloramphenicol

(5 µg/ml), kanamycin (5 µg/ml), erythromycin (1 µg/ml), benzyl alcohol (BnOH, 0.1 %) or

magnesium sulphate (MgSO$_4$, 6-20 mM) when necessary.

Growth curves

Growth experiments were performed either manually or automatically with a PowerWave 340 microplate reader (BioTek® Instruments, U.S.A). Strains were pre-cultured overnight in 3 ml LB or SMM medium at 37 °C with shaking at 200 rpm. Next, stationary or late-exponentially cells were diluted with fresh LB or SMM medium (supplemented when necessary), and cell densities (OD_{600}) were measured every 1 hour when monitored manually or every 10 min when monitored automatically, for a total time of 7-22 hours.

Fluorescence microscopy

For standard fluorescence microscopy, exponentially growing cells were immobilized on microscope slides covered with a thin film of 1 % agarose (w/v) in water or the appropriate medium. For TIRFM, agarose pads were mounted using Gene Frames (1.7 x 2.8 cm chamber, 0.25 mm thickness, 125 μL volume) from ThermoScientific. Standard fluorescence microscopy was carried out using an Axio Zeiss Imager M1 fluorescence microscope (EC Plan-Neofluar 100x/1.30 Oil Ph3 objective) equipped with an AxioCam HRm camera and an Nikon-Ti-E microscope (Nikon Instruments, Tokyo, Japan) equipped with Hamamatsu Orca Flash 4.0 camera.

For Laurdan and TIRFM experiments, a Delta Vision Elite microscope (Applied Precision, GE Healthcare) equipped with an Insight SSI™ Illumination, an X4 Laser module, a CoolSnap HQ^2 CCD camera and a temperature-controlled chamber set up at 37 °C was used. Laurdan images were taken with an Olympus UplanSApo 100x/1.4 oil objective. TIRFM image series were taken using an Olympus UAPO N 100X/1.49 TIRF objective and a 561 nm laser (50 mW, 100 % power). Data processing was performed with softWoRx Suite 2.0 Software.

Visualization of cell wall synthesis

Peptidoglycan (PG) synthesis was assessed by labelling the cells with HADA (7-

hydroxycoumarin 3-carboxylic acid-amino-D-alanine)[57], Van-FL[33] or D-Ala-D-Pra[38].

HADA: synthesized as described[34]. Overnight cultures of *B. subtilis* strains were diluted 1:100

into fresh LB medium or LB medium supplemented with 0.1 % (w/v) of benzyl alcohol

(BnOH), a membrane fluidizer. Cells were grown until exponential phase, a sample of 1 ml of

culture was spun down for 30 sec, 5000 × g and the cell pellet was resuspended in 25 µl of

fresh pre-warmed LB or LB containing 0.1% (w/v) BnOH. HADA was added to a 50 µM final

concentration. Cells were incubated for 10 min in the dark (37 ^{o}C, 200 rpm) and then washed

twice in 1 ml PBS buffer (58 mM Na_2HPO_4, 17 mM NaH_2PO_4, 68 mM NaCl, pH 7.3) to remove

the excess of unbounded HADA. Cells were spun down again and resuspended in 25 µl of the

appropriate medium and 2 µl of cells were mounted on 1% agarose slides before visualization.

Visualization of HADA patterns (excitation: 358 nm/emission: 461 nm) under fluorescence

microscopy from two biological replicates and cell length measurements were taken from at

least 100 cells each strain/treatment.

Van-FL: a 1:1 mixture of vancomycin (Sigma Aldrich) and BODIPY®FL Vancomycin

(Molecular Probes) at a final concentration 1 µg/ml was used to label cells for 5-10 min at room

temperature.

D-Ala-D-Pra: synthesized as described[38]. 1 ml of cell cultures was pelleted and resuspended in

50 µl of PBS buffer. Dipeptide was added to a final concentration of 0.5 mM following with 5

min incubation at room temperature. Cells were fixed by adding 70 % ethanol and incubated

for minimum 2h in -20 °C. Next, cells were washed twice with PBS in order to remove

unattached peptides, and resuspended in 50 µl of PBS. The D-Ala-D-Pra was subsequently

labelled via a click reaction with fluorescent azide (20 µM) that was incubated for 15 min at

room   temperature   with   addition   of   copper   sulphate   (CuSO4,   1   mM),   tris-

hydroxypropyltriazolylmethylamine (THPTA, 125 μM) and ascorbic acid (1.2 mM). The

sample was washed twice with PBS and resuspended in 50 µl of the same buffer.

Laurdan staining and GP calculations

Laurdan (6-Dodecanoyl-N, N-dymethyl2-naphthylamine, Sigma-Aldrich) was used to detect

the liquid ordering in the membrane, as decribed[24], with modifications. Cells were grown in

LB or SMM medium until late exponential phase. Laurdan, dissolved in dimethyformamide

(DMF), was added at 10 μM final concentration and cells were incubated for 10 min in the

dark at 37 °C, 200 rpm. Cells were then washed twice in PBS buffer supplemented with 0.2 %

(w/v) glucose and 1% (w/v) DMF, and resuspended in fresh prewarmed appropriate medium.

Laurdan was excited at 360 ± 20 nm, and fluorescence emission was captured at 460 ± 25 nm

(exposure time: 500 ms) and at 535 ± 25 nm (exposure time: 1 sec)[13]. The image analysis

including the generation of GP maps was carried out using Fiji Software[58] in combination with

the macro tool CalculateGP designed by Norbert Vischer

(http://sils.fnwi.uva.nl/bcb/objectj/examples/CalculateGP/MD/gp.html). The GP values were

measured for at least 100 individual cells after background subtraction, from two biological

replicates.

Other fluorescent membrane probes

B. subtilis cell membranes were probed with Nile Red (0.5 µg.ml^{-1}), FM4-64 (0.5 µg.ml^{-1}) or

DiI-C12 (2.5 µg.ml^{-1}). To this end, an overnight culture was grown in appropriate antibiotic,

diluted 1:100 in LB medium supplemented with DiI-C12 followed by growth until exponential

phase. Membranes were probed with Nile Red or FM4-64 for 5 min at room temperature after

reaching exponential phase. The stained cells were washed three times in prewarmed LB

medium supplemented with 1 % DMSO before visualization under fluorescence microscopy.

TIRF time lapse microscopy

Time-lapse TIRFM movies were taken in two independent experiments for each strain and

condition. To this end, overnight cultures of strains grown in LB medium supplemented with

the appropriate antibiotic were diluted 1:100 in medium containing 0.5 % (w/v) xylose and

grown until exponential phase. All experiments were performed inside the incubation chamber

set to 37 $^\circ$C, no longer than 10 min after taking the sample. The cells were imaged over 30 s

with 1 s inter-frame intervals in a continuous illumination and ultimate focus correction mode.

The single particle tracking analyses and kymographs were done using Fiji Software[58] in

combination with the MTrackJ[59] and MicrobeJ plugins[60].

Bocillin labelling

Cells were grown until an OD600 of 0.4 - 0.5, and washed twice with PBS. Next, samples were

resuspended in 50 μl PBS containing Bocillin-FL (5 μg.ml^{-1}) and incubated at room

temperature for 10 min. Subsequently cells were harvested, lysed by sonication and cell-free

extracts were prepared. Samples, equalized for culture OD, were prepared with SDS-PAGE

sample buffer and run on a 12 % SDS-PAGE gel. Fluorescent bands were visualized using a

Typhoon Trio (GE Healthcare) scanner.

Isolation of membranes

Membrane isolation was adapted from[61]. Briefly, cells were grown until an OD600 of 0.4 - 0.5,

cell fractions were collected and resuspended in PBS with Lysozyme (1 μg.ml^{-1}), EDTA (5

mM), 1/10 tablet cOmplete protease inhibitor (Roche), and DNAse (5 μg.ml^{-1}) and incubated

for 30 min on ice. Samples were sonicated, cell which did not lyze were spun down (8000 rpm,

2 min, 4°C), and the supernatant fraction was centrifuged at 4°C and 40000 rpm for 1 h. The

membrane pellet was dissolved in ACA750 buffer (750 mM aminocaproic acid, 50 mM Bis-

Tris, pH 7.0) to a final protein concentration of 1 μg/μl. Membranes were solubilized overnight

at 4°C in 1 % (w/v) dodecylmaltoside (DDM) and either used directly or stored at -20°C.

Blue native PAGE (BN-PAGE)

The experiment was performed as described[36]. Samples were prepared by mixing sample buffer

(0.1 % Ponceau S, 42.5 % Glycerol) with solubilized membranes in a 1:3 ratio. Samples were

resolved on a mini-PROTEAN™ TGX Stain-Free™ gradient gel (4-15%, BioRad) using

cathode (50 mM Tricine and 15 mM BisTris), and anode (50 mM BisTris pH 7.0) buffers. The

Novex® NativeMark™ Unstained Protein Standard® marker was used as a Mw marker.

Second dimension SDS PAGE (2D SDS-PAGE)

A lane of interest was excised from the Native-PAGE gel and immobilized horizontally on top

of a SDS-PAGE gel (5% stacking, 12% resolving). The excised fragment was flanked with a

piece of Whatman™ paper soaked with PageRuler™ Prestained Protein Ladder. The gel

fragment to be resolved in the second dimension was topped with a mix of 1 % (w/v)

LowTemperature agarose, 0.5 % (w/v) SDS and bromophenol blue. After the agarose had

solidified, standard SDS-PAGE electrophoresis was performed.

TEM

Cultures were harvested by centrifugation and a small amount of pellet was placed on a copper

dish. A 400 copper mesh grid and a 75 μm aperture grid was placed on top of the cells to create

a thin layer. The sandwiched cells were plunged rapidly into liquid propane. Sandwiches were

then disassembled and placed on frozen freeze-substitution medium containing 1 % osmium

tetroxide, 0.5 % uranyl acetate and 5 % water in acetone. Cells were dehydrated and fixed using

the rapid freeze substitution method[62]. Samples were embedded in epon and ultrathin sections

were collected on formvar coated and carbon evaporated copper grids and inspected using a

CM12 (Philips) transmission electron microscope. For each strain 70 random septa were

imaged with pixel resolution of 1.2 nm. Using ImageJ the cell wall thickness for each septum

was measured at 4 places from which the average was taken.

Statistical Analysis

Each set of micrographs to be analysed was imaged with the same exposure time. For the

septum intensity analysis of HADA, FM4-64 and Nile Red, the wildtype strain (expressing

GFP) and the Δ*floAT strains* were mixed, labelled and imaged on the same agarose pad.

Intensity of the fluorescently labelled septa was measured using the ObjectJ macro tool

PeakFinder   (https://sils.fnwi.uva.nl/bcb/objectj/examples/PeakFinder/peakfinder.html)[63].   A

perpendicular line was drawn across the septal plane, the background intensity was removed

resulting in a maximum peak intensity. The number of septa compared was indicated for every

individual experiment. Populations were compared using the non-parametric Mann-Whitney

test. The null hypothesis was tested with the *p* value of 0.05. The statistical analyses and their

graphical representation (box plots) were generated with GraphPad Prism 8.1 (San Diego,

California, USA). Box plots show the median and the interquartile range (box), the 5th and

95th percentile (whiskers). Laurdan fluorescence generalized polarization, cell length and

MreB speed statistical analyses were performed using Kruskal-Wallis with Dunn's multiple

comparison *post-hoc* test.

Fatty acid composition analysis

The fatty acid composition of *B. subtilis* wild-type cells and the flotillin/PBP mutants was

analyzed with gas chromatography as fatty acid methyl esters. Cells for the analyses were

grown at 37 °C in LB or SMM until mid-exponential ($OD_{600} \sim 0.5$), harvested (6000 rpm, 10

min, 4 °C) and washed with 100 mM NaCl. Next, the cells were freeze dried at -50 ºC, 0.012

mbar for a minimum of 18 h. All analyses were carried out on biological duplicates by the

Identification Service of the DSMZ, Braunschweig, Germany.

Sample preparation for solid-state NMR

FloT was essentially purified as described[24], in solubilized form, and stored in buffer A (50

mM Tris HCl pH 7.5, 150 mM NaCl, 5 mM $MgCl_2$) supplemented with 0.05 % Triton X-100.

Liposomes containing POPC-d31 were prepared by mixing appropriate lipid powders in

organic solvents (chloroform/methanol, 2:1 ratio). Solvents were evaporated under a flow of

N_2 to obtain a thin lipid film. Lipids were rehydrated with ultrapure water before lyophilisation

over night. The lipid powder was hydrated with an appropriate amount of buffer A with 10%

glycerol and homogenized by three cycles of vortexing, freezing (liquid nitrogen, -196°C, 1

min) and thawing (40°C in a water bath, 10 min). This protocol generated a milky suspension

of micrometer-sized multilamellar vesicles. FloT was solubilized in Buffer A supplemented

with 0.05% Triton X-100 and added to preformed liposomes and incubated for 1h at room

temperature. A dialysis step was then performed against Buffer A at 4°C under agitation to

remove the detergent. Samples were centrifuged at 100,000g at 4°C for 1h to pellet the

proteoliposomes. ^2H solid-state NMR spectra were recorded of liposomes in the presence or

absence of FloT at a lipid/protein ratio of 25:1 at 298 K.

Solid-state NMR

^{2}H NMR spectroscopy experiments were performed using a Bruker Avance III 500 MHz WB

(11.75 T) spectrometer. They were recorded on ^{2}H-labeled POPC at 76.77 MHz with a phase-

cycled quadrupolar echo pulse sequence (90°x-t-90°y-t-acq). Acquisition parameters were as

follows: spectral window of 500 kHz for 2H NMR spectroscopy, p/2 pulse width of 3.90 ms

for ^{2}H, interpulse delays (t) were of 40 µs, recycled delays of 1.3 s for 2H; 3000 and 8000 scans

were used for ^{2}H NMR spectroscopy on liposomes and liposomes with FloT, respectively.

Spectra were processed using a Lorentzian line broadening of 300 Hz for ^{2}H NMR spectra

before Fourier transformation from the top of the echo. Samples were equilibrated for 30 min

at a given temperature before data acquisition. All spectra were processed and analyzed using

Bruker Topspin 3.2 software. Spectral moments were calculated for each temperature using the

NMR Depaker 1.0rc1 software [Copyright (C) 2009 Sébastien Buchoux]. Orientational order

parameters (SCD) were calculated from experimental quadrupolar splittings (DnQ) as

described in Huster 2014. For ^{31}P ssNMR, we applied a static Hahn spin echo sequence at the

^{31}P frequency of 162 MHz on a 400 MHz (9.4T) Bruker Avance III HD spectrometer, with a

90◦ pulse of 8 µs, a delay of 40 µs, a recycle delay of 5 s, a spectral window of 400 ppm and a

number of scans of 4000 and 3400 was used on liposomes and liposomes with FloT,

respectively. Spectra were processed using a Lorentzian line broadening of 100 Hz.

Acknowledgements:

We thank Henrik Strahl for discussions and sharing of unpublished data, Rut Carballido-Lopez

for strain RWBS5 and Luiza Morawska and Oscar Kuipers for the P*rrnB*-*gfp* plasmid.

This work was funded by NWO grant 864.09.010 (DJS), DFG grants BR 2915/4-1; INST

86/1452-1 (MB), ERC starting grant 757913; NWO grant 721.014.008 (AKHH); PhD

fellowships DAAD-GSSP to AS; and SFRH/BD/78061/2011- POPH/FSE/FCT to ASB.

Contributions:

AZ, AS, ASB, BH, MBr, DJS designed experiments. AZ, AS, ASB, MBe, DM, JR, RB

performed experiments. AZ, AS, ASB, MBe, DM, JR, RB, IK, BH, MBr, DJS analysed

experiments. AH, AKHH contributed reagents. AZ, AS, BH, MBr, DJS wrote the manuscript,

all authors read and commented on the manuscript.

Competing interests

The authors declare that they have no competing interests.

References

1   Typas, A., Banzhaf, M., Gross, C. A. & Vollmer, W. From the regulation of
peptidoglycan synthesis to bacterial growth and morphology. *Nat Rev Microbiol* **10**,
123-136, doi:10.1038/nrmicro2677 (2012).
2   Zhao, H., Patel, V., Helmann, J. D. & Dorr, T. Don't let sleeping dogmas lie: new views
of peptidoglycan synthesis and its regulation. *Mol Microbiol* **106**, 847-860,
doi:10.1111/mmi.13853 (2017).
3   Cho, H. *et al.* Bacterial cell wall biogenesis is mediated by SEDS and PBP polymerase
families functioning semi-autonomously. *Nat Microbiol*, 16172,
doi:10.1038/nmicrobiol.2016.172 (2016).
4   Meeske, A. J. *et al.* SEDS proteins are a widespread family of bacterial cell wall
polymerases. *Nature* **537**, 634-638, doi:10.1038/nature19331 (2016).
5   Taguchi, A. *et al.* FtsW is a peptidoglycan polymerase that is functional only in
complex with its cognate penicillin-binding protein. *Nat Microbiol*,
doi:10.1038/s41564-018-0345-x (2019).
6   Morales Angeles, D. & Scheffers, D. J. in *Bacillus: Cellular and Molecular Biology* (ed
P. L. Graumann) Ch. 10, 295-332 (Caister Academic Press, U.K., 2017).
7   Strahl, H. & Errington, J. Bacterial Membranes: Structure, Domains, and Function.
Annual Review of Microbiology **71**, 519-538, doi:10.1146/annurev-micro-102215-
095630 (2017).
8   Hussain, S. *et al.* MreB filaments align along greatest principal membrane curvature
to orient cell wall synthesis. *eLife* **7**, doi:10.7554/eLife.32471 (2018).
9   Ursell, T. S. *et al.* Rod-like bacterial shape is maintained by feedback between cell
curvature and cytoskeletal localization. *Proc Natl Acad Sci U S A* **111**, E1025-1034,
doi:10.1073/pnas.1317174111 (2014).
10   Dominguez-Escobar, J. *et al.* Processive movement of MreB-associated cell wall
biosynthetic complexes in bacteria. *Science* **333**, 225-228,
doi:10.1126/science.1203466 (2011).
11   Garner, E. C. *et al.* Coupled, circumferential motions of the cell wall synthesis
machinery and MreB filaments in B. subtilis. *Science* **333**, 222-225,
doi:10.1126/science.1203285 (2011).

12 Billaudeau, C. *et al.* Contrasting mechanisms of growth in two model rod-shaped
bacteria. *Nat Commun* **8**, 15370, doi:10.1038/ncomms15370 (2017).
13 Strahl, H., Burmann, F. & Hamoen, L. W. The actin homologue MreB organizes the
bacterial cell membrane. *Nat Commun* **5**, 3442, doi:10.1038/ncomms4442 (2014).
14 Ganchev, D. N., Hasper, H. E., Breukink, E. & de Kruijff, B. Size and orientation of the
lipid II headgroup as revealed by AFM imaging. *Biochemistry* **45**, 6195-6202,
doi:10.1021/bi051913e (2006).
15 Witzke, S., Petersen, M., Carpenter, T. S. & Khalid, S. Molecular Dynamics
Simulations Reveal the Conformational Flexibility of Lipid II and Its Loose Association
with the Defensin Plectasin in the Staphylococcus aureus Membrane. *Biochemistry*
55, 3303-3314, doi:10.1021/acs.biochem.5b01315 (2016).
16 Muchova, K., Wilkinson, A. J. & Barak, I. Changes of lipid domains in Bacillus subtilis
cells with disrupted cell wall peptidoglycan. *FEMS Microbiol Lett* **325**, 92-98,
doi:10.1111/j.1574-6968.2011.02417.x (2011).
17 Schirner, K. *et al.* Lipid-linked cell wall precursors regulate membrane association of
bacterial actin MreB. *Nat Chem Biol* **11**, 38-45, doi:10.1038/nchembio.1689 (2015).
18 Bramkamp, M. & Lopez, D. Exploring the existence of lipid rafts in bacteria. *Microbiol*
Mol Biol Rev **79**, 81-100, doi:10.1128/MMBR.00036-14 (2015).
19 Lopez, D. & Koch, G. Exploring functional membrane microdomains in bacteria: an
overview. *Current Opinion in Microbiology* **36**, 76 - 84,
doi:https://doi.org/10.1016/j.mib.2017.02.001 (2017).
20 García-Fernández, E. *et al.* Membrane Microdomain Disassembly Inhibits MRSA
Antibiotic Resistance. *Cell* **171**, 1354-1367.e1320, doi:10.1016/j.cell.2017.10.012
(2017).
21 Lopez, D. & Kolter, R. Functional microdomains in bacterial membranes. *Genes Dev*
24, 1893-1902, doi:10.1101/gad.1945010 (2010).
22 Mielich-Suss, B. & Lopez, D. Molecular mechanisms involved in Bacillus subtilis
biofilm formation. *Environmental microbiology* **17**, 555-565, doi:10.1111/1462-
2920.12527 (2015).
23 Mielich-Suss, B., Schneider, J. & Lopez, D. Overproduction of flotillin influences cell
differentiation and shape in Bacillus subtilis. *mBio* **4**, e00719-00713,
doi:10.1128/mBio.00719-13 (2013).
24 Bach, J. N. & Bramkamp, M. Flotillins functionally organize the bacterial membrane.
Mol Microbiol **88**, 1205-1217, doi:10.1111/mmi.12252 (2013).
25 Dempwolff, F., Moller, H. M. & Graumann, P. L. Synthetic motility and cell shape
defects associated with deletions of flotillin/reggie paralogs in Bacillus subtilis and
interplay of these proteins with NfeD proteins. *J Bacteriol* **194**, 4652-4661,
doi:10.1128/JB.00910-12 (2012).
26 Yepes, A. *et al.* The biofilm formation defect of a Bacillus subtilis flotillin-defective
mutant involves the protease FtsH. *Mol Microbiol* **86**, 457-471, doi:10.1111/j.1365-
2958.2012.08205.x (2012).
27 Schneider, J. *et al.* Spatio-temporal remodeling of functional membrane
microdomains organizes the signaling networks of a bacterium. *PLoS genetics* **11**,
e1005140, doi:10.1371/journal.pgen.1005140 (2015).
28 Huang, X., Gaballa, A., Cao, M. & Helmann, J. D. Identification of target promoters
for the Bacillus subtilis extracytoplasmic function sigma factor, sigma W. *Mol*
Microbiol **31**, 361-371 (1999).

29 Nicolas, P. *et al.* Condition-dependent transcriptome reveals high-level regulatory
architecture in Bacillus subtilis. *Science* **335**, 1103-1106,
doi:10.1126/science.1206848 (2012).
30 Dempwolff, F. *et al.* Super Resolution Fluorescence Microscopy and Tracking of
Bacterial Flotillin (Reggie) Paralogs Provide Evidence for Defined-Sized Protein
Microdomains within the Bacterial Membrane but Absence of Clusters Containing
Detergent-Resistant Proteins. *PLoS genetics* **12**, e1006116,
doi:10.1371/journal.pgen.1006116 (2016).
31 Mielich-Suss, B. *et al.* Flotillin scaffold activity contributes to type VII secretion
system assembly in Staphylococcus aureus. *PLoS pathogens* **13**, e1006728,
doi:10.1371/journal.ppat.1006728 (2017).
32 Kuru, E. *et al.* In Situ probing of newly synthesized peptidoglycan in live bacteria with
fluorescent D-amino acids. *Angew Chem Int Ed Engl* **51**, 12519-12523,
doi:10.1002/anie.201206749 (2012).
33 Daniel, R. A. & Errington, J. Control of cell morphogenesis in bacteria: two distinct
ways to make a rod-shaped cell. *Cell* **113**, 767-776 (2003).
34 Morales Angeles, D. *et al.* Pentapeptide-rich peptidoglycan at the Bacillus subtilis
cell-division site. *Mol Microbiol* **104**, 319-333, doi:10.1111/mmi.13629 (2017).
35 Kahan, F. M., Kahan, J. S., Cassidy, P. J. & Kropp, H. The mechanism of action of
fosfomycin (phosphonomycin). *Ann N Y Acad Sci* **235**, 364-386, doi:10.1111/j.1749-
6632.1974.tb43277.x (1974).
36 Trip, E. N. & Scheffers, D. J. A 1 MDa protein complex containing critical components
of the Escherichia coli divisome. *Sci Rep* **5**, 18190, doi:10.1038/srep18190 (2015).
37 Scheffers, D. J. & Errington, J. PBP1 Is a Component of the Bacillus subtilis Cell
Division Machinery. *Journal of Bacteriology* **186**, 5153-5156 (2004).
38 Sarkar, S., Libby, E. A., Pidgeon, S. E., Dworkin, J. & Pires, M. M. In Vivo Probe of Lipid
II-Interacting Proteins. *Angew Chem Int Ed Engl* **55**, 8401-8404,
doi:10.1002/anie.201603441 (2016).
39 Dajkovic, A. *et al.* Hydrolysis of peptidoglycan is modulated by amidation of meso-
diaminopimelic acid and Mg(2+) in Bacillus subtilis. *Mol Microbiol* **104**, 972-988,
doi:10.1111/mmi.13673 (2017).
40 Konopasek, I., Strzalka, K. & Svobodova, J. Cold shock in Bacillus subtilis: different
effects of benzyl alcohol and ethanol on the membrane organisation and cell
adaptation. *Biochim Biophys Acta* **1464**, 18-26 (2000).
41 Molugu, T. R., Lee, S. & Brown, M. F. Concepts and Methods of Solid-State NMR
Spectroscopy Applied to Biomembranes. *Chemical Reviews* **117**, 12087-12132,
doi:10.1021/acs.chemrev.6b00619 (2017).
42 Legrand, A. *et al.* Nanodomain Clustering of the Plant Protein Remorin by Solid-State
NMR. *Frontiers in Molecular Biosciences* **6**, doi:10.3389/fmolb.2019.00107 (2019).
43 Scheidt, H. A. & Huster, D. Structure and Dynamics of the Myristoyl Lipid
Modification of Src Peptides Determined by 2H Solid-State NMR Spectroscopy.
Biophysical Journal **96**, 3663-3672, doi:https://doi.org/10.1016/j.bpj.2009.02.028
(2009).
44 Huster, D. Solid-state NMR spectroscopy to study protein–lipid interactions.
Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids **1841**,
1146-1160, doi:https://doi.org/10.1016/j.bbalip.2013.12.002 (2014).

45 Bach, J. N. & Bramkamp, M. Dissecting the molecular properties of prokaryotic
flotillins. *PLoS One* **10**, e0116750, doi:10.1371/journal.pone.0116750 (2015).
46 Dion, M. F. *et al.* Bacillus subtilis cell diameter is determined by the opposing actions
of two distinct cell wall synthetic systems. *Nat Microbiol*, doi:10.1038/s41564-019-
0439-0 (2019).
47 Calvez, P., Jouhet, J., Vie, V., Durmort, C. & Zapun, A. Lipid Phases and Cell Geometry
During the Cell Cycle of Streptococcus pneumoniae. *Frontiers in microbiology* **10**,
351, doi:10.3389/fmicb.2019.00351 (2019).
48 Gohrbandt, M. *et al.* Low membrane fluidity triggers lipid phase separation and
protein segregation in vivo. *bioRxiv*, 852160, doi:10.1101/852160
(2019).
49 Pazos, M. *et al.* Z-ring membrane anchors associate with cell wall synthases to
initiate bacterial cell division. *Nat Commun* **9**, 5090, doi:10.1038/s41467-018-07559-
2 (2018).
50 Varma, A. & Young, K. D. In Escherichia coli, MreB and FtsZ direct the synthesis of
lateral cell wall via independent pathways that require PBP 2. *J Bacteriol* **191**, 3526-
3533, doi:10.1128/JB.01812-08 (2009).
51 Aaron, M. *et al.* The tubulin homologue FtsZ contributes to cell elongation by guiding
cell wall precursor synthesis in Caulobacter crescentus. *Molecular Microbiology* **64**,
938-952, doi:doi:10.1111/j.1365-2958.2007.05720.x (2007).
52 Szenk, M., Dill, K. A. & de Graff, A. M. R. Why Do Fast-Growing Bacteria Enter
Overflow Metabolism? Testing the Membrane Real Estate Hypothesis. *Cell Syst* **5**, 95-
104, doi:10.1016/j.cels.2017.06.005 (2017).
53 Donovan, C. & Bramkamp, M. Characterization and subcellular localization of a
bacterial flotillin homologue. *Microbiology* **155**, 1786-1799,
doi:10.1099/mic.0.025312-0 (2009).
54 Harwood, C. R. & Cutting, S. M. *Molecular Biological Methods for Bacillus*. (John
Wiley and Sons, 1990).
55 Lennox, E. S. Transduction of linked genetic characters of the host by bacteriophage
P1. *Virology* **1**, 190-206, doi:10.1016/0042-6822(55)90016-7 (1955).
56 Anagnostopoulos, C. & Spizizen, J. Requirements for transformation in Bacillus
subtilis. *Journal of Bacteriology* **81**, 741-746 (1961).
57 Kuru, E., Tekkam, S., Hall, E., Brun, Y. V. & Van Nieuwenhze, M. S. Synthesis of
fluorescent D-amino acids and their use for probing peptidoglycan synthesis and
bacterial growth in situ. *Nat Protoc* **10**, 33-52, doi:10.1038/nprot.2014.197 (2015).
58 Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat
Methods* **9**, 676-682, doi:10.1038/nmeth.2019 (2012).
59 Meijering, E., Dzyubachyk, O. & Smal, I. Methods for cell and particle tracking.
Methods Enzymol **504**, 183-200, doi:10.1016/B978-0-12-391857-4.00009-4 (2012).
60 Ducret, A., Quardokus, E. M. & Brun, Y. V. MicrobeJ, a tool for high throughput
bacterial cell detection and quantitative analysis. *Nat Microbiol* **1**, 16077,
doi:10.1038/nmicrobiol.2016.77 (2016).
61 Schneider, J., Mielich-Suss, B., Bohme, R. & Lopez, D. In vivo characterization of the
scaffold activity of flotillin on the membrane kinase KinC of Bacillus subtilis.
Microbiology **161**, 1871-1887, doi:10.1099/mic.0.000137 (2015).

62 McDonald, K. L. Out with the old and in with the new: rapid specimen preparation
procedures for electron microscopy of sectioned biological material. *Protoplasma*
251, 429-448, doi:10.1007/s00709-013-0575-y (2014).
63 Vischer, N. O. *et al.* Cell age dependent concentration of Escherichia coli divisome
proteins analyzed with ImageJ and ObjectJ. *Frontiers in microbiology* **6**, 586,
doi:10.3389/fmicb.2015.00586 (2015).
64 Scheffers, D.-J., Jones, L. J. F. & Errington, J. Several distinct localization patterns for
penicillin-binding proteins in Bacillus subtilis. *Mol Microbiol* **51**, 749-764 (2004).
65 Lages, M. C., Beilharz, K., Morales Angeles, D., Veening, J. W. & Scheffers, D. J. The
localization of key Bacillus subtilis penicillin binding proteins during cell growth is
determined by substrate availability. *Environmental microbiology* **15**, 3272-3281,
doi:10.1111/1462-2920.12206 (2013).
66 Veening, J. W., Murray, H. & Errington, J. A mechanism for cell cycle regulation of
sporulation initiation in Bacillus subtilis. *Genes Dev* **23**, 1959-1970,
doi:10.1101/gad.528209 (2009).
756
757
758
759

760 **Table 1. Strain list**

Strain	Genotype	Source
168	*trpC2*	Laboratory collection
BB001	*trpC2 yqfA::tet*	[24]
BB003	*trpC2 yuaG::pMUTIN4 yqfA::tet*	[24]
DB003	*trpC2 yuaG::pMUTIN4*	[53]
2082	*trpC2 pbpD::cat Pxyl–gfp–pbpD $^{1–510}$*	[64]
2083	*trpC2 ponA::cat Pxyl–gfp–ponA $^{1–394}$*	[64]
2085	*trpC2 dacA::cat Pxyl–gfp–dacA $^{1–423}$*	[64]
3105	*trpC2 pbpC::cat Pxyl–gfp–pbpC $^{1–768}$*	[64]
3122	*trpC2 pbpB::cat Pxyl-gfp-pbpB $^{1–825}$*	[64]
3511	*trpC2 ponA::spc*	[37]
4042	*trpC2 pbpA::cat Pxyl-mkate2-pbpA $^{1–804}$*	[65]
4056	*trpC2 dacA::kan*	[34]
RWBS5	*amyE::spc Pxyl mrfpruby-mreB*	[10]
4059	*trpC2 dacA::cat Pxyl–gfp–dacA $^{1–423}$ yuaG::pMUTIN4 yqfA::tet*	This work
4064	*trpC2 dacA::kan yuaG::pMUTIN4 yqfA::tet*	This work
4090	*trpC2 ponA::spc yuaG::pMUTIN4*	This work
4091	*ponA::spc yqfA::tet*	This work
4092	*trpC2 ponA::spc yuaG::pMUTIN4 yqfA::tet*	This work
4095	*trpC2 ponA::cat Pxyl–gfp–ponA $^{1–394}$ yuaG::pMUTIN4 yqfA::tet*	This work

4099	*trpC2 pbpB::cat Pxyl-gfp-pbpB* ^{1–825} *yuaG::pMUTIN4 yqfA::tet*	This work
4102	*trpC2 pbpA::cat Pxyl-mkate2-pbpA* ^{1–804} *yuaG::pMUTIN4 yqfA::tet*	This work
4108	*trpC2 pbpD::cat Pxyl–gfp–pbpD* ^{1–510} *yuaG::pMUTIN4 yqfA::tet*	This work
4122	*trpC2 pbpC::cat Pxyl–gfp–pbpC* ^{1–768} *yuaG::pMUTIN4 yqfA::tet*	This work
4128	*trpC2 ponA::spc pbpD::cat Pxyl–gfp–pbpD* ^{1–510}	This work
4129	*trpC2 ponA::spc yuaG::pMUTIN4 pbpD::cat Pxyl–gfp–pbpD* ^{1–510}	This work
4070	*trpC2 mreB::kan amyE::spc Pxyl mrfpruby-mreB*	This work
4076	*trpC2 mreB::kan amyE::spc Pxyl mrfpruby-mreB yuaG::pMUTIN4 yqfA::tet*	This work
4259	*trpC2 amyE::P_{rrnB}-GFP*	This work, [66]