Membrane fluidity controls peptidoglycan synthesis and MreB movement.

Aleksandra Zielińska[#,1], Abigail Savietto[#,2,7], Anabela de Sousa Borges[1], Joël R. Roelofsen[1], Alwin M. Hartman[3,4,5], Rinse de Boer[6], Ida J. van der Klei[6], Anna K. H. Hirsch[3,4,5], Marc Bramkamp[*,2,7], Dirk-Jan Scheffers[*,1]

[#]equal contribution; [*]corresponding author

[1]Molecular Microbiology, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, the Netherlands; [2]Biozentrum, Ludwig-Maximilians-Universität München, Germany; [3] Department of Drug Design and Optimization (DDOP), Helmholtz-Institute for Pharmaceutical Research Saarland (HIPS) - Helmholtz Centre for Infection Research (HZI), Saarbrücken, Germany; [4]Department of Pharmacy, Saarland University, Saarbrücken, Germany; [5]Strathingh Institute for Chemistry, University of Groningen, The Netherlands; [6]Molecular Cell Biology, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, the Netherlands. [7]Institute for General Microbiology, Christian-Albrechts-University Kiel, Germany.

[*]correspondence: marc.bramkamp@lmu.de; d.j.scheffers@rug.nl

Abstract

The bacterial plasma membrane is an important cellular compartment. In recent years it has become obvious that protein complexes and lipids are not uniformly distributed within membranes. Current hypotheses suggest that flotillin proteins are required for the formation of complexes of membrane proteins including cell-wall synthetic proteins. We show here that bacterial flotillins are important factors for membrane fluidity homeostasis. Loss of flotillins leads to changes in membrane fluidity that in turn lead to alterations in MreB dynamics and, as a consequence, in peptidoglycan synthesis. Our data support a model in which flotillins are required for growth-rate dependent control of membrane fluidity rather than for the formation of protein complexes via direct protein-protein interactions.

Introduction

The shape of a bacterium is predominantly defined by the structure of its peptidoglycan. Although there is a great variety in bacterial shapes, the overall chemistry of peptidoglycan is very similar between bacteria and thus the shape of peptidoglycan is primarily determined by the temporal and spatial regulation of peptidoglycan synthesis. In rod-shaped bacteria, peptidoglycan synthesis is thought to be mediated by two protein assemblies, the elongasome and the divisome, that synthesize peptidoglycan along the long axis and across the division plane of the cell, respectively[1,2]. These complexes contain a set of proteins required for the final steps of synthesis and translocation of the peptidoglycan precursor, LipidII, from the inner to the outer leaflet of the cytoplasmic membrane, and proteins that incorporate LipidII into peptidoglycan. These include SEDS (Shape, Elongation, Division and Sporulation) proteins that can perform glycosyl transferase reactions[3-5], and Penicillin Binding Proteins (PBPs) that are divided in class A PBPs (aPBPs) that catalyse both glycosyl transferase and transpeptidase reactions, class B PBPs (bPBPs) that only catalyse transpeptidase reactions and low molecular weight PBPs that modify peptidoglycan, as well as hydrolases[2,6].

Coordination of these complexes is linked to cytoskeletal elements, MreB (-like proteins) for the elongasome and FtsZ for the divisome. In models, the cytoplasmic membrane is often depicted as a passive environment in which these machineries are embedded. However, it is becoming clear that the structure of the membrane plays a critical role in the coordination of peptidoglycan synthesis[7]. Inward membrane curvature serves as a localization trigger for MreB and the elongasome, and enhanced local synthesis at bulges straightens out the membrane sufficient to convey a rod shape to spherical cells[8,9]. Interestingly, MreB localizes to and organizes regions of increased membrane fluidity (RIF)[10], which in turn is linked to the presence of LipidII, which favours a more fluid membrane and promotes local membrane disorder[11,12]. Inhibition of LipidII synthesis by genetic or chemical means results in a

dissolution of membrane structures observed with the dye FM 4-64 and release of MreB from the membrane[13-16]. In *Bacillus subtilis*, MreB interacts with the aPBP PBP1[17] and both proteins require an intact membrane potential for localization[18,19].

Next to RIFs, membrane regions of decreased fluidity have been identified in bacteria[7,20,21]. These so-called functional membrane microdomains (FMMs) are thought to be organized by the bacterial flotillin proteins, are enriched in isoprenoid lipids [22,23], and can be found in so-called Detergent Resistant Membrane (DRM) fractions of the membrane. Since the formulation of the FMM hypothesis, FMMs have been linked to many processes, such as protein secretion, biofilm formation, competence and cell morphology[24-27]. Cell morphology effects are linked to cell wall synthesis, and analysis of the protein content of *Bacillus subtilis* DRMs identified several PBPs, MreC and other proteins involved in cell wall metabolism as well as the two flotillins, FloA and FloT[23,26,28]. FloA is constitutively expressed, whereas FloT is expressed primarily during stationary growth and sporulation[29]. Super resolution microscopy showed that the flotillins and other proteins in DRMs do not colocalize and have different dynamics[30], so it is unlikely that FMMs are regions in the membrane that offer a favourable environment for membrane protein activity. Recently, the hypothesis has been put forward that FMMs/flotillins form a platform for the formation of functional protein oligomers, as work in *Staphylococcus aureus* showed that multimerization of Type 7 secretion systems and PBP2a depends on FMMs[21,22,31].

Here, we have analysed the role of flotillins in peptidoglycan synthesis in *B. subtilis.* Our results show that, at high growth rates, flotillins control membrane fluidity in a manner that is critical for peptidoglycan synthesis and MreB dynamics, but have no effect on PBP oligomerization. This results in a new model for flotillin function in the physical organization of membranes during fast growth.

78   Results

79   Absence of flotillins leads to increased peptidoglycan synthesis and membrane fluidity at

80   division septa

In previous studies, a double deletion of *floA/floT* was either reported to suffer severe shape

defects and perturbed membrane structure[27] or not strong shape defects but a change in the

overall lipid ordering of the membrane[26]. We grew wild-type and Δ*floAT* strains and analysed

exponentially growing cells. We did not observe striking shape defects but did see an increase

in median cell length and distribution of cell lengths in the absence of flotillins (**Fig**. 1A, B).

To look at effects on peptidoglycan synthesis and membrane structure, we labelled cells with

HADA, a fluorescent D-Alanine analogue that reports on sites of active peptidoglycan

synthesis[32], fluorescent vancomycin (Van-FL), which labels LipidII and peptidoglycan

containing pentapeptide side chains[33,34], as well as FM4-64, Nile-Red and DiI-C12, which are

lipid dyes that accumulate in zones enriched in fluid lipids[10]. This revealed a significant

accumulation of both peptidoglycan synthesis stains and lipid dyes at division septa in the

Δ*floAT* strain (**Fig**. 1A, C-G). The Δ*floAT* strain also showed a stronger accumulation of DiI-

C12 in patches, suggesting that the more fluid regions of the membrane are coalescing into

larger regions. Inspection of the septa by electron microscopy revealed that there was no

difference between the thicknesses of the septa between the wild type and Δ*floAT* strain, ruling

out that the increase in signal was due to formation of thicker septa (**Fig**. S1). We also ruled

out that the observed difference was due to a change in the expression, folding or complex

formation by PBPs in the absence of flotillins, using combinations of SDS-PAGE and Native-

PAGE (**Fig.** S2). Also, none of five functional GFP-PBPs examined changed their localization

in the Δ*floAT* strain (**Fig.** S2). Overall, the data indicate that in the absence of flotillins,

peptidoglycan synthesis and membrane fluidity are increased at division septa.

Figure. 1 Accumulation of peptidoglycan synthesis and membrane material at division

sites in a flotillin mutant.

A. Morphology of the exponentially growing wild type (WT) and Δ*floAT* strains labelled with

HADA, fluorescent Vancomycin (Van-FL), Nile Red, FM 4-64, and DiIC12. Scale bar 5 μm.

B. Distribution of the cell length of the strains analyzed in (A). Statistical analysis of the data

(n=100) was performed with Prism, resulting in box plot graphs. **C-G**. Peak intensity of HADA

(C), Van-FL (D), Nile Red (E), FM4-64 (F) and DiI-C12 (G) labelled division sites of the cells

shown in (A). Cells from each strain (n ≥ 100, except E, n = 60) were analysed using the ObjectJ

macro tool PeakFinder followed with statistical analysis with Prism. Significant differences are

based on the two-tailed Mann-Whitney test (* p<0.05; ** p<0.001).

The absence of both flotillins and PBP1 causes a severe phenotype, linked to a loss of

membrane fluidity.

We reasoned that a non-lethal defect in septal peptidoglycan synthesis could reveal more about

the role of flotillins and constructed a flotillin mutant that lacks PBP1, a bifunctional glycosyl

transferase/transpeptidase that is required for efficient cell division[35]. Simultaneous deletion of

pbp1, *floA*, and *floT* resulted in strong filamentation and delocalization of peptidoglycan synthesis as well as membrane dyes to patches **(Fig. 2A, Fig. S3A)**. Deletion of single flotillin genes and PBP1 had similar, albeit less severe effects **(Fig. S3B, C)**. To exclude the possibility that an alteration of peptidoglycan modification resulted in the delocalization of HADA and Van-FL, we used D-Alanine-D-Propargylglycine (D-Ala-D-Pra), a clickable dipeptide analogue which is exclusively incorporated into peptidoglycan via LipidII[36]. D-Ala-D-Pra incorporation was delocalized in the Δ*pbp1*Δ*floAT* strain, indicating that peptidoglycan synthesis itself is delocalized (**Fig. S3D**). So far, our experiments were done with fast growing cells and Lysogeny Broth (LB) as the growth medium. Strikingly, all same strains had no apparent phenotype when cultivated in Spizizen's minimal medium (SMM, **Fig. 2B**), and peptidoglycan synthesis and membrane material were no longer accumulating at division sites in the Δ*floAT* strain (**Fig. S4**). SMM has a higher Mg^{2+} concentration, which is known to rescue various cell shape mutations by inhibition of cell wall hydrolysis[37]. However, the increase in Mg^{2+} was not sufficient to explain the reversal of phenotype as cells grown on LB supplemented with Mg^{2+} (6 mM, concentration in SMM, or 20 mM) still displayed the elongated phenotype with delocalized peptidoglycan synthesis **(Fig. S5)**. This indicated that the phenotypes associated with the lack of flotillins are growth-rate related.

Figure 2. Cell morphology and cell wall synthesis localization is dependent on growth conditions. A. Morphology of the *WT*, Δ*floAT*, Δ*pbp1*, and Δ*pbp1*Δ*floAT* strains grown in rich (LB), minimal (SMM) medium, and in rich medium with membrane fluidizing conditions (0.1 % benzyl alcohol, LB+BnOH). Cells were labelled with HADA, and aberrant cell shape and peptidoglycan synthesis are indicated with arrowheads. **B.** Corresponding cell length distributions are shown (n ≥ 100). Distributions were analysed using Dunn's multiple comparison tests after Kruskal–Wallis. Statistically significant cell length distribution classes (*p < 0.001*) are represented as letters above each graph. Scale bar: 4 μm.

Next, we determined lipid packing order in the different strains using the fluorescent dye Laurdan, a reporter for flotillin-mediated lipid ordering[26]. LB-grown cells lacking flotillins displayed an increased generalized polarization (GP)[26], indicative of an overall increase in ordered lipid packing in the membrane, but the effect of flotillins on membrane ordering

completely disappeared when cells were grown on SMM **(Fig. 3)**. Overall ordered lipid

packing was increased in cells grown on SMM compared to LB **(Fig. 3)**, and the absence of

PBP1 alone had a slight effect on membrane ordering in LB-grown cells, but did not contribute

to the increase in GP when combined with flotillin deletions **(Fig. 3)**. The changes in lipid

ordering were not due to changes in the overall fatty acid composition of the membranes - the

ratios of C17/C15 side chains and *iso/anteiso* fatty acids, which are indicative of fluidity[10],

were identical for wild type and Δ*floAT* strains (**Fig. S6**).

Figure 3. Flotillins increase overall membrane fluidity at high growth rate. Changes in
overall membrane fluidity were assessed by Laurdan microscopy in cells grown on LB (**A**),
SMM (**B**) and LB+BnOH (**C**). Micrographs show colour-coded generalized polarization (GP)
maps in which red indicates regions of decreased fluidity (scale bar: 4 µm). Correspondent
theoretical GP measurements in the graphs vary from -1 (more fluid) to 1 (less fluid).
Significant statistical differences according to Dunn's multiple comparison tests after Kruskal–
Wallis are represented as letters above each graph (*p < 0.001;* n ≥ 150, two biological
replicates).

Restoring membrane fluidity rescues normal peptidoglycan synthesis.

The GP values indicated that membranes are more ordered when cells are grown on minimal

medium, and this suggests that the flotillin-associated increase in overall membrane fluidity is

important for cell shape control at high growth rates. This was tested by growing the strains

lacking flotillins and PBP1 on LB in the presence of the membrane fluidizer benzyl alcohol[38].

Indeed, increasing fluidity (see Fig. 3C) restored normal cell length and normal peptidoglycan

synthesis patterns to the *pbp1/floA/floT* strain **(Fig. 2C)**.

In *B. subtilis*, the rate of growth and of peptidoglycan synthesis is linked to the speed of MreB

movement – in minimal media, the speed of MreB patches is reduced compared to the speed

in rich media[39]. Analysis of the movement of a fully functional mrfp-ruby-MreB fusion[14] by

time lapse TIRF (Total Internal Reflection Fluorescence) microscopy, confirmed that MreB

patch mobility is higher in cells grown on LB than in cells grown on SMM, with MreB speeds

similar to those reported previously[39] **(Fig. 4, Movie S1, 2)**. Strikingly, in the absence of

flotillins, MreB patch mobility was notably decreased in cells grown on LB, while in SMM

grown cells MreB patch mobility was independent of the presence of flotillins (**Fig. 4, Movie**

S3,4). Fluidizing the membrane with benzyl alcohol almost completely restored MreB mobility

in LB grown cells **(Fig. 4, Movie S5, 6)**. Notably, the growth rates of the wildtype and the

floA/floT strains were identical, also in the presence of benzyl alcohol (**Fig. S7**). This result

suggests that the synthesis of peptidoglycan, that is mediated by the MreB-associated

elongasome, is reduced in fast growing cells in the absence of flotillins, and that fluidizing the

membrane can restore this synthesis. Moreover, the results indicate that the MreB patch

mobility is, in part, controlled by membrane fluidity.

Figure 4. MreB speed is linked to membrane fluidity. (A) The MreB speed in different strain backgrounds and growth conditions was analysed by time-lapse TIRF microscopy. Scatter plot of the speed of patches obtained from individual tracks in 5 different cells are represented per fusion and condition. Average speeds are shown; error bars indicate the standard deviation. Significant statistical differences according to Dunn's multiple comparison tests after Kruskal–Wallis are represented (*p < 0.001*). **(B)** Representative kymographs showing fast and slow moving patches of mrfRuby-MreB in *B. subtilis* cells lacking endogenous *mreB* (WT) or *mreB* and *floAT* (Δ*floAT*). See supplementary movies S1-S6 for corresponding raw image series.

Discussion

Our data provide an alternative explanation for the role of flotillins in peptidoglycan synthesis which does not require that flotillins act as a platform for PBP oligomerization as suggested before[40]. We propose that flotillin-mediated control of membrane fluidity at high growth rates is critical and sufficient to allow essential membrane bound processes, such as peptidoglycan synthesis, to occur. Given the identical fatty acid composition of wild-type and flotillin-mutant strains, we propose that this control is direct, through efficient separation of states of liquid ordered and disordered lipid domains in the membrane bilayer[26]. The predominantly physical role in membrane organization for flotillins fits with our observation that adding a chemical fluidizer is sufficient to restore MreB dynamics and cell shape to fast growing cells that lack

flotillins. We also note that membrane fluidity is not solely a function of temperature, but also

of growth conditions.

A sufficiently fluid membrane is necessary for the efficient recruitment and movement of

MreB, and provides a more favorable environment for the peptidoglycan precursor LipidII[8,9,16].

It has recently been shown that modulation of either MreBCD or PBP1 levels is sufficient to

alter the shape of *B. subtilis* cells[41], underscoring the importance of both systems. In the

absence of flotillins, the activity of the MreBCD component is strongly reduced – as evidenced

by the reduction of MreB speed – and the overall rigidity of the membrane is increased which

results in a less favourable environment for the peptidoglycan precursor LipidII, which prefers

more liquid, disordered membrane phases[11,12,42]. To compensate, membrane fluidity and

peptidoglycan synthesis activity is increased at division sites in flotillin mutants, which is

sufficient to keep the overall cell shape intact, although cells are elongated. The increased

fluidity at division sites is in line with a recent study that showed phases of different fluidity in

Streptococcus pneumoniae membranes, with more fluid membranes and LipidII localizing at

midcell[42] where the membrane is most bent. It may seem paradoxical that cells elongate when

elongasome activity is reduced, but one has to remember that a large amount of the

peptidoglycan synthesis contributing to elongation of bacterial cells is actually taking place at

midcell, before the ingrowth of the septum[43-45]. The shift to peptidoglycan synthesis at future

division sites makes the activity of PBP1 critical and explains why its deletion has such a

dramatic effect in cells lacking flotillins. Restoring fluidity using a chemical fluidizer allows

the MreBCD component to again efficiently drive peptidoglycan synthesis during elongation,

which is sufficient to suppress the phenotype. At low growth rates, there is no difference

between wild type cells and cells lacking flotillins with respect to membrane fluidity, and the

speed of MreB is similar between the two cell types - the net effect of this is that cells lacking

PBP1 and flotillins behave as cells that only lack PBP1, which is similar to wild type. An

overall rigidification of the membrane may also lead to retardation of processes which require

membrane modifications such as division and sporulation, which is indeed observed in *B.*

subtilis flotillin mutants[27,46].

One of the proposed roles for flotillin proteins is that they form a 'platform' that transiently

interacts with membrane proteins that need to oligomerize into functional complexes[21]. We

tested this hypothesis for *B. subtilis* PBPs by comparing their localization and oligomerization

in wild type and flotillin mutant strains. Although we were capable of detecting a high MW

complex containing various PBPs (notably PBP1, 2a, 2b, 3 and 4), the complex was not

dependent on the presence of flotillins. We also note that PBP1 was present in the complex, as

well as present in a large smear in the first dimension native gel, which would explain why

PBP1 was detected by mass spectrometry analysis of a native PAGE band containing FloA[29].

PBP5, on the other hand, was not part of the high MW complex, which fits with its role in

processing of the terminal D-Ala from stem-peptides that have not been cross-linked, which it

exercises over the entire surface of the cell[32]. Although we cannot exclude that flotillins may

affect PBPs that are not easily detected by Bocillin-FL, our results do not provide any evidence

for a role for flotillins in the oligomerization of PBPs in *B. subtilis*. This extends the finding of

the Graumann lab that found either transient or no colocalization between flotillins and other

proteins present in DRM fractions[30]. Although it is obvious that peptidoglycan synthesis is

altered in the absence of flotillins, our data strongly suggest that the basis for this alteration is

in the physical organisation of the membrane rather than inefficient formation of divisome or

elongasome complexes in the absence of flotillins, as flotillin mutants strains show no synthetic

phenotype on minimal medium, and the defects on rich medium can be reverted by chemically

fluidizing the membrane.

In conclusion, our data provide a new model for flotillin function in the physical organization

of membranes during fast growth. The observation that flotillins differentially affect the

membrane at different growth rates also explains the diversity of phenotypes described for

flotillin mutants in the literature.

Methods

B. subtilis strains and growth conditions

All *B. subtilis* strains used in this study are derived from strain 168 and are listed in **Table 1**.

Construction of new strains was based on natural competence of *B. subtilis*[47]. Gene integration

or deletion was validated by colony PCR whereas the expression and localization of the

fluorescent fusions was additionally validated by microscopy. Cells were grown either in LB[48]

or Spizizen minimal medium (SMM)[49] supplemented with 1% glucose, at 37°C and 200 rpm,

unless indicated otherwise. Induction of the P_{xyl} promoter was triggered by addition of 0.2 -

0.5 % xylose. Cell cultures were supplemented with spectinomycin (50 μg/ml), tetracycline

(10 μg/ml), chloramphenicol (5 μg/ml), kanamycin (5 μg/ml), erythromycin (1 μg/ml), benzyl

alcohol (BnOH, 0.1 %) or magnesium sulphate ($MgSO_4$, 6-20 mM) when necessary.

Fluorescence microscopy

For standard fluorescence microscopy, exponentially growing cells were immobilized on

microscope slides covered with a thin film of 1 % agarose (w/v) in water or the appropriate

medium. For TIRFM, agarose pads were mounted using Gene Frames (1.7 x 2.8 cm chamber,

0.25 mm thickness, 125 μL volume) from ThermoScientific. Standard fluorescence

microscopy was carried out using an Axio Zeiss Imager M1 fluorescence microscope (EC Plan-

Neofluar 100x/1.30 Oil Ph3 objective) equipped with an AxioCam HRm camera and an Nikon-

Ti-E microscope (Nikon Instruments, Tokyo, Japan) equipped with Hamamatsu Orca Flash 4.0

camera..

For Laurdan and TIRFM experiments, a Delta Vision Elite microscope (Applied Precision, GE

Healthcare) equipped with an Insight SSI[TM] Illumination, an X4 Laser module, a CoolSnap

HQ^2 CCD camera and a temperature-controlled chamber set up at 37 ^{0}C was used. Laurdan

images were taken with an Olympus UplanSApo 100x/1.4 oil objective. TIRFM image series

were taken using an Olympus UAPO N 100X/1.49 TIRF objective and a 561 nm laser (50 mW,

100 % power). Data processing was performed with softWoRx Suite 2.0 Software.

Visualization of cell wall synthesis

Peptidoglycan (PG) synthesis was assessed by labelling the cells with HADA (7-

hydroxycoumarin 3-carboxylic acid-amino-D-alanine)[50], Van-FL[33] or D-Ala-D-Pra[36].

HADA: synthesized as described[34]. Overnight cultures of *B. subtilis* strains were diluted 1:100

into fresh LB medium or LB medium supplemented with 0.1 % (w/v) of benzyl alcohol

(BnOH), a membrane fluidizer. Cells were grown until exponential phase, a sample of 1 ml of

culture was spun down for 30 sec, 5000 × g and the cell pellet was resuspended in 25 µl of

fresh pre-warmed LB or LB containing 0.1% (w/v) BnOH. HADA was added to a 50 µM final

concentration. Cells were incubated for 10 min in the dark (37 ^{0}C, 200 rpm) and then washed

twice in 1 ml PBS buffer (58 mM Na_2HPO_4, 17 mM NaH_2PO_4, 68 mM NaCl, pH 7.3) to remove

the excess of unbounded HADA. Cells were spun down again and resuspended in 25 µl of the

appropriate medium and 2 µl of cells were mounted on 1% agarose slides before visualization.

Visualization of HADA patterns (excitation: 358 nm/emission: 461 nm) under fluorescence

microscopy from two biological replicates and cell length measurements were taken from at

least 100 cells each strain/treatment.

Van-FL: a 1:1 mixture of vancomycin (Sigma Aldrich) and BODIPY®FL Vancomycin

(Molecular Probes) at a final concentration 1 µg/ml was used to label cells for 5-10 min at room

temperature.

D-Ala-D-Pra: synthesized as described[36]. 1 ml of cell cultures was pelleted and resuspended in

50 µl of PBS buffer. Dipeptide was added to a final concentration of 0.5 mM following with 5

min incubation at room temperature. Cells were fixed by adding 70 % ethanol and incubated

for minimum 2h in -20 °C. Next, cells were washed twice with PBS in order to remove

unattached peptides, and resuspended in 50 µl of PBS. The D-Ala-D-Pra was subsequently

labelled via a click reaction with fluorescent azide (20 µM) that was incubated for 15 min at

room temperature with addition of copper sulphate ($CuSO_4$, 1 mM), tris-

hydroxypropyltriazolylmethylamine (THPTA, 125 µM) and ascorbic acid (1.2 mM). The

sample was washed twice with PBS and resuspended in 50 µl of the same buffer.

Laurdan staining and GP calculations

Laurdan (6-Dodecanoyl-N, N-dymethyl2-naphthylamine, Sigma-Aldrich) was used to detect

the liquid ordering in the membrane, as decribed[26], with modifications. Cells were grown in

LB or SMM medium until late exponential phase. Laurdan, dissolved in dimethyformamide

(DMF), was added at 10 µM final concentration and cells were incubated for 10 min in the

dark at 37 °C, 200 rpm. Cells were then washed twice in PBS buffer supplemented with 0.2 %

(w/v) glucose and 1% (w/v) DMF, and resuspended in fresh prewarmed appropriate medium.

Laurdan was excited at 360 ± 20 nm, and fluorescence emission was captured at 460 ± 25 nm

(exposure time: 500 ms) and at 535 ± 25 nm (exposure time: 1 sec)[10]. The image analysis

including the generation of GP maps was carried out using Fiji Software[51] in combination with

the macro tool CalculateGP designed by Norbert Vischer

(http://sils.fnwi.uva.nl/bcb/objectj/examples/CalculateGP/MD/gp.html). The GP values were

measured for at least 100 individual cells after background subtraction, from two biological

replicates.

Other fluorescent membrane probes

B. subtilis cell membranes were probed with Nile Red (0.5 µg.ml^{-1}), FM4-64 (0.5 µg.ml^{-1}) or DiI-C12 (2.5 µg.ml^{-1}). To this end, an overnight culture was grown in appropriate antibiotic, diluted 1:100 in LB medium supplemented with DiI-C12 followed by growth until exponential phase. Membranes were probed with Nile Red or FM4-64 for 5 min at room temperature after reaching exponential phase. The stained cells were washed three times in prewarmed LB medium supplemented with 1 % DMSO before visualization under fluorescence microscopy.

TIRF time lapse microscopy

Time-lapse TIRFM movies were taken in two independent experiments for each strain and condition. To this end, overnight cultures of strains grown in LB medium supplemented with the appropriate antibiotic were diluted 1:100 in medium containing 0.5 % (w/v) xylose and grown until exponential phase. All experiments were performed inside the incubation chamber set to 37 $^{\circ}$C, no longer than 10 min after taking the sample. The cells were imaged over 30 s with 1 s inter-frame intervals in a continuous illumination and ultimate focus correction mode. The single particle tracking analyses and kymographs were done using Fiji Software[51] in combination with the MTrackJ[52] and MicrobeJ plugins[53].

Bocillin labelling

Cells were grown until an OD600 of 0.4 - 0.5, and washed twice with PBS. Next, samples were resuspended in 50 µl PBS containing Bocillin-FL (5 µg.ml^{-1}) and incubated at room temperature for 10 min. Subsequently cells were harvested, lysed by sonication and cell-free extracts were prepared. Samples, equalized for culture OD, were prepared with SDS-PAGE sample buffer and run on a 12 % SDS-PAGE gel. Fluorescent bands were visualized using a Typhoon Trio (GE Healthcare) scanner.

Isolation of membranes

Membrane isolation was adapted from[54]. Briefly, cells were grown until an OD600 of 0.4 - 0.5, cell fractions were collected and resuspended in PBS with Lysozyme (1 µg.ml^{-1}), EDTA (5 mM), 1/10 tablet cOmplete protease inhibitor (Roche), and DNAse (5 µg.ml^{-1}) and incubated for 30 min on ice. Samples were sonicated, cell which did not lyze were spun down (8000 rpm, 2 min, 4°C), and the supernatant fraction was centrifuged at 4°C and 40000 rpm for 1 h. The membrane pellet was dissolved in ACA750 buffer (750 mM aminocaproic acid, 50 mM Bis-Tris, pH 7.0) to a final protein concentration of 1 µg/µl. Membranes were solubilized overnight at 4°C in 1 % (w/v) dodecylmaltoside (DDM) and either used directly or stored at -20°C.

Blue native PAGE (BN-PAGE)

The experiment was performed as described[55]. Samples were prepared by mixing sample buffer (0.1 % Ponceau S, 42.5 % Glycerol) with solubilized membranes in a 1:3 ratio. Samples were resolved on a mini-PROTEAN™ TGX Stain-Free™ gradient gel (4-15%, BioRad) using cathode (50 mM Tricine and 15 mM BisTris), and anode (50 mM BisTris pH 7.0) buffers. The Novex® NativeMark™ Unstained Protein Standard® marker was used as a Mw marker.

Second dimension SDS PAGE (2D SDS-PAGE)

A lane of interest was excised from the Native-PAGE gel and immobilized horizontally on top of a SDS-PAGE gel (5% stacking, 12% resolving). The excised fragment was flanked with a piece of Whatman™ paper soaked with PageRuler™ Prestained Protein Ladder. The gel fragment to be resolved in the second dimension was topped with a mix of 1 % (w/v) LowTemperature agarose, 0.5 % (w/v) SDS and bromophenol blue. After the agarose had solidified, standard SDS-PAGE electrophoresis was performed.

TEM

Cultures were harvested by centrifugation and a small amount of pellet was placed on a copper dish. A 400 copper mesh grid and a 75 μm aperture grid was placed on top of the cells to create a thin layer. The sandwiched cells were plunged rapidly into liquid propane. Sandwiches were then disassembled and placed on frozen freeze-substitution medium containing 1 % osmium tetroxide, 0.5 % uranyl acetate and 5 % water in acetone. Cells were dehydrated and fixed using the rapid freeze substitution method[56]. Samples were embedded in epon and ultrathin sections were collected on formvar coated and carbon evaporated copper grids and inspected using a CM12 (Philips) transmission electron microscope. For each strain 70 random septa were imaged with pixel resolution of 1.2 nm. Using ImageJ the cell wall thickness for each septum was measured at 4 places from which the average was taken.

Statistical Analysis

Each set of micrographs to be analysed was imaged with the same exposure time. Intensity of the fluorescently labelled septa was measured using the ObjectJ macro tool PeakFinder (https://sils.fnwi.uva.nl/bcb/objectj/examples/PeakFinder/peakfinder.html)[57]. A perpendicular line was drawn across the septal plane, the background intensity was removed resulting in a maximum peak intensity. The number of septa compared was indicated for every individual experiment. The Kolgomorov-Smirnoff test was used to test for normal distribution of values, and non-parametric tests were employed. The null hypothesis was tested with the *p* value of 0.05 or 0.001. The statistical analyses and their graphical representation (box plots) were generated with GraphPad Prism 8.1 (San Diego, California, USA). Box plots show the median and the interquartile range (box), the 5th and 95th percentile (whiskers). Laurdan fluorescence

generalized polarization, cell length and MreB speed statistical analyses were performed using

Kruskal-Wallis with Dunn's multiple comparison *post-hoc* test.

Fatty acid composition analysis

The fatty acid composition of *B. subtilis* wild-type cells and the flotillin/PBP mutants was

analyzed with gas chromatography as fatty acid methyl esters. Cells for the analyses were

grown at 37 °C in LB or SMM until mid-exponential ($OD_{600} \sim 0.5$), harvested (6000 rpm, 10

min, 4 °C) and washed with 100 mM NaCl. Next, the cells were freeze dried at -50 ºC, 0.012

mbar for a minimum of 18 h. All analyses were carried out on biological duplicates by the

Identification Service of the DSMZ, Braunschweig, Germany.

Acknowledgements:

We thank R. Carballido-Lopez for strain RWBS5.

This work was funded by NWO grant 864.09.010 (DJS), DFG grants BR 2915/4-1; INST

86/1452-1 (MB), ERC starting grant 757913; NWO grant 721.014.008 (AKHH); PhD

fellowships DAAD-GSSP to AS; and SFRH/BD/78061/2011- POPH/FSE/FCT to ASB.

Contributions:

AZ, AS, ASB, MB, DJS designed experiments. AZ, AS, ASB, JR, RB performed experiments.

AZ, AS, ASB, JR, RB, IK, MB, DJS analysed experiments. AH, AKHH contributed reagents.

AZ, AS, MB, DJS wrote the manuscript, all authors read and commented on the manuscript.

Competing interests

The authors declare that they have no competing interests.

References

1   Typas, A., Banzhaf, M., Gross, C. A. & Vollmer, W. From the regulation of
peptidoglycan synthesis to bacterial growth and morphology. *Nat Rev Microbiol* **10**,
123-136, doi:10.1038/nrmicro2677 (2012).
2   Zhao, H., Patel, V., Helmann, J. D. & Dorr, T. Don't let sleeping dogmas lie: new views
of peptidoglycan synthesis and its regulation. *Mol Microbiol* **106**, 847-860,
doi:10.1111/mmi.13853 (2017).
3   Cho, H. *et al.* Bacterial cell wall biogenesis is mediated by SEDS and PBP polymerase
families functioning semi-autonomously. *Nat Microbiol*, 16172,
doi:10.1038/nmicrobiol.2016.172 (2016).
4   Meeske, A. J. *et al.* SEDS proteins are a widespread family of bacterial cell wall
polymerases. *Nature* **537**, 634-638, doi:10.1038/nature19331 (2016).
5   Taguchi, A. *et al.* FtsW is a peptidoglycan polymerase that is functional only in
complex with its cognate penicillin-binding protein. *Nat Microbiol*,
doi:10.1038/s41564-018-0345-x (2019).
6   Morales Angeles, D. & Scheffers, D. J. in *Bacillus: Cellular and Molecular Biology* (ed
P. L. Graumann) Ch. 10, 295-332 (Caister Academic Press, U.K., 2017).
7   Strahl, H. & Errington, J. Bacterial Membranes: Structure, Domains, and Function.
Annual Review of Microbiology **71**, 519-538, doi:10.1146/annurev-micro-102215-
095630 (2017).
8   Hussain, S. *et al.* MreB filaments align along greatest principal membrane curvature
to orient cell wall synthesis. *eLife* **7**, doi:10.7554/eLife.32471 (2018).
9   Ursell, T. S. *et al.* Rod-like bacterial shape is maintained by feedback between cell
curvature and cytoskeletal localization. *Proc Natl Acad Sci U S A* **111**, E1025-1034,
doi:10.1073/pnas.1317174111 (2014).
10 Strahl, H., Burmann, F. & Hamoen, L. W. The actin homologue MreB organizes the
bacterial cell membrane. *Nat Commun* **5**, 3442, doi:10.1038/ncomms4442 (2014).
11 Ganchev, D. N., Hasper, H. E., Breukink, E. & de Kruijff, B. Size and orientation of the
lipid II headgroup as revealed by AFM imaging. *Biochemistry* **45**, 6195-6202,
doi:10.1021/bi051913e (2006).
12 Witzke, S., Petersen, M., Carpenter, T. S. & Khalid, S. Molecular Dynamics
Simulations Reveal the Conformational Flexibility of Lipid II and Its Loose Association
with the Defensin Plectasin in the Staphylococcus aureus Membrane. *Biochemistry*
55, 3303-3314, doi:10.1021/acs.biochem.5b01315 (2016).
13 Muchova, K., Wilkinson, A. J. & Barak, I. Changes of lipid domains in Bacillus subtilis
cells with disrupted cell wall peptidoglycan. *FEMS Microbiol Lett* **325**, 92-98,
doi:10.1111/j.1574-6968.2011.02417.x (2011).
14 Dominguez-Escobar, J. *et al.* Processive movement of MreB-associated cell wall
biosynthetic complexes in bacteria. *Science* **333**, 225-228,
doi:10.1126/science.1203466 (2011).
15 Garner, E. C. *et al.* Coupled, circumferential motions of the cell wall synthesis
machinery and MreB filaments in B. subtilis. *Science* **333**, 222-225,
doi:10.1126/science.1203285 (2011).
16 Schirner, K. *et al.* Lipid-linked cell wall precursors regulate membrane association of
bacterial actin MreB. *Nat Chem Biol* **11**, 38-45, doi:10.1038/nchembio.1689 (2015).
17 Kawai, Y., Daniel, R. A. & Errington, J. Regulation of cell wall morphogenesis in
Bacillus subtilis by recruitment of PBP1 to the MreB helix. *Molecular Microbiology*
71, 1131-1144 (2009).

18   Lages, M. C., Beilharz, K., Morales Angeles, D., Veening, J. W. & Scheffers, D. J. The
localization of key Bacillus subtilis penicillin binding proteins during cell growth is
determined by substrate availability. *Environmental microbiology* **15**, 3272-3281,
doi:10.1111/1462-2920.12206 (2013).

19   Strahl, H. & Hamoen, L. W. Membrane potential is important for bacterial cell
division. *Proc Natl Acad Sci U S A* **107**, 12281-12286, doi:10.1073/pnas.1005485107
(2010).

20   Bramkamp, M. & Lopez, D. Exploring the existence of lipid rafts in bacteria. *Microbiol*
Mol Biol Rev **79**, 81-100, doi:10.1128/MMBR.00036-14 (2015).

21   Lopez, D. & Koch, G. Exploring functional membrane microdomains in bacteria: an
overview. *Current Opinion in Microbiology* **36**, 76 - 84,
doi:https://doi.org/10.1016/j.mib.2017.02.001 (2017).

22   García-Fernández, E. *et al.* Membrane Microdomain Disassembly Inhibits MRSA
Antibiotic Resistance. *Cell* **171**, 1354-1367.e1320, doi:10.1016/j.cell.2017.10.012
(2017).

23   Lopez, D. & Kolter, R. Functional microdomains in bacterial membranes. *Genes Dev*
24, 1893-1902, doi:10.1101/gad.1945010 (2010).

24   Mielich-Suss, B. & Lopez, D. Molecular mechanisms involved in Bacillus subtilis
biofilm formation. *Environmental microbiology* **17**, 555-565, doi:10.1111/1462-
2920.12527 (2015).

25   Mielich-Suss, B., Schneider, J. & Lopez, D. Overproduction of flotillin influences cell
differentiation and shape in Bacillus subtilis. *mBio* **4**, e00719-00713,
doi:10.1128/mBio.00719-13 (2013).

26   Bach, J. N. & Bramkamp, M. Flotillins functionally organize the bacterial membrane.
Mol Microbiol **88**, 1205-1217, doi:10.1111/mmi.12252 (2013).

27   Dempwolff, F., Moller, H. M. & Graumann, P. L. Synthetic motility and cell shape
defects associated with deletions of flotillin/reggie paralogs in Bacillus subtilis and
interplay of these proteins with NfeD proteins. *J Bacteriol* **194**, 4652-4661,
doi:10.1128/JB.00910-12 (2012).

28   Yepes, A. *et al.* The biofilm formation defect of a Bacillus subtilis flotillin-defective
mutant involves the protease FtsH. *Mol Microbiol* **86**, 457-471, doi:10.1111/j.1365-
2958.2012.08205.x (2012).

29   Schneider, J. *et al.* Spatio-temporal remodeling of functional membrane
microdomains organizes the signaling networks of a bacterium. *PLoS genetics* **11**,
e1005140, doi:10.1371/journal.pgen.1005140 (2015).

30   Dempwolff, F. *et al.* Super Resolution Fluorescence Microscopy and Tracking of
Bacterial Flotillin (Reggie) Paralogs Provide Evidence for Defined-Sized Protein
Microdomains within the Bacterial Membrane but Absence of Clusters Containing
Detergent-Resistant Proteins. *PLoS genetics* **12**, e1006116,
doi:10.1371/journal.pgen.1006116 (2016).

31   Mielich-Suss, B. *et al.* Flotillin scaffold activity contributes to type VII secretion
system assembly in Staphylococcus aureus. *PLoS pathogens* **13**, e1006728,
doi:10.1371/journal.ppat.1006728 (2017).

32   Kuru, E. *et al.* In Situ probing of newly synthesized peptidoglycan in live bacteria with
fluorescent D-amino acids. *Angew Chem Int Ed Engl* **51**, 12519-12523,
doi:10.1002/anie.201206749 (2012).

526	33	Daniel, R. A. & Errington, J. Control of cell morphogenesis in bacteria: two distinct ways to make a rod-shaped cell. *Cell* **113**, 767-776 (2003).
528	34	Morales Angeles, D. *et al.* Pentapeptide-rich peptidoglycan at the Bacillus subtilis cell-division site. *Mol Microbiol* **104**, 319-333, doi:10.1111/mmi.13629 (2017).
530	35	Scheffers, D. J. & Errington, J. PBP1 Is a Component of the Bacillus subtilis Cell Division Machinery. *Journal of Bacteriology* **186**, 5153-5156 (2004).
532	36	Sarkar, S., Libby, E. A., Pidgeon, S. E., Dworkin, J. & Pires, M. M. In Vivo Probe of Lipid II-Interacting Proteins. *Angew Chem Int Ed Engl* **55**, 8401-8404, doi:10.1002/anie.201603441 (2016).
535	37	Dajkovic, A. *et al.* Hydrolysis of peptidoglycan is modulated by amidation of meso-diaminopimelic acid and Mg(2+) in Bacillus subtilis. *Mol Microbiol* **104**, 972-988, doi:10.1111/mmi.13673 (2017).
538	38	Konopasek, I., Strzalka, K. & Svobodova, J. Cold shock in Bacillus subtilis: different effects of benzyl alcohol and ethanol on the membrane organisation and cell adaptation. *Biochim Biophys Acta* **1464**, 18-26 (2000).
541	39	Billaudeau, C. *et al.* Contrasting mechanisms of growth in two model rod-shaped bacteria. *Nat Commun* **8**, 15370, doi:10.1038/ncomms15370 (2017).
543	40	Garcia-Fernandez, E. *et al.* Membrane Microdomain Disassembly Inhibits MRSA Antibiotic Resistance. *Cell* **171**, 1354-1367 e1320, doi:10.1016/j.cell.2017.10.012 (2017).
546	41	Dion, M. F. *et al.* Bacillus subtilis cell diameter is determined by the opposing actions of two distinct cell wall synthetic systems. *Nat Microbiol*, doi:10.1038/s41564-019-0439-0 (2019).
549	42	Calvez, P., Jouhet, J., Vie, V., Durmort, C. & Zapun, A. Lipid Phases and Cell Geometry During the Cell Cycle of Streptococcus pneumoniae. *Frontiers in microbiology* **10**, 351, doi:10.3389/fmicb.2019.00351 (2019).
552	43	Pazos, M. *et al.* Z-ring membrane anchors associate with cell wall synthases to initiate bacterial cell division. *Nat Commun* **9**, 5090, doi:10.1038/s41467-018-07559-2 (2018).
555	44	Varma, A. & Young, K. D. In Escherichia coli, MreB and FtsZ direct the synthesis of lateral cell wall via independent pathways that require PBP 2. *J Bacteriol* **191**, 3526-3533, doi:10.1128/JB.01812-08 (2009).
558	45	Aaron, M. *et al.* The tubulin homologue FtsZ contributes to cell elongation by guiding cell wall precursor synthesis in Caulobacter crescentus. *Molecular Microbiology* **64**, 938-952, doi:doi:10.1111/j.1365-2958.2007.05720.x (2007).
561	46	Donovan, C. & Bramkamp, M. Characterization and subcellular localization of a bacterial flotillin homologue. *Microbiology* **155**, 1786-1799, doi:10.1099/mic.0.025312-0 (2009).
564	47	Harwood, C. R. & Cutting, S. M. *Molecular Biological Methods for Bacillus*. (John Wiley and Sons, 1990).
566	48	Lennox, E. S. Transduction of linked genetic characters of the host by bacteriophage P1. *Virology* **1**, 190-206, doi:10.1016/0042-6822(55)90016-7 (1955).
568	49	Anagnostopoulos, C. & Spizizen, J. Requirements for transformation in Bacillus subtilis. *Journal of Bacteriology* **81**, 741-746 (1961).
570	50	Kuru, E., Tekkam, S., Hall, E., Brun, Y. V. & Van Nieuwenhze, M. S. Synthesis of fluorescent D-amino acids and their use for probing peptidoglycan synthesis and bacterial growth in situ. *Nat Protoc* **10**, 33-52, doi:10.1038/nprot.2014.197 (2015).

51 Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*
Methods **9**, 676-682, doi:10.1038/nmeth.2019 (2012).
52 Meijering, E., Dzyubachyk, O. & Smal, I. Methods for cell and particle tracking.
Methods Enzymol **504**, 183-200, doi:10.1016/B978-0-12-391857-4.00009-4 (2012).
53 Ducret, A., Quardokus, E. M. & Brun, Y. V. MicrobeJ, a tool for high throughput
bacterial cell detection and quantitative analysis. *Nat Microbiol* **1**, 16077,
doi:10.1038/nmicrobiol.2016.77 (2016).
54 Schneider, J., Mielich-Suss, B., Bohme, R. & Lopez, D. In vivo characterization of the
scaffold activity of flotillin on the membrane kinase KinC of Bacillus subtilis.
Microbiology **161**, 1871-1887, doi:10.1099/mic.0.000137 (2015).
55 Trip, E. N. & Scheffers, D. J. A 1 MDa protein complex containing critical components
of the Escherichia coli divisome. *Sci Rep* **5**, 18190, doi:10.1038/srep18190 (2015).
56 McDonald, K. L. Out with the old and in with the new: rapid specimen preparation
procedures for electron microscopy of sectioned biological material. *Protoplasma*
251, 429-448, doi:10.1007/s00709-013-0575-y (2014).
57 Vischer, N. O. *et al.* Cell age dependent concentration of Escherichia coli divisome
proteins analyzed with ImageJ and ObjectJ. *Frontiers in microbiology* **6**, 586,
doi:10.3389/fmicb.2015.00586 (2015).
58 Scheffers, D.-J., Jones, L. J. F. & Errington, J. Several distinct localization patterns for
penicillin-binding proteins in Bacillus subtilis. *Mol Microbiol* **51**, 749-764 (2004).

597 **Table 1. Strain list**

Strain	Genotype	Source
168	*trpC2*	Laboratory collection
BB001	*trpC2 yqfA::tet*	[26]
BB003	*trpC2 yuaG::pMUTIN4 yqfA::tet*	[26]
DB003	*trpC2 yuaG::pMUTIN4*	[46]
2082	*trpC2 pbpD::cat Pxyl–gfp–pbpD* $^{1-510}$	[58]
2083	*trpC2 ponA::cat Pxyl–gfp–ponA* $^{1-394}$	[58]
2085	*trpC2 dacA::cat Pxyl–gfp–dacA* $^{1-423}$	[58]
3105	*trpC2 pbpC::cat Pxyl–gfp–pbpC* $^{1-768}$	[58]
3122	*trpC2 pbpB::cat Pxyl-gfp-pbpB* $^{1-825}$	[58]
3511	*trpC2 ponA::spc*	[35]
4042	*trpC2 pbpA::cat Pxyl-mkate2-pbpA* $^{1-804}$	[18]
4056	*trpC2 dacA::kan*	[34]
RWBS 5	*amyE::spc Pxyl mrfpruby-mreB*	[14]
4059	*trpC2 dacA::cat Pxyl–gfp–dacA* $^{1-423}$ *yuaG::pMUTIN4 yqfA::tet*	This work
4064	*trpC2 dacA::kan yuaG::pMUTIN4 yqfA::tet*	This work

4090	*trpC2 ponA::spc yuaG::pMUTIN4*	This work
4091	*ponA::spc yqfA::tet*	This work
4092	*trpC2 ponA::spc yuaG::pMUTIN4 yqfA::tet*	This work
4095	*trpC2 ponA::cat Pxyl–gfp–ponA $^{1–394}$ yuaG::pMUTIN4 yqfA::tet*	This work
4099	*trpC2 pbpB::cat Pxyl-gfp-pbpB $^{1–825}$ yuaG::pMUTIN4 yqfA::tet*	This work
4102	*trpC2 pbpA::cat Pxyl-mkate2-pbpA $^{1–804}$ yuaG::pMUTIN4 yqfA::tet*	This work
4108	*trpC2 pbpD::cat Pxyl–gfp–pbpD $^{1–510}$ yuaG::pMUTIN4 yqfA::tet*	This work
4122	*trpC2 pbpC::cat Pxyl–gfp–pbpC $^{1–768}$ yuaG::pMUTIN4 yqfA::tet*	This work
4128	*trpC2 ponA::spc pbpD::cat Pxyl–gfp–pbpD $^{1–510}$*	This work
4129	*trpC2 ponA::spc yuaG::pMUTIN4 pbpD::cat Pxyl–gfp–pbpD $^{1–510}$*	This work
4070	*trpC2 mreB::kan amyE::spc Pxyl mrfpruby-mreB*	This work
4076	*trpC2 mreB::kan amyE::spc Pxyl mrfpruby-mreB yuaG::pMUTIN4 yqfA::tet*	This work