Evolved differences in *cis* and *trans* regulation between the maternal and zygotic mRNA complements in the *Drosophila* embryo

Emily L. Cartwright and Susan E. Lott

Department of Evolution and Ecology, University of California, Davis CA 95616

Data availability: The raw sequencing reads and processed data are available at GEO/NCBI at accession number GSE136646.

Running title: Regulatory evolution in embryogenesis

Keywords: gene expression regulation, maternal mRNA deposition, zygotic

transcription, *cis* and *trans* regulation

Corresponding Author Information:

Emily L. Cartwright

Department of Evolution and Ecology

2320 Storer Hall

University of California, Davis

Davis, CA 95616

(530)-752-1112

elcartwright@ucdavis.edu

Susan E. Lott

Department of Evolution and Ecology

2320 Storer Hall

University of California, Davis

Davis, CA 95616

(530) 752-2939

selott@ucdavis.edu

ABSTRACT

How gene expression can evolve depends on the mechanisms driving gene

expression. Gene expression is controlled in different ways in different developmental

stages; here we ask whether different developmental stages show different patterns of

regulatory evolution. To explore the mode of regulatory evolution, we used the early

stages of embryonic development controlled by two different genomes, that of the

mother and that of the zygote. During embryogenesis in all animals, initial

developmental processes are driven entirely by maternally provided gene products

deposited into the oocyte. The zygotic genome is activated later, when developmental

control is handed off from maternal gene products to the zygote during the maternal-to-

zygotic transition. Using hybrid crosses between sister species of *Drosophila* (*D.*

simulans, *D. sechellia*, and *D. mauritiana*) and transcriptomics, we find that the

regulation of maternal transcript deposition and zygotic transcription evolve through

different mechanisms. We find that patterns of transcript level inheritance in hybrids,

relative to parental species, differ between maternal and zygotic transcripts, and

maternal transcript levels are more likely to be conserved. Changes in transcript levels

occur predominantly through differences in *trans* regulation for maternal genes, while

changes in zygotic transcription occur through a combination of both *cis* and *trans*

regulatory changes. Differences in the underlying regulatory landscape in the mother

and the zygote are likely the primary determinants for how maternal and zygotic

transcripts evolve.

INTRODUCTION

Since the proposal that regulation of gene expression may be one of the primary

drivers of morphological evolution was introduced (King and Wilson 1975; Carroll 1995),

much research has been directed toward elucidating mechanisms underlying the

evolution of gene expression. The mechanistic basis of regulatory control determines

the substrate for evolution of gene expression. At the transcriptional level, gene

expression is controlled by *cis*-regulatory elements (regulatory DNA sequences, such as

enhancers or promoters) and by *trans* acting factors (such as transcription factors or

miRNAs that bind the regulatory DNA of many genes). Due to their modular structure

(Britten and Davidson 1969; Davidson and Peter 2015), changes in *cis*-regulatory

elements can lead to the evolution of altered gene expression and thus new traits

(Prud'homme *et al.* 2006; Chan *et al.* 2010; Kvon *et al.* 2016) without the pleiotropic

consequences for other critical traits controlled by the same gene. On the other hand,

transcription factors that bind to enhancers often have higher pleiotropy and thus can

affect the expression of many genes. This poses a fundamental mechanistic question

as to how gene regulatory evolution occurs: whether changes are more likely to occur in

cis-regulatory elements or *trans*-acting factors.

In order to determine the relative contributions of *cis* and *trans* elements to the

evolution of gene expression genome-wide, previous studies implemented the use of

genetic hybrids and methods of detecting allele-specific expression (Wittkopp *et al.*

2004; Landry *et al.* 2005; Graze *et al.* 2009; McManus *et al.* 2010; Coolon *et al.* 2014;

León-Novelo *et al.* 2014). Several studies point to differences in *cis* regulation as the

primary mechanism of change in transcript abundance within or between species

(Graze *et al.* 2009; Mack *et al.* 2016), while other studies indicate that *trans* changes

are more widespread (McManus *et al.* 2010; Glaser-Schmitt *et al.* 2018). Previous

research has proposed that the difference in the abundance of *cis* versus *trans* changes

affecting gene expression can be explained by the divergence times of the strains or

species being compared (Coolon *et al.* 2014), or the particular tissue type examined

(Buchberger *et al.* 2019). Given this body of previous work, it is surprising that, to our

knowledge, no previous study has compared *cis* and *trans* contributions to gene

expression evolution across developmental stages in a model organism.

In this study, we ask whether contributions of *cis* and *trans* changes to gene

regulatory evolution differ across developmental stages. We chose stages during

embryogenesis that are close together in developmental time, yet are likely to broadly

differ in the mechanistic basis of regulation. The general regulatory architecture of a

particular stage likely affects how regulation at this stage can evolve. The early stages

of development utilized here are under regulatory control of entirely different genomes:

that of the mother, and that of the zygote. The earliest developmental processes in

embryogenesis are regulated by maternally provided RNA and protein, which lay the

foundation for the rest of development (Tadros and Lipshitz 2009; Vastenhouw *et al.*

2019). These maternally derived gene products carry out all initial developmental events

because at the time of fertilization, the zygotic genome is transcriptionally silent.

Because the zygotic genome is not yet transcriptionally active, post-transcriptional

mechanisms also play an important role in regulating the amount of maternal gene

products present (Tadros *et al.* 2007; Rouget *et al.* 2010; Barckmann and Simonelig

2013). As the zygotic genome is activated, control of developmental processes is

handed off from the maternally deposited factors to those derived from the zygotic

genome in a process known as the maternal-to-zygotic transition (MZT). The MZT is a

highly conserved and regulated process during early development that occurs in all

animals and in some species of flowering plants (Baroux *et al.* 2008; Tadros and

Lipshitz 2009; Vastenhouw *et al.* 2019).

Regulation of early zygotic gene expression has been extensively studied over

many years (Mannervik 2014; Schulz and Harrison 2019), while regulatory control of

maternal genes is not well understood. Studies of maternal transcripts have largely

focused on their transport into the oocyte (Mische *et al.* 2007; Kugler and Lasko 2009),

their localization and movement within the oocyte and embryo (Theurkauf and Hazelrigg

1998; Kugler and Lasko 2009), their post-transcriptional regulation (Salles *et al.* 1994),

and their degradation (Tadros *et al.* 2007; Bushati *et al.* 2008; Laver *et al.* 2015).

Zygotic gene expression is carefully controlled, with classic examples such as the *even-*

skipped gene having multiple enhancer elements along with multiple transcription

factors responsible for producing complex expression patterns in developmental time

and embryonic space (Small *et al.* 1992; Perry *et al.* 2011; Mannervik 2014). In contrast,

maternal transcripts are produced by support cells called nurse cells during oogenesis,

which are polyploid and highly transcriptionally active (Kugler and Lasko 2009; Lasko

2012), rapidly producing large amounts of transcripts. Roughly 50-75% of the genome is

maternally deposited (Tadros *et al.* 2007; De Renzis *et al.* 2007; Thomsen *et al.* 2010;

Lott *et al.* 2011; Vastenhouw *et al.* 2019) and there is considerable post-transcriptional

regulation of maternal factors (Tadros *et al.* 2007; Bushati *et al.* 2008; Laver *et al.*

2015). For these reasons, regulation at the transcript level may not need to be as

precise. While the mechanisms behind maternal transcription are not well understood,

the regulatory environments driving the production of the maternal and zygotic

transcriptomes are quite different.

To study the regulatory basis of differences in transcript levels at the maternal

and zygotic stages of early development between species, we focused on three closely

related species of *Drosophila* (*D. simulans*, *D. sechellia*, and *D. mauritiana*). Despite

having a relatively close divergence time of 250,000 years (McDermott and Kliman

2008), these sister species have differences in the pools of transcripts present in the

developing embryo both at a stage where only maternal transcripts are present, and at

a stage after zygotic genome activation (ZGA; Atallah and Lott 2018). By comparing

hybrids and parental lines of the species *D. simulans*, *D. sechellia*, and *D. mauritiana*,

we asked whether, at each of these two developmental stages, changes in gene

expression between species occurred due to changes in *cis*, in *trans*, or in a

combination of the two.

We found that patterns of gene regulatory changes between species are distinct

across developmental stages for maternally deposited transcripts and for genes with

primarily zygotic expression (see Methods) in early embryogenesis. Differences in

maternal transcripts occur much more frequently due to *trans* as opposed to *cis*

regulatory changes, while differences in zygotic gene transcription occur through a mix

of *cis*, *trans*, and the combined action of *cis* and *trans* regulatory changes. The complex

pattern of changes found in our study at the zygotic stage speaks to what has been

known about regulation at this stage for some time, that both *cis* and *trans* elements are

necessary for the intricate control of gene expression in time and space at this stage of

embryogenesis. The large proportion of *trans* regulatory signal found at the maternal

stage may reveal fundamental properties of the regulatory architecture during

oogenesis. *Trans* regulators can affect a large number of genes at the same time, as

might be necessary to maximize mRNA production to load sufficient numbers of

transcripts into the egg. We also identified motifs associated with *trans* regulation at the

level of chromatin at the maternal stage, which lends evidence to an emerging picture of

how gene expression might be regulated during oogenesis. Overall, we find distinct

patterns of gene regulatory changes at the two embryonic timepoints, before and after

ZGA, indicating evolved changes in gene regulation differ based on the developmental

context.

128   **MATERIALS AND METHODS**

Crossing scheme and sample acquisition

Three *Drosophila* species were used for this study: *D. sechellia* (Dsec/wild-

type;14021-0248.25) and *D. simulans* (Dsim/w[501]; 14021-0251.011) from the 12

Genomes study (Clark *et al.* 2007) and *D. mauritiana* (Dmau/[w1];14021-0241.60). For

interspecific crosses, each vial was set up using 7-12 virgin females from one species

and 7-10 males from another. We did not cross *D. sechellia* females and *D. simulans*

males, as this combination is known to be incompatible (Lachaise *et al.* 1986). Two

types of hybrid crosses were established from which embryos were collected: 1) to

determine the regulatory basis of changes in zygotic gene expression, hybrid F1

embryos were collected; and 2) to determine the regulatory basis of changes in

maternal gene expression, embryos produced by hybrid F1 mothers were collected. To

investigate the regulatory basis of changes in zygotic gene expression, we collected

hybrid F1 embryos at the very end of blastoderm stage, stage 5 (Bownes' stages;

Bownes 1975; Campos-Ortega and Hartenstein 2013), a timepoint after the zygotic

genome is activated. We define late stage 5 by morphology; it is the point when

cellularization is complete, but gastrulation has not yet begun. In order to determine the

regulatory basis of changes in maternal gene expression, similar crosses were

established with hybrid females from the F1 generations of the initial crosses and males

that were the same species as the maternal species in the parental cross. We set up

crosses in this manner in order to establish consistency amongst crosses, although the

male genotype is unlikely to affect our data. The contribution of sperm mRNA to the

zygote is debated but known contributions are small (Fischer *et al.* 2012) and likely not

detectable via RNA-sequencing (Ali-Murthy *et al.* 2013). This second set of crosses was

used to collect stage 2 embryos (Bownes' stages; Bownes 1975; Campos-Ortega and

Hartenstein 2013), during which time only maternal gene products are present. At this

point in development, the cytoplasm has retracted from the vitelline membrane at the

anterior and posterior poles of the embryo but pole cells have not yet migrated to the

posterior (Ashburner 1989). As is conventional in *Drosophila* genetics, we denote our

crosses by listing the female genotype first and the male genotype second. For

example, in a cross between *D. mauritiana* and *D. simulans*, we write the genotype of

the resulting stage 5 hybrid embryo as *mau* x *sim*. We describe the hybrid genotype of

stage 2 embryos from backcrosses of hybrid F1 females and males of the maternal

species in the initial cross in a similar way, e.g. (*mau* x *sim*) x *mau* (also see Figure 1 for

cross diagram).

All flies were raised in vials on standard cornmeal media at 25°C. Flies were

allowed to lay eggs for ~2 hours (for collecting stage 2 embryos) and ~3 hours (for

collecting stage 5 embryos) before they were transferred to a new vial so that the eggs

could be harvested. Eggs were collected from 4-14 day old females, dechorionated

using 50% bleach and moved into halocarbon oil on a microscope slide for staging.

Embryos were staged at the appropriate developmental time point under a microscope

(Zeiss AxioImager M2), imaged, and promptly collected at stage 2 or at the end of stage

5 (Bownes' stages; Bownes 1975; Campos-Ortega and Hartenstein 2013), of embryonic

development.

To collect the samples after staging, the embryos were quickly transferred with a

paintbrush to Parafilm (Bemis) and rolled (to remove excess halocarbon oil) into a drop

of TRIzol (Ambion). The embryos were ruptured with a needle so that the contents

dissolved in the TRIzol and were transferred to a tube to be frozen at -80°C until

extraction. RNA was extracted using glycogen as a carrier (as per manufacturer

instructions) in a total volume of 1mL TRIzol. Approximately 80-120ng total RNA was

extracted from individual embryos, measured using a Qubit 2.0 fluorimeter (Invitrogen).

The quality of the RNA was validated on RNA Pico Chips using an Agilent Bioanalyzer.

Genotyping was performed to determine embryo sex for stage 5 samples, as

dosage compensation is not complete and transcript levels for genes on the X

chromosome may differ for males and females at this time in development (Lott *et al.*

2014). DNA was extracted from each sample along with the RNA as per manufacturer

instructions and amplified using a whole genome amplification kit (illustra GenomePhi

v2, GE Healthcare). Sex-specific primers (Table S1) designed for use with all three

species, two sets for the Y chromosome (to the *ORY* and *kl2* genes) and one control set

(to *ftz*), were used to genotype the single embryos after genome amplification. For the

stage 5 samples, a total of three male and three female embryos from each cross were

used for sequencing. One noted exception is that in the *sim* x *mau* cross, a total of four

female and two male embryos were collected, as determined from the transcriptomic

data. A total of three stage 2 embryos were collected. We did not perform genotyping

for embryo sex on the stage 2 embryos because the zygotic genome is not yet active at

this stage in development.

Library preparation and sequencing

The RNA from single embryos was treated with DNase (TurboDNA-free, Life

AM1907) using manufacturer instructions and RNA sequencing libraries were

constructed with Illumina TruSeq v2 kits following the manufacturer's low sample

protocol. The Illumina protocol uses oligo (dT) beads to enrich for polyadenylated

transcripts. Because poly(A) tail length is important in determining many post-

transcriptional processes during early development, including translational efficiency, it

is important to ensure that the method used for mRNA selection does not produce a

biased set of poly(A) tail lengths. Previous datasets report poly(A) length distributions

for transcripts during oogenesis and early development (Lim *et al.* 2016; Eichhorn *et al.*

2016). We could not directly compare our data to previous reports, as these studies

were done using *D. melanogaster*, which may have a different poly(A) tail length

distribution than the species used in our analysis. However, previous studies comparing

distributions of poly(A) tail lengths of all genes to poly(A) tail lengths of transcripts

recovered through poly(A) selection in *D. melanogaster* have demonstrated that poly(A)

selection with commonly used methods does not bias which transcripts are recovered

from the total pool of transcripts present (Eichhorn *et al.* 2016). This includes studies

that used the same single embryo approaches utilized here (Crofton *et al.* 2018; Atallah

and Lott 2018). Therefore, it seems unlikely that poly(A) selection heavily biases the

extracted RNA relative to the RNAs present at these developmental stages. cDNA

libraries were quantified using a Qubit 2.0 fluorimeter (dsDNA BR Assay Kits) and the

quality of the libraries were assessed on High Sensitivity DNA chips on an Agilent

Bioanalyzer. The libraries were pooled (11-12 samples per lane) and sequenced

(100bp, paired-end) in four lanes on an Illumina HiSeq2500 sequencer. Sequencing

was done at the Vincent J. Coates Genomics Sequencing Laboratory at UC Berkeley.

Data Processing

Raw reads were processed to remove adapter sequences and gently trimmed

(PHRED Q<5; Macmanes 2014) using Cutadapt (version 1.7.1; Martin 2011). TopHat

(version 2.0.13; Trapnell *et al.* 2012) was used to align reads to the *D. simulans* (version

r2.02) and *D. sechellia* (version r1.3) genome assemblies (from the twelve species

project, downloaded from Flybase) and to the *D. mauritiana* MS17 assembly (Nolte *et*

al. 2013). Because the *D. mauritiana* line used for sequencing and the line used to

construct the genome assembly differed, variant sites from the lab line, called using

Genome Analysis Toolkit's (GATK) Haplotypecaller, were incorporated into the MS17

assembly using Pseudoref (http://yangjl.com/pseudoRef/; Xu *et al.* 2020). Additionally,

an updated annotation file for the MS17 assembly (Torres-Oliva *et al.* 2016) was used

during alignment and in subsequent processing steps. Annotation files for *D. simulans*

and *D. sechellia* were obtained from the same versions of the genome release of each

species. Read alignment, mismatches, edit distance, and gap length were all set to

three when using TopHat (version 2.0.13; Trapnell *et al.* 2012) to allow for a higher rate

of read alignment.

In order to differentiate reads derived from each parental species, variant sites

were called between the genomes of the species used in this analysis. RNA-seq reads

from parental species samples (from previous data from Atallah and Lott, 2018, GEO

accession GSE112858) were aligned to every other parental genome in each pairwise

comparison using TopHat (version 2.0.13; Trapnell *et al.* 2012). The BAM files from the

TopHat output were sorted and indexed using Samtools (version 1.2; Li *et al.* 2009).

Picard tools (version 2.7.1) and GATK tools (Van der Auwera *et al.* 2013) were then

used to identify variant sites by using the following programs:

AddorReplaceReadGroups, MarkDuplicates, ReorderSam, SplitNCigarReads, and

HaplotypeCaller. Additionally, indels were excluded and sites with single variants

selected using the SelectVariants tool. The variants were ordered using a Pysam script

(Python version 2.7.10) and read assignments to parental genomes were subsequently

organized with custom R scripts using the variant sites that exist between the parental

genomes (R version 3.4.1; R Core Team 2017) (Files S1, S2, S3 and S4). This pipeline

was also used to update the *D. mauritiana* MS17 assembly (Nolte *et al.* 2013) with

variants present in the line we used in the lab (Dmau/[w1];14021-0241.60).

Normalization to the upper quantile was performed across all samples in each set of

pairwise species comparisons. This was used to account for differences in the number

of reads for each sample as a result of sequencing.

Read counts represent the number of reads mapping to variant sites within a

gene. A cutoff of 5 or more reads mapped to any given gene was set to determine if a

gene was expressed. Genes with read counts <5 in both species in any pairwise

comparison were not considered to be expressed in either species so were removed

from the analysis. This cutoff was tested empirically and was set to exclude genes with

low count numbers that had a higher frequency of mapping in a biased manner to both

parental genomes. Genes analyzed in this analysis were also limited to those with

annotated orthologs in both species in any pairwise comparison. An orthologs table

from Atallah and Lott, 2018, was updated using the annotations available on FlyBase

(v2017) and an updated set of annotations from Torres-Oliva, *et al*. 2016. This revised

orthologs table (Table S2) was used to compare genes between each species and in

each direction of mapping.

Mapping bias due to differing genome quality may occur when using two different

reference genomes. In our study, mapping bias can result when a higher proportion of

reads from one allele map to the genome of a species used in the cross than reads from

the other allele. In order to alleviate mapping bias that may occur when mapping the

parental and hybrid samples to each parental reference genome, Poisson Gamma (PG)

models (León-Novelo *et al.* 2014) were employed to calculate mapping bias for every

set of mappings, in each pairwise comparison of species. We compared the 95%

confidence intervals (CIs) from PG models (with fixed bias parameter, q = 0.5) in each

direction of mapping. We set a slightly conservative standard for classifying allelic

imbalance where genes with CIs below 0.49 or above 0.51 were called differentially

expressed. Genes with CIs close to 0.50 did not appear differentially expressed when

looking at the count data, so we used a more conservative cutoff. Genes that appeared

differentially expressed in one direction of mapping (CIs fell outside of the range of 0.49

- 0.51 when comparing the expression levels of parental alleles in each replicate) but

not in the other direction of mapping were removed from the analysis, as this was

determined to be a result of mapping bias between the two genomes. We also removed

genes that had disparate confidence intervals in the two mapping directions (i.e. one

mapping direction yielded a CI that fell above the 0.49 - 0.51 range and the other

direction of mapping yielded a CI that fell below the 0.49 - 0.51 range).

We found that between 9.6% and 10.9% of genes expressed at stage 2 (with a

count >5) mapped in a biased way to parental genomes when compared to the total

number of orthologous genes between any pair of species. Each between-species

comparison has a different number of orthologous genes so the proportion of biased

genes varies based on the pair of species compared in a cross. In contrast to the

maternal stage, we found that between 5.0% and 6.2% of genes expressed at stage 5

mapped in a biased way to parental genomes when compared to the total number of

orthologous genes between a pair of species. Overall, when looking at the total

proportion of biased genes, not just those that were called "expressed" in our analysis,

we found that between 24.8% and 28.3% of genes at stage 2 and between 20.0% and

21.2% of genes at stage 5 mapped in a biased manner when compared to the total

number of orthologous genes in any comparison between species in a cross. All the

genes that mapped in a biased manner were removed from our analysis. Genes that

were not biased in their mapping and had a read count of >5 reads were retained for

analysis.

Genes used for stage 5 analysis

To focus on the gene regulation from the zygotic genome after ZGA, we removed

genes with high levels of maternal transcript deposition from our analysis. We limited

the pool of genes analyzed to those that are mostly zygotic because roughly half of

maternal transcripts are not entirely degraded by stage 5 (although studies are

somewhat variable in the percent reported; Tadros *et al.* 2007; De Renzis *et al.* 2007;

Thomsen *et al.* 2010; Lott *et al.* 2014) and we wanted to examine only those genes that

have a larger contribution to expression from the zygotic genome. For this analysis, we

included genes with "zygotic-only" expression (those that are not maternally deposited)

and genes that are "mostly zygotic" (those with 8-fold higher expression at stage 5

relative to stage 2, a $\log_2$ difference greater than three). We added a count of 1 to the

transcript count for genes at the maternal and zygotic stages when calculating the $\log_2$

difference to avoid errors associated with taking the $\log_2$ of zero. We tested several

cutoffs but chose the 8-fold threshold because at this conservative cutoff, most genes

with high maternal transcript deposition are removed from the analysis. Additionally, for

this analysis we used confidence intervals and averages generated from only female

samples for genes on the X chromosome because dosage compensation is not

complete at stage 5 (Lott *et al.* 2011).

Correlation analysis and PCA

We performed correlation analysis (Figure 2, Table S3) between single embryos across replicates, stages, and genotypes in R (R Core Team 2017) using the Spearman option within the *corr* function. Principal component analysis (PCA) was also performed in R using the *prcomp* function (Figure S1).

***Cis/Trans* analysis**

To identify evolved regulatory changes between species, we first determined which genes showed differential expression between alleles using the 95% CIs from PG models (León-Novelo *et al.* 2014) that were also used to interpret mapping bias. We compared expression levels of alleles for individual genes within the hybrids and also between parental samples using this model. Genes with CIs that fell outside of the 49% - 51% range were defined as differentially expressed, either within the hybrid or between parental samples, while those falling within the 49% - 51% range were identified as having the same level of expression. Genes were then categorized using *cis*, *trans*, *cis + trans*, *cis x trans*, *compensatory*, and *conserved* categories as described in Landry, *et al*. 2005; McManus, *et al*. 2010; and Coolon, *et al*. 2014 (Figures 3, S2 and S3). We assigned the following categories for regulatory change based on the CIs generated from PG models for individual genes (see Figure 4 for individual examples):

cis: Genes categorized as having changes in *cis* are those that are differentially expressed (CIs do not overlap 49% - 51%) between the parental species and in

the hybrids. (CIs for parental species and hybrids overlap each other for changes

purely in *cis*. To determine this, we used the CIs generated from mapping to the

D. simulans genome for *D. simulans/D. mauritiana* and *D. simulans/ D. sechellia*

comparisons and CIs generated from mapping to the *D. sechellia* genome for *D.*

sechellia/D. mauritiana comparisons.)

trans: Genes that are differentially expressed between the parental species (CI

does not overlap 49% - 51%) but are not differentially expressed in the hybrid (CI

overlaps 49% - 51%).

cis + *trans*: Genes that are differentially expressed in the hybrids and between

the parental species (CI does not overlap 0.49% - 0.51%) and the CI is in the

same direction for both the parents and the hybrid (i.e. both are greater than 51%

but the CIs for the parents and hybrid do not overlap. For this comparison, we

used the CIs generated from mapping to the *D. simulans* genome for *D.*

simulans/D. mauritiana and *D. simulans/ D. sechellia* comparisons and CIs

generated from mapping to the *D. sechellia* genome for *D. sechellia/D.*

mauritiana comparisons.)

cis x *trans*: Genes that are differentially expressed in the hybrids and between

the parental species (CI does not overlap 49% - 51%) and the CI is in opposite

directions for the parents and the hybrid (i.e. one is greater than 51%, the other is

less than 49%)

compensatory: Genes that are not differentially expressed between the parental

species (CI overlaps 49% - 51%) but are differentially expressed in the hybrids

(CI does not overlap 49% - 51%).

conserved: Genes are not differentially expressed between the parental species

or within the hybrids (CIs overlap 49% - 51%).

Inheritance Patterns

Previous studies from Gibson, *et al*. 2004 and McManus, *et al*. 2010 identified

and outlined ways to classify inheritance patterns of transcript abundance in hybrids in

relation to parental samples. We used these methods in our study to compare the

averages of total expression levels in the hybrids relative to those of parental samples.

Gene expression was considered conserved if the expression level between parental

samples and the total expression in the hybrid (sum of the expression of the two

species-specific alleles in the hybrid) were within 1.25-fold of one another, a $\log_2$-fold

change of 0.32. Overdominant genes were expressed at least 1.25-fold more in the

hybrid than in either parent while underdominant genes were expressed at least 1.25-

fold lower in the hybrid than in either parent. Genes that were expressed at an

intermediate level in the hybrid in comparison to the parental species samples involved

in the cross were defined as additive. Dominance was determined when the hybrid had

expression within 1.25-fold of one of the parental species such that total transcript levels

in the hybrid was more similar to transcript levels in one parental species than in the

other parental species.

Candidate transcription factor identification

We took a computational approach to identify potential transcription factors that

may change in *trans* regulation between the species in our analysis. We used motif

enrichment programs to find potential binding sites in the upstream regions of genes

changing in regulation in *D. sechellia* and *D. simulans*. We omitted *D. mauritiana* from

this analysis because the *D. mauritiana* genome is not as well annotated as the

genomes for *D. simulans* and *D. sechellia*. We used the Differential Enrichment mode in

MEME (Bailey and Elkan 1994) as well as the findMotifs.pl script in HOMER (Heinz *et*

al. 2010) to identify overrepresented motifs in the regions 500bp upstream of the

annotated starting location for genes changing in regulation or with conserved

regulation between species in every set of comparisons at stage 2. We utilized a 500bp

region as this was empirically determined to give the highest enrichment of signal for

motifs in target genes relative to background (see Omura and Lott 2020 for more

information). In MEME, we used options to find motifs with any number of repetitions

and a motif width of 8-12. We used default options for HOMER and supplied a

background FASTA file for enrichment analysis. The background lists supplied were

500bp upstream regions from all annotated genes in the species except for those that

were in the target set (either those genes with conserved or changing regulation in any

set of comparisons). The 500bp regions were extracted from FASTA files (versions

were the same as ones used for mapping) for each species using BEDTools (Quinlan

and Hall 2010). Significantly overrepresented motifs in the target lists relative to the

background supplied were then compared against databases of known transcription

factor binding sites using Tomtom (MEME suite) and HOMER. All enriched motifs that

appeared in both HOMER and MEME analyses are included in Table S4. All potential

targets of discovered motifs with significant E-values (MEME) or high Match Rank

scores in HOMER (>0.8) are also listed in Table S4 (see Figure S4 for transcript levels

of differentially maternally deposited targets in embryos of parental species).

Gene Ontology

Gene ontology (GO) analysis was done with the statistical overrepresentation

test in PANTHER (Mi *et al.* 2019) using the default settings. We looked at the GO

complete annotations for biological processes and molecular function but did not find

any significant terms represented in the cellular component categories. For this

analysis, we set a cutoff of Bonferroni adjusted p-value < 0.05. We searched for

enrichment of GO categories amongst genes that change in *trans* in each cross,

compared to the background of genes that are expressed (having a count >5) in each

cross. We used REVIGO (Supek *et al.* 2011) to reduce the number of redundant GO

categories and used the small (0.5) level of similarity as a cutoff for redundant GO

terms. GO categories shared between two or more crosses at stage 5 are represented

in Figure 5 and GO categories unique to a cross are shown in Figure S5. All enriched

categories are listed in Table S5.

Data Availability

All sequencing data and processed data files from this study are available at NCBI/GEO

at accession number: GSE136646.

439   **RESULTS**

In order to compare the regulatory basis of evolved changes in gene expression

at different stages of early embryogenesis, one stage where all the transcripts are

maternally provided and the other after zygotic genome activation (ZGA), we performed

a series of crosses between closely related species followed by RNA-seq on resulting

embryos (Figure 1). We used the sister species *D. simulans*, *D. sechellia*, and *D.*

mauritiana, all of which may be crossed reciprocally (with the exception of *D. sechellia*

females to *D. simulans* males; Lachaise *et al.* 1986). As transcripts in the two early

embryonic stages of interest are produced by different genomes, that of the mother and

that of the zygote, we performed crosses to produce a hybrid genome in the appropriate

generation (that of the mother or that of the zygote; Figure 1). To investigate regulatory

changes in zygotic gene expression, the three species were crossed pairwise (with the

noted exception), to produce F1 hybrid embryos, which were collected at a stage after

zygotic genome activation (end of blastoderm stage or the end of stage 5, Bownes'

stages; Bownes 1975; Campos-Ortega and Hartenstein 2013). While the zygotic

genome is fully activated at this developmental stage, maternal transcripts are not yet

entirely degraded so we limited our analysis to those genes that are expressed at a

much higher level after ZGA than before the zygotic genome is activated (see Methods).

To discover the regulatory basis of changes in maternal transcript deposition, the F1

females were crossed to males of the same species as the maternal species in the

initial cross. Resulting embryos were collected at stage 2 (Bownes' stages; Bownes

1975; Campos-Ortega and Hartenstein 2013), when all the transcripts in the egg are

maternal in origin (Figure 1). Three replicate samples were obtained for each cross at

stage 2, and since stage 5 features incomplete X chromosomal dosage compensation

(Lott *et al.* 2011), 6 replicates were obtained for each cross at late stage 5 (3 female

and 3 male embryos with one noted exception, see Methods). mRNA-sequencing

libraries were constructed from each embryo sample using poly(A) selection. Libraries

were sequenced paired-end, 100bp, on an Illumina HiSeq2500.

Reproducibility of Single Embryo RNA Sequencing Data

Previous studies have shown that single-embryo RNA-seq data is highly

reproducible, despite replicate samples representing both biological and technical

replicates (Lott *et al.* 2011, 2014; Paris *et al.* 2015; Atallah and Lott 2018). Our current

study extends this to include replicates of F1 crosses and subsequent backcrosses

between closely related species, which are as reproducible as the within-species

replicates. Spearman's rank correlation coefficients are high between replicate samples

of the same species or cross at the same developmental stage (Figure 2, A,B,D,E,

Table S3). For example, when comparing the RNA-seq data from stage 2 samples of

the (*mau* x *sim*) x *mau* and (*sim* x *mau*) x *sim* hybrid crosses, correlation coefficients

range from 0.965 to 0.995 (Table S3). Stage 5 hybrids from the *mau* x *sim* cross have

equally high correlation coefficients, ranging from 0.980 to 0.996 (Table S3). Similarly,

correlation coefficients for the RNA-seq data from *D. simulans* stage 5 embryos, when

compared with other *D. simulans* stage 5 embryos, range from 0.985 to 0.990. The high

correlation coefficients between replicates may be due, in part, to the removal of genes

with differential mapping to either parental genome and those genes with very low

transcript abundances (see Methods) from this analysis.

Transcript levels for embryos of the same stage but different genotypes (parental

lines and hybrids) are highly similar, as indicated by their Spearman's rank correlation

coefficients (Table S3), with one notable exception. When we compare RNA-seq

profiles from stage 5 hybrids to stage 5 embryos of the paternal species in the cross, we

see more divergent patterns of gene expression than when we compare stage 5 hybrids

to stage 5 embryos of the maternal species in the cross. This is due to the fact that

many maternal transcripts are still present at the zygotic stage, and thus the hybrid

zygotic embryo has many remaining maternal transcripts from the maternal species, not

the paternal species, in the cross. For example, comparisons between *D. simulans*

stage 5 embryos and stage 5 embryos of the *sim* x *mau* cross, where *D. simulans* is the

maternal species in the cross, yield high correlation coefficients, ranging from 0.955 to

0.972. In contrast, correlation coefficients are much lower when comparing *sim* x *mau*

stage 5 hybrid embryos to stage 5 embryos of the paternal species in the cross, *D.*

mauritiana, ranging from 0.863 to 0.887. In this particular comparison, the lower

correlation coefficients are likely due to having *D. simulans* as the maternal species in

the hybrid cross for the *sim x mau* embryos. Remaining maternal transcripts are from

the *D. simulans* alleles and likely explain why these hybrid embryos correlate more

highly with *D. simulans* stage 5 embryos.

In contrast to highly correlated samples within a stage, comparing the transcript

abundance of different stages yields strikingly lower correlation coefficients (Figure 2C

and F, Table S3), emphasizing the turnover of transcripts between these stages. For

example, when comparing stage 2 hybrids from crosses with *D. mauritiana* and *D.*

simulans to stage 5 hybrids from the same cross, correlation coefficients range from

0.483 to 0.573 (Table S3). The correlation coefficients are lower when comparing

transcript abundances of embryos of different stages than when comparing embryos

within a stage, indicating that the pool of transcripts present at the maternal stage is

different from that at the zygotic stage of development.

The above finding is reinforced by principal component analysis (PCA; Figure

S1) for RNA expression profiles of the samples in each set of pairwise comparisons

between species. We found that the first principal component corresponds to

developmental stage and explains between 80.65% and 81.86% of the variance in the

three sets of comparisons. The second principal component of this PCA accounts for

between 6.94-8.44% of the variance in the three sets of pairwise comparisons between

species and corresponds to genotype. This indicates that there is a more substantial

difference between pools of transcripts at the maternal and the zygotic stage of

development than there is between the pools of transcripts present in embryos of

different genotypes (parental species and hybrids) at the same developmental stage.

Regulatory changes at the maternal stage of development

Changes in gene expression can occur at many levels of regulation:

transcriptional, post-transcriptional, translational or post-translational. Here, we address

whether at the transcriptional level, changes in gene expression between species

occurred due to changes in *cis* or in *trans* and whether the pattern of regulatory

changes differs based on developmental stage. Changes in *cis* regulation can occur

through changes in the DNA of regulatory regions proximal to the gene that they

regulate. These types of regulatory changes have an allele-specific effect on gene

expression. In contrast, changes in *trans* regulation typically occur via changes in

factors that bind to the DNA, such as transcription factor proteins. Changes in *trans*

regulation affect the expression of both alleles.

In order to determine regulatory changes in *cis* and in *trans* that lead to

differences in maternal transcript deposition between *D. simulans*, *D. sechellia*, and *D.*

mauritiana, we used Poisson Gamma (PG) models (León-Novelo *et al.* 2014). This

allowed us to determine mapping bias as well as differential expression between alleles,

between parental lines and within hybrid embryos (see Methods). We identified

differential maternal transcript deposition between the parental lines as well as between

species-specific alleles in the stage 2 embryos produced by hybrid mothers. We then

compared the two sets of analyses to determine the proportion of *cis* and *trans*

regulatory changes underlying differential maternal transcript deposition between

species. We used the logic of Landry, *et al.* 2005 to classify genes as having changed in

cis or in *trans* regulation, by comparing confidence intervals of the bias parameters

generated through the PG models (see Methods).

We found that most regulatory changes underlying differentially maternally

deposited transcripts occurred in *trans* between each pair of species examined (Figure

3A and C, Figure S2), where a change in a transcription factor or other *trans*-acting

regulatory element affects both alleles equally (shown in Figure 4A and B). In all pairs of

comparisons between species, the proportion of *trans* changes was higher than any

other category of changes. Comparisons between *D. simulans* and *D. mauritiana* had

the highest percentage of *trans*-only regulatory changes (between 49.4% and 50.8%)

while comparisons between *D. simulans* and *D. sechellia* had a lower percentage of

regulatory changes solely in *trans* (32.9%). The second highest proportion (between

15.0% and 26.7%) of regulatory changes between species at the maternal stage

occurred only in *cis* regulation. Slightly fewer regulatory changes occurred due to a

combination of *cis* and *trans* acting factors (between 13.4% and 15.8% in all

comparisons). Most genes that change in *cis* and in *trans* regulation are assigned to the

cis + *trans* category, which indicates that the allele with higher expression in the

parental lines is also preserved as the allele with higher expression in the hybrid (the

changes in *cis* and in *trans* affect gene expression in the same direction; Landry *et al.*

2005; McManus *et al.* 2010; Coolon *et al.* 2014). We found a smaller proportion of

genes changed in regulation through *cis* x *trans* interactions, where changes in *cis* and

in *trans* have opposing effects on gene expression and the allele with the lower level of

expression in the hybrid is from the parental line with the higher level of expression

(Landry *et al.* 2005; McManus *et al.* 2010; Coolon *et al.* 2014). We also found a

percentage of genes with conserved levels of maternal transcript deposition between

species, between 16.4% and 25.1% in all crosses. *D. simulans* and *D. sechellia* have

the highest percentage of conserved genes while *D. simulans* and *D. mauritiana* have

the lowest percentage of conserved genes. We also found a small proportion of genes,

between 4.2% and 4.7% in all comparisons, that have evolved compensatory

mechanisms of regulation, where the genes are not differentially expressed between the

parental samples but are differentially expressed in hybrids. This implies that while

transcript levels are the same between species, regulatory changes have occurred,

which then become visible in the environment of the hybrid.

Genes with expression differences due to *trans* regulatory changes include

regulators with critical functions in important processes governed by maternal gene

products, such as *Cdk1*, a cell-cycle regulator necessary for the rapid cleavage cycles

in early development (Farrell and O'Farrell 2014). Examples of individual genes with

changes attributed to *trans* regulatory differences at the maternal stage are represented

in Figure 4A and B.

Binding sites for chromatin modifiers are enriched in the regulatory regions of

maternally deposited genes

As *trans* regulatory changes can affect numerous genetic loci, we asked whether

there are *trans* regulatory factors that may affect the differential deposition of a number

of maternal transcripts between the species studied. For this, we identified binding sites

in the predicted *cis*-regulatory regions of all differentially expressed genes and

compared them to identified binding sites in the *cis*-regulatory regions of genes with

conserved expression between species at the maternal stage. In the pool of genes with

altered expression between species, we included not only those with differences in

trans regulation, but also those with differences in other regulatory categories (*trans*,

cis, *cis* + *trans*, *cis* x *trans*, and compensatory). This is because genes with changes in

cis regulation may have had changes that affect the binding of the same *trans*

regulators. For genes with differential expression, and separately for genes with

conserved expression between species, we took a computational approach. We used

both HOMER and MEME (Bailey and Elkan 1994; Heinz *et al.* 2010), to search for

overrepresented motifs within 500 base pairs upstream of transcription start sites, as

compared to the rest of the genome (see Methods). We used the upstream regions of

genes in *D. simulans* and *D. sechellia* because the *D. mauritiana* genome is not as well

annotated, as compared to the other two species in this study.

Interestingly, we found that the *cis*-regulatory regions from both genes with

conserved and genes with differential transcript levels between species are enriched in

motifs associated with insulator binding (Table S4). Specifically, we found that the

Dref/BEAF-32 binding site (BEAF-32 and Dref bind overlapping DNA sequences; Hart

et al. 1999) is the most significantly enriched (Table S4). These factors are annotated

as insulators (Matzat and Lei 2014; Ali *et al.* 2016) and known to be associated with

topologically associated domains (TADs) (Liang *et al.* 2014; Ramírez *et al.* 2018). The

binding site for M1BP also appeared significantly enriched in both sets of genes that

change in regulation and in ones that are conserved in regulation across species (Table

S4). M1BP is involved in transcriptional regulation and RNA polymerase II pausing at

the promoter of genes (Li and Gilmour 2013), which may also be associated with

regulating chromatin state (Ramírez *et al.* 2018). Our findings are consistent with

previous studies that identified the enrichment of binding sites for M1BP, BEAF-32 and

Dref in the promoter regions of genes that are maternally deposited (Chen *et al.* 2013;

Omura and Lott 2020). These binding sites have also been associated with

housekeeping genes (Zabidi *et al.* 2015), and the pool of maternal transcripts is

enriched with housekeeping genes (Liu *et al.* 2014). However, maternal genes that are

not housekeeping genes are even more highly enriched for these binding sites (Omura

and Lott 2020), thus the involvement of insulators or other chromatin regulators in

maternal regulation is unlikely due only to the inclusion of housekeeping genes among

maternal genes. Transcript abundance data from our study indicates that *Dref*, *BEAF-*

32, and *M1BP* are differentially maternally deposited in several between-species

comparisons (Figure S4), although in certain crosses, hybrid reads mapped in a biased

way to *Dref*, *BEAF-32* and *M1BP*, and thus they were excluded from our regulatory

analysis. As the motifs for these *trans*-acting factors are enriched in the upstream

regions of all maternal genes, relative to upstream regions of all annotated genes, they

are likely important regulators of transcription during oogenesis, and therefore also likely

targets of regulatory evolution between species.

Evolution of regulation for zygotically expressed genes

To determine the regulatory basis of changes in zygotic transcript abundance

between species, we compared expression levels in late stage 5 parental species

samples to late stage 5 hybrid samples and used PG models to identify *cis* and *trans*

regulatory changes, similar to our maternal analysis (see Methods). We limited our

analysis at the zygotic stage to those genes that are mostly-zygotic: zygotically

expressed but not maternally deposited (zygotic-only) or expressed at the zygotic stage

at an 8-fold higher level when compared to the maternal stage (we will refer to these as

mostly zygotic genes, see Methods).

While we found that most gene expression changes at the maternal stage of

development are due to changes in *trans* regulation between the three sister species,

we see strikingly different patterns of regulatory changes after ZGA. At the zygotic

stage, differences in gene expression between the three species examined occur

mostly due to regulatory changes in both *cis* and *trans*, either by *cis* + *trans* or *cis* x

trans interactions (Figure 3B and D, Figure S3). Changes in both *cis* and *trans*

regulatory elements (either *cis* + *trans* or *cis* x *trans* interactions) account for expression

differences in 39% to 47% of zygotic genes at stage 5 in our between-species

comparisons. We also see a higher proportion of these interactions occurring in a *cis* +

trans pattern (between 29% and 35% of all genes) as opposed to a *cis* x *trans* pattern

(between 9% and 12% of all genes) of regulatory interactions. In contrast, *cis*-only and

trans-only changes account for a smaller number of differences in gene expression

levels at this stage in development. In all comparisons, we found between 15% and

21% of genes changing only in *trans* regulation. There are between 16% and 30% of

genes that change only in *cis* regulation between each pair of species compared at this

stage in development. Compared to the maternal stage, we found a larger proportion of

genes with compensatory changes (between 7% and 10% of all genes) in gene

regulation and a smaller proportion of genes that are conserved (between 6% and 8%

of all genes) between each pair of species comparisons. The smaller proportion of

genes with conserved transcript levels at the zygotic stage compared to the maternal

stage is consistent with earlier findings showing maternal transcripts to be more highly

conserved between species than zygotic transcripts (Atallah and Lott 2018). Examples

of evolved changes include regulators critical to important early zygotic processes, such

as gap gene *Kruppel* and pair-rule gene *sloppy paired 1*, which are required for

segmentation along the anterior-posterior axis (Nüsslein-Volhard and Wieschaus 1980;

Grossniklaus *et al.* 1992) (Figure 4C and D).

Transcriptional regulation at the maternal stage may be broadly determined by

regulation at the level of chromatin, as evidenced in this work and by another study

(Omura and Lott 2020). In contrast, regulation at the zygotic stage can be gene or

pathway specific and involve transcription only in a spatially localized subset of cells

(Jäckle *et al.* 1986; Johnston and Nüsslein-Volhard 1992). As such, if a *trans* regulator

changed at the zygotic stage, it may affect genes involved in a specific developmental

process. For these reasons, we wanted to ask if genes whose zygotic expression

differed between species due to changes solely in *trans* regulation had a specific

molecular function or were part of a particular biological process. We used PANTHER

(Mi *et al.* 2019) to perform gene ontology (GO) analysis on genes changing only in *trans*

regulation in each pairwise comparison of species at stage 5 (see Methods). Identifying

GO categories over multiple crosses identifies the types of genes that evolve changes

repeatedly over evolution. Shared categories were broad, and included those related to

DNA binding, positive regulation of transcription by RNA polymerase II, cell fate

determination, and several developmental categories (Figure 5). The range in GO

categories represented, while broadly important at this developmental timepoint,

demonstrate how genes changing in *trans* are distributed across developmental

processes. As may be expected for zygotic genes at this stage in development, this

finding suggests that changes in *trans* regulators of zygotic genes can affect a broad

range of molecular and developmental processes. We also investigated biological

process categories unique to each specific cross, these are represented in Figure S5

and Table S5.

Modes of Inheritance in Hybrids

Misexpression in hybrid offspring has been used to examine regulatory

incompatibilities that may contribute to speciation (Michalak 2003; Ortíz-Barrientos *et al.*

2006; Moehring *et al.* 2007; Mack *et al.* 2016). As the maternal and zygotic transcripts

examined during embryogenesis showed different patterns of gene expression

evolution, we also asked whether there were more hybrid incompatibilities present at

one stage than the other by looking at whether these two developmental stages showed

different levels of transcript misexpression. One way to identify misexpression in hybrids

is to compare the inheritance of transcript levels in hybrids to transcript levels in each

parental species. Here, we quantify the total transcript abundance for a gene by

summing the levels of both species-specific alleles in the hybrid and comparing this

level to the transcript abundance in both parental lines. We used methods developed by

Gibson, *et al*., 2004 to define modes of inheritance in our hybrids; as in previous studies

(Gibson *et al.* 2004; Landry *et al.* 2005; McManus *et al.* 2010; Coolon *et al.* 2014), we

used a conservative fold change of 1.25 ($\log_2$-fold change of 0.32) to define those

genes that do not change in the total transcript abundance between genotypes

(conserved). Genes with transcript levels that are higher (overdominant) or lower

(underdominant) in the hybrid relative to either parental species are categorized as

misexpressed. The total transcript abundance in the hybrids can also be more similar to

one parent versus the other. Here, we categorize the parental line with expression most

similar to the hybrid as the dominant parent. Expression in the hybrid can also have a

level intermediate to both parental species (additive).

Both the maternal and zygotic stages show a significant proportion of genes

where one species' allele is dominant (Figure 6, Table S6). In both stages, the species

that is dominant in each set of crosses is consistent. A higher proportion of genes that

are dominant have *D. simulans*-like expression (in any cross involving *D. simulans*) in

comparison to the proportion that have expression more like the other parental line in

the cross (see Table S6 for percentages). We found that *D. mauritiana* has the least

dominance in any cross involving this species. Taken together, our findings indicate that

D. simulans has the most dominant effect on gene expression at both developmental

stages, while *D. mauritiana* has the least dominant effect, with dominance in *D.*

sechellia falling between the other two species. While there has been previous work

proposing relationships between the proportion of dominance and the physiology of

unique species (McManus *et al.* 2010), it is difficult to determine any known factors

between these three species that would predict this pattern of relative dominance

(species range, effective population size, egg size/maternal investment). It is, however,

interesting that while the proportion of changes in *cis* and *trans* vary considerably

between stages that both stages have dominance among the largest categories of

modes of inheritance, and that the relative patterns of which species are dominant is

conserved.

Strikingly, while many genes show conservation of expression levels between

parental species and in the hybrids at both developmental stages, we found a much

higher percentage of conserved transcript levels between parents and hybrids for genes

that are maternally deposited (Figure 6, Table S6). We found a high proportion of genes

with conserved transcript levels at stage 2 in all crosses, between 15.4% and 31.4% of

all genes. In contrast, in stage 5 crosses we found conserved transcript abundance in

between 4% and 8% of all genes that are either zygotic-only or are mostly zygotic (see

Methods for definitions). While there is a large difference in the percentage of

conserved genes between the two stages, our stage 5 analysis is limited to those genes

with much higher expression at the zygotic stage in comparison to the maternal stage of

development. There may be more genes that are mostly zygotic or zygotic-only that are

misregulated at this stage in development relative to all of the genes that are expressed

at stage 5.

In contrast to maternal expression patterns, we found more genes that have an

additive mode of inheritance or that are misexpressed in the hybrids for zygotic genes

(Figure 6, Table S6). Previous studies indicate that additive inheritance is associated

with *cis* regulatory divergence (Lemos *et al.* 2008; McManus *et al.* 2010). This is

consistent with our findings that a larger proportion of genes at the zygotic stage have

expression divergence due, in part or wholly, to *cis* regulatory changes and that more

zygotic genes show an additive pattern of inheritance. Higher levels of misexpression at

the zygotic stage, taken together with lower conservation of transcript levels at the

zygotic stage, suggests that zygotic genes may contribute more to genetic

incompatibilities than maternal genes.

754   **DISCUSSION**

In this study, we asked whether evolution of gene regulation differs at different

developmental stages. We found striking differences in the proportions of *cis* and *trans*

regulatory changes between the stage of embryogenesis where all transcripts are

maternally derived, and a stage just a few hours later after the zygotic genome has

been activated. Between the species examined, we uncovered an overwhelming

number of *trans* regulatory changes resulting in differential maternal transcript levels,

whereas a complex mix of *cis*, *trans*, and the combination of the two were responsible

for changes in zygotic transcription of mostly zygotic genes (see Methods). Here, we

propose that the differences in the patterns of gene regulatory evolution between the

stages we examined may be due to fundamental differences in the biological context

and regulatory architecture producing the transcriptomes present at these stages.

Maternal transcripts are produced by support cells called nurse cells during

oogenesis and are either transported by microtubule-dependent mechanisms or

dumped into the oocyte along with the cytoplasmic contents of the nurse cells upon

apoptosis (Kugler and Lasko 2009). Many aspects of maternal provisioning have been

well-studied in *D. melanogaster*, including transport of transcripts into the oocyte

(Mische *et al.* 2007), localization of transcripts within the oocyte (Theurkauf and

Hazelrigg 1998), post-transcriptional regulation of maternal gene products (Salles *et al.*

1994), and subsequent degradation of maternal transcripts (Tadros *et al.* 2007; Bushati

et al. 2008; Laver *et al.* 2015). Surprisingly, how transcription is regulated in the nurse

cells is not well understood. Nurse cells are polyploid, and are able to rapidly transcribe

a large quantity of RNA that represents a large proportion of the genome (Tadros *et al.*

2007; De Renzis *et al.* 2007; Thomsen *et al.* 2010; Lott *et al.* 2011; Vastenhouw *et al.*

2019) to provide the oocyte with the large stock of transcripts needed. The oocyte itself

is thought to be largely transcriptionally silent (Navarro-Costa *et al.* 2016). What we

found here, and what was also found in another study investigating binding motifs in

maternal factors across the *Drosophila* genus using computational methods (Omura

and Lott 2020), is that maternal transcription is associated with *trans* factors annotated

to be insulators and that interact with topologically associated domains (TADs). This

provides evidence that maternal transcription may be controlled broadly at the level of

chromatin state. In this context, we predict that changes in only a few *trans* factors can

be responsible for the bulk of the between-species changes in maternal transcription.

Thus, changes in the levels of *trans* regulators at this stage may easily be responsible

for changes in transcription level for a number of genes.

In contrast to the large proportion of regulatory changes in *trans* at the maternal

stage, differences in zygotic gene transcription for genes that are mostly zygotic (see

Methods) between these species is predominantly explained by a combination of

changes in *cis*, *trans*, *cis + trans*, and *cis* x *trans*. Zygotic gene transcription for genes

without a maternal complement is fundamentally different than maternal gene

transcription. Unlike the bulk transcription that takes place in the nurse cells, zygotic

gene transcription is precisely regulated at the spatial and temporal level across the

embryo with enhancer regions playing a large role in where, when and how much genes

are expressed (Haines and Eisen 2018). Due to these fundamental differences in gene

regulation, the embryo at the zygotic stage may be more sensitive to changes in gene

expression than at the maternal stage. Specifically, changes in *trans* regulation, which

can affect the expression of many genes, may be detrimental to the developing

organism at this stage. In contrast, changes in *cis* regulation are gene-specific and may

only affect gene expression in a subset of the embryo, which might be a more suitable

way of fine-tuning zygotic expression. We propose that fundamental differences in the

regulatory landscape and developmental purpose of the maternal versus the zygotic

stage likely explain why the evolution of gene expression occurs through different

mechanisms for transcripts that are maternally deposited and genes that are primarily

zygotic.

While this study was directed at understanding the regulatory basis of evolution

in gene expression at the maternal and zygotic stages of embryogenesis, it also

provides insight into the relative conservation of gene expression, both between species

and between parent and hybrid offspring. Here, in both the analysis of regulatory

changes and the analysis of modes of inheritance, we found more genes with

conserved transcript levels among those that are maternally deposited relative to those

that are zygotically transcribed. This is in agreement with previous studies that identified

high conservation of maternally deposited transcripts relative to those transcribed

zygotically between species (Heyn *et al.* 2014; Atallah and Lott 2018) and indicates that

the maternal stage is highly conserved. We observe lower conservation of transcript

levels at the zygotic stage. A caveat our gene expression analysis at the zygotic stage

is that we had to remove genes that still have a large maternal component at this stage

(roughly 50% of total transcript pool at late stage 5 is maternally derived; Lott *et al.*

2014). Thus, our finding is best viewed as genes whose transcripts are primarily zygotic

at stage 5, have a higher rate of evolutionary change. Additionally, the large proportion

of genes with conserved transcript levels at the maternal stage may be unexpected

considering that there is substantial post-transcriptional regulation of maternally

deposited factors (Tadros *et al.* 2007; Rouget *et al.* 2010; Barckmann and Simonelig

2013), so it is not clear that a high degree of conservation at the transcript level should

be necessary to maintaining conservation at the protein level. Alternatively, if the

maternal genome is primarily regulated at the level of chromatin state, this may be a

mechanistic constraint on evolution at the level of gene expression. It may be

functionally difficult for a gene located in a region of open chromatin to be repressed, or

for a gene in a region of heterochromatin to gain expression. Thus, it may be easier to

evolve differences in expression over evolutionary time via post-transcriptional

mechanisms for maternal genes. Further study is needed to disentangle conservation at

the transcript and protein levels of maternal factors across species.

In addition to the differences in conservation between stages, we also found

differences in the patterns of inheritance of gene expression between species at the

maternal and zygotic stages of embryogenesis. The zygotic stage has a larger

proportion of additive differences, which some previous theory (Gibson *et al.* 2004) and

empirical studies (Lemos *et al.* 2008; McManus *et al.* 2010) have suggested may be

more likely to be changes in *cis* regulation. This would be consistent both with the

increased relative role of *cis* changes at the zygotic stage compared to the maternal

stage found here, as well as what is known about zygotic gene regulation more

generally (Mannervik 2014). In addition, a larger proportion of changes at the zygotic

stage fall into the broad category of misregulation (underdominant, overdominant),

which have been proposed to increase with divergence time (Coolon *et al.* 2014) and

may be a potential source of hybrid incompatibility between species (Michalak 2003;

Ortíz-Barrientos *et al.* 2006; Moehring *et al.* 2007; Mack *et al.* 2016).

In this study, we found that differences between species in levels of maternally

deposited transcripts and zygotically transcribed genes evolve via different patterns of

regulatory change. We found that maternal transcript abundance is more conserved but

when changes do occur, they occur more frequently through *trans* regulation in

comparison to zygotic complements. Regulatory organization, constraints, and

developmental processes that are specific to each developmental stage likely play a

large role in determining how gene regulation can evolve at these two embryonic

timepoints. Further study is needed to characterize the molecular basis of evolved

changes in transcript level on a single gene level, and more generally to determine what

is controlling the regulatory landscape at each stage in development.

861   **ACKNOWLEDGMENTS**

We would like to thank members of the Lott lab for comments on the manuscript, as well

as members of the UC Davis fly community for their contributions. We would also like to

thank Amanda Crofton and Anthony Le for their assistance with genotyping and RNA

extraction and Graham Coop for helpful discussions. This work was supported by the

National Institute of General Medical Sciences of the National Institutes of Health grant

R01GM111362 and the Floyd and Mary Schwall Fellowship.

REFERENCES

Ali T., R. Renkawitz, and M. Bartkuhn, 2016 Insulators and domains of gene

expression. Current Opinion in Genetics & Development 37: 17–26.

https://doi.org/10.1016/j.gde.2015.11.009

Ali-Murthy Z., S. E. Lott, M. B. Eisen, and T. B. Kornberg, 2013 An essential role for

zygotic expression in the pre-cellular Drosophila embryo. PLoS genetics 9:

e1003428. https://doi.org/10.1371/journal.pgen.1003428

Ashburner M., 1989 *Drosophila: A Laboratory Handbook*(C. spring harbor laboratory

press, Ed.).

Atallah J., and S. E. Lott, 2018 Evolution of maternal and zygotic mRNA complements

in the early Drosophila embryo, (K. A. Dyer, Ed.). PLoS genetics 14: e1007838.

https://doi.org/10.1371/journal.pgen.1007838

Bailey T. L., and C. Elkan, 1994 Fitting a mixture model by expectation maximization to

discover motifs in bipolymers

Barckmann B., and M. Simonelig, 2013 Control of maternal mRNA stability in germ cells

and early embryos. Biochimica et Biophysica Acta (BBA) - Gene Regulatory

Mechanisms 1829: 714–724. https://doi.org/10.1016/j.bbagrm.2012.12.011

Baroux C., D. Autran, C. S. Gillmor, D. Grimanelli, and U. Grossniklaus, 2008 The

Maternal to Zygotic Transition in Animals and Plants. Cold Spring Harbor Symposia

on Quantitative Biology 73: 89–100. https://doi.org/10.1101/sqb.2008.73.053

Bownes M., 1975 A photographic study of development in the living embryo of

Drosophila melanogaster. Development 33: 789–801.

Britten R. J., and E. H. Davidson, 1969 Gene Regulation for Higher Cells - a Theory.

Science (New York, N.Y.) 165: 349-+. https://doi.org/10.1126/science.165.3891.349

Buchberger, Reis, Lu, and Posnien, 2019 Cloudy with a Chance of Insights: Context

Dependent Gene Regulation and Implications for Evolutionary Studies. Genes 10:

492–21. https://doi.org/10.3390/genes10070492

Bushati N., A. Stark, J. Brennecke, and S. M. Cohen, 2008 Temporal reciprocity of

miRNAs and their targets during the maternal-to-zygotic transition in Drosophila.

Current Biology 18: 501–506. https://doi.org/10.1016/j.cub.2008.02.081

Campos-Ortega J. A., and V. Hartenstein, 2013 *The Embryonic Development of*

Drosophila melanogaster. Springer Science & Business Media.

Carroll S. B., 1995 Homeotic Genes and the Evolution of Arthropods and Chordates.

Nature 376: 479–485. https://doi.org/10.1038/376479a0

Chan Y. F., M. E. Marks, F. C. Jones, G. Villarreal, M. D. Shapiro, *et al.*, 2010 Adaptive

Evolution of Pelvic Reduction in Sticklebacks by Recurrent Deletion of a Pitx1

Enhancer. Science (New York, N.Y.) 327: 302–305.

https://doi.org/10.1126/science.1182213

Chen K., J. Johnston, W. Shao, S. Meier, C. Staber, *et al.*, 2013 A global change in

RNA polymerase II pausing during the Drosophila midblastula transition. Elife 2:

e00861. https://doi.org/10.7554/elife.00861

Clark A. G., M. B. Eisen, D. R. Smith, C. M. Bergman, B. Oliver, *et al.*, 2007 Evolution of

genes and genomes on the Drosophila phylogeny. Nature 450: 203–218.

https://doi.org/10.1038/nature06341

Coolon J. D., C. J. McManus, K. R. Stevenson, B. R. Graveley, and P. J. Wittkopp,

2014 Tempo and mode of regulatory evolution in Drosophila. Genome Research 24:

797–808. https://doi.org/10.1101/gr.163014.113

Crofton A. E., E. L. Cartwright, A. A. Feitzinger, and S. E. Lott, 2018 Effect of Larval

Nutrition on Maternal mRNA Contribution to the Drosophila Egg. G3:

Genes|Genomes|Genetics 8: 1933–1941. https://doi.org/10.1534/g3.118.200283

Davidson E. H., and I. S. Peter, 2015 *Genomic control process*.

De Renzis S. D., O. Elemento, S. Tavazoie, and E. F. Wieschaus, 2007 Unmasking

Activation of the Zygotic Genome Using Chromosomal Deletions in the Drosophila

Embryo. PLOS Biology 5: e117. https://doi.org/10.1371/journal.pbio.0050117

Eichhorn S. W., A. O. Subtelny, I. Kronja, and J. K. Elife, 2016 mRNA poly (A)-tail

changes specified by deadenylation broadly reshape translation in Drosophila

oocytes and early embryos. cdn.elifesciences.org.

https://doi.org/10.7554/elife.16955.001

Farrell J. A., and P. H. O'Farrell, 2014 From Egg to Gastrula: How the Cell Cycle Is

Remodeled During the Drosophila Mid-Blastula Transition. Annual Review of

Genetics, Vol 48 48: 269–294. https://doi.org/10.1146/annurev-genet-111212-

133531

Fischer B. E., E. Wasbrough, L. A. Meadows, O. Randlet, S. Dorus, *et al.*, 2012

Conserved properties of Drosophila and human spermatozoal mRNA repertoires.

Proc Royal Soc B Biological Sci 279: 2636–2644.

https://doi.org/10.1098/rspb.2012.0153

Gibson G., R. Riley-Berger, L. Harshman, A. Kopp, S. Vacha, *et al.*, 2004 Extensive

Sex-Specific Nonadditivity of Gene Expression in Drosophila melanogaster. Genetics

167: 1791–1799. https://doi.org/10.1534/genetics.104.026583

Glaser-Schmitt A., A. Zečić, and J. Parsch, 2018 Gene Regulatory Variation in

Drosophila melanogaster Renal Tissue. Genetics genetics.301073.2018-46.

https://doi.org/10.1534/genetics.118.301073

Graze R. M., L. M. McIntyre, B. J. Main, M. L. Wayne, and S. V. Nuzhdin, 2009

Regulatory Divergence in Drosophila melanogaster and D. simulans, a Genomewide

Analysis of Allele-Specific Expression. Genetics 183: 547–561.

https://doi.org/10.1534/genetics.109.105957

Grossniklaus U., R. K. Pearson, and W. J. Gehring, 1992 The Drosophila sloppy paired

locus encodes two proteins involved in segmentation that show homology to

mammalian transcription factors. Genes & Development 6: 1030–1051.

https://doi.org/10.1101/gad.6.6.1030

Haines J. E., and M. B. Eisen, 2018 Patterns of chromatin accessibility along the

anterior-posterior axis in the early Drosophila embryo, (J. M. Vaquerizas, Ed.). PLoS

genetics 14: e1007367-25. https://doi.org/10.1371/journal.pgen.1007367

Hart C. M., O. Cuvier, and U. K. Laemmli, 1999 Evidence for an antagonistic

relationship between the boundary element-associated factor BEAF and the

transcription factor DREF. Chromosoma 108: 375–383.

https://doi.org/10.1007/s004120050389

Heinz S., C. Benner, N. Spann, E. Bertolino, Y. C. Lin, *et al.*, 2010 Simple Combinations

of Lineage-Determining Transcription Factors Prime cis-Regulatory Elements

Required for Macrophage and B Cell Identities. Molecular cell 38: 576–589.

https://doi.org/10.1016/j.molcel.2010.05.004

Heyn P., M. Kircher, A. Dahl, J. Kelso, P. Tomancak, *et al.*, 2014 The Earliest

Transcribed Zygotic Genes Are Short, Newly Evolved, and Different across Species.

CellReports 6: 285–292. https://doi.org/10.1016/j.celrep.2013.12.030

Jäckle H., D. Tautz, R. Schuh, E. Seifert, and R. Lehmann, 1986 Cross-regulatory

interactions among the gap genes of Drosophila. Nature 324: 668–670.

https://doi.org/10.1038/324668a0

Johnston D. S., and C. Nüsslein-Volhard, 1992 The Origin of Pattern and Polarity in the

Drosophila Embryo. Cell 68: 201–219. https://doi.org/10.1016/0092-8674(92)90466-p

King M. C., and A. C. Wilson, 1975 Evolution at two levels in humans and chimpanzees.

Science (New York, N.Y.) 188: 107–116. https://doi.org/10.1126/science.1090005

Kugler J.-M., and P. Lasko, 2009 Localization, anchoring and translational control of

oskar, gurken, bicoid and nanos mRNA during Drosophila oogenesis. Fly 3: 15–28.

https://doi.org/10.4161/fly.3.1.7751

Kvon E. Z., O. K. Kamneva, U. S. Melo, I. Barozzi, M. Osterwalder, *et al.*, 2016

Progressive Loss of Function in a Limb Enhancer during Snake Evolution. Cell 167:

633-642.e11. https://doi.org/10.1016/j.cell.2016.09.028

Lachaise D., J. R. David, F. Lemeunier, L. Tsacas, and M. Ashburner, 1986 The

Reproductive Relationships of Drosophila sechellia with D. mauritiana, D. simulans,

and D. melanogaster from the Afrotropical Region. Evolution 40: 262–271.

https://doi.org/10.1111/j.1558-5646.1986.tb00468.x

983 Landry C. R., P. J. Wittkopp, C. H. Taubes, J. M. Ranz, A. G. Clark, *et al.*, 2005

984  Compensatory cis-trans evolution and the dysregulation of gene expression in

985  interspecific hybrids of Drosophila. Genetics 171: 1813–1822.

986  https://doi.org/10.1534/genetics.105.047449

987 Lasko P., 2012 mRNA Localization and Translational Control in Drosophila Oogenesis.

988  Cold Spring Harbor Perspectives in Biology 4: a012294–a012294.

989  https://doi.org/10.1101/cshperspect.a012294

990 Laver J. D., X. Li, D. Ray, K. B. Cook, N. A. Hahn, *et al.*, 2015 Brain tumor is a

991  sequence-specific RNA-binding protein that directs maternal mRNA clearance during

992  the Drosophila maternal-to-zygotic transition. Genome biology 16: 94.

993  https://doi.org/10.1186/s13059-015-0659-4

994 Lemos B., L. O. Araripe, P. Fontanillas, and D. L. Hartl, 2008 Dominance and the

995  evolutionary accumulation of cis- and trans-effects on gene expression. Proceedings

996  of the National Academy of Sciences 105: 14471–14476.

997  https://doi.org/10.1073/pnas.0805160105

998 León-Novelo L. G., L. M. McIntyre, J. M. Fear, and R. M. Graze, 2014 A flexible

999  Bayesian method for detecting allelic imbalance in RNA-seq data. BMC Genomics

1000  15: 920. https://doi.org/10.1186/1471-2164-15-920

1001 Li H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, *et al.*, 2009 The Sequence

1002  Alignment/Map format and SAMtools. Bioinformatics 25: 2078–2079.

1003  https://doi.org/10.1093/bioinformatics/btp352

Li J., and D. S. Gilmour, 2013 Distinct mechanisms of transcriptional pausing

orchestrated by GAGA factor and M1BP, a novel transcription factor. The EMBO

Journal 32: 1829–1841. https://doi.org/10.1038/emboj.2013.111

Liang J., L. Lacroix, A. Gamot, S. Cuddapah, S. Queille, *et al.*, 2014 Chromatin

Immunoprecipitation Indirect Peaks Highlight Long-Range Interactions of Insulator

Proteins and Pol II Pausing. Molecular cell 53: 672–681.

https://doi.org/10.1016/j.molcel.2013.12.029

Lim J., M. Lee, A. Son, H. Chang, and V. N. Kim, 2016 mTAIL-seq reveals dynamic

poly(A) tail regulation in oocyte-to-embryo development. Genes & Development 30:

1671–1682. https://doi.org/10.1101/gad.284802.116

Liu M. M., J. W. Davey, D. J. Jackson, M. L. Blaxter, and A. Davison, 2014 A conserved

set of maternal genes? Insights from a molluscan transcriptome. Int J Dev Biology

58: 501–511. https://doi.org/10.1387/ijdb.140121ad

Lott S. E., J. E. Villalta, G. P. Schroth, S. Luo, L. A. Tonkin, *et al.*, 2011 Noncanonical

Compensation of Zygotic X Transcription in Early Drosophila melanogaster

Development Revealed through Single-Embryo RNA-Seq. PLOS Biology 9:

e1000590. https://doi.org/10.1371/journal.pbio.1000590

Lott S. E., J. E. Villalta, Q. Zhou, D. Bachtrog, and M. B. Eisen, 2014 Sex-specific

embryonic gene expression in species with newly evolved sex chromosomes., (H. S.

Malik, Ed.). PLoS genetics 10: e1004159.

https://doi.org/10.1371/journal.pgen.1004159

Mack K. L., P. Campbell, and M. W. Nachman, 2016 Gene regulation and speciation in

house mice. Genome Research gr.195743.115.

https://doi.org/10.1101/gr.195743.115

Macmanes M. D., 2014 On the optimal trimming of high-throughput mRNA sequence

data. Frontiers in genetics 5: 13. https://doi.org/10.3389/fgene.2014.00013

Mannervik M., 2014 Control of Drosophila embryo patterning by transcriptional co-

regulators. Experimental Cell Research 321: 47–57.

https://doi.org/10.1016/j.yexcr.2013.10.010

Martin M., 2011 Cutadapt removes adapter sequences from high-throughput

sequencing reads. EMBnet.journal 17: 10–12. https://doi.org/10.14806/ej.17.1.200

Matzat L. H., and E. P. Lei, 2014 Surviving an identity crisis: A revised view of

chromatin insulators in the genomics era. Biochimica et Biophysica Acta (BBA) -

Gene Regulatory Mechanisms 1839: 203–214.

https://doi.org/10.1016/j.bbagrm.2013.10.007

McDermott S. R., and R. M. Kliman, 2008 Estimation of Isolation Times of the Island

Species in the Drosophila simulans Complex from Multilocus DNA Sequence Data.

PLoS ONE 3: e2442. https://doi.org/10.1371/journal.pone.0002442

McManus C. J., J. D. Coolon, M. O. Duff, J. Eipper-Mains, B. R. Graveley, *et al.*, 2010

Regulatory divergence in Drosophila revealed by mRNA-seq. Genome Research 20:

816–825. https://doi.org/10.1101/gr.102491.109

Mi H., A. Muruganujan, X. Huang, D. Ebert, C. Mills, *et al.*, 2019 Protocol Update for

large-scale genome and gene function analysis with the PANTHER classification

system (v.14.0). Nature Protocols 1–21. https://doi.org/10.1038/s41596-019-0128-8

Michalak P., 2003 Genome-Wide Patterns of Expression in Drosophila Pure Species

and Hybrid Males. Molecular Biology and Evolution 20: 1070–1076.

https://doi.org/10.1093/molbev/msg119

Mische S., M. Li, M. Serr, and T. S. Hays, 2007 Direct Observation of Regulated

Ribonucleoprotein Transport Across the Nurse Cell/Oocyte Boundary, (S. Wente,

Ed.). Molecular Biology of the Cell 18: 2254–2263. https://doi.org/10.1091/mbc.e06-

10-0959

Moehring A. J., K. C. Teeter, and M. A. F. Noor, 2007 Genome-Wide Patterns of

Expression in Drosophila Pure Species and Hybrid Males. II. Examination of Multiple-

Species Hybridizations, Platforms, and Life Cycle Stages. Molecular Biology and

Evolution 24: 137–145. https://doi.org/10.1093/molbev/msl142

Navarro-Costa P., A. McCarthy, P. Prudêncio, C. Greer, L. G. Guilgur, *et al.*, 2016 Early

programming of the oocyte epigenome temporally controls late prophase I

transcription and chromatin remodelling. Nat Commun 7: 12331.

https://doi.org/10.1038/ncomms12331

Nolte V., R. V. Pandey, R. Kofler, and C. Schlötterer, 2013 Genome-wide patterns of

natural variation reveal strong selective sweeps and ongoing genomic conflict in

Drosophila mauritiana. Genome Research 23: 99–110.

https://doi.org/10.1101/gr.139873.112

Nüsslein-Volhard C., and E. Wieschaus, 1980 Mutations affecting segment number and

polarity in Drosophila. Nature 287: 795–801. https://doi.org/10.1038/287795a0

Omura C. S., and S. E. Lott, 2020 The conserved regulatory basis of mRNA

contributions to the early Drosophila embryo differs between the maternal and

zygotic genomes. Plos Genet 16: e1008645.

https://doi.org/10.1371/journal.pgen.1008645

Ortíz-Barrientos D., B. A. Counterman, and M. A. F. Noor, 2006 Gene expression

divergence and the origin of hybrid dysfunctions. Genetica 129: 71–81.

https://doi.org/10.1007/s10709-006-0034-1

Paris M., J. E. Villalta, M. B. Eisen, and S. E. Lott, 2015 Sex Bias and Maternal

Contribution to Gene Expression Divergence in Drosophila Blastoderm Embryos., (A.

G. Clark, Ed.). PLoS genetics 11: e1005592.

https://doi.org/10.1371/journal.pgen.1005592

Perry M. W., A. N. Boettiger, and M. Levine, 2011 Multiple enhancers ensure precision

of gap gene-expression patterns in the Drosophila embryo. Proceedings of the

National Academy of Sciences 108: 13570–13575.

https://doi.org/10.1073/pnas.1109873108

Prud'homme B., N. Gompel, A. Rokas, V. A. Kassner, T. M. Williams, *et al.*, 2006

Repeated morphological evolution through cis -regulatory changes in a pleiotropic

gene. Nature 440: 1050–1053. https://doi.org/10.1038/nature04597

Quinlan A. R., and I. M. Hall, 2010 BEDTools: a flexible suite of utilities for comparing

genomic features. Bioinformatics 26: 841–842.

https://doi.org/10.1093/bioinformatics/btq033

R Core Team, 2017 *R: A language and environment for statistical computing*.

Ramírez F., V. Bhardwaj, L. Arrigoni, K. C. Lam, B. A. Grüning, *et al.*, 2018 High-

resolution TADs reveal DNA sequences underlying genome organization in flies.

Nature Communications 9: 1–15. https://doi.org/10.1038/s41467-017-02525-w

Rouget C., C. Papin, A. Boureux, A.-C. Meunier, B. Franco, *et al.*, 2010 Maternal mRNA

deadenylation and decay by the piRNA pathway in the early Drosophila embryo.

Nature 467: 1128-U144. https://doi.org/10.1038/nature09465

Salles F. J., M. E. Lieberfarb, C. Wreden, J. P. Gergen, and S. Strickland, 1994

Coordinate initiation of Drosophila development by regulated polyadenylation of

maternal messenger RNAs. Science (New York, N.Y.) 266: 1996–1999.

https://doi.org/10.1126/science.7801127

Schulz K. N., and M. M. Harrison, 2019 Mechanisms regulating zygotic genome

activation. Nature Reviews Genetics 1–14. https://doi.org/10.1038/s41576-018-0087-

x

Small S., A. Blair, and M. Levine, 1992 Regulation of Even-Skipped Stripe-2 in the

Drosophila Embryo. Embo Journal 11: 4047–4057. https://doi.org/10.1002/j.1460-

2075.1992.tb05498.x

Supek F., M. Bošnjak, N. Škunca, and T. Šmuc, 2011 REVIGO Summarizes and

Visualizes Long Lists of Gene Ontology Terms, (C. Gibas, Ed.). PLoS ONE 6:

e21800-9. https://doi.org/10.1371/journal.pone.0021800

Tadros W., A. L. Goldman, T. Babak, F. Menzies, L. Vardy, *et al.*, 2007 SMAUG Is a

Major Regulator of Maternal mRNA Destabilization in Drosophila and Its Translation

Is Activated by the PAN GU Kinase. Developmental Cell 12: 143–155.

https://doi.org/10.1016/j.devcel.2006.10.005

Tadros W., and H. D. Lipshitz, 2009 The maternal-to-zygotic transition: a play in two

acts. Development 136: 3033–3042. https://doi.org/10.1242/dev.033183

Theurkauf W. E., and T. I. Hazelrigg, 1998 In vivo analyses of cytoplasmic transport and

cytoskeletal organization during Drosophila oogenesis: characterization of a multi-

step anterior localization pathway. Dev Camb Engl 125: 3655–66.

Thomsen S., S. Anders, S. C. Janga, W. Huber, and C. R. Alonso, 2010 Genome-wide

analysis of mRNA decay patterns during early Drosophila development. Genome

biology 11: R93. https://doi.org/10.1186/gb-2010-11-9-r93

Torres-Oliva M., I. Almudi, A. P. McGregor, and N. Posnien, 2016 A robust (re-

)annotation approach to generate unbiased mapping references for RNA-seq-based

analyses of differential expression across closely related species. BMC Genomics

17: 392. https://doi.org/10.1186/s12864-016-2646-x

Trapnell C., A. Roberts, L. Goff, G. Pertea, D. Kim, *et al.,* 2012 Differential gene and

transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks.

Nature Protocols 7: 562. https://doi.org/10.1038/nprot.2012.016

Van der Auwera G. A. V. der, M. O. Carneiro, C. Hartl, R. Poplin, G. del Angel, *et al.*,

2013 Current Protocols in Bioinformatics. Curr Protoc Bioinform Ed Board Andreas D

Baxevanis Et Al 43: 11.10.1-11.10.33.

https://doi.org/10.1002/0471250953.bi1110s43

Vastenhouw N. L., W. X. Cao, and H. D. Lipshitz, 2019 The maternal-to-zygotic

transition revisited. Development 146: dev161471-20.

https://doi.org/10.1242/dev.161471

Wittkopp P. J., B. K. Haerum, and A. G. Clark, 2004 Evolutionary changes in cis and

trans gene regulation. Nature 430: 85–88. https://doi.org/10.1038/nature02698

Xu G., J. Lyu, Q. Li, H. Liu, D. Wang, *et al.*, 2020 Adaptive evolution of DNA methylation

reshaped gene regulation in maize. Biorxiv 2020.03.13.991117.

https://doi.org/10.1101/2020.03.13.991117

Zabidi M. A., C. D. Arnold, K. Schernhuber, M. Pagani, M. Rath, *et al.*, 2015 Enhancer–

core-promoter specificity separates developmental and housekeeping gene

regulation. Nature 518: 556–559. https://doi.org/10.1038/nature13994

1144

Figure 1: Crosses to produce hybrid embryos for the zygotic and maternal stages. To look at changes in regulation for zygotic genes, hybrid stage 5 embryos (left) were produced by crossing two parental species and collecting their eggs at the appropriate stage (late stage 5). To look at regulatory changes in maternal transcript deposition, F1 hybrid mothers were were mated to males and stage 2 embryos were collected (right). In both cases, transcription is coming from a F1 hybrid genome, either that of the zygote (left) which is measured after zygotic genome activation (late stage 5) or the mother (right) which is measured when all the transcripts in the embryo are maternally deposited (stage 2).

Figure 2: Hybrid and parental species single embryo transcript levels are highly reproducible. (A,B,D,E) Spearman's rank correlation coefficients are high when counts from replicate transcriptomes of the same stage and genotype are compared. Correlation coefficients are similarly high in parental species (D,E) and when comparing replicates from hybrid crosses (A,B). (C,F) Samples from different stages and the same genotype have much lower correlations, indicating a large difference in transcriptomes between the maternal and zygotic stages.

Figure 3: Different types of evolved regulatory changes dominate in maternal transcript deposition vs. zygotic transcription. Proportion of genes that fall into categories of regulatory change for each cross are shown for both the maternal transcript deposition (A) and zygotic gene transcription (B), for mostly-zygotic genes. Transcript level ratios between parental lines and within hybrids at stage 2 (C) and stage 5 (D) describe regulatory changes between *D. mauritiana* and *D. simulans* in one direction of crosses (for the rest of the crosses, see Figures S2 and S3).

Figure 4: Examples of the type of regulatory changes observed, for individual genes. Transcript abundance, shown in counts, for each gene is plotted for the total mRNA abundance in both parental lines and for each parental allele within the hybrid; error bars shown represent the standard deviation. Total transcript abundance in the hybrid (the summation of levels from parental alleles in the hybrid) is shown as the last bar on the right in each graph. (A) Maternal transcript deposition of *Cyclin-dependent kinase 1 (Cdk1),* a critical cell cycle regulator in early development, changes in *trans* regulation between *D. mauritiana* and *D. sechellia*. Hybrid mRNA abundance is from the (*mau* x *sec*) x *mau* cross. (*Cdk1* also changes in *trans* regulation in the reciprocal cross comparison, (*sec* x *mau*) x *sec.*) (B) Maternal transcript deposition of *CG11241*, a gene of currently unknown function, changes in *trans* regulation between *D. sechellia* and *D. simulans*. Hybrid mRNA abundance is from the (*sim* x *sec*) x *sim* cross. (C) At stage 5 in development, *sloppy paired 1,* a critical pair-rule segmentation gene, changes in regulation through a combination of *cis* and *trans* regulatory changes (*cis* + trans) between *D. simulans* and *D. sechellia*. Hybrid expression is shown for the *sim* x *sec* cross. *Sloppy paired 2* also changes in *cis* regulation between these two species. (D) At stage 5, *Kruppel*, a gap gene crucial to segmentation changes in regulation through a combination of *cis* and *trans* regulatory elements (*cis* x *trans*) between *D. mauritiana* and *D. simulans*. Here, expression in the hybrid is from the *sim* x *mau* cross but in the reciprocal cross (*mau* x *sim*), *Kruppel* also changes in *cis* x *trans* regulation.

Figure 5: Gene ontology (GO) analysis identifies transcription factors that act in developmental processes as types of genes that change zygotically. Significantly enriched GO terms are listed for zygotically transcribed genes that change in *trans* regulation between each pair of species compared. Genes represented in this analysis are categorized as mostly zygotic (see Methods). Terms are listed for Biological Processes and Molecular Function categories and only terms that appear in more than one cross are shown in this figure. Terms unique to a specific cross are listed in Figure S5. Biological process categories identified relate to development, molecular function categories identify functions consistent with DNA binding and regulation.

Figure 6: Patterns of inheritance show dominance of particular parental genomes at both stages. A) Shows patterns of inheritance for stage 2, over all genes and all crosses. B) Shows patterns of inheritance for stage 5, for mostly zygotic genes (see Methods) and all crosses. The maternal stage (A) shows a higher proportion of conserved genes than the zygotic stage (B). Both stages show a high degree of dominance for *D. simulans* for crosses involving that species, and for *D. sechellia* in crosses with *D. mauritiana*, forming the general dominance pattern of *D. simulans* > *D. sechellia* > *D. mauritiana.* There is a greater proportion of additive inheritance for the zygotic stage (B) than the maternal stage (A).

Figure S1: PCA plots for transcript abundance in all crosses. Samples of each stage, 2 or 5, cluster together. Samples of each genotype also cluster together, parental samples and hybrids. Proportion of variance explained by each principal component is listed on each axis.

Figure S2: Regulatory changes in all pairwise comparisons for maternally deposited transcripts. Transcript level ratios between parental lines and within hybrids at stage 2 describe regulatory changes between species in each set of crosses. The number of genes in each category of regulatory change (n=) is listed in each plot. For definitions of categories of changes and criteria, see Methods.

Figure S3: Regulatory changes in all pairwise comparisons for mostly zygotic genes. Transcript level ratios between parental lines and within hybrids at for mostly zygotic genes (see Methods) at stage 5 describe regulatory changes between species in each set of crosses. The number of genes in each category of regulatory change (n=) is listed in each plot. For definitions of categories of changes and criteria, see Methods.

Figure S4: Transcript abundance from parental lines at stage 2 demonstrates differential maternal deposition of *M1BP*, *Dref* and *BEAF-32*. Counts for *D. simulans*/*D. mauritiana* and *D. simulans*/*D. sechellia* comparisons are averages across replicates from alignment to the *D. simulans genome*. Counts for *D. sechellia*/*D. mauritiana* comparison are averages across replicates from alignments to the *D. sechellia* genome. Error bars represent standard deviations. Count data for the same species and gene may differ across comparisons due to the genome used for alignment in each comparison and normalization of counts within a comparison.

Figure S5: Gene ontology (GO) analysis for categories unique to a specific cross show enrichment for specific developmental processes. Significantly enriched GO terms are listed for zygotically transcribed genes that change in *trans* regulation between each pair of species compared. Again, zygotically transcribed genes are limited to those that are mostly zygotic (see Methods), in comparison to the maternal stage of development. Terms are listed for the biological processes category. Gene categories identified uniquely in a single cross primarily represent specific types of developmental processes, and may indicate evolved differences in parental genomes in these processes.

Target Gene	Sequence
L_ORY	aatacaactcaggagcgggacaatttt
R_ORY	tcgtaccatttgcaatccgactag
L_kl3	gaacgcgcatccattttattct
R_kl3	tcgaaaagcccacgacaggtattt
L_ftz	accaaccccgtgaagaagctgaagtaca
R_ftz	cgtgtgtgatgcctacctgatgccaaagt

Table S1: Primers for genes *ORY*, *kl3* (both on the Y chromosome) and *ftz* (control locus, on 3R) that were used for genotyping stage 5 embryos as male or female.

Motif	Predicted Binding Proteins	Enriched in upstream regions of *D. sechellia* genes that are conserved in expression (E-value)	Enriched in upstream regions of *D. sechellia* genes that are changing in expression (E-value)	Enriched in upstream regions of *D. simulans* genes that are conserved in expression (E-value)	Enriched in upstream regions of *D. simulans* genes that are changing in expression (E-value)
	Dref, BEAF-32	(*sec* x *mau*) x *sec* (1.7e-009) (*sim* x *sec*) x *sim* (9.6e-011)	(*mau* x *sec*) x *mau* (2.2e-023) (*sec* x *mau*) x *sec* (3.5e-024) (*sim* x *sec*) x *sim* (2.7e-011)	(*sim* x *sec*) x *sim* (1.7e-006)	(*mau* x *sim*) x *mau* (2.5e-010) (*sim* x *mau*) x *sim* (2.8e-032) (*sim* x *sec*) x *sim* (2.2e-032)
	M1BP	(*mau* x *sec*) x *mau* (2.8e-006) (*sec* x *mau*) x *sec* (3.0e-009) (*sim* x *sec*) x *sim* (1.1e-006)	(*mau* x *sec*) x *mau* (7.7e-008) (*sec* x *mau*) x *sec* (5.5e-028) (*sim* x *sec*) x *sim* (1.8e-015)		(*sim* x *mau*) x *sim* (2.3e-008)
	crp, salivary gland-expressed bHLH, similar to Deadpan, E-box, nau (only in *sec* x *mau* and *sim* x *sec* comparison in *D. sechellia* upstream regions)	(*mau* x *sec*) x *mau* (6.6e-007) (*sec* x *mau*) x *sec* (6.6e-003) (*sim* x *sec*) x *sim* (7.5e-004)			

Table S4: Enriched motifs found upstream of maternally deposited genes. Sequences 500bp upstream were extracted for genes in *D. simulans* and *D. sechellia* that change in regulation or that are conserved in each pairwise comparison. Motifs that were significantly enriched in analysis using MEME and HOMER are listed in the table and predicted binding proteins discovered using Tomtom and Homer are also described. E-values generated by MEME indicating the enrichment of each motif compared to background in each cross are also listed. The position weight matrix represented is a representative example of the discovered motifs.

Table: Number of genes with each pattern of inheritance in hybrids							
	Additive	*sim* dominant	*sec* dominant	*mau* dominant	Underdominant	Overdominant	Conserved
(*sim* x *mau*) x *sim*	88	735		182	197	270	376
(*mau* x *sim*) x *mau*	97	733		210	168	260	412
(*sec* x *mau*) x *sec*	237		540	292	280	371	322
(*mau* x *sec*) x *mau*	220		489	302	329	382	313
(*sim* x *sec*) x *sim*	180	450	215		141	187	536
sim x *mau*	51	68		20	71	27	17
mau x *sim*	77	57		31	40	40	11
sec x *mau*	70		62	30	39	44	17
mau x *sec*	58		69	35	34	49	19
sim x *sec*	49	59	42		51	24	19

Table S6: Number of genes within each category of inheritance pattern in the hybrids. Stage 2 hybrids are shown in the first five rows and stage 5 hybrids are shown in the last five rows of the table.