Title: Acute feeding disorder after oral RNAi exposure results in efficient pest control

Authors: Yasuhiko Chikami[a,b], Haruka Kawaguchi[a], Takamasa Suzuki[c], Hirofumi

Yoshioka[d], Yutaka Sato[d,e,f], Toshinobu Yaginuma[d] and Teruyuki Niimi[a,b,d, 1]

Author affiliations:

[a] Division of Evolutionary Developmental Biology, National Institute for Basic Biology,

Nishigonaka 38, Myodaiji, Okazaki, Aichi 444-8585, Japan

[b] Department of Basic Biology, School of Life Science, SOKENDAI (The Graduate

University for Advanced Studies), Nishigonaka 38, Myodaiji, Okazaki, Aichi 444-8585,

Japan

[c]Department of Biological Chemistry, College of Bioscience and Biotechnology, Chubu

University, Matsumoto-cho 1200, Kasugai, Aichi 487-8501, Japan.

[d]Graduate School of Bioagricultural Sciences, Nagoya University, Furo-cho, Chikusa,

Nagoya, Aichi 464-8601, Japan

[e] Plant Genetics Laboratory, Genetic Strains Research Center, National Institute of

Genetics, Yata 1111, Mishima, Shizuoka 411-8540, Japan

[f]Department of Genetics, School of Life Science, SOKENDAI (The Graduate University

for Advanced Studies), Yata 1111, Mishima, Shizuoka 411-8540, Japan

[1] **Corresponding author:**

Teruyuki Niimi

E-mail: niimi@nibb.ac.jp

Division of Evolutionary Developmental Biology, National Institute for Basic

Biology, Nishigonaka 38, Myodaiji, Okazaki, Aichi 444-8585, Japan

TEL: +81 564 55 7606

ABSTRACT

Oral feeding of RNA interference (f-RNAi) is an innovative technology in the protection of crops from herbivorous pests. The strength of this technology lies in choosing optimal f-RNAi targets. Conventional targets have been chosen on due to their induction of lethality or infertility in target pests. However, conventional targets are slow-acting (c.a. one week) and crop damage can progress until the induction of lethality due to f-RNAi. Here, we show that f-RNAi of *death-associated inhibitor of apoptosis protein 1* (*diap1*) induces acute feeding cessation. In the solanaceous pest, *Henosepilachna vigintioctopunctata*, f-RNAi of *diap1* caused a decrease in feeding leaf area during the first 24 hours after treatment. Feeding cessation in f-RNAi individuals was achieved during 24-48 hours. We compared our targets with conventional targets by analyzing the influence of frequently used target genes, *vacuolar ATPase subunits A* and *E*, and showed that unlike *diap1* f-RNAi, conventional targets did not induce acute feeding disorder. Feeding experiments of *diap1* double-stranded RNA (dsRNA) from other insects against *H. vigintioctopunctata* did not induce feeding cessation and showed high species-specificity of *diap1* dsRNA. We also showed that 1.6 ng *diap1* dsRNA is sufficient to induce acute feeding cessation in *H. vigintioctopunctata*. These results suggest that *diap1* is an optimal target gene for rapid reduction of crop damage by herbivorous insects. In addition, our findings suggest that acute feeding cessation should be assessed as well as conventional criteria in selecting target genes for pest control by f-RNAi.

KEYWORDS: RNAi, pest control, Inhibitor of apoptosis protein

INTRODUCTION

Techniques for manipulating endogenous essential genes in pests using the RNA interference (RNAi)[1,2,3] or genome editing such as CRISPR/Cas9 system[4,5] have recently attracted attention alternative strategies for pest management to conventional chemical pesticides. Pest control mediated by endogenous gene manipulation can be achieved by inducing lethality[6,7] or infertility[5,8], which are essential for eradication of pests. However, since pests continue to feed on crops until the effects of lethality or infertility appears, such target genes are not necessarily optimal for preventing crop damage. Therefore, it would be more desirable for pest control to identify genes that can inducing acute feeding cessation in pests.

We found that the *death-associated inhibitor of apoptosis protein 1* (*diap1*) gene can be an effective target gene for pest control meeting the criterion. Diap1 is one of insect paralogues of *inhibitor of apoptosis protein* genes of vertebrates and essential for maintaining cell survival by inhibiting the apoptotic pathway through inactivating caspase proteins[9,10]. For an optimal target of pest control by oral feeding of RNAi (f-RNAi), *diap1* has ideal properties such as an evolutionarily conserved protein with moderate diverged sequences, an extremely short half-life of protein molecules in cells (40 minutes)[11] and higher expression in the midgut (Fig. S1). We investigated the effects of f-RNAi for *diap1* in the 28-spotted ladybird, *Henosepilachna vigintioctopunctata,* which is a pest for Solanaceae (Fig. S2) and compared its pest control efficacy with those of *vacuolar ATPase subunit* (*v-ATPase*) *A* and *E* which were previously reported as effective target genes[6,12] and frequently used.

RESULTS AND DISCUSSIONS

diap1, v-ATPase A and *v-ATPase E* orthologs cloned from *H. vigintioctopunctata* (Figs. S3, S4) were expressed in whole body including the gut (Fig. 1a). For f-RNAi, we fed 50 ng of the double-stranded (ds) RNA targeting each gene to the 3rd instar larvae (Fig. S5). Quantitative RT-PCR analysis revealed that the expression of each target gene in the larvae at two-days after f-RNAi treatment significantly decreased as compared to the negative control larvae fed with *egfp* dsRNA (Fig. 1b). This result indicated that the knockdown of each target gene by f-RNAi was effective in *H. vigintioctopunctata*.

We found that the body weight significantly decreased in the larvae fed with 50 ng of *diap1* dsRNA as compared to the larvae fed with *egfp* dsRNA (Fig. 1c and Table S1). We further found that potato leaves were hardly eaten by *diap1* f-RNAi treated larvae (Fig. 1d). In order to evaluate the influence on crop damage after RNAi treatment, we measured the area of potato leaves consumed by the larvae during 0-24 and 24-48 hours after dsRNA ingestion (Fig. 1e). *diap1* RNAi treated larvae showed a significantly narrower area of consumed leaves than negative control larvae within 48 hours after dsRNA ingestion (Fig. 1f and Tables S1, S2), whereas no significant difference was observed in the *v-ATPase A* and *E* RNAi treatment. These results indicate that the knockdown of *diap1* by f-RNAi induces rapid feeding cessation in the larvae after RNAi treatment, although all larvae treated with f-RNAi in our experiments did not show lethality at least until 48 hours after f-RNAi. Previous reports evaluated lethality and growth-inhibition in pest treated with injection- or f-RNAi targeting several genes including *v-ATPase*[6,12] and *diap1*[13] in longer periods (c.a., a week). However, the acute effects of f-RNAi on feeding cessation have not been evaluated thus far. Our findings provide the first report on effective target genes which

induce rapid feeding cessation, an essential phenotype for crop protection. Since *diap1* f-

RNAi induced almost complete feeding cessation that is not observed in conventional

target gene RNAi treatments (Table S1), *diap1* seems to be the most desirable target gene

for the alleviation of crop damage.

The moderately diverged sequence in the *diap1* gene is useful for designing species-

specific RNAi pesticides. The nucleotide sequences of *diap1* when compared among 10

insect species showed 37.3-53.7% identities (Fig. 2a). Notably, the nucleotide sequences

between *H. vigintioctopunctata* and *Harmonia axyridis*, which belong to the same family

showed lower identity in *diap1* (Fig. 2a) than those of *v-ATPase A* and *E* (Fig. S6). This

finding suggested that dsRNA targeting the homologous region in the *diap1* ortholog genes

can use for species-specific RNAi pesticide reagents, and allowed us to examine RNAi

specificity on *H. vigintioctopunctata* larvae using 320 ng dsRNAs targeting *diap1*

sequences cloned from 4 other insects (Fig. 2b). Of the RNAi treatments, only *H.*

vigintioctopunctata diap1 RNAi treatment showed reduction in feeding damage in 0-24

hours after RNAi treatment, and feeding cessation in 24-48 hours after treatment (Fig. 2c).

In contrast, f-RNAi effects under the cross-species treatment of the *v-ATPase* genes were

previously reported[6]. These results indicate that *diap1* is also an optimal target for

designing highly species-specific RNAi reagents.

We also investigated the dose-sensitivity of *diap1* f-RNAi. Larvae fed with 1.6 and 8 ng

of *diap1* dsRNA showed significant decreases in the area of consumed leaves (Fig. 2d).

This result suggests that 1.6 ng dsRNA is sufficient to induce feeding cessation in the 3rd

instar larva of *H. vigintioctopunctata*. Since many previous reports of f-RNAi for pest

management require an amount ranging from several hundred nano-grams to a few micro-

grams of dsRNA[14], *diap1* is also the most efficient dsRNA target for f-RNAi to date.

The technologies for application of RNAi pesticide have been developed[3]. It has also been

reported that combination of RNAi and Bt pesticide produces a stronger insecticide effect

than either alone[15]. However, these studies did not investigate genes with either acute

affects or high species- and dose-sensitivity, which are essential to the practice of RNAi as

a pesticide. In the future, targeting *diap1* may allow the application of highly efficient

RNAi pesticide treatments.

Furthermore, our results indicate the representative target genes in terms of lethality and

growth-inhibition are not necessarily the optimal target for stopping crop damage. We

propose the quantitative evaluation of acute feeding disorders should be required for

selecting target genes in the application of RNAi pesticides, in addition to the conventional

criteria.

MATERUALS AND METHODS

Insects

Henosepilachna vigintioctopunctata were collected from potato leaves at either Nagoya

University or National Institute for Basic Biology, in Japan and used these animals to

establish a laboratory line. Laboratory stocks of *Harmonia axyridis* were derived from field

collections in Aichi, Japan. They were reared as described by Niimi *et al.*[17] *Helicoverpa*

armigera was kindly provided by Dr. Chie Goto (NARO Agricultural Research Center).

Blattela lateralis were purchased from Remix. The larvae of *Oxya yezoensis* were collected

from rice field at Nagoya University.

Cloning of partial *diap1* sequences from insects

Total RNA was extracted from the gonad of *H. vigintioctopunctata*, the embryos of *H. axyridis*, the anterior wing primordia of *H. armigera*, the posterior wing primordia of *B. lateralis* or the larval posterior legs of *O. yezoensis* using TRIzol (Invitrogen, California, USA) according to the manufactural protocol. The first-strand cDNA was synthesized using the SuperScript II Reverse Transcriptase (Life Technologies Japan Ltd., Tokyo, Japan) with SMART RACE cDNA Amplification kit (Clontech, Mountain View, California, USA) from 1 μg total RNA. *diap1* cDNA fragment of each species was amplified using degenerated primer sets designed from conserved amino acid sequences among insects (diap1_F1 or diap1_F2/diap1_R in Table S7). The RT-PCR were performed using AmpliTaq Gold DNA polymerase (Perkin Elmer, Boston, USA). PCR fragments were separated with electrophoresis using 1% agarose gel in tris-borate-EDTA (TBE) buffer and extracted using MagExtractor (TOYOBO Co. Ltd., Osaka, Japan). Each PCR fragment was subcloned into the EcoRV recognition site of pBluescript KS (+) vector (Stratagene, La Jolla, CA, USA) using DNA Ligation kit Ver. 2 (TaKaRa bio. Inc., Shiga, Japan). The vector was transformed into XL1-Blue *Escherichia coli* competent cells (GMbiolab Co., Ltd, Taichung, Taiwan). The plasmid was extracted from the transformed colony using FlexiPrep Kit (Amersham Pharmacia Biotech Inc., New Jersey, USA). The nucleotide sequences of the PCR fragment inserted into vectors were confirmed using the dideoxy chain-termination method using an automatic DNA sequencer (CEQ 2000XL; Beckman Coulter, California, USA). Sequence analysis was carried out using a DNASIS system (Hitachi Software Engineering, Tokyo, Japan). The partial length *diap1* nucleotide sequences of the insects were deposited in DNA Data Bank of Japan (DDBJ) (accession

numbers: LC473085 in *H. axyridis*, LC473088 in *H. armigera*, LC473087 in *B. lateralis*

and LC473086 in *O. yezoensis*).

Rapid amplification of cDNA end (RACE) of *diap1* in *H. vigintioctopunctata*

To obtain a full-length *diap1* cDNA in *H. vigintioctopunctata*, the 5'- and 3'- rapid

amplification of cDNA end (RACE) were performed according to the manufacturer's

protocol of the SMART RACE cDNA Amplification kit (Clontech, Mountain View,

California, USA). The gene-specific primers are described in Table S7 (1st PCR:

Hvig_diap1_1 for 5'-RACE and Hvig_diap1_3 for 3'-RACE; nested PCR: Hvig_diap1_2

for 5' RACE and Hvig_diap1_4 for 3'RACE). These primers were designed from the *diap1*

cDNA fragment cloned above. The cloning and confirmation of nucleotide sequences of

RACE fragment were performed according to the same way as the above method. The full

length *diap1* nucleotide sequences of *H. vigintioctopunctata* was deposited in DDBJ

(accession numbers: LC473084). The *diap1* full-length nucleotide sequence is shown in

Supplementary Sequences.

Transcriptomic analysis

In order to obtain *diap1*, *v-ATPase A* and *v-ATPase E* sequences of *H. vigintioctopunctata*,

transcriptomic analysis was performed using total RNA from the whole body of *H.*

vigintioctopunctata 3rd instar larvae. Total RNAs were extracted using the RNeasy Mini

kit (QIAGEN, Tokyo, Japan). The integrity of extracted RNAs were confirmed using a

Bioanalyzer 2000 (Agilent Technologies) and these RNAs were used to construct cDNA

libraries using the Truseq™ RNA Sample Prep Kit (Illumina) according to the

manufacture's instruction. Pair-end sequences with 200 bp were obtained from these cDNA

libraries through Hiseq2000 (Illumina). The raw reads were deposited in DDBJ Sequence

Read Archive (accession number: #### [the deposition of data is now proceeding, and the

accession number will be obtained during editorial process]). We rearranged pair-end reads

in order and excluded unpaired reads using cmpfastq_pe

(http://compbio.brc.iop.kcl.ac.uk/software/cmpfastq_pe.php). Low quality reads and

adapter sequences were trimmed from obtained short-reads using Cutadapt v1.15[18]. Then,

de novo assembly of trimmed short-read sequence was performed using Trinity-v4.2.0

(https://github.com/trinityrnaseq/trinityrnaseq/)[19]. We also analyzed the RNA-seq data of

H. axyridis obtained from National Center for Biotechnology Information (NCBI)

Sequence Read Archive database (run number.: ERR1309559) in the same way as *H.*

vigintioctopunctata.

BLAST search and phylogenetic analysis of target genes

BLAST databases of the assembled transcriptomic sequences in *H. vigintioctopunctata* and

H. axyridis were established by makeblastdb program in BLAST+ ver. 2.7.1[20]. We searched

homologous genes of *v-ATPase A, v-ATPase E* and *diap1* in the *H. vigintioctopunctata*

BLAST database with the orthologues in *Drosophila melanogaster* (accession number:

Diap1, NP_001261916.1; V-ATPase A, NP_652004.2; V-ATPase E, NP_524237.1)

obtained from NCBI database as query sequences using the tblastn program in BLAST+

ver. 2.7.1. Then, we confirmed the orthology of these obtained sequences by molecular

phylogenetic analysis (Fig. S4). The homologous sequences of V-ATPase A*,* V-ATPase E

and Diap1 in other animals were obtained from NCBI database (Table S6). All sequences

including *H. vigintioctopunctata* homologues were aligned using MAFFT ver. 7[21] with the

L-INS-i program. The using sites for phylogenetic construction were selected by trimAl

v1.2[22] with a gap threshold value of 0.7. The multiple alignments were shown in Figs. S7-

S9. The selection of best-hit substitution model and construction of maximal-likelihood

phylogeny were performed using MEGA X[23]. Bootstrap values were calculated after 1000

replications. BLAST search and multiple alignment were performed under Super Computer

Facilities of National Institute of Genetics. The *diap1*, *v-ATPase A* and *v-ATPase E*

nucleotide sequences of *H. vigintioctopunctata* and *H. axyridis* are shown in

Supplementary Sequences.

Expression profile analysis of target genes in larval tissues

Total RNA was extracted from nerve cord, legs, wing discs fat body, Malpighian tubules,

gut and carcass of 4th instar larvae of *H. vigintioctopunctata* which is the dissectable stage

for each tissue using the RNeasy Mini kit (QIAGEN, Tokyo, Japan). The first-strand cDNA

was synthesized from the 1 µg total RNA using SuperScript III Reverse Transcriptase (Life

Technologies Japan Ltd., Tokyo, Japan). RT-PCR for the target genes and the internal

control (*ribosomal protein 49* (*rp49*), accession number: AB480201) was performed using

the first-strand cDNA and Q5 High-Fidelity DNA Polymerase (New England Biolabs

Japan Inc., Tokyo, Japan) with 30 cycles. Primers for RT-PCR were designed by

Primer3web version 4.1.0[24]   and are shown in Table S7 (Hvig_diap1_RT-PCR_F/

Hvig_diap1_RT-PCR_R, Hvig_v-ATPase_A_RT-PCR_F/Hvig_v-ATPase_A_RT-PCR_R,

Hvig_v-ATPase_E_RT-PCR_F/ Hvig_v-ATPase_E_RT-PCR_R and Hvig_rp49_RT-

PCR_F/Hvig_rp49_RT-PCR_R). The annealing temperatures for RT-PCR were calculated

by NEB Tm Calculator (https://tmcalculator.neb.com/). The expression pattern was confirmed by agarose gel electrophoresis of the PCR products with 2% Agarose S (NIPPON GENE Co. Ltd., Toyama, Japan).

Double strand RNA synthesis

<u>Comparison analysis of RNAi efficiency of target genes</u>

The total RNA was extracted from 3rd instar larvae of *H. vigintioctopunctata* using TRI Reagent (Molecular Research Center Inc., Ohio, USA) according to the manufactural protocol. The first-strand cDNA was synthesized from 1 µg total RNA using SuperScript III Reverse Transcriptase (Life Technologies Japan Ltd., Tokyo, Japan). The templates of dsRNA were initially amplified from the first-strand cDNA and secondary amplified from the initial PCR products by RT-PCR using Q5 High-Fidelity DNA Polymerase (New England Biolabs Japan Inc., Tokyo, Japan). The initial PCR primers are described as Hvig_diap1_F/Hvig_diap1_R, Hvig_v-ATPase_A_F/Hvig_v-ATPase_A_R and Hvig_v-ATPase_E_F/ Hvig_v-ATPase_E_R in Table S7. The nested primers were flanked with T7 promoter sequences on the 5' ends (Hvig_T7-diap1_F/Hvig_T7-diap1_R, Hvig_T7-v-ATPase_A_F/Hvig_T7-v-ATPase_A_R and Hvig_T7-v-ATPase_E_F/Hvig_T7-v-ATPase_E_R in Table S7). The nested PCR products were purified using MagExtractor (TOYOBO Co. Ltd., Osaka, Japan). The sequences of interest were confirmed by sanger DNA sequencing service at FASMAC Co. Ltd. (Kanagawa, Japan). The dsRNAs of target genes were synthesized from the purified PCR products using AmpliScribe T7-Flash Transcription Kit (Epicentre Technologies, Co., Wisconsin, USA).

<u>Species-specificity and dose analysis of *diap1* in *H. vigintioctopunctata*</u>

The templates of dsRNA were amplified from the above pBluescript KS (+) vectors

inserted *diap1* sequences in each insect by PCR using AmpliTaq Gold (Perkin Elmer,

Boston, USA). The PCR primers are described in Table S7 (T7-KS/T7-SK). The PCR

products were purified by the same ways to the above method. The dsRNAs were

synthesized using MEGAscript T7 Transcription Kit (Ambion, Texas, USA).

Feeding RNAi assay

Synthesized dsRNA was artificially fed to 3rd instar larvae of *H. vigintioctopunctata* within

24 hours after molting (Fig. S5). The 0.5 µl solution including dsRNA of each gene was

dropped on slide glasses in front of the larvae which had been starved after molting to the

3rd instar. Larvae starved for this amount of time readily consumed the dsRNA solution.

After the larvae had consumed the entire droplet, each larva was put on a fresh potato leaf

and kept in a plastic container at room temperature. Body weight of the larvae after 48

hours were measured using a precision electronic balance (Table S1). The potato leaves

were collected and exchanged for new potato leaves up to 48 hours every 12 or 24 hours

(Fig. 1e). dsRNA of *enhanced green fluorescent protein* (*egfp*) were fed to 3rd instar larvae

as the negative control.

Evaluation of knockdown of mRNAs expression by quantitative (q) RT-PCR

The total RNA was extracted from the whole-body of *H. vigintioctopunctata* 3rd instar

larvae 48-hours after f-RNAi treatment using TRI Reagent (Molecular Research Center

Inc., Ohio, USA). The first-strand cDNA was synthesized from 1 µg total RNA using

SuperScript III Reverse Transcriptase (Life Technologies Japan Ltd., Tokyo, Japan). qRT-

PCR was performed on LightCycler 96 instrument (Roche, Basel, Swizerland) using

THUNDERBIRD SYBR qPCR Mix (TOYOBO Co. Ltd., Osaka, Japan) according to the

manufactural protocol with the first-strand cDNA as the template. The same primer sets as

the above expression profile analysis of each gene were used. The expression level of target

genes relative to *rp49* was calculated by the $2^{-\Delta\Delta Ct}$ method[25].

Measurements of leaf area eaten by feeding RNAi larvae

We measured the potato leaf area eaten by f-RNAi larvae using the following method with

the digital microscope system (VHX-5000, KEYENCE, Osaka, Japan). A leaf was put on

the glass plate on the stage and was covered with a plastic wrap. Then, the region of leaf

eaten was manually bordered with a black pen. After focusing on the leaf using 10x

magnification, we automatically binarized the image and measured the area of the region

eaten by larvae with the digital microscope system.

Statistical analysis

We performed the Welch's *t*-test to evaluate the differences of effects of f-RNAi assay. We

adjusted *P*-value with the Holm's methods for the multiple-comparisons. The significance

level used in our analysis was *P*-value < 0.05. All statistical analyses were performed by

R-3.5.0 (https://cran.r-project.org/).

Sequence analysis of target genes among insects

The nucleotide and amino acid sequences of target genes in 9 insect species were obtained

from NCBI's database (Table S6). Then, multiple alignments were performed for each pair

of species using MAFFT ver.7 with L-INS-i program. The proportions of identical sites for

length of coding sequence region of each gene were manually calculated. The identical

sites of partial sequences of *diap1* between *H. vigintioctopunctata* and other insects using

for f-RNAi assay were identified in the same way (alignment: Fig. S10).

Acknowledgements

We would like to thank to R. A. Zinna (Mars Hill University) for his critical read of this

manuscript. We also thank J. Yatomi for helping in the part of this work and T. Ando, T.

Nakamura and H. Sakai (National Institute for Basic Biology, Japan) for helpful

discussions. We express our gratitude to C. Goto (NARO Agricultural Research Center,

Japan) for providing *Helicoverpa armigera* and T. Konagaya (National Institute for Basic

Biology, Japan) for the critical advice to statistical analysis. We thank the Model Plant

Research Facility, NIBB BioResource Center for their technical support. This work was

supported in part by the Center for the Promotion of Integrated Sciences (CPIS) of

SOKENDAI (to T.N. and Y.S.).

Contributions

Y.C., H.Y., Y.S., T.Y. and T.N. conceived and designed the study. Y.C., H.K. and T.N.

performed the majority of experiments. Y.C., T.S. and Y.S. analyzed RNA-seq data. H.K.

maintained *H. vigintioctopunctata* and potato in laboratory. T.N. supervised the study. Y.C.

and T.N. wrote the manuscript. All authors discussed the manuscripts.

Competing interests

The authors declare that no competing interests exist.

References

1. Y.-B. Mao, W.-J. Cai, J.-W. Wang, G.-J. Hong, X.-Y. Tao, L.-J. Wang, Y.-P. Huang, X.-Y. Chen, Silencing a cotton bollworm P450 monooxygenase gene by plant-mediated RNAi impairs larval tolerance of gossypol. *Nat. Biotechnol.* **25,** 1307–1313 (2007).

2. Y. Wang , H. Zhang , H. Li, X. Miao, Second-generation sequencing supply an effective way to screen RNAi targets in large scale for potential application in pest insect control. *PloS one*. **6,** e18644 (2011).

3. J. Zhang, S. A. Khan, C. Hasse, S. Ruf,.D. G. Heckel, R. Bock, Full crop protection from an insect pest by expression of long double-stranded RNAs in plastids. *Science*. **347,** 991-994 (2015).

4. Y. Wang, Z. Li, J. Xu, B. Zeng, L. Ling, L. You, Y. Chen, Y. Huang, A. Tan, The CRISPR/Cas System mediates efficient genome engineering in *Bombyx mori. Cell Res*. **23,** 1414-1416 (2013).

5. K. Kyrou, A. M. Hammond, R. Galizi, N. Kranjc, A. Burt, A. K. Beaghton, T. Nolan, A. Crisanti, A CRISPR–Cas9 gene drive targeting doublesex causes complete population suppression in caged *Anopheles gambiae* mosquitoes. *Nat. Biotechnol*. **36,** 1062-1066 (2018).

6. J. A. Baum, T. Bogaert, W. Clinton, G. R. Heck, P. Feldmann, O. Ilagan, S. Johnson, G. Plaetinck, T. Munyikwa, M. Pleau, T. Vaughn, J. Roberts, Control of coleopteran insect pests through RNA interference. *Nat. Biotechnol.* **25,** 1322-1326 (2007).

7. J. Ulrich, V. A. Dao, U. Majumdar, C. Schmitt-Engel, J. Schwirz, D. Schultheis, N.

Ströhlein, N. Troelenberg, D. Grossmann, T. Richter, J. Dönitz, L. Gerischer, G. Leboulle, A. Vilcinskas, M. Stanke, G. Bucher, Large scale RNAi screen in *Tribolium* reveals novel target genes for pest control and the proteasome as prime target. *BMC Genomics*. **16,** 674 (2015).

8. A. Hammond, R. Galizi, K. Kyrou, A. Simoni, C. Siniscalchi, D. Katsanos, M. Gribble, D. Baker, E. Marois, S. Russell, A. Burt, N. Windbichler, A. Crisanti, T. Nolan, A CRISPR-Cas9 gene drive system targeting female reproduction in the malaria mosquito vector *Anopheles gambiae*. *Nat. Biotechnol.* **34,** 78–83 (2016).

9. G. S. Salverson, C. S. Duchett, Apoptosis: IAP proteins: blocking the road to death's door. *Nat. Rev. Mol. Cell Biol.* **3,** 401–410 (2002).

10. D. Vasudevan, H. D. Ryoo, Regulation of cell death by IAPs and their antagonists. *Curr. Topic. Dev. Biol*. **114,** 185–208 (2015).

11. S, J, Yoo, J. R. Huh, I. Muro, H. Yu, L. Wang, S. L. Wang, R. M. R. Feldman, R. J. Clem, H.-A. J. Müller, B. A. Hay, Hid, Rpr and Grim negatively regulate DIAP1 levels through distinct mechanisms. *Nat. Cell Biol.* **4,** 416–424 (2002).

12. D. R. G. Price, J. A. Gatehouse, RNAi-mediated crop protection against insects. *Trends Biotechnol.* **26,** 393-400 (2008).

13. T. B. Rodrigues, L. K. Rieske, J. J. Duan, K. Mogilicherla, S. R. Palli, Development of RNAi method for screening candidate genes to control emerald ash borer, *Agrilus planipennis*. *Sci. Rep.* **7,** 7379 (2017).

14. R. Katoch, A. Sethi, N. Thakur, L. L. Murdock, RNAi for insect control: current perspective and future challenges. *Appl. Biochem. Biotechnol.* **171,** 847–873 (2013).

15. B. E. Tabashnik, Y. Carrière, Surge in insect resistance to transgenic crops and prospects

for sustainability. *Nat. Biotechnol.* **35,** 926-935 (2017).

16. B. Misof, S. Liu, K. Meusemann, R. S. Peters, A. Donath, C. Mayer, P. B. Frandsen, J.

Ware, T. Flouri, R. G. Beutel, O. Niehuis, M. Petersen, F. Izquierdo-Carrasco, T. Wappler,

J Rust10, A. J. Aberer, U. Aspöck, H. Aspöck, D. Bartel, A. Blanke, S. Berger, A. Böhm,

T. R. Buckley, B. Calcott, J. Chen, F. Friedrich, M. Fukui, M. Fujita, C. Greve, P. Grobe,

S. Gu, Y. Huang, L. S. Jermiin, A. Y. Kawahara, L. Krogmann, M. Kubiak, R. Lanfear, H.

Letsch, Y. Li, Z. Li, J. Li, H. Lu, R. Machida, Y. Mashimo, P. Kapli, D. D. McKenna, G.

Meng, Y. Nakagaki, J. L. Navarrete-Heredia, M. Ott, Y. Ou, G. Pass, L. Podsiadlowski, H.

Poh, B. M. von Reumont, K. Schütte, K. Sekiya, S. Shimizu, A. Slipinski, A. Stamatakis,

W. Song, X. Su, N. U. Szucsich, M. Tan, X. Tan, M. Tang, J. Tang, G. Timelthaler, S.

Tomizuka, M. Trautwein, X. Tong, T. Uchifune, M. G. Walzl, B. M. Wiegmann, J.

Wilbrandt, B. Wipfler, T. K. F. Wong, Q. Wu, G. Wu, Y. Xie, S. Yang, Q. Yang, D. K. Yeates,

K. Yoshizawa, Q. Zhang, R. Zhang, W. Zhang, Y. Zhang, J. Zhao, C. Zhou, L. Zhou, T.

Ziesmann, S. Zou, Y. Li, X. Xu, Y. Zhang, H. Yang, J. Wang, J. Wang, K. M. Kjer, X. Zhou,

Phylogenomics resolves the timing and pattern of insect evolution. *Science*. **346,** 763–767

(2014).

17. T. Niimi, H. Kuwayama, T. Yaginuma, Larval RNAi applied to the analysis of

postembryonic development in the ladybird beetle, *Harmonia axyridis*. *J. Insect Biotechnol.*

Sericol. **74,** 95-102 (2005).

18. M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet. journal.* **17,** 10-12 (2011).

19. M. G. Grabherr, B. J. Haas, M. Yassour, J. Z. Levin, D. A. Thompson, I. Amit, X.

Adiconis, L. Fan, R. Raychowdhury, Q. Zeng, Z. Chen, E. Mauceli, N. Hacohen, A. Gnirke,

N. Rhind, F. di Palma, B. W. Birren, C. Nusbaum, K. Lindblad-Toh, N. Friedman, A. Regev,

Trinity: reconstructing a full-length transcriptome without a genome from RNA-Seq data.

Nat. Biotechnol. **29,** 644-652 (2011).

20. C. Camacho, G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer, T. L.

Madden, BLAST+: architecture and applications. *BMC Bioinform*. **10,** 421 (2009).

21. Katoh, K. & Standley, D.M. *Mol. Biol. Evol*. **30**, 772-780 (2013).

22. S. Capella-Gutiérrez, J. M. Silla-Martínez, T. Gabaldón, trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*. **25**, 1972-1973

(2009).

23. S. Kumar, G. Stecher, M. Li, C. Knyaz, K. Tamura, MEGA X: molecular evolutionary

genetics analysis across computing platforms. *Mol. Biol. Evol.* **35**, 1547-1549 (2018).

24. A. Untergasser, I. Cutcutache, T. Koressaar, J. Ye, B. C. Faircloth, M. Remm, S. G.

Rozen, Primer3—new capabilities and interfaces. *Nucleic Acids Res*. **40,** e115 (2012).

25. K. J. Livak, T. D. Schmittgen, Analysis of relative gene expression data using real-time

quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods*. 25, 402-408 (2001).

26. V. R. Chintapalli, J. Wang, J. A. Dow, Using FlyAtlas to identify better *Drosophila*

melanogaster models of human disease. *Nat. Genet.* 39, 715-720 (2007).

Figure Legends

Figure 1: Effects of f-RNAi of *diap1, v-ATPase A* and *v-ATPase E* in *Henosepilachna*

vigintioctopunctata. (a) Tissue expression profile in 4th instar larva. nc, nerve cord; l, legs;

wd, wing discs; fb, fat body; mp, Malpighian tube; g, gut; c, carcass. (b) Quantitative RT-

PCR in the larvae two-days after f-RNAi. (c) Body-weight of the larvae two-days after f-

RNAi. (d) The leaves consumed by larvae two-days after f-RNAi. Scale, 1 cm. (e) Scheme

of f-RNAi experiment to evaluate feeding disorders. (f) Leaf area consumed by the larvae

for 0-24 hours (dark gray) and 24-48 hours (right gray) after f-RNAi. *P*-values were

calculated by Welch's *t*-test and adjusted by Holm's method for multiple comparisons.

Adjusted *P*-values were shown in c and f, and non-significances were not shown. Data of

b, c and f are means $\pm$ s.d., *n*=7 each gene RNAi larvae. Information are provided in Table

S1 for c and f.

Figure 2: Species-specificity and efficiency of *diap1* dsRNA. (a) Identity of *diap1*

nucleotide among insects. Numbers in each column indicates the identity between species.

Branches show the phylogenetic relationship[16]. (b) Image of sequence identity of *diap1*

dsRNAs among insects. Blue bars indicate the identical nucleotide sites between *H.*

vigintioctopunctata and each species. UTR, untranslated region; CDS, coding sequence;

BIR, baculovirus inhibitior of apoptosis protein repeat; RING, really interesting new gene.

(c) Area consumed by the larvae during 12-24 (upper) and 24-48 (lower) hours after various

insect-derived *diap1* dsRNAs ingestion. (d) Area consumed by the *diap1* f-RNAi larvae

treated with different amounts of dsRNA. Data of c and d are means $\pm$ s.d., *n*=3 in c and

n=4 in d. *P*-values were calculated by Welch's *t*-test and adjusted by Holm's method.

Adjusted *P*-values were shown in c and d, and non-significances were not shown. Amel,

Apis melifera; Blat, *Blatta lateraris*; Bmor, *Bombyx mori*; Caqu, *Catajapyx aquilonaris*;

Dmel, *Drosophila melanogaster*; Hvig, *Henosepilachna vigintioctopunctata*; Harm,

Helicoverpa armigera; Haxy, *Harmonia axyridis*; Ldec, *Leptinotarsa decemlineata*; Lmig,

Locusta migratoria; Ofas, *Oncopeltus fasciatus*; Oyez, *Oxya yezoensis*. Information for a,

c and d is provided in Tables S3, 4 and 5, respectively.

Figure 1

Figure 2