Testosterone-mediated behavior shapes the emergent properties of social networks

Roslyn Dakin[1,2], Ignacio T. Moore[3], Brent M. Horton[4], Ben J. Vernasco[3], T. Brandt Ryder[1,5]

[1]Migratory Bird Center, Smithsonian Conservation Biology Institute, National Zoological Park, Washington, DC 20013, USA
[2]Current address: Department of Biology, Carleton University, Ottawa, Ontario K1S 5B6, Canada
[3]Department of Biological Sciences, Virginia Tech, Blacksburg, Virginia 24061, USA
[4]Department of Biology, Millersville University, Millersville, Pennsylvania 17551, USA
[5]Current address: Bird Conservancy of the Rockies, Fort Collins, Colorado, 80525, USA

Correspondence: rydert@si.edu

Abstract

1. Social networks can vary in their organization and dynamics, with implications for ecological and evolutionary processes. Understanding the mechanisms that drive social network dynamics requires integrating individual-level biology with comparisons across multiple social networks.

2. Testosterone is a key mediator of vertebrate social behavior and can influence how individuals interact with social partners. Although the effects of testosterone on individual behavior are well established, no study has examined whether hormone-mediated behavior can scale up to shape the emergent properties of social networks.

3. We investigated the relationship between testosterone and social network dynamics in the wire-tailed manakin, a lekking bird species in which male-male social interactions form complex social networks. We used an automated proximity system to longitudinally monitor several leks and we quantified the social network structure at each lek. Our analysis examines three emergent properties of the networks: social specialization (the exclusivity of partnerships), network stability (the overall persistence of partnerships through time), and behavioral assortment (the tendency for like to associate with like). All three properties are expected to promote the evolution of cooperation. As the predictor, we analyzed the collective testosterone of males within each social network.

4. Social networks that were composed of high-testosterone dominant males were less specialized, less stable, and had more negative behavioral assortment, after accounting for other factors. These results support our main hypothesis that individual-level hormone physiology can predict group-level network dynamics. We also observed that larger leks

38   with more interacting individuals had more positive behavioral assortment, suggesting

that small groups may constrain the processes of homophily and behavior-matching.

40   5. Overall, these results provide evidence that hormone-mediated behavior can shape the

broader architecture of social groups. Groups with high average testosterone exhibit

42   social network properties that are predicted to impede the evolution of cooperation.

44   **Keywords:** androgens, behavioral endocrinology, collective behavior, cooperation, dynamic

networks, social networks, testosterone

Introduction

Behavioral interactions are the foundation of social network structures that can vary through time, among populations, and across species. Network structures play an important role in many ecological and evolutionary processes, including the spread of diseases (Sah *et al.* 2018), the transmission of information and resources (Aplin *et al.* 2012), and selection on individual behavior (Ohtsuki *et al.* 2006). A major challenge is understanding how individual-level factors, such as physiological and behavioral mechanisms, scale up to drive the emergent structural properties of social groups (Krause & Ruxton 2002). Linking these two levels of analysis is difficult because it requires integrating individual-level data with repeated measures of entire social groups (Sah *et al.* 2017).

Here, we use a comparison of multiple social networks through time to investigate how hormone-mediated behavior shapes the higher-order structure of social networks. Testosterone is a steroid hormone that is well known for its influence on social behavior and its sensitivity to changes in the social environment (Wingfield *et al.* 1990; Adkins-Regan 2005; Goymann 2009). Testosterone often promotes physical aggression and other behaviors associated with social dominance (Oyegbile & Marler 2005; Fuxjager *et al.* 2010). Testosterone can also promote status-seeking behaviors in a non-aggressive context, including cooperative and gregarious behavior (Eisenegger *et al.* 2011; Boksem *et al.* 2013; Ryder *et al.* 2020). Overall, these hormone-signaling pathways are essential for the development and modulation of complex behavioral phenotypes (Cohen *et al.* 2012).

To investigate how testosterone is associated with social network dynamics, we studied the wire-tailed manakin (*Pipra filicauda*), a bird species in which the males engage in coordinated displays with each other at sites known as leks. In *P. filicauda*, males exhibit two

social status classes: the dominant males who hold territories on the leks, and subordinate,

70   "floater" males who must acquire a territory on a lek before they can mate (Heindl 2002; Ryder

et al. 2011a). Male social partnerships in wire-tailed manakins can be remarkably stable and

72   typically occur between two unrelated males, most commonly, but not exclusively, between a

territory-owner and a floater (Ryder *et al.* 2011a; Dakin & Ryder 2020). Two features of the

74   manakin leks make them especially well-suited to studying the relationship between hormones

and group-level social structure. First, testosterone is known to affect the social behavior of

76   individual male wire-tailed manakins (Ryder *et al.* 2020). Second, these partnerships among

males form the basis of complex and dynamic social networks that are replicated across leks,

78   facilitating a comparative approach (Dakin & Ryder 2018, 2020). Given this background, we

sought to test whether testosterone and its effects on male behavioral phenotype could drive the

80   emergent properties of the social network.

The broader function of male-male social behavior and coordinated displays in the

82   manakin family has been the subject of considerable study (e.g., Prum 1994; DuVal 2007;

McDonald 2007; Ryder *et al.* 2008, 2009; Díaz-Muñoz *et al.* 2014). One possible function is that

84   dyadic displays may be competitive and/or they may serve to maintain an individual's position in

a dominance hierarchy (Prum 1994; Heindl 2002). Social interactions may also represent

86   cooperative coalitions that provide benefits to both parties (Ryder *et al.* 2008, 2009). The

potential competitive and cooperative functions of male-male social behavior are not mutually

88   exclusive, and function may be context- and/or status-specific (Ryder *et al.* 2008). Coordinated

displays may also be a vestige of ancestral cooperative behavior (i.e., the behavior may have

90   been directly beneficial to both parties in the past, and it persists today, even if it no longer has

adaptive benefits) (Prum 1994). Given this background and recent evidence that testosterone

92   modulates male social behavior in wire-tailed manakins (Ryder *et al.* 2020), we focused this

study on three emergent properties of the social networks formed by male-male interactions that

94   can influence the evolution and maintenance of cooperation (Fig. 1).

The first property we examined, social specialization, seeks to capture the exclusivity of

96   the relationships between partners in a social network. In the context of manakin behavior,

highly specialized networks exhibit a high frequency of exclusive social partnerships between

98   males of different status classes, as illustrated in Fig. 1A. In humans, social specialization has

been found to maximize the ability of a team to successfully perform a challenging task (Jehn &

100   Shah 1997). In manakins, we expect specialization to improve the familiarity of social partners

and the behavioral coordination of their displays. Greater specialization is also expected to

102   minimize conflict over mating and territorial ascension opportunities (Schjelderup-Ebbe 1922;

McDonald 1993).

104   The second property, network stability (Fig. 1B), quantifies the average persistence of

social partnerships through time (Poisot *et al.* 2012; Dakin & Ryder 2020). Coordinated displays

106   in manakins require the synchronization of complex behaviors, and previous empirical work

indicates that longer partnership tenure has a positive effect on display coordination (Trainer &

108   McDonald 1995; Trainer *et al.* 2002). Greater temporal stability of partnerships also increases

the opportunity for familiarity and reciprocity within a social network (Trivers 1971; Roberts &

110   Sherratt 1998; Croft *et al.* 2006).

The third property, behavioral assortment (Fig. 1C), captures the extent to which males

112   interact with partners who express similar behaviors (i.e., is like associated with like? (Croft *et*

al. 2006; Farine 2014)). At the proximate level, positive assortment may represent the outcome

114   of generalized reciprocity and/or partner choice (Fowler & Christakis 2010; Dakin & Ryder

2018). At the ultimate level, positive assortment has also been shown to promote the evolution of

116   cooperation (Ohtsuki *et al.* 2006). To quantify the overall behavioral assortment of the manakin

networks, we focused on two correlated metrics of social behavior within the network: "strength"

118   (a male's frequency of daily social interactions) and "degree" (his daily number of social

partnerships; (Dakin & Ryder 2018)). We used a composite measure of assortment that averaged

120   the assortativity indices of these two phenotypes.

As a potential predictor of specialization, stability, and assortment, we quantified the

122   collective testosterone of the manakin leks (Fig. 2; (Akinola *et al.* 2016)). Because hormone-

behavior relationships are status-specific in manakins and many other species (Eisenegger *et al.*

124   2011; Boksem *et al.* 2013; Ryder *et al.* 2020), we analyzed collective testosterone for each of the

two status classes separately. High individual testosterone levels are associated with reduced

126   sociality in dominant males, but increased sociality in subordinate males (Ryder *et al.* 2020). We

therefore predicted that the average (or collective) testosterone of territorial males in a group

128   would be negatively associated with its specialization, stability, and assortment. In contrast, we

predicted that the collective testosterone of floater males in a group would be positively

130   associated with these three emergent properties of the social network.

132   **Materials and Methods**

Study Population

134   We studied wire-tailed manakins (*Pipra filicauda*) at the Tiputini Biodiversity Station in

Orellana Province, Ecuador (0º 38' S, 76º 08' W). This population of *P. filicauda* has been

136   studied and individuals color-banded annually since 2002 (e.g., Ryder *et al.* 2008, 2009). The

present study was conducted on 11 leks during peak breeding activity (December-March) across

138   three field seasons: 2015-16, 2016-17 and 2017-18. All research was approved by the

Smithsonian ACUC (protocols #12-23, 14-25, and 17-11) and the Ecuadorean Ministry of the

140   Environment (MAE-DNB-CM-2015-0008).

142   ***Testosterone Assessment***

Male manakins were captured using mist-nets on the leks as described in (Ryder *et al.* 2020). We

144   deployed up to 16 mist-nets simultaneously at a given lek, with the intention of capturing every

male on the lek. We rotated the nets between leks with the goal of capturing each male up to

146   three times per field season. Each mist-net was checked on a 30-minute schedule with variation

resulting from capture rate (i.e., multiple birds being caught on the same net run; Vernasco *et al.*

148   2019; Ryder *et al.* 2020). The amount of time a male spends in the net has a subtle, but

significant, negative effect on his circulating testosterone (Vernasco et al. 2019). Therefore, we

150   used video monitoring to determine the duration of time that each bird was in the mist-net prior

to blood sampling, so that it could be accounted for in further analyses (mean = 17.5 minutes, SD

152   = 10.3 minutes, range = 1–72 minutes). Following removal of a male from the mist-net, a small

blood sample (< 125uL) was taken from the brachial vein and stored on ice prior to being

154   centrifuged at 10,000 rpm for 5 min, as described in previous studies (Ryder *et al.* 2011b;

Vernasco *et al.* 2019; Ryder *et al.* 2020). Plasma volume was measured to the nearest 0.25 ul and

156   stored in 0.75 ml of 100% ethanol (Goymann *et al.* 2007). Testosterone was double extracted

from the plasma using dichloromethane. Following extraction, a direct radioimmunoassay was

158   used to measure the total plasma androgen concentration (ng/ml) adjusted by the extraction

efficiency and plasma volume of each sample (Eikenaar *et al.* 2011; Ryder *et al.* 2011b).

160   Hormone assays were conducted annually, and the detection limits were 0.12, 0.08, and 0.09

ng/ml for 2015-16, 2016-17 and 2017-18, respectively; any sample that fell below the assay-

162   specific limit of detection was assigned that limit as its testosterone concentration as a most

conservative estimate. The extraction efficiency for all samples was between 62-73%, and the

164   intra-assay coefficients of variation were 6.6%, 11.6%, and 9.2% for 2015-16, 2016-17 and

2017-18, respectively; the inter-assay coefficient of variation was 19.5%.

Behavioral Recording

168   We used an automated data-logging system to monitor male-male interactions on the display

territories of the leks (Ryder *et al.* 2012; Dakin & Ryder 2018; Ryder *et al.* 2020). The territories

170   on these leks are specifically used for male-male coordinated social displays (as described in

Schwartz & Snow 1978). At the beginning of each field season, male manakins were outfitted

172   with coded nano-tags (NTQB-2, Lotek Wireless; 0.35 g). The tags transmitted a unique VHF

signal ping once per 20 s for three months. In total, 296 tag deployments were performed on 180

174   individuals (mean 1.7 field seasons per male ± SD 0.7), 178 of whom also had hormone data

(mean number of hormone samples per male = 3 ± SD 1.5). Approximately 10 days (± SD 7)

176   after tagging and sampling was completed at a given lek, a proximity data-logger (SRX-DL800,

Lotek Wireless) was deployed within each display territory at the lek to record tagged males

178   within a detection radius of 30 m (a distance that corresponds to the typical diameter of a

manakin display territory (Heindl 2002; Dakin & Ryder 2018)).

180   Proximity recording sessions ran from 06:00 to 16:00 for ~6 consecutive days (± SD 1

day) and were performed ~3 times per field season at a given lek. Occasionally, the length of a

182   recording session was extended due to extenuating circumstances such as inclement weather.

The recording sessions were scheduled to be distributed evenly throughout each field season at

184   each lek, to minimize any confounding seasonal effects. Each recording session represents an

observation of the social network at a given lek. In total, we conducted 86 recording sessions

186   (29,760 data-logger hours) representing repeated measures of the social activity at 11 leks during

three field seasons (see Fig. S1 in the supplement for additional details on the sampling regime).

188   Prior to data-logger deployment, each territory was also observed on non-recording days

to identify the territory-holder based on his color-bands, following previous studies (Ryder *et al.*

190   2008, 2009). These status assignments were subsequently verified in the proximity data.

192   ***Data Processing and Statistical Analysis***

All computational and statistical analyses were performed in R (R Core Team 2018). Network

194   illustrations were made using the igraph package (Csardi *et al.* 2018).

196   ***Social Interactions***

Social networks were constructed by first defining interactions between two males that occurred

198   on the display territories. To do this, a computational algorithm was used to identify joint

detections, wherein two males were located at the same display territory within a pre-defined

200   spatial and temporal threshold (Ryder *et al.* 2008, 2012; Dakin & Ryder 2018). For the temporal

threshold, the two males had to occur < 45 s apart. This temporal threshold was chosen to allow

202   for the fact that each tag pinged with a 20 second pulse rate, such that overlapping individuals

could have up to a 40 s gap between their respective pings. For the spatial threshold, the two

204   males had to have a difference in received signal strength values (Δ RSSI) < 10. This threshold

corresponded to a typical distance between 0 to 5 m apart in a ground-truthing experiment

206   (Dakin & Ryder 2018). Hence, according to our definition, a social interaction is initiated only

after males come within this approximate spatial threshold. We chose this spatial threshold

208 because it is close enough to permit visual and acoustic contact during typical social behaviors

(such as those described in Schwartz & Snow 1978). After completing our study, we also

210 performed a sensitivity analysis to verify that our main results were robust to alternative spatial

threshold definitions (see supplement for details).

212 Any repeated co-occurrence of the two males within 5 minutes was considered to be part

of the same social interaction, but after a gap of $\geq$ 5 minutes, it was considered to be a new

214 interaction between those two males. The average duration of social interactions defined by this

method was 5.1 min $\pm$ SD 13.3, 8.8 min $\pm$ SD 20.1, and 7.0 min $\pm$ SD 20.5 in the three respective

216 field seasons, further indicating that these were sustained social interactions, rather than random

encounters.

218 In total, we identified 36,885 social interactions over the three field seasons of study.

These interactions were used to define a weighted social network for each lek recording session.

220 The nodes in the network were the individual males, and the edges were weighted by the number

of social interactions between each pair of males. An earlier validation study compared social

222 interactions that were detected by the proximity system with those that were directly observed

for 11 males (Ryder *et al.* 2012), and confirmed that all of the interactions detected by the

224 proximity system were also directly observed.

226 ***Null Model Validation of the Social Networks***

Two broad classes of methods have been described for building social networks: (i) networks

228 that are built based purely on the proximity of animals unrelated to their behavioral context

("gambit of the group"), and (ii) networks that are built based on specific behavioral criteria that

230   are directly observed (Franks *et al*. 2010; Croft *et al*. 2011; Farine 2015). Although the

interactions in our study were not directly observed, they were recorded at specialized display

232   perches that have a known function in male-male social interactions (Schwartz & Snow 1978;

Ryder *et al*. 2011a). Hence, they do not qualify as the gambit of the group. To further

234   demonstrate this point, we performed a pre-network permutation of the raw data to determine

whether the observed network edges occurred more often than expected by chance (Farine *et al*.

236   2017). This analysis was highly conservative in that it preserved key features of the data

including male visit rates to specific territories within each recording session and lek; it is

238   described in detail in the supplement and Fig. S2. The results demonstrated that 95% of the

observed network edges had a greater edge weight than expected under stringent permutation

240   conditions. Moreover, these preferred edges had edge weights that were 10-fold to 50-fold

greater than expected by chance under these stringent conditions (Fig. S2). This provides an

242   additional validation to our approach, because it indicates that the observed network edges were

nearly always preferred, even relative to other possible interactions within the same lek. These

244   results are also consistent with a previous validation study indicating that these methods capture

male-male coalition partnerships that are directly observed (Ryder *et al*. 2012).

246   Although pre-network data permutations are sometimes used to derive adjusted

association indices, or to prune networks prior to further analysis, we did not take this approach

248   for several reasons. First, the statistical rarity of a relationship in our system does not *a priori*

define the importance of any one social interaction, especially given that all of the interactions

250   occurred on display territories with specialized function. A single interaction between two rarely

interacting males may have been highly consequential (e.g., if one of those males had a highly

252   influential hormone-behavioral phenotype). Conversely, a single interaction between two

frequently interacting males may have been relatively unimportant. It would be unwarranted to

254   assume that partnership rarity indicates the importance of any single interaction in this context.

Second, and perhaps more importantly, the goal of this study was a comparison across multiple

256   networks. Some networks are genuinely less preferential, and more random, than others. If we

modified the network edges based on permutation-based indices, it would disproportionately

258   prune the truly random networks, introducing a source of bias that would be contrary to our main

goal. Hence, all further analyses are based on networks where the edge weights are given by the

260   observed number of interactions, as this method is most appropriate for our study system and

aims. Below, we also describe a separate node-label permutation that provided an additional

262   check on our final statistical analyses.

264   ***Social Specialization***

To quantify social specialization, we sought a metric that would capture the extent to which a

266   network was partitioned into exclusive social relationships (as opposed to a network made up of

non-specific or non-exclusive partnerships). To do this, we used a network metric of

268   specialization that is commonly used in community ecology called H_2' (Blüthgen *et al.* 2006).

An advantage of H_2' is that it is standardized against a theoretical maximum, based on the

270   overall activity levels of different nodes and Shannon entropy (Blüthgen *et al.* 2006); this makes

it possible to compare the extent of specialization across different bipartite networks in a

272   standardized way. To apply this metric to our manakin data, we converted each lek's social

network into its bipartite adjacency matrix (Fig. 1A), with floaters along one axis, and territory-

274   holders on the other, and then calculated social specialization as H_2' using the bipartite package

(Dormann *et al.* 2019). Higher values of specialization indicate that the network is made up of

276   more exclusive relationships, as illustrated in Fig. 1A. We chose to focus on floater-territorial

specialization because these two social classes are well-defined and floater-territorial

278   partnerships tend to be the most common in this species (Fig. S3 and (Ryder *et al.* 2011a)).

280   *Network Stability*

To quantify the stability of social relationships, we compared each lek's social network from a

282   given recording session to its subsequent recording session within the same field season (Fig.

1B). Network stability was then calculated as the number of male-male partnerships (binary

284   network edges) shared by both time points divided by the number of partnerships at either time

point (Dakin & Ryder 2020). Higher values of stability indicate greater persistence of social

286   relationships within the network, independent of any changes in the representation of particular

males (nodes) (Poisot *et al.* 2012). To focus this measure on the persistence of strong

288   relationships, we computed network stability of partnerships that occurred at least six times

following (Dakin & Ryder 2020). The threshold of six was chosen because it corresponds to an

290   average rate of one social interaction per day in our data. We conducted an additional sensitivity

analysis to verify that alternative thresholds for the stability calculation (greater or less than six)

292   did not change our main results (see supplement for details). Previous work using this metric of

stability has shown that the wire-tailed manakin social networks are far more stable than

294   expected by chance (Dakin & Ryder 2020).

296   *Behavioral Assortment*

Assortment refers to the extent to which individuals associate with similar partners (Fig. 1C); it

298   can be due to partner choice (homophily), shared environments, or the social transmission of

behavior (Dakin & Ryder 2018). Assortment was quantified using Newman's assortativity index,

300   which is a correlation coefficient for the statistical association among linked nodes within a

network. It ranges from –1 (a negative association), through 0 (no association), and up to +1 (a

302   positive association). To quantify the assortment of social behaviors, we first determined the

daily frequency of two behaviors for each male: his number of social interactions per day

304   (strength), and his number of unique social partnerships per day (degree). Strength and degree

are both repeatable measures of a male's social behavior in our study population (Dakin & Ryder

306   2018). We used the average log-transformed values of each male's strength and degree within

the recording session, and then calculated the assortativity coefficient for the entire social

308   network using the algorithm for weighted networks in the assortnet package (Farine 2016).

Because assortativity values for strength and degree were highly correlated (Pearson's $r = 0.78$, p

310   < 0.0001, $n = 86$ networks), we took the average of these two values as the measure of overall

behavioral assortment within the social network. Note that we used log-transformed values of

312   strength and degree because these two variables are strongly positively skewed (Dakin & Ryder

2018; Ryder *et al.* 2020), and assortativity is based on a Pearson's correlation. Finally, we also

314   computed the assortativity of the two discrete status classes (floater and territory-holder), to

ensure that our analysis of behavioral assortment was not solely driven by status-assortment.

Collective Testosterone

318   To understand how hormones might predict network properties, we derived a measure of

collective testosterone of each social network. This was based on the hormonal trait that was the

320   best predictor of social behavior in our previous study, referred to as "mean testosterone" (Ryder

et al. 2020). A male's mean testosterone is his average residual circulating testosterone. It is

322   calculated using a linear regression of log-transformed testosterone to statistically account for the

        effects of field season, Julian date, time of day when captured, and duration of restraint, all of

324   which may influence point estimates of a male's baseline hormone level (Vernasco *et al.* 2019).

        Hence, mean testosterone represents a standardized measure of a male's circulating testosterone,

326   independent of his capture conditions (Fig. 2A). Next, to determine collective testosterone, we

        took the average mean testosterone for each social network, weighted by the interaction

328   frequency (strength) of the males within the network (Fig. 2B). Collective testosterone is thus a

        group-level characteristic that is weighted towards the males that made the greatest contribution

330   to group social structure. In other words, networks with low collective testosterone are made up

        of mostly low-testosterone individuals, whereas networks with high collective testosterone are

332   made up of mostly high-testosterone individuals. We calculated collective testosterone separately

        for each status class, because the effects of hormones on social behavior are status-dependent

334   (Eisenegger *et al.* 2011; Boksem *et al.* 2013; Ryder *et al.* 2020).

336   ***Statistical Analysis***

        To evaluate the hypothesis that collective testosterone predicts network properties, we analyzed

338   mixed-effects models of the social network properties in the lme4 package (Bates *et al.* 2018).

        The three response variables were social specialization, network stability, and behavioral

340   assortment (Fig. 1). We used Akaike's Information Criterion (AIC) to compare four candidate

        models for each response variable, as follows: (1) collective testosterone of territory-holders +

342   collective testosterone of floaters; (2) collective testosterone of territory-holders; (3) collective

        testosterone of floaters, and (4) no testosterone predictors. All of the models included additional

344   fixed effects to account for field season (a categorical variable with three levels), the average

Julian date of the recording session, the average number of recorded hours per territory, and the

346   size of the social network (number of individuals), as well as a random effect of lek to account

        for repeated measures.

348   Model comparison was performed on models fit with maximum likelihood, and the best-

        fit models were re-fit using restricted estimation of maximum likelihood (REML) to derive

350   parameter estimates (Zuur *et al.* 2009). We used the lmerTest package to compute p-values for

        parameter estimates in the mixed-effects models based on Satterthwaite's method (Kuznetsova *et*

352   *al.* 2018). We also report Nakagawa and Schielzeth's R^2_{LMM} values as an estimate of effect size

        (Nakagawa & Schielzeth 2013). We verified that all models met the assumptions of linear

354   regression analyses. First, we checked that the Pearson residuals met the assumption of

        normality. We visually inspected the partial residual plots for each fixed and random effect, to

356   confirm that there were no outliers or departures from the assumption of homoscedasticity. To

        check for multicollinearity, we used the performance package (Lüdecke *et al.* 2019) to calculate

358   variance inflation factors (VIFs), and we verified that all VIFs were between $1 - 1.8$.

360   ***Sample Sizes and Exclusions***

        In two field seasons (2016-17 and 2017-18), we performed an experiment as part of a separate

362   study to test the influence of transiently-elevated testosterone on individuals (n = 5 individuals in

        2016-17 and n = 4 in 2017-18 (Ryder *et al.* 2020)). The results of that experiment demonstrated

364   that elevated testosterone caused a temporary decrease in the frequency and the number of social

        partnerships in the altered males (Ryder *et al.* 2020). It is important to note that this experiment

366   was not designed to test emergent properties at the network level, because it was conducted on a

        limited scale with only one or two individuals temporarily altered within each lek. We therefore

368   excluded the 6 post-manipulation networks from the main analysis in this study. We verified that

when we included these manipulated leks, all of our main conclusions were unchanged.

370   After excluding the 6 post-manipulation observations, 80 of the original 86 recording

sessions remained. Table 1 summarizes the sample sizes for the network-level analyses. In 10

372   cases, specialization could not be calculated because the bipartite network did not have sufficient

data to determine H_2'. Stability could not be calculated in 26 cases, when either the recording

374   session occurred at the end of a field season, or when there were insufficient partnerships that

met the criteria for the stability calculation.

Node-label Permutation Analysis

378   To evaluate the possibility that our results could be influenced by other properties of the leks that

were independent of hormonal traits, we also performed a statistical permutation of the post-

380   network data (Farine *et al.* 2017). The purpose of this analysis was to verify that the results were

driven by (and sensitive to) the relative contribution and social position of different males. To do

382   this, we performed a statistical permutation that randomized the node labels (male IDs) within

each of the social networks, retaining network topology, and leaving each male's testosterone

384   traits unchanged. Hence, this analysis preserved which males were present in which recording

session, but it randomized the relative position and contribution of each male. After generating

386   1,000 of these node-label permutation datasets, we recalculated specialization, stability, and

assortment, and then refit the top models from our mixed-model analysis. We then compared the

388   slope estimates from the observed data to those derived from 1,000 node-label permutations. As

a one-sided p-value, we calculated the proportion of slope estimates from the node-label

390   permutations that were more negative than the corresponding estimate from the observed data.

Results

392 The lek social networks had 14 males on average, but there was considerable variation among

leks ranging from 3 to 43 individuals (Fig. S1). Additional descriptive statistics are provided in

394 Table 1. Although network size has a lognormal distribution (Fig. S1), all other emergent

properties of the social structure were approximately normally distributed (Fig. S4).

Table 1. Descriptive statistics for manakin social networks. Means and standard deviations
398 (SD) are provided for each field season. The bottom row provides the sample sizes for the
number of social networks analyzed (n_{obs}) at each lek (N_{lek}). Additional data is provided in the
400 supplement in Fig. S1 and Fig. S4.

Field season	Network size (# birds)	Social specialization	Network stability	Behavioral assortment	Collective T (terr.)	Collective T (floa.)
15-16	14.5 (8.9)	0.50 (0.22)	0.43 (0.25)	0.01 (0.36)	0.44 (0.42)	–0.31 (0.56)
16-17	17.8 (10.5)	0.30 (0.24)	0.47 (0.20)	0.06 (0.30)	0.49 (0.24)	–0.22 (0.37)
17-18	10.3 (4.5)	0.42 (0.30)	0.42 (0.27)	0.02 (0.37)	0.43 (0.40)	–0.70 (0.46)
Sample size	$n_{obs} = 80$	$n_{obs} = 70$	$n_{obs} = 54$	$n_{obs} = 80$	$n_{obs} = 80$	$n_{obs} = 80$
	$N_{lek} = 11$	$N_{lek} = 11$	$N_{lek} = 11$	$N_{lek} = 11$	$N_{lek} = 11$	$N_{lek} = 11$

402 We found that the collective testosterone of territorial males could predict all three emergent

properties of the social networks (specialization, stability, and assortment). The leks with greater

404 representation of high-testosterone territorial males were less specialized, less stable, and more

negatively assorted (Fig. 3; all $p \leq 0.03$ in mixed-effects models). The slope coefficients for these

406 three relationships were also all greater than expected under a node-label permutation (inset

panels, Fig. 3; all $p < 0.02$). Model selection results for the observed data indicated considerable

408 uncertainty in the best-fit model for each of the three network properties analyzed (Table S1).

However, the collective testosterone of territory-holders was a significant predictor in all of the

410 best-supported models (Table S2). In contrast, the collective testosterone of floater males was

not a significant predictor of network properties in any of the best-fit models (Table S2).

412   We did not detect any significant effects of Julian date on network properties within our

study period, but we did observe some significant year-to-year differences (e.g., social

414 specialization and behavioral assortment in Table S2). Assortment was the only property that

was significantly related to network size and recording effort (Table S2). All else being equal,

416 behavioral assortment was more positive in larger networks, and it was more negative in

networks that had longer recording sessions.

418   $R^2_{LMM(m)}$ provides an estimate of the proportion of variance explained by the fixed effects

in a model (Table S1). The $R^2_{LMM(m)}$ for assortment in the best-fit model was 0.52. This indicates

420 that about 52% the variation in behavioral assortment could be explained by the combined

associations with collective testosterone, network size, sampling effort, and annual variation

422 (Table S2). The $R^2_{LMM(m)}$ for stability was 0.12, indicating that collective testosterone and the

other predictors (network size, sampling effort, year) could explain about 12% the variation in

424 that metric (Table S2). Finally, for specialization, the $R^2_{LMM(m)}$ indicated that the combined

effects of testosterone and these other sources of variation together explained about 20% of the

426 variation (Table S2).

428 **Discussion**

The collective testosterone of the dominant, territory-holding males within a lek was associated

430 with multiple emergent properties of the social network (Fig. 3). Variation in collective

testosterone is a function of both the number of high-testosterone males and their frequency of

432 social interactions. Our results indicate that the hormone-mediated behavior of these individuals

may affect all three social network properties of specialization, stability and assortment. This

434 indicates that the effects of testosterone on dominant males may mediate an extended phenotype

with the power to shape social structure (Dawkins 1982). In contrast, the collective testosterone

436   of floater males was not significantly associated with any of the emergent network properties

(Table S2).

438   How can the relationship between hormone levels and network properties be explained in

terms of individual mechanisms? Given that testosterone has antagonistic effects on the sociality

440   of territorial males (Ryder *et al.* 2020; Vernasco *et al.* 2020), we hypothesize that the behavior of the

high-testosterone individuals can cause several features of the social organization to break down.

442   Our previous results showed that high-testosterone dominant individuals have a reduced ability

to attract and maintain social partners (Ryder *et al.* 2020). We propose that this weakening of

444   social relationships may cause floater males to prospect elsewhere for new partners, negatively

affecting both the stability and specialization of the lek network as a whole. Likewise, high-

446   testosterone dominant males may inhibit the processes of social contagion, reciprocity, and/or

behavioral matching that can cause positive behavioral assortment (Dakin & Ryder 2018).

448   Testing these ideas will require direct observation of the individual social behaviors that occur

within dyads, and how these behaviors change through time. Because our current data cannot

450   assess the fine-scale valence of social interactions, further studies are needed that combine direct

observation with high-throughput data on social network dynamics.

452   We included several additional parameters in our analyses to account for social network

size and sampling effort. Although it was not one of our main hypotheses, we noted that network

454   size was positively associated with behavioral assortment (Table S2). In other words, males were

more likely to associate with behaviorally similar partners in larger leks, whereas they were more

456   likely to associate with dissimilar partners in smaller leks. Effects of group size on assortment

have been noted in a few other studies, although the form of this relationship varies (Griffiths &

458   Magurran 1997; Ilmarinen *et al.* 2017; McDonald *et al.* 2017). In manakins and other lekking

systems, larger leks are known to have heightened display activity and higher female visitation

460   rates (Lank & Smith 1992; Durães *et al.* 2009). This raises the possibility that social facilitation

and heightened activity may be associated with increased homophily and/or behavioral matching

462   (either through partner choice, contagion, or reciprocity). Another plausible explanation is that

smaller social groups may constrain these behavioral processes, by making it more difficult to

464   find or match a suitable partner. This hypothesis could be explored in experiments on captive

systems and simulation models.

466   Behavioral assortment was also more positive in networks that were recorded for less

time in our study (in other words, leks where males associated with behaviorally similar partners

468   tended to be recorded for fewer days). Recording time was designed to be approximately even

among leks (Fig. S1), with the most common reason for an extended recording duration being

470   inclement weather that extended the recording session. Therefore, we speculate that inclement

weather may have reduced the amount of behavioral matching (assortment) while also affecting

472   recording time. Although we did not collect data on weather at each lek as this was not our main

goal, the question of how inclement weather affects group-level social dynamics is an interesting

474   one that merits further study.

       Conducting experimental tests of causation at the level of whole social networks remains

476   a major challenge in ecology (Pinter-Wollman *et al.* 2014; James *et al.* 2009). Although our

node-label permutation analysis provides evidence that our results are not merely due to

478   structural differences among networks, we cannot rule out the possibility that high-testosterone

individuals chose to participate in certain networks due to other factors that may also influence

480   the emergent properties of the network (e.g., environmental quality and/or female activity). It is

important to note that in this system, as in many other wild animals, controlled experimental

482   manipulations of the broader social network structure are not yet possible (Zyphur *et al.* 2009;

Akinola *et al.* 2016). Nevertheless, our data here indicate that the increased prevalence of

484   dominant, high-testosterone individuals can predict changes social dynamics and subsequent

higher-order network structure. These findings establish that hormone-behavior relationships are

486   not limited to one individual, but instead hormones have population-level consequences

(McClintock 1981; Robinson 1992). Taken together with previous studies, we propose that

488   testosterone-mediated behavior can alter social network dynamics in ways that impede the

evolution of stable social relationships and cooperation.

Figure Legends

Figure 1. Example social networks illustrating social specialization, network stability, and behavioral assortment. (A) Social specialization is measured using the bipartite form of the
social network. In the example on the left, the associations between floater males and territory-holders are poorly partitioned, creating a network with a relatively low specialization. On the
right, there is greater partitioning, such that each floater male maintains a smaller number of associations with specific territory-holding males. (B) Network stability is measured by
evaluating the persistence of partnerships from one recording session (t_1) to the next (t_2). The example at the top of (B) has a lower stability than the example on the bottom. (C) Behavioral
assortment measures the tendency of like to associate with like. In (C), nodes are shaded to indicate a continuous gradient of daily behavior (in this case, each male's average daily strength
is shown). In the example network on the left, the males tend to associate with behaviorally dissimilar partners, yielding a negative behavioral assortment. In the example on the right, the
males tend to associate with behaviorally similar partners, yielding a positive assortment. See Table 1 for additional data.

        Figure 2. Collective testosterone of manakin social networks. (A) Circulating testosterone
508   varies among individual males. This graph shows repeated measures from 210 individual male manakins, sorted along the x-axis by mean testosterone, a standardized measure of a male's
510   average circulating testosterone level (Ryder *et al.* 2020). Opacity is used to denote the two male status classes (with subordinate floaters colored semi-transparent, and dominant territory-holders
512   colored opaque). Color ramping is used to denote each individual's hormone phenotype. (B-D)

These data were used to define collective testosterone in the present study. Node colors indicate an individual's mean testosterone following the scale in (A). The collective testosterone of each social network is calculated as the average of the individual hormone phenotypes, weighted by strength, and is shown below each example. Collective testosterone was determined for each status class separately. See Table 1 for additional data.

Figure 3. Collective testosterone predicts the structure of the social network. The networks with higher-testosterone dominant males were less specialized, less stable over time, and had more negative behavioral assortment. Each plot shows the partial residuals from a statistical analysis that also accounts for field season, the average Julian date of the recording session, the number of recording hours, network size, and lek identity. Because floater and territorial males can differ in behavior, the analysis of behavioral assortment also accounted for status assortment within each network. Different symbols are used to indicate repeated measures of 11 different leks, colored according to the collective testosterone scale in Fig. 2. Inset panels show the results of node-label permutation tests. In each case, the slope in the best-fit model (solid vertical line) is significantly more negative than expected based on the distribution of permuted estimates (grey distribution, dotted line is at 0). See Tables S1-S2 for additional data.

Acknowledgments

We thank Camilo Alfonso, Brian Evans, David and Consuelo Romo, Kelly Swing, Diego

Mosquera, Gabriela Vinueza, and the Tiputini Biodiversity Station of the Universidad San

Francisco de Quito. Funding was provided by the National Science Foundation (NSF) IOS

1353085 and the Smithsonian Migratory Bird Center.

Author Contributions

RD, ITM, BMH, and TBR designed the study. ITM, BMH, BJV, and TBR collected the data.

RD, ITM, and TBR analyzed the data. RD and TBR wrote the manuscript. All authors edited the

manuscript.

Declaration of Interests

The authors declare no competing interests.

Data Availability

All data and R scripts necessary to reproduce this study are available for download at:

https://figshare.com/s/13a311662fee686fa4f3

Supporting Information

Supplemental text, Figures S1-S5, and Table S1-S3 in the attached PDF

References

552   Adkins-Regan, E. (2005). *Hormones and animal social behavior*. Princeton University Press.

Akinola, M., Page-Gould, E., Mehta, P.H. & Lu, J.G. (2016). Collective hormonal profiles predict
554   group performance. *PNAS*, 113, 9774–9779.

Aplin, L.M., Farine, D.R., Morand-Ferron, J. & Sheldon, B.C. (2012). Social networks predict
556   patch discovery in a wild population of songbirds. *Proceedings of the Royal Society B: Biological Sciences*, 279, 4199–4205.

558   Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R.H.B., Singmann, H., *et al.* (2018). *lme4 1.1-18-1: linear mixed-effects models using "Eigen" and S4*.

560   Blüthgen, N., Menzel, F. & Blüthgen, N. (2006). Measuring specialization in species interaction networks. *BMC Ecology*, 6, 9.

562   Boksem, M.A.S., Mehta, P.H., Van den Bergh, B., van Son, V., Trautmann, S.T., Roelofs, K., *et al.* (2013). Testosterone Inhibits trust but promotes reciprocity. *Psychol Sci*, 24, 2306–2314.

564   Cohen, A.A., Martin, L.B., Wingfield, J.C., McWilliams, S.R. & Dunne, J.A. (2012). Physiological regulatory networks: ecological roles and evolutionary constraints. *Trends in Ecology & Evolution*, 27, 428–435.
566

Croft, D.P., James, R., Thomas, P.O.R., Hathaway, C., Mawdsley, D., Laland, K.N., *et al.* (2006).
Social structure and co-operative interactions in a wild population of guppies (*Poecilia reticulata*). *Behav Ecol Sociobiol*, 59, 644–650.

Croft, D.P., Madden, J.R., Franks, D.W. & James, R. (2011). Hypothesis testing in animal social networks. *Trends in Ecology & Evolution*, 26, 502–507.

Csardi, G., Nepusz, T. & and 84 other coauthors. (2018). *igraph 1.2.2: network analysis and visualization.*

Dakin, R. & Ryder, T.B. (2018). Dynamic network partnerships and social contagion drive cooperation. *Proceedings of the Royal Society B: Biological Sciences*, 285, 20181973.

Dakin, R. & Ryder, T.B. (2020). Reciprocity and behavioral heterogeneity govern the stability of social networks. *PNAS*, 117, 2993–2999.

Dawkins, R. (1982). *The Extended Phenotype: The Gene as the Unit of Selection*. W. H. Freeman and Company, San Francisco.

Díaz-Muñoz, S.L., DuVal, E.H., Krakauer, A.H. & Lacey, E.A. (2014). Cooperating to compete: altruism, sexual selection and causes of male reproductive cooperation. *Animal Behaviour*, 88, 67–78.
582

Dormann, C.F., Fruend, J. & Gruber, B. (2019). *bipartite 2.13: visualising bipartite networks and*
584   *calculating some (ecological) indices*.

Durães, R., Loiselle, B.A., Parker, P.G. & Blake, J.G. (2009). Female mate choice across spatial
586   scales: influence of lek and male attributes on mating success of blue-crowned manakins. *Proceedings of the Royal Society B: Biological Sciences*, 276, 1875–1881.

588   DuVal, E.H. (2007). Adaptive advantages of cooperative courtship for subordinate male lance-tailed manakins. *The American Naturalist*, 169, 423–432.

590   Eikenaar, C., Whitham, M., Komdeur, J., van der Velde, M. & Moore, I.T. (2011). Endogenous testosterone is not associated with the trade-off between paternal and mating effort.
592   *Behav Ecol*, 22, 601–608.

Eisenegger, C., Haushofer, J. & Fehr, E. (2011). The role of testosterone in social interaction. *Trends in Cognitive Sciences*, 15, 263–271.

Farine, D.R. (2014). Measuring phenotypic assortment in animal social networks: weighted associations are more robust than binary edges. *Animal Behaviour*, 89, 141–153.

Farine, D.R. (2015). Proximity as a proxy for interactions: issues of scale in social network analysis. *Animal Behaviour*, 104, e1–e5.

Farine, D.R. (2016). *assortnet 0.12: calculate the assortativity coefficient of weighted and binary networks*.

Farine, D.R., Freckleton, R. & Rands, S. (2017). A guide to null models for animal social network analysis. *Methods in Ecology and Evolution*, 8, 1309–1320.

Fowler, J.H. & Christakis, N.A. (2010). Cooperative behavior cascades in human social networks. *PNAS*, 107, 5334–5338.

Franks, D.W., Ruxton, G.D. & James, R. (2010). Sampling animal association networks with the gambit of the group. *Behav Ecol Sociobiol*, 64, 493–503.

Fuxjager, M.J., Forbes-Lorman, R.M., Coss, D.J., Auger, C.J., Auger, A.P. & Marler, C.A. (2010). Winning territorial disputes selectively enhances androgen sensitivity in neural pathways related to motivation and social aggression. *PNAS*, 107, 12393–12398.

Goymann, W. (2009). Social modulation of androgens in male birds. *General and Comparative Endocrinology*, 9th International Symposium on Avian Endocrinology, 163, 149–157.

Goymann, W., Schwabl, I., Trappschuh, M. & Hau, M. (2007). Use of ethanol for preserving steroid and indoleamine hormones in bird plasma. *General and Comparative Endocrinology*, 150, 191–195.

Griffiths, S.W. & Magurran, A.E. (1997). Schooling preferences for familiar fish vary with group size in a wild guppy population. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 264, 547–551.

Heindl, M. (2002). Social organization on leks of the wire-tailed manakin in Southern Venezuela. *The Condor*, 104, 772–779.

Ilmarinen, V.-J., Vainikainen, M.-P., Verkasalo, M.J. & Lönnqvist, J.-E. (2017). Homophilous friendship assortment based on personality traits and cognitive ability in middle childhood: the moderating effect of peer network size. *European Journal of Personality*, 31, 208–219.

James, R., Croft, D.P. & Krause, J. (2009). Potential banana skins in animal social network analysis. *Behav Ecol Sociobiol*, 63, 989–997.

Jehn, K.A. & Shah, P.P. (1997). Interpersonal relationships and task performance: an examination of mediation processes in friendship and acquaintance groups. *Journal of Personality and Social Psychology*, 72, 775–790.

Krause, J. & Ruxton, G.D. (2002). *Living in Groups*. OUP Oxford.

Kuznetsova, A., Brockhoff, P.B. & Christensen, R.H.B. (2018). *lmerTest 3.0-1: tests in linear mixed effects models*.

Lank, D.B. & Smith, C.M. (1992). Females prefer larger leks: field experiments with ruffs (*Philomachus pugnax*). *Behav Ecol Sociobiol*, 30, 323–329.

Lüdecke, D., Makowski, D. & Waggoner, P. (2019). *performance 0.4.2: assessment of regression mdoels performance*.

McClintock, M.K. (1981). Social control of the ovarian cycle and the function of estrous synchrony. *Integr Comp Biol*, 21, 243–256.

McDonald, D.B. (1993). Delayed plumage maturation and orderly queues for status: a manakin mannequin experiment. *Ethology*, 94, 31–45.

McDonald, D.B. (2007). Predicting fate from early connectivity in a social network. *PNAS*, 104, 10910–10914.

McDonald, G.C., Farine, D.R., Foster, K.R. & Biernaskie, J.M. (2017). Assortment and the analysis of natural selection on social traits. *Evolution*, 71, 2693–2702.

Nakagawa, S. & Schielzeth, H. (2013). A general and simple method for obtaining R2 from generalized linear mixed-effects models. *Methods Ecol Evol*, 4, 133–142.

Ohtsuki, H., Hauert, C., Lieberman, E. & Nowak, M.A. (2006). A simple rule for the evolution of cooperation on graphs and social networks. *Nature*, 441, 502–505.

Oyegbile, T.O. & Marler, C.A. (2005). Winning fights elevates testosterone levels in California mice and enhances future ability to win fights. *Hormones and Behavior*, 48, 259–267.

Pinter-Wollman, N., Hobson, E.A., Smith, J.E., Edelman, A.J., Shizuka, D., de Silva, S., *et al.* (2014). The dynamics of animal social networks: analytical, conceptual, and theoretical
advances. *Behav Ecol*, 25, 242–255.

Poisot, T., Canard, E., Mouillot, D., Mouquet, N. & Gravel, D. (2012). The dissimilarity of species
654 interaction networks. *Ecology Letters*, 15, 1353–1361.

Prum, R.O. (1994). Phylogenetic analysis of the evolution of alternative social behavior in the
656 manakins (Aves: Pipridae). *Evolution*, 48, 1657–1675.

R Core Team. (2018). *R 3.5.1: A Language and Environment for Statistical Computing*. R
658 Foundation for Statistical Computing, Vienna, Austria.

Roberts, G. & Sherratt, T.N. (1998). Development of cooperative relationships through
660 increasing investment. *Nature*, 394, 175.

Robinson, G.E. (1992). Regulation of division of labor in insect societies. *Annual Review of*
662 *Entomology*, 37, 637–665.

Ryder, T.B., Blake, J.G., Parker, P.G. & Loiselle, B.A. (2011a). The composition, stability, and
664 kinship of reproductive coalitions in a lekking bird. *Behav Ecol*, 22, 282–290.

Ryder, T.B., Dakin, R., Vernasco, B.J., Evans, B.S., Horton, B.M. & Moore, I.T. (2020).
666 Testosterone modulates status-specific patterns of cooperation in a social network. *The American Naturalist*.

668 Ryder, T.B., Horton, B.M. & Moore, I.T. (2011b). Understanding testosterone variation in a tropical lek-breeding bird. *Biology Letters*, 7, 506–509.

670 Ryder, T.B., Horton, B.M., Tillaart, M. van den, Morales, J.D.D. & Moore, I.T. (2012). Proximity data-loggers increase the quantity and quality of social network data. *Biology Letters*, 8,
672 917–920.

Ryder, T.B., McDonald, D.B., Blake, J.G., Parker, P.G. & Loiselle, B.A. (2008). Social networks in
674 the lek-mating wire-tailed manakin (*Pipra filicauda*). *Proceedings of the Royal Society B*, 275, 1367–1374.

676 Ryder, T.B., Parker, P.G., Blake, J.G. & Loiselle, B.A. (2009). It takes two to tango: reproductive skew and social correlates of male mating success in a lek-breeding bird. *Proceedings of*
678 *the Royal Society B*, 276, 2377–2384.

Sah, P., Leu, S.T., Cross, P.C., Hudson, P.J. & Bansal, S. (2017). Unraveling the disease consequences and mechanisms of modular structure in animal social networks. *Proc Natl Acad Sci USA*, 114, 4165–4170.

Sah, P., Mann, J. & Bansal, S. (2018). Disease implications of animal social network structure: A synthesis across social systems. *J Anim Ecol*, 87, 546–558.

Schjelderup-Ebbe, T. (1922). Beiträge zur sozialpsychologie des haushuhns. [Observation on the social psychology of domestic fowls.]. *Zeitschrift für Psychologie und Physiologie der Sinnesorgane. Abt. 1. Zeitschrift für Psychologie*, 88, 225–252.

Schwartz, P. & Snow, D.W. (1978). Display and related behavior of the wire-tailed manakin. *Living Bird*, 17, 51–78.

Trainer, J.M. & McDonald, D.B. (1995). Singing performance, frequency matching and courtship success of long-tailed manakins (*Chiroxiphia linearis*). *Behav Ecol Sociobiol*, 37, 249–254.

Trainer, J.M., McDonald, D.B. & Learn, W.A. (2002). The development of coordinated singing in cooperatively displaying long-tailed manakins. *Behav Ecol*, 13, 65–69.

Trivers, R.L. (1971). The evolution of reciprocal altruism. *The Quarterly Review of Biology*, 46, 35–57.

Vernasco, B.J., Horton, B.M., Moore, I.T. & Ryder, T.B. (2020). Reduced cooperative behavior as a cost of high testosterone in a lekking passerine bird. *Behav Ecol*.

Vernasco, B.J., Horton, B.M., Ryder, T.B. & Moore, I.T. (2019). Sampling baseline androgens in free-living passerines: Methodological considerations and solutions. *General and Comparative Endocrinology*, Endocrinology of Neotropical Vertebrates, 273, 202–208.

Wingfield, J.C., Hegner, R.E., Dufty, Alfred M. & Ball, G.F. (1990). The "challenge hypothesis": theoretical implications for patterns of testosterone secretion, mating systems, and breeding strategies. *The American Naturalist*, 136, 829–846.

Zuur, A., Ieno, E.N., Walker, N., Saveliev, A.A. & Smith, G.M. (2009). *Mixed effects models and extensions in ecology with R*. Springer Science & Business Media.

Zyphur, M.J., Narayanan, J., Koh, G. & Koh, D. (2009). Testosterone–status mismatch lowers collective efficacy in groups: evidence from a slope-as-predictor multilevel structural equation model. *Organizational Behavior and Human Decision Processes*, Biological Basis of Business, 110, 70–79.

Figure 1

A Social specialization

Low
0.08

High
0.83

Territory
-holders

Floaters

B Network stability

t_1　　　　t_2

Low
0.33

High
0.88

C Behavioral assortment

Negative
-0.52

Positive
0.66

Avg. daily behavior

Figure 2

Figure 3