Testosterone-mediated behavior shapes the emergent properties of social networks

Roslyn Dakin[1,2], Ignacio T. Moore[3], Brent M. Horton[4], Ben J. Vernasco[3], T. Brandt Ryder[1,5]

[1]Migratory Bird Center, Smithsonian Conservation Biology Institute, National Zoological Park, Washington, DC 20013, USA
[2]Current address: Department of Biology, Carleton University, Ottawa, Ontario K1S 5B6, Canada
[3]Department of Biological Sciences, Virginia Tech, Blacksburg, Virginia 24061, USA
[4]Department of Biology, Millersville University, Millersville, Pennsylvania 17551, USA
[5]Current address: Bird Conservancy of the Rockies, Fort Collins, Colorado, 80525, USA

Correspondence: rydert@si.edu

Abstract

1. Social networks can vary in their organization and dynamics, with implications for ecological and evolutionary processes. Understanding the mechanisms that drive these dynamics requires integrating individual-level biology with comparisons across multiple social networks.

2. Testosterone is a key regulator of vertebrate social behavior. Previous studies have demonstrated that testosterone can influence how individuals interact with social partners. Although the effects of testosterone on individual behaviors are well established, no study has examined whether hormone-mediated behavior can scale up to shape the emergent structure of social groups.

3. We tested this hypothesis in the wire-tailed manakin, a bird species in which the males cooperate at specialized courtship sites known as leks. We used an automated proximity system to longitudinally monitor multiple leks and quantify the social network interactions at each site. Our analysis examines three emergent properties of the social networks that are expected to enhance the benefits of cooperative behavior: social specialization (the exclusivity of relationships), network stability (the overall persistence of partnerships through time), and behavioral assortment (the tendency for like to associate with like).

4. The manakin leks with higher-testosterone dominant males were less specialized, less stable, and had more negative behavioral assortment, after accounting for other factors. These results support our main hypothesis that individual-level hormone physiology can predict group-level network dynamics. We also observed that larger leks with more

38   interacting individuals had more positive behavioral assortment. This suggests that small groups may constrain processes of homophily and behavior-matching.

40   5. Overall, these results provide evidence that hormones can act as an extended phenotype that shapes the broader architecture of social groups. High-testosterone groups exhibit

42   collective behaviors that are predicted to impede the evolution of cooperation.

44   **Keywords:** androgens, behavioral endocrinology, collective behavior, cooperation, dynamic networks, social networks, testosterone

46 **Introduction**

Behavioral interactions are the foundation of social network structures that can vary through

48 time, among populations, and among species. Network structures play an important role in many

ecological and evolutionary processes, including the spread of diseases (Sah *et al.* 2018), the

50 transmission of information and resources (Aplin *et al.* 2012), and selection on individual

behavior (Ohtsuki *et al.* 2006). A major challenge is understanding how individual-level factors,

52 such as physiological and behavioral mechanisms, scale up to drive the emergent structural

properties of social groups (Krause & Ruxton 2002). Linking these two levels of analysis is

54 difficult because it requires integrating individual-level data with repeated measures of entire

social groups (Sah *et al.* 2017).

56 Here, we use a comparison of multiple social networks through time to investigate the

hypothesis that testosterone destabilizes social structure. Testosterone is a steroid hormone that is

58 well known for its influence on the social behavior of individuals (Wingfield *et al.* 1990;

Oyegbile & Marler 2005; Adkins-Regan 2005; Goymann 2009; Fuxjager *et al.* 2010; Eisenegger

60 *et al.* 2011). These hormone-signaling pathways are essential for the development and expression

of complex behavioral phenotypes (Cohen *et al.* 2012). Recent work has shown that testosterone

62 can influence the cooperative behavior of individuals (Boksem *et al.* 2013; Ryder *et al.* 2020).

We consequently hypothesize that hormone-mediated behaviors may also shape higher-order

64 social network structure.

To investigate this hypothesis, we studied the wire-tailed manakin (*Pipra filicauda*), a

66 bird species in which the males engage in cooperative display coalitions at specialized courtship

sites known as leks. In *P. filicauda*, males can represent two status classes: dominant, territory-

68 holding males, and subordinate, "floater" males who must acquire a territory on a lek before they

can mate (Heindl 2002; Ryder *et al.* 2011a). Cooperative coalitions occur between two unrelated

70   males, most commonly a territory-owner and a floater (Ryder *et al.* 2011a). A hallmark of

reproductive cooperation is that, on average, both partners gain some benefit, although the

72   magnitude of these benefits need not be equal (e.g., partners can receive delayed benefits; DuVal

2007; Díaz-Muñoz *et al.* 2014). Consistent with this definition, long-term studies of *P. filicauda*

74   have shown that coalition partnerships are mutually beneficial. First, subordinate males with

more social partners are more likely to acquire a territory in subsequent years (Ryder *et al.*

76   2008), similar to other cooperative systems (McDonald 2007). Second, the territory-holders with

more social partners sire more offspring, as compared to their direct competitors (Ryder *et al.*

78   2009).

It is impossible to manipulate long-term social relationships in the wild to experimentally

80   test the association between sociality and fitness. Nevertheless, further evidence also supports the

hypothesis that partnerships are beneficial in *P. filicauda*. For example, when a male has a social

82   partner present, he displays at a higher frequency and for a longer period of time (Ryder *et al.*

2011a; Vernasco *et al.* 2020). Given that these conspicuous displays are often a target of female

84   mate choice, this provides a mechanism by which coalitions may increase male attractiveness

and subsequent reproductive success (Lank & Smith 1992; Trainer & McDonald 1995; Durães *et*

86   *al.* 2009). Territory-holders also use calls to solicit visitation by coalition partners (Schwartz &

Snow 1978; Ryder and Vernasco, personal observations), a behavior that would be unlikely if

88   these partnerships were not directly beneficial. Finally, territory-holders who visit and perform

displays with males on neighboring territories may also benefit, as a result of reciprocity and

90   behavioral contagion (Dakin & Ryder 2018).

Two features of the manakin leks make them especially well-suited to study the

92   relationship between hormones and group-level social structure. First, we recently established

that testosterone has status-specific effects on the coalition behavior of individual male wire-

94   tailed manakins (Ryder *et al.* 2020). Second, these coalition partnerships among males form the

basis of complex and dynamic social networks (Dakin & Ryder 2018). Given this background,

96   we sought to test whether testosterone and its effects on male behavioral phenotype could drive

the emergent properties of the social network that facilitate cooperation. Previous theoretical

98   studies have shown that cooperation in social networks is favored when individuals interact with

a limited set of familiar partners, when these relationships are temporally stable, and when social

100 relationships are reciprocated by immediate neighbors in the group (Trivers 1971; Roberts &

Sherratt 1998; Ohtsuki *et al.* 2006). Accordingly, we used repeated measures of the manakin

102 networks to analyze whether testosterone was related to three emergent properties of the social

network, described below (Fig. 1).

104   The first property, social specialization, seeks to capture the exclusivity of the

relationships between coalition partners. In the context of manakin behavior, highly specialized

106 networks exhibit a high frequency of exclusive social partnerships between males of different

status classes, as illustrated in Fig. 1A. In other species, social specialization has been found to

108 maximize the quality and coordination of different types of behaviors (Jehn & Shah 1997;

Laskowski *et al.* 2016). In manakins, we expect specialization to improve the signal quality of

110 male-male courtship displays (e.g., as a result of improved behavioral coordination). Greater

specialization is also expected to minimize conflict over mating and territorial ascension

112 opportunities (Schjelderup-Ebbe 1922; McDonald 1993).

The second property, network stability (Fig. 1B), quantifies the average persistence of

114   social partnerships through time (Poisot *et al.* 2012; Dakin & Ryder 2020). Coalition

partnerships require the coordination of complex behaviors, and previous empirical work

116   indicates that longer partnership tenure has a positive effect on display coordination (Trainer &

McDonald 1995; Trainer *et al.* 2002). Greater temporal stability of partnerships also increases

118   the opportunity for sustained reciprocity within the network (Trivers 1971; Roberts & Sherratt

1998; Croft *et al.* 2006).

120   The third property, behavioral assortment (Fig. 1C), captures the extent to which males

interact with partners who express similar behaviors (i.e., is like associated with like? (Croft *et*

122   *al.* 2006; Farine 2014)). At the proximate level, positive assortment may represent the outcome

of generalized reciprocity and/or partner choice (Fowler & Christakis 2010; Dakin & Ryder

124   2018). At the ultimate level, positive assortment has also been shown to promote the evolution of

cooperation (Ohtsuki *et al.* 2006). To quantify the overall behavioral assortment of the manakin

126   networks, we used a composite measure that averaged the assortativity of "strength" (a male's

frequency of daily social interactions) with that of "degree" (his daily number of social

128   partnerships; (Dakin & Ryder 2018)).

As a potential predictor of specialization, stability, and assortment, we quantified the

130   collective testosterone of the manakin leks (Fig. 2; (Akinola *et al.* 2016)). Because hormone-

behavior relationships are status-specific in manakins and many other species (Eisenegger *et al.*

132   2011; Boksem *et al.* 2013; Ryder *et al.* 2020), we analyzed collective testosterone for each of the

two status classes separately. High testosterone levels are associated with reduced cooperative

134   behavior in dominant males, but increased cooperative behavior in subordinate males (Ryder *et*

al. 2020). We consequently predicted that the collective testosterone for territorial males would

136   be negatively associated with specialization, stability, and assortment, whereas the collective

testosterone for floater males would be positively associated with these three emergent

138   properties.

140   **Materials and Methods**

Study Population

142   We studied wire-tailed manakins (*Pipra filicauda*) at the Tiputini Biodiversity Station in

Orellana Province, Ecuador (0º 38' S, 76º 08' W). This population of *P. filicauda* has been

144   studied and individuals color-banded annually since 2002 (Ryder *et al.* 2008, 2009). The present

study was conducted on 11 leks during peak breeding activity (December-March) across three

146   field seasons: 2015-16, 2016-17 and 2017-18 (Ryder *et al.* 2020). All research was approved by

the Smithsonian ACUC (protocols #12-23, 14-25, and 17-11) and the Ecuadorean Ministry of the

148   Environment (MAE-DNB-CM-2015-0008).

150   *Testosterone Assessment*

Male manakins were captured using mist-nets on the display leks as described in (Ryder *et al.*

152   2020). We deployed up to 16 mist-nets simultaneously at a given lek, with the goal of capturing

each male up to three times per field season. Each mist-net was checked on a 30-minute schedule

154   (Vernasco *et al.* 2019; Ryder *et al.* 2020). The amount of time a male spends in the net has a

subtle, but significant, effect on his circulating testosterone (Vernasco et al. 2019). Therefore, we

156   used video monitoring to determine the duration of time that each bird was in the mist-net prior

to blood sampling, so that it could be accounted for in further analyses (mean = 17.5 minutes, SD

158   = 10.3 minutes, range = 1–72 minutes). Following removal of a male from the mist-net, a small

blood sample (< 125uL) was taken from the brachial vein and stored on ice prior to being

160   centrifuged at 10,000 rpm for 5 min, as described in previous studies (Ryder *et al.* 2011b;

Vernasco *et al.* 2019; Ryder *et al.* 2020). Plasma volume was measured to the nearest 0.25 ul and

162   stored in 0.75 ml of 100% ethanol (Goymann *et al.* 2007). Testosterone was double extracted

from the plasma using dichloromethane. Following extraction, direct radioimmunoassay was

164   used to measure the total plasma androgen concentration (ng/ml) adjusted by the extraction

efficiency and plasma volume of each sample (Eikenaar *et al.* 2011; Ryder *et al.* 2011b).

166   Hormone assays were conducted annually, and the detection limits were 0.12, 0.08, and 0.09

ng/ml for 2015-16, 2016-17 and 2017-18, respectively; any sample that fell below the assay-

168   specific limit of detection was assigned that limit as its testosterone concentration as a most

conservative estimate. The extraction efficiency for all samples was between 62-73%, and the

170   intra-assay coefficients of variation were 6.6%, 11.6%, and 9.2% for 2015-16, 2016-17 and

2017-18, respectively; the inter-assay coefficient of variation was 19.5%.

Behavioral Recording

174   We used an automated data-logging system to monitor male-male interactions on the display

territories of 11 different leks (Ryder *et al.* 2012; Dakin & Ryder 2018; Ryder *et al.* 2020). The

176   territories on these leks are specifically used for courtship displays and male-male social

interactions. At the beginning of each field season, male manakins were outfitted with coded

178   nano-tags (NTQB-2, Lotek Wireless; 0.35 g). The tags transmitted a unique VHF signal ping

once per 20 s for three months. In total, 296 tag deployments were performed on 180 individuals

180   (mean 1.7 field seasons per male ± SD 0.7), 178 of whom also had hormone data (mean number

of hormone samples per male = 3 ± SD 1.5). Approximately 10 days (± SD 7) after tagging and

182   sampling was completed at a given lek, a proximity data-logger (SRX-DL800, Lotek Wireless)

was deployed within each display territory at the lek to record tagged males within a detection

184   radius of 30 m (a distance that corresponds to the typical diameter of a manakin display territory

(Heindl 2002; Dakin & Ryder 2018)).

186   Proximity recording sessions ran from 06:00 to 16:00 for ~6 consecutive days (± SD 1

day) and were performed ~3 times per field season at a given lek. Occasionally, the length of a

188   recording session was extended due to extenuating circumstances such as inclement weather.

The recording sessions were scheduled to be distributed evenly throughout each field season at

190   each lek, to minimize any confounding seasonal effects. Each recording session represents an

observation of the social network at a given lek. In total, we conducted 86 recording sessions

192   (29,760 data-logger hours) representing repeated measures of the social activity at 11 different

leks over the course of three field seasons (see Fig. S1 in the supplement for additional details on

194   the sampling regime).

          Prior to data-logger deployment, each territory was also observed on non-recording days

196   to identify the territory-holder based on his color-bands, following previous studies (Ryder *et al.*

2008, 2009). These status assignments were subsequently verified in the proximity data.

Data Processing and Statistical Analysis

200   All computational and statistical analyses were performed in R (R Core Team 2018). Network

illustrations were made using the igraph package (Csardi & coathors 2018).

Social Interactions

206   Social networks were constructed by first defining interactions between two males that occurred

on the display territories. To do this, a computational algorithm was used to identify joint

208   detections, wherein two males were located at the same display territory within a pre-defined

spatial and temporal threshold (Ryder *et al.* 2008, 2012; Dakin & Ryder 2018). For the temporal

210   threshold, the two males had to occur < 45 s apart. This temporal threshold was chosen to allow

for the fact that each tag pinged with a 20 second pulse rate, such that overlapping individuals

212   could have up to a 40 s gap between their respective pings. For the spatial threshold, the two

males had to have a difference in received signal strength values (Δ RSSI) < 10. This threshold

214   corresponded to a typical distance < 5 m apart in a ground-truthing experiment (Dakin & Ryder

2018). Hence, according to our definition, a social interaction is initiated only after males come

216   within this approximate spatial threshold. We chose this spatial threshold because it is close

enough to permit visual and acoustic contact during typical social behaviors, such as those

218   described in (Schwartz & Snow 1978). Any repeated co-occurrence of the two males within 5

minutes was considered to be part of the same social interaction, but after a gap of ≥ 5 minutes, it

220   was considered to be a new interaction between those two males. The average duration of social

interactions defined by this method was 5.1 min $\pm$ SD 13.3, 8.8 min $\pm$ SD 20.1, and 7.0 min $\pm$

222   SD 20.5 in the three respective field seasons, further indicating that these were sustained social

interactions, rather than random encounters.

224   In total, we identified 36,885 social interactions over the three field seasons of study.

These interactions were used to define a weighted social network for each lek recording session.

226   The nodes in the network were the individual males, and the edges were weighted by the number

of social interactions between each pair of males. An additional validation study confirmed that

228   the social interactions defined by this method corresponded to male-male partnerships that were

directly observed (Ryder *et al.* 2012).

Pre-network Permutation Analysis

232   Two broad classes of methods have been described for building social networks: (i) networks

that are built based purely on the proximity of animals unrelated to their behavioral context

234   ("gambit of the group"), and (ii) networks that are built based on specific behavioral criteria that

are directly observed (Franks *et al.* 2010; Farine 2015). Although the interactions in our study

236   were not directly observed, they were recorded at specialized display perches that have a known

function in male-male reproductive coalitions (Ryder *et al.* 2011a). Hence, they do not qualify as

238   the gambit of the group. To further demonstrate this point, we performed a pre-network

permutation of the raw data to validate that most of the observed network edges occurred more

240   often than expected by chance, even within lek recording sessions (Farine *et al.* 2017). This

analysis preserved key features of the data including male visit rates to specific territories within

242   each recording session; it is described in detail in the supplement and Fig. S2. The results

demonstrated that 95% of the observed network edges had a greater edge weight than expected

244   under stringent permutation conditions. This provides additional validation to our approach,

because it indicates that the observed network edges were nearly always preferred, even relative

246   to other possible interactions within the same lek. These results are also consistent with a

previous validation study indicating that these methods capture male-male coalition partnerships

248   that are directly observed (Ryder *et al.* 2012).

       Although pre-network data permutations are sometimes used to derive adjusted

250   association indices, or to prune networks prior to further analysis, we did not take this approach

for several reasons. First, the statistical rarity of a relationship in our system does not *a priori*

252   define the importance of any one social interaction, especially given that all of the interactions

occurred on display territories with specialized function. A single interaction between two rarely

254   interacting males may have been highly consequential (e.g., if one of those males had a highly

influential hormone-behavioral phenotype). Conversely, a single interaction between two

256   frequently interacting males may have been relatively unimportant. It would be unwarranted to

assume that partnership rarity indicates the importance of any single interaction in this context.

258   Second, and perhaps more importantly, the goal of this study was a comparison across multiple

networks. Some networks are genuinely less preferential, and more random, than others. If we

260   modified the network edges based on permutation-based indices, it would disproportionately

prune the truly random networks, introducing a source of bias that would be contrary to our main

262   goal. Hence, all further analyses are based on networks where the edge weights are given by the

observed number of interactions, as this method is most appropriate for our study system and

264   aims.

266   ***Social Specialization***

To quantify social specialization, we sought a metric that would capture the extent to which a

268   network was partitioned into exclusive social relationships (as opposed to a network made up of

non-specific or non-exclusive partnerships). To do this, we used a network metric of

270   specialization that is commonly used in community ecology called H_2' (Blüthgen *et al.* 2006).

An advantage of H_2' is that it is standardized against a theoretical maximum, based on the

272   overall activity levels of different nodes and Shannon entropy (Blüthgen *et al.* 2006); this makes

it possible to compare the extent of specialization across different bipartite networks in a

274   standardized way. To apply this metric to our manakin data, we converted each lek social

network into its bipartite adjacency matrix (Fig. 1A), with floaters along one axis, and territory-

276   holders on the other, and then calculated social specialization as H_2' using the bipartite package

(Dormann *et al.* 2019). Higher values of specialization indicate that the network is made up of

278   more exclusive relationships, as illustrated in Fig. 1A. We chose to focus on floater-territorial

specialization because these two classes are well-defined and floater-territorial partnerships tend

280   to be the most common in this species (Fig. S3 and (Ryder *et al.* 2011a)).

282   ***Network Stability***

To quantify the stability of social relationships, we compared each lek social network from a

284   given recording session to its subsequent recording session within the same field season (Fig.

1B). Network stability was then calculated as the number of male-male partnerships (network

286   edges) shared by both time points divided by the number of partnerships at either time point

(Dakin & Ryder 2020). Higher values of stability indicate greater persistence of social

288   relationships within the network, independent of any changes in the representation of particular

males (nodes) (Poisot *et al.* 2012). To focus this measure on the persistence of strong

290   relationships, we computed network stability of partnerships that occurred at least six times (i.e.,

at least once per day, on average) following (Dakin & Ryder 2020). Previous work using this

292   metric shows that the wire-tailed manakin social networks are more stable than expected by

chance (Dakin & Ryder 2020).

Behavioral Assortment

298   Assortment refers to the extent to which individuals associate with similar partners (Fig. 1C); it

can be due to partner choice (homophily), shared environments, or the social transmission of

300   behavior (Dakin & Ryder 2018). Assortment was quantified using Newman's assortativity,

which is a correlation coefficient for the statistical association among linked nodes within a

302   network. It ranges from –1 (a negative association), through 0 (no association), and up to +1 (a

positive association). To quantify the assortment of social behaviors, we first determined the

304   daily frequency of two behaviors for each male: his number of social interactions per day

(strength), and his number of unique social partnerships per day (degree). Strength and degree

306   are both repeatable measures of a male's social behavior in our study population (Dakin & Ryder

2018). We used the average log-transformed values of each male's strength and degree within

308   the recording session, and then calculated the assortativity coefficient for the entire social

network using the algorithm for weighted networks in the assortnet package (Farine 2016).

310   Because assortativity values for strength and degree were highly correlated (Pearson's $r = 0.78$, p

< 0.0001, $n = 86$ networks), we took the average of these two values as the measure of overall

312   behavioral assortment within the social network. Note that we used log-transformed values of

strength and degree because these two variables are strongly positively skewed (Dakin & Ryder

314   2018; Ryder *et al.* 2020), and assortativity is based on a Pearson's correlation. Finally, we also

computed the assortativity of the two discrete status classes (floater and territory-holder), to

316   ensure that our analysis of behavioral assortment was not solely driven by status-assortment.

320 *Collective Testosterone*

To understand how hormones might predict network properties, we derived a measure of

322 collective testosterone of each social network. This was based on the hormonal trait that was the

best predictor of cooperative behavior in our previous study, referred to as "mean T" (Ryder *et*

324 *al.* 2020). A male's mean T is his average residual circulating testosterone. It is calculated using

a linear regression of log-transformed testosterone to statistically account for the effects of field

326 season, Julian date, time of day when captured, and duration of restraint, all of which may

influence point estimates of a male's baseline hormone level (Vernasco *et al.* 2019). Hence,

328 mean T represents a standardized measure of a male's circulating testosterone, independent of

his capture conditions (Fig. 2A). Next, to determine collective testosterone, we took the average

330 mean T for each social network, weighted by the interaction frequency (strength) of the males

within the network (Fig. 2B). Collective testosterone is thus a group-level characteristic that is

332 weighted towards the males that made the greatest contribution to group social structure. In other

words, networks with low collective testosterone are made up of mostly low-testosterone

334 individuals, whereas networks with high collective testosterone are made up of mostly high-

testosterone individuals. We calculated collective testosterone separately for each status class,

336 because the effects of hormones on social behavior are status-dependent (Eisenegger *et al.* 2011;

Boksem *et al.* 2013; Ryder *et al.* 2020).

Statistical Analysis

340 To evaluate the hypothesis that collective testosterone predicts network properties, we analyzed

mixed-effects models of the social network properties in the lme4 package (Bates *et al.* 2018).

342 The three response variables were social specialization, network stability, and behavioral

assortment. We used Akaike's Information Criterion (AIC) to compare four candidate models for each response variable, as follows: (1) collective testosterone of territory-holders + collective testosterone of floaters; (2) collective testosterone of territory-holders; (3) collective testosterone of floaters, and (4) no testosterone predictors. All of the models included additional fixed effects to account for field season (a categorical variable with three levels), the average Julian date of the recording session, the average number of recorded hours per territory, and the size of the social network (number of individuals), as well as a random effect of lek to account for repeated measures.

Model comparison was performed on models fit with maximum likelihood, and the best-fit models were re-fit using restricted estimation of maximum likelihood (REML) to derive parameter estimates (Zuur *et al.* 2009). We used the lmerTest package to compute p-values for parameter estimates in the mixed-effects models based on Satterthwaite's method (Kuznetsova *et al.* 2018). We also report Nakagawa and Schielzeth's R^2_{LMM} values as an estimate of effect size (Nakagawa & Schielzeth 2013). We verified that all models met the assumptions of linear regression analyses. First, we checked that the Pearson residuals met the assumption of normality. We visually inspected the partial residual plots for each fixed and random effect, to confirm that there were no outliers or departures from the assumption of homoscedasticity. To check for multicollinearity, we used the performance package (Lüdecke *et al.* 2019) to calculate variance inflation factors (VIFs), and we verified that all VIFs were between $1 - 1.8$.

Sample Sizes and Exclusions

In two field seasons (2016-17 and 2017-18), we performed an experiment as part of a separate study to test the influence of transiently-elevated testosterone on individuals (n = 5 individuals in

366   2016-17 and n = 4 in 2017-18 (Ryder *et al.* 2020)). The results of that experiment demonstrated

that elevated testosterone caused a temporary decrease in the frequency and the number of social

368   partnerships in the altered males (Ryder *et al.* 2020). It is important to note that this experiment

was not designed to test emergent properties at the network level, because it was conducted on a

370   limited scale with only one or two individuals temporarily altered within each lek. We therefore

excluded the 6 post-manipulation networks from the main analysis in this study. We verified that

372   when we included these manipulated leks, all of our main conclusions were unchanged.

After excluding the 6 post-manipulation observations, 80 of the original 86 recording

374   sessions remained. Table 1 summarizes the sample sizes for the network-level analyses. In 10

cases, specialization could not be calculated because the bipartite network did have sufficient

376   data to determine H_2'. Stability could not be calculated in 26 cases when either the recording

session occurred at the end of a field season, or there were insufficient partnerships that met the

378   criteria for the stability calculation.

380   *Node-label Permutation Analysis*

To evaluate the possibility that our results could be influenced by other properties of the leks that

382   were independent of hormonal traits, we also performed a statistical permutation of the post-

network data (Farine *et al.* 2017). The purpose of this analysis was to verify that the results were

384   driven by (and sensitive to) the relative contribution and social position of different males. To do

this, we performed a statistical permutation that randomized the node labels (male IDs) within

386   each of the social networks, retaining network topology, and leaving each male's testosterone

traits unchanged. Hence, this analysis preserved which males were present in which recording

388   session, but it randomized the relative position and contribution of each male. After generating

1,000 of these node-label permutation datasets, we recalculated specialization, stability, and

390   assortment, and then refit the top models from our mixed-model analysis. We then compared the

slope estimates from the observed data to those derived from 1,000 node-label permutations. As

392   a one-sided p-value, we calculated the proportion of slope estimates from the node-label

permutations that were more negative than the corresponding estimate from the observed data.

Results

396   The manakin social networks had 14 males on average, but there was considerable variation

among leks ranging from 3 to 43 individuals (Fig. S1). Additional descriptive statistics are

398   provided in Table 1. Although network size has a lognormal distribution (Fig. S1), all other

emergent properties of the social structure were approximately normally distributed (Fig. S4).

402   **Table 1. Descriptive statistics for manakin social networks.** Means and standard deviations
(SD) are provided for each field season. The bottom row provides sample sizes for the number of
404   social networks analyzed (n_{obs}) at each lek (N_{lek}). Additional data is provided in the supplement
in Fig. S1 and Fig. S4.

Field season	Network size (# birds)	Social specialization	Network stability	Behavioral assortment	Collective T (terr.)	Collective T (floa.)
15-16	14.5 (8.9)	0.50 (0.22)	0.43 (0.25)	0.01 (0.36)	0.44 (0.42)	–0.31 (0.56)
16-17	17.8 (10.5)	0.30 (0.24)	0.47 (0.20)	0.06 (0.30)	0.49 (0.24)	–0.22 (0.37)
17-18	10.3 (4.5)	0.42 (0.30)	0.42 (0.27)	0.02 (0.37)	0.43 (0.40)	–0.70 (0.46)
Sample size	$n_{obs} = 80$	$n_{obs} = 70$	$n_{obs} = 54$	$n_{obs} = 80$	$n_{obs} = 80$	$n_{obs} = 80$
	$N_{lek} = 11$	$N_{lek} = 11$	$N_{lek} = 11$	$N_{lek} = 11$	$N_{lek} = 11$	$N_{lek} = 11$

We found that the collective testosterone of territorial males could predict all three emergent

408   properties of the networks (specialization, stability, and assortment). The leks with a greater

contribution from high-testosterone territorial males were less specialized, less stable, and more

410   negatively assorted (Fig. 3; all p ≤ 0.03 in mixed-effects models). The slope coefficients for these

three relationships were also all greater than expected under a node-label permutation (inset

412   panels, Fig. 3; all p < 0.02). Model selection results for the observed data indicated considerable

uncertainty in the best-fit model for each of the three network properties analyzed (Table S1).

414   However, the collective testosterone of territory-holders was a significant predictor in all of the

best-supported models (Table S2). In contrast, the collective testosterone of floater males was

416   not a significant predictor of network properties in any of the best-fit models (Table S2).

We did not detect any significant effects of Julian date on network properties within our

418   study period, but we did observe some significant year-to-year differences (e.g., social

specialization and behavioral assortment in Table S2). Assortment was the only property that

420   was significantly related to network size and recording effort, but in opposite ways (Table S2).

All else being equal, behavioral assortment was more positive in larger networks, but it was

422   more negative in networks that had longer recording sessions.

$R^2_{LMM(m)}$ provides an estimate of the proportion of variance explained by the fixed effects

424   in a model (Table S1). The $R^2_{LMM(m)}$ for assortment in the best-fit model was 0.52. This indicates

that around 52% the variation in behavioral assortment could be explained by the combined

426   associations with collective testosterone, network size, sampling effort, and annual variation

(Table S2). The $R^2_{LMM(m)}$ for stability was 0.13, indicating that collective testosterone could

428   explain about 13% the variation in that metric (Table S2). Finally, for specialization, the

$R^2_{LMM(m)}$ indicated that the combined effects of testosterone and year-to-year variation together

430   explained about 20% of the variation (Table S2).

Discussion

434   The collective testosterone of dominant, territory-holding males was associated with multiple

features of the social network (Fig. 3). This indicates that the effects of testosterone on dominant

436   males may represent an extended phenotype with the power to shape emergent social structure

(Dawkins 1982). In contrast, the collective testosterone of floater males was not significantly

438   associated with any of the emergent network properties (Table S2).

How can the relationship between hormone levels and network properties be explained in

440   terms of individual mechanisms? Given that testosterone has antagonistic effects on the

cooperative behavior of territorial males (Ryder *et al.* 2020), we hypothesize that the behavior of

442   high-testosterone individuals causes several features of the social organization to break down.

Our previous results showed that high-testosterone dominant individuals have a reduced ability

444   to attract and maintain coalition partners (Ryder *et al.* 2020). We propose that this weakening of

coalition bonds may cause floater males to prospect elsewhere for new partnerships, negatively

446   affecting both the stability and specialization of the lek network as a whole. Likewise, high-

testosterone dominant males may inhibit the processes of social contagion, reciprocity, and/or

448   behavioral matching that cause positive behavioral assortment (Dakin & Ryder 2018).

We included several additional parameters in our analysis to account for network size and

450   sampling effort. Although not one of our main hypotheses, we noted that network size was

positively associated with behavioral assortment (Table S2). This result indicates that males are

452   more likely to associate with behaviorally-similar partners in larger leks, whereas they are more

likely to associate with dissimilar partners in smaller leks. Effects of group size on assortment

454   have been noted in a few other studies, although the form of this relationship varies (Griffiths &

Magurran 1997; Ilmarinen *et al.* 2017; McDonald *et al.* 2017). In manakins and other lekking

456   systems, larger leks are known to have heightened display activity and higher female visitation

rates (Lank & Smith 1992; Durães *et al.* 2009). This raises the possibility that social facilitation

458   and heightened activity may be associated with increased homophily and/or behavioral matching

(either through partner choice, contagion, or reciprocity). Another plausible explanation is that

460   smaller networks may constrain these behavioral processes, by making it more difficult to find or

match a suitable partner. This hypothesis could be explored in experiments on captive systems

462   and simulation models.

Behavioral assortment was also more positive in networks that were recorded for less

464   time in our study. Recording time was designed to be approximately even among leks, with the

most common reason for departure being inclement weather that extended the recording session

466   (Fig. S1). Therefore, we speculate that inclement weather may have reduced the amount of

behavioral matching (assortment) while also affecting recording time. Although we did not

468   collect data on weather at the leks as this was not our main goal, the question of how inclement

weather affects group-level social dynamics is an interesting one that merits further study.

470   Conducting experimental tests of causation at the level of whole social networks remains

a major challenge in ecology (Pinter-Wollman *et al.* 2014; James *et al.* 2009). Although our

472   node-label permutation analysis provides evidence that our results are not merely due to

structural differences among networks, we cannot rule out the possibility that high-testosterone

474   individuals chose to participate in certain networks due to other factors that may also influence

the emergent properties of the network (e.g., environmental quality and/or female activity). It is

476   important to note that in this system, as in many other wild animals, controlled experimental

manipulations of the broader social network structure are not yet possible (Zyphur *et al.* 2009;

478   Akinola *et al.* 2016). Nevertheless, our data here indicate that the increased prevalence of

dominant, high-testosterone individuals can predict changes to higher-order social structure.

480   These findings establish that hormone-behavior relationships are not limited to one individual,

but instead hormones may act as extended phenotypes that have population-level consequences

482   (McClintock 1981; Dawkins 1982; Robinson 1992). Taken together with previous studies, we

propose that testosterone can alter social network dynamics in ways that impede the evolution of

484   cooperation.

Figure Legends

Figure 1. Example social networks illustrating social specialization, network stability, and behavioral assortment. (A) Social specialization is measured using the bipartite form of the

social network. In the example on the left, the coalitions between floater males and territory-holders are poorly partitioned, creating a network with a relatively low specialization. On the

right, there is greater partitioning, such that each floater male maintains a smaller number of partnerships with specific territory-holding males. (B) Network stability is measured by

evaluating the persistence of partnerships from one recording session (t_1) to the next (t_2). The

example at the top of (B) has a lower stability than the example on the bottom. (C) Behavioral assortment measures the tendency of like to associate with like. In (C), nodes are shaded to

indicate a continuous gradient of daily behavior (in this case, each male's average daily strength is shown). In the example network on the left, the males tend to associate with behaviorally

dissimilar partners, yielding a negative behavioral assortment. In the example on the right, the males tend to associate with behaviorally similar partners, yielding a positive assortment. See

Table 1 for additional data.

Figure 2. Collective testosterone of manakin social networks. (A) Circulating testosterone varies among individual males. This graph shows repeated measures from 210 individual male

manakins, sorted along the x-axis by mean T, a standardized measure of a male's average testosterone level (Ryder *et al.* 2020). Opacity is used to denote the two male status classes (with

subordinate floaters colored semi-transparent, and dominant territory-holders colored opaque). Color ramping is used to denote each individual's hormone phenotype. (B-D) These data were

508 used to define collective testosterone in the present study. Node colors indicate an individual's

mean T following the scale in (A). The collective testosterone of each social network is

510 calculated as the average of the individual hormone phenotypes, weighted by strength, and is

shown below each example. Collective testosterone was determined for each status class

512 separately. See Table 1 for additional data.

514 **Figure 3. Collective testosterone predicts the structure of the social network.** The networks

with higher-testosterone dominant individuals were less specialized, less stable over time, and

516 had more negative behavioral assortment. Each plot shows the partial residuals from a statistical

analysis that also accounts for field season, the average Julian date of the recording session, the

518 number of recording hours, network size, and lek identity. Because floater and territorial males

can differ in behavior, the analysis of behavioral assortment also accounted for status assortment

520 within each network. Different symbols are used to indicate repeated measures of 11 different

leks, colored according the collective testosterone scale in Fig. 2. Inset panels show the results of

522 the node-label permutation tests. In each case, the slope in the best-fit model (solid vertical line)

is significantly more negative than expected based on the distribution of permuted estimates

524 (grey distribution). See Tables S1-S2 for additional data.

Acknowledgments

526   We thank Camilo Alfonso, Brian Evans, David and Consuelo Romo, Kelly Swing, Diego

Mosquera, Gabriela Vinueza, and Tiputini Biodiversity Station of the Universidad San Francisco

528   de Quito. Funding was provided by the National Science Foundation (NSF) IOS 1353085 and

the Smithsonian Migratory Bird Center.

Author Contributions

532   RD, ITM, BMH, and TBR designed the study. ITM, BMH, BJV, and TBR collected the data.

RD, ITM, and TBR analyzed the data. RD and TBR wrote the manuscript. All authors edited the

534   manuscript.

Declaration of Interests

The authors declare no competing interests.

Data Availability

540   All data and R scripts necessary to reproduce this study are available for download at:

https://figshare.com/s/13a311662fee686fa4f3

Supporting Information

544   Supplemental text, Figures S1-S4, and Table S1-S2 in the attached PDF

546   **References**

Adkins-Regan, E. (2005). *Hormones and animal social behavior*. Princeton University Press.
548   Akinola, M., Page-Gould, E., Mehta, P.H. & Lu, J.G. (2016). Collective hormonal profiles predict
            group performance. *PNAS*, 113, 9774–9779.
550   Aplin, L.M., Farine, D.R., Morand-Ferron, J. & Sheldon, B.C. (2012). Social networks predict
            patch discovery in a wild population of songbirds. *Proceedings of the Royal Society B:*
552   *Biological Sciences*, 279, 4199–4205.
Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R.H.B., Singmann, H., *et al.* (2018).
554   *lme4 1.1-18-1: linear mixed-effects models using "Eigen" and S4*.
Blüthgen, N., Menzel, F. & Blüthgen, N. (2006). Measuring specialization in species interaction
556   networks. *BMC Ecology*, 6, 9.
Boksem, M.A.S., Mehta, P.H., Van den Bergh, B., van Son, V., Trautmann, S.T., Roelofs, K., *et al.*
558   (2013). Testosterone Inhibits trust but promotes reciprocity. *Psychol Sci*, 24, 2306–2314.
Cohen, A.A., Martin, L.B., Wingfield, J.C., McWilliams, S.R. & Dunne, J.A. (2012). Physiological
560   regulatory networks: ecological roles and evolutionary constraints. *Trends in Ecology &*
            Evolution, 27, 428–435.
562   Croft, D.P., James, R., Thomas, P.O.R., Hathaway, C., Mawdsley, D., Laland, K.N., *et al.* (2006).
            Social structure and co-operative interactions in a wild population of guppies (*Poecilia*
564   *reticulata*). *Behav Ecol Sociobiol*, 59, 644–650.
Csardi, G. & coauthors. (2018). *igraph 1.2.2: network analysis and visualization.*
566   Dakin, R. & Ryder, T.B. (2018). Dynamic network partnerships and social contagion drive
            cooperation. *Proceedings of the Royal Society B: Biological Sciences*, 285, 20181973.
568   Dakin, R. & Ryder, T.B. (2020). Reciprocity and behavioral heterogeneity govern the stability of
            social networks. Preprint. https://doi.org/10.1101/694166. *Proceedings of the National*
570   *Academy of Sciences*.
Dawkins, R. (1982). *The Extended Phenotype: The Gene as the Unit of Selection*. W. H. Freeman
572   and Company, San Francisco.
Díaz-Muñoz, S.L., DuVal, E.H., Krakauer, A.H. & Lacey, E.A. (2014). Cooperating to compete:
574   altruism, sexual selection and causes of male reproductive cooperation. *Animal*
            Behaviour, 88, 67–78.
576   Dormann, C.F., Fruend, J. & Gruber, B. (2019). *bipartite 2.13: visualising bipartite networks and*
            calculating some (ecological) indices.
578   Durães, R., Loiselle, B.A., Parker, P.G. & Blake, J.G. (2009). Female mate choice across spatial
            scales: influence of lek and male attributes on mating success of blue-crowned
580   manakins. *Proceedings of the Royal Society B: Biological Sciences*, 276, 1875–1881.
DuVal, E.H. (2007). Adaptive advantages of cooperative courtship for subordinate male lance-
582   tailed manakins. *The American Naturalist*, 169, 423–432.
Eikenaar, C., Whitham, M., Komdeur, J., van der Velde, M. & Moore, I.T. (2011). Endogenous
584   testosterone is not associated with the trade-off between paternal and mating effort.
            Behav Ecol, 22, 601–608.
586   Eisenegger, C., Haushofer, J. & Fehr, E. (2011). The role of testosterone in social interaction.
            Trends in Cognitive Sciences, 15, 263–271.

588   Farine, D.R. (2014). Measuring phenotypic assortment in animal social networks: weighted
           associations are more robust than binary edges. *Animal Behaviour*, 89, 141–153.
590   Farine, D.R. (2015). Proximity as a proxy for interactions: issues of scale in social network
           analysis. *Animal Behaviour*, 104, e1–e5.
592   Farine, D.R. (2016). *assortnet 0.12: calculate the assortativity coefficient of weighted and binary
           networks*.
594   Farine, D.R., Freckleton, R. & Rands, S. (2017). A guide to null models for animal social network
           analysis. *Methods in Ecology and Evolution*, 8, 1309–1320.
596   Fowler, J.H. & Christakis, N.A. (2010). Cooperative behavior cascades in human social networks.
           PNAS, 107, 5334–5338.
598   Franks, D.W., Ruxton, G.D. & James, R. (2010). Sampling animal association networks with the
           gambit of the group. *Behav Ecol Sociobiol*, 64, 493–503.
600   Fuxjager, M.J., Forbes-Lorman, R.M., Coss, D.J., Auger, C.J., Auger, A.P. & Marler, C.A. (2010).
           Winning territorial disputes selectively enhances androgen sensitivity in neural
602   pathways related to motivation and social aggression. *PNAS*, 107, 12393–12398.
       Goymann, W. (2009). Social modulation of androgens in male birds. *General and Comparative
604   Endocrinology*, 9th International Symposium on Avian Endocrinology, 163, 149–157.
       Goymann, W., Schwabl, I., Trappschuh, M. & Hau, M. (2007). Use of ethanol for preserving
606   steroid and indoleamine hormones in bird plasma. *General and Comparative
           Endocrinology*, 150, 191–195.
608   Griffiths, S.W. & Magurran, A.E. (1997). Schooling preferences for familiar fish vary with group
           size in a wild guppy population. *Proceedings of the Royal Society of London. Series B:
610   Biological Sciences*, 264, 547–551.
       Heindl, M. (2002). Social organization on leks of the wire-tailed manakin in southern venezuela.
612   *The Condor*, 104, 772–779.
       Ilmarinen, V.-J., Vainikainen, M.-P., Verkasalo, M.J. & Lönnqvist, J.-E. (2017). Homophilous
614   friendship assortment based on personality traits and cognitive ability in middle
           childhood: the moderating effect of peer network size. *European Journal of Personality*,
616   31, 208–219.
       James, R., Croft, D.P. & Krause, J. (2009). Potential banana skins in animal social network
618   analysis. *Behav Ecol Sociobiol*, 63, 989–997.
       Jehn, K.A. & Shah, P.P. (1997). Interpersonal relationships and task performance: an
620   examination of mediation processes in friendship and acquaintance groups. *Journal of
           Personality and Social Psychology*, 72, 775–790.
622   Krause, J. & Ruxton, G.D. (2002). *Living in Groups*. OUP Oxford.
       Kuznetsova, A., Brockhoff, P.B. & Christensen, R.H.B. (2018). *lmerTest 3.0-1: tests in linear
624   mixed effects models*.
       Lank, D.B. & Smith, C.M. (1992). Females prefer larger leks: field experiments with ruffs
626   (*Philomachus pugnax*). *Behav Ecol Sociobiol*, 30, 323–329.
       Laskowski, K.L., Montiglio, P.-O. & Pruitt, J.N. (2016). Individual and group performance suffers
628   from social niche disruption. *The American Naturalist*, 187, 776–785.
       Lüdecke, D., Makowski, D. & Waggoner, P. (2019). *performance 0.4.2: assessment of regression
630   mdoels performance*.

McClintock, M.K. (1981). Social control of the ovarian cycle and the function of estrous
632   synchrony. *Integr Comp Biol*, 21, 243–256.

McDonald, D.B. (1993). Delayed plumage maturation and orderly queues for status: a manakin
634   mannequin experiment. *Ethology*, 94, 31–45.

McDonald, D.B. (2007). Predicting fate from early connectivity in a social network. *PNAS*, 104,
636   10910–10914.

McDonald, G.C., Farine, D.R., Foster, K.R. & Biernaskie, J.M. (2017). Assortment and the analysis
638   of natural selection on social traits. *Evolution*, 71, 2693–2702.

Nakagawa, S. & Schielzeth, H. (2013). A general and simple method for obtaining R2 from
640   generalized linear mixed-effects models. *Methods Ecol Evol*, 4, 133–142.

Ohtsuki, H., Hauert, C., Lieberman, E. & Nowak, M.A. (2006). A simple rule for the evolution of
642   cooperation on graphs and social networks. *Nature*, 441, 502–505.

Oyegbile, T.O. & Marler, C.A. (2005). Winning fights elevates testosterone levels in California
644   mice and enhances future ability to win fights. *Hormones and Behavior*, 48, 259–267.

Pinter-Wollman, N., Hobson, E.A., Smith, J.E., Edelman, A.J., Shizuka, D., de Silva, S., *et al.*
646   (2014). The dynamics of animal social networks: analytical, conceptual, and theoretical
       advances. *Behav Ecol*, 25, 242–255.

648 Poisot, T., Canard, E., Mouillot, D., Mouquet, N. & Gravel, D. (2012). The dissimilarity of species
       interaction networks. *Ecology Letters*, 15, 1353–1361.

650 R Core Team. (2018). *R 3.5.1: A Language and Environment for Statistical Computing*. R
       Foundation for Statistical Computing, Vienna, Austria.

652 Roberts, G. & Sherratt, T.N. (1998). Development of cooperative relationships through
       increasing investment. *Nature*, 394, 175.

654 Robinson, G.E. (1992). Regulation of division of labor in insect societies. *Annual Review of
       Entomology*, 37, 637–665.

656 Ryder, T.B., Blake, J.G., Parker, P.G. & Loiselle, B.A. (2011a). The composition, stability, and
       kinship of reproductive coalitions in a lekking bird. *Behav Ecol*, 22, 282–290.

658 Ryder, T.B., Dakin, R., Vernasco, B.J., Evans, B.S., Horton, B.M. & Moore, I.T. (2020).
       Testosterone modulates status-specific patterns of cooperation in a social network. *The
660   American Naturalist*.

Ryder, T.B., Horton, B.M. & Moore, I.T. (2011b). Understanding testosterone variation in a
662   tropical lek-breeding bird. *Biology Letters*, 7, 506–509.

Ryder, T.B., Horton, B.M., Tillaart, M. van den, Morales, J.D.D. & Moore, I.T. (2012). Proximity
664   data-loggers increase the quantity and quality of social network data. *Biology Letters*, 8,
       917–920.

666 Ryder, T.B., McDonald, D.B., Blake, J.G., Parker, P.G. & Loiselle, B.A. (2008). Social networks in
       the lek-mating wire-tailed manakin (*Pipra filicauda*). *Proceedings of the Royal Society B*,
668   275, 1367–1374.

Ryder, T.B., Parker, P.G., Blake, J.G. & Loiselle, B.A. (2009). It takes two to tango: reproductive
670   skew and social correlates of male mating success in a lek-breeding bird. *Proceedings of
       the Royal Society B*, 276, 2377–2384.

672 Sah, P., Leu, S.T., Cross, P.C., Hudson, P.J. & Bansal, S. (2017). Unraveling the disease
       consequences and mechanisms of modular structure in animal social networks. *Proc
Natl Acad Sci USA*, 114, 4165–4170.

Sah, P., Mann, J. & Bansal, S. (2018). Disease implications of animal social network structure: A synthesis across social systems. *J Anim Ecol*, 87, 546–558.

Schjelderup-Ebbe, T. (1922). Beiträge zur sozialpsychologie des haushuhns. [Observation on the social psychology of domestic fowls.]. *Zeitschrift für Psychologie und Physiologie der Sinnesorgane. Abt. 1. Zeitschrift für Psychologie*, 88, 225–252.

Schwartz, P. & Snow, D.W. (1978). Display and related behavior of the wire-tailed manakin. *Living Bird*, 17, 51–78.

Trainer, J.M. & McDonald, D.B. (1995). Singing performance, frequency matching and courtship success of long-tailed manakins (*Chiroxiphia linearis*). *Behav Ecol Sociobiol*, 37, 249–254.

Trainer, J.M., McDonald, D.B. & Learn, W.A. (2002). The development of coordinated singing in cooperatively displaying long-tailed manakins. *Behav Ecol*, 13, 65–69.

Trivers, R.L. (1971). The evolution of reciprocal altruism. *The Quarterly Review of Biology*, 46, 35–57.

Vernasco, B.J., Horton, B.M., Moore, I.T. & Ryder, T.B. (2020). Reduced cooperative behavior as a cost of high testosterone in a lekking passerine bird. *Behav Ecol*.

Vernasco, B.J., Horton, B.M., Ryder, T.B. & Moore, I.T. (2019). Sampling baseline androgens in free-living passerines: Methodological considerations and solutions. *General and*
Comparative Endocrinology, Endocrinology of Neotropical Vertebrates, 273, 202–208.

Wingfield, J.C., Hegner, R.E., Dufty, Alfred M. & Ball, G.F. (1990). The "challenge hypothesis":
694 theoretical implications for patterns of testosterone secretion, mating systems, and breeding strategies. *The American Naturalist*, 136, 829–846.

696 Zuur, A., Ieno, E.N., Walker, N., Saveliev, A.A. & Smith, G.M. (2009). *Mixed effects models and extensions in ecology with R*. Springer Science & Business Media.

698 Zyphur, M.J., Narayanan, J., Koh, G. & Koh, D. (2009). Testosterone–status mismatch lowers collective efficacy in groups: evidence from a slope-as-predictor multilevel structural
700 equation model. *Organizational Behavior and Human Decision Processes*, Biological Basis of Business, 110, 70–79.

Figure 1

A Social specialization

Low
0.08

High
0.83

Territory
-holders

Floaters

B Network stability

t_1

t_2

Low
0.33

High
0.88

C Behavioral assortment

Negative
-0.52

Positive
0.66

Avg. daily behavior

Figure 2

A

Log T (ng/ml)

Floaters
Territory-holders

Low T male ← Mean T → High T male

B Low T group

C

D High T group

Collective testosterone

Figure 3