A modular cloning toolkit for genome editing in plants

Florian Hahn[1], Andrey Korolev[1§], Laura Sanjurjo Loures[1] and Vladimir Nekrasov[1*]

[1]Plant Sciences Department, Rothamsted Research, Harpenden AL5 2JQ, United Kingdom
[§]Present address: John Innes Centre, Norwich Research Park, Norwich NR4 7UH, United Kingdom
*Correspondence: vladimir.nekrasov@rothamsted.ac.uk
Tel.: +44 (0)1582 938 292
FH ORCID: 0000-0002-0215-3209; VN ORCID: 0000-0001-9386-1683

Keywords: CRISPR, Cas9, plant, genome editing, Golden Gate, MoClo

Abstract

The modular cloning (MoClo), based on the Golden Gate (GG) method, has enabled development of cloning systems with standardised genetic parts, e.g. promoters, coding sequences or terminators, that can be easily interchanged and assembled into expression units, which in their own turn can be further assembled into higher order multigene constructs. Here we present an expanded cloning toolkit that contains modules encoding a variety of CRISPR/Cas-based nucleases and their corresponding guide RNA backbones. Among other components, the toolkit includes a number of promoters that allow expression of CRISPR/Cas nucleases (or any other coding sequences) and their guide RNAs in monocots and dicots. As part of the toolkit, we present a set of modules that enable quick and facile assembly of tRNA-sgRNA polycistronic units without a PCR step involved. We believe the toolkit will contribute towards wider adoption of the CRISPR/Cas genome editing technology and modular cloning by researchers across the plant science community.

Introduction

The CRISPR/Cas technology has recently become an easily accessible genome editing tool for many organisms, including plants (Belhaj et al., 2015). Generating gene knockouts has become a rather straightforward CRISPR/Cas application in many plant systems (Hahn et al., 2017; Nekrasov et al., 2013, 2017), while more sophisticated applications, such as allele replacements or targeted gene insertions, still remain a challenge due to low efficiency of homology-directed repair (HDR) in plants (Hahn et al., 2018).

In its conventional form, the CRISPR/Cas system includes a DNA nuclease, such as Cas9, which is guided to a specific genomic location by the guide RNA. Therefore, in order to perform targeted mutagenesis *in planta*, one needs to co-express both the CRISPR/Cas nuclease and its cognate guide RNA. Usually, the gene encoding the CRISPR/Cas nuclease is expressed

using an RNA polymerase II (Pol II) promoter (e.g. 35Sp), while the guide RNA is expressed under an RNA polymerase III (Pol III) promoter (e.g. U6p or U3p), which has a defined transcription start nucleotide ('G' for U6p or 'A' for U3p). One of the advantages of CRISPR/Cas is multiplexing i.e. one can target DNA at multiple genomic locations by co-expressing multiple guide RNAs specific to those loci. Guide RNAs can be expressed either as individual transcriptional units, each under its own Pol III promoter (Nekrasov et al., 2017), or as a tRNA-sgRNA polycistronic transcript (Xie et al., 2015). In the latter case, guide RNAs are interspaced with tRNAs in a single transcript that gets processed into individual guide RNAs by the highly conserved tRNA processing machinery inside the plant cell (Xie et al., 2015).

As genome editing applications in plants often rely on delivering multiple expression units into plant cells, including a selectable marker, a CRISPR/Cas nuclease-encoding gene and one or more guide RNAs, it is important to be able to assemble DNA constructs encoding such expression units easily and rapidly. The modular cloning (MoClo) system based on the Golden Gate (GG) cloning method (Weber et al., 2011) is highly flexible and versatile, and provides a means for quick and facile assembly of multi-expression unit constructs using standard genetic parts, such as promoters, terminators, coding sequences etc. The system has already been successfully used for genome editing applications in plants (Brooks et al., 2014; Castel et al., 2019; Nekrasov et al., 2013, 2017; Raitskin et al., 2019) but lacks modules encoding many of the newest genome editing reagents. Here we report on an expanded GG cloning toolkit for genome editing applications in monocot and dicot plants. We believe the toolkit will become a valuable addition to already existing GG-based tools for plant genome editing and be widely used by plant researchers across the community.

Materials and methods

DNA construct assembly

All PCR amplifications were performed using Q5® DNA Polymerase (New England Biolabs) according to the manufacturer's instruction. All GG cut-ligation reactions were performed according to the protocol described in the Supplementary methods.

All ligations were transformed into One Shot™ TOP10 chemically competent *E. coli* (Thermo Fisher Scientific) and constructs were verified by sequencing (Eurofins Genomics).

Specific details related to assembly of all GG modules reported in this study are provided in the Supplementary methods. All PCR primers used in the study are listed in the Table S4. All DNA constructs generated as part of this study were deposited with Addgene (www.addgene.org) with

Addgene IDs indicated for each construct in the Table S5. Sequence information of all constructs can be found in GenBank (.gb) supplementary files.

Protoplast assay

Protoplasts were isolated from 10 day old, etiolated wheat seedlings (cv. Cadenza) as previously described (Shan et al., 2014) with some modifications. Cellulase R10 and Macerozyme R10 were obtained from Duchefa Biochemie (Haarlem, the Netherlands) and the enzymatic digestion was performed at 26 °C for 4 h. Subsequently, 50,000 protoplasts in a volume of 100 µL were transformed with 20 µg of each plasmid (2 µg/µL) purified using the Maxi kit (Qiagen, Germany). One batch of protoplasts was treated with an equivalent amount of water and used as the negative (untransformed) control. Then, protoplasts were cultured in 6-well plates for 60 hours. DNA was isolated by pelleting the protoplasts for 4 min at 16,000 g, resuspending the pellet in 300 µL TPS buffer (100 mM Tris-HCl, pH 8, 10 mM EDTA, 1 M KCl) and then incubating for 3 min at 50 C. At the following step, DNA was precipitated by adding 300 µL of isopropanol and incubating at room temperature for 20 min. The precipitated DNA was pelleted by centrifuging at 16,000 g for 20 min and the DNA pellet was washed with 70% ethanol. After a 5 min centrifugation at 16,000 g and drying the pellet, DNA was resuspended in 20 µL of the TE buffer (10 mM Tris-HCl, pH 7.5, 1 mM EDTA). CRISPR/Cas induced mutations were detected by PCR amplification of the target genes using the DreamTaq DNA polymerase (ThermoFisher Scientific) and primer pairs FH432/FH433 (Target gene 1), FH436/FH437 (Target gene 2) and FH440/FH441 (Target gene 3) (Table S4).

Results

During this study we have generated a set of GG modules that enable one to perform genome editing in both monocot and dicot plant species (Table S5). The cloning toolkit is an addition to previously published GG modules (Castel et al., 2019; Engler et al., 2014; Raitskin et al., 2019; Weber et al., 2011) and includes new sequence-specific CRISPR/Cas nucleases (codon-optimised for both monocots and dicots), Pol II and Pol III promoters, as well as guide RNA backbone modules. The latter enable insertion of the guide sequence by cloning in a pair of annealed complimentary oligos without a PCR amplification step involved. The toolkit enables assembly of CRISPR/Cas constructs that target a single as well as multiple targets with guide RNAs expressed either under individual Pol III promoters (Figure 1) or using a polycistronic tRNA-sgRNA construct (Figure 4).

CRISPR/Cas nuclease modules

The CRISPR/Cas nuclease level 0 modules include *Streptococcus pyogenes* Cas9 (SpCas9; pFH13, pFH24 and pFH25) as well as Cas9 variants coming from other bacterial species: *Staphylococcus aureus* (SaCas9; pFH14), *Streptococcus thermophilus* Cas9 (StCas9; pFH15) and *Streptococcus canis* (ScCas9; pFH76) (Table S5). SpCas9 (Jinek et al., 2012) with the 'NGG' protospacer adjacent motif (PAM) is the most commonly used Cas9 variant for genome editing applications in various organisms, including plants. SaCas9 ('NNGRRT' PAM) and StCas9 ('NNRGAA' PAM) are less common but have also been successfully used in rice, tobacco (Kaya et al., 2016) and Arabidopsis (Schmidt et al., 2019; Steinert et al., 2015; Wolter et al., 2018). ScCas9 (pFH76), SpCas9-NG (pFH32) and SpCas9-derived xCas9 (pFH22) are characterised by broadened PAM motif requirements: 'NNG' for ScCas9 (Chatterjee et al., 2018), and 'NG' for SpCas9-NG (Nishimasu et al., 2018) and xCas9 (Hu et al., 2018). We have also included modules with Cas12a (Cpf1) CRISPR nucleases from *Francisella novicida* (FnCas12a; pFH16 and pFH46) and *Lachnospiraceae bacterium* (LbCas12a; pFH17 and pFH47) as well as with four related Cms1 nucleases (pFH18-21) (Table S5). LbCas12a (Tang et al., 2017), FnCas12a (Endo et al., 2016) and Cms1 (Begemann et al., 2017b) have all been shown to work in plants.

Base editors are a rather recent addition to the range of available genome editing tools and allow targeted conversion of DNA base pairs as following: C-G to T-A (Komor et al., 2016) and A-T to G-C (Gaudelli et al., 2017) without introducing a double-strand break (DSB). The former base editor is based on the cytidine deaminase while the latter – on the adenosine deaminase. Both base editors have now been shown to be functional in various plants, including wheat, rice and tomato (Hua et al., 2019; Li et al., 2018; Shimatani et al., 2017; Zong et al., 2017, 2018). We have therefore generated level 0 modules encoding cytidine deaminase (pFH55 and pFH79) and adenosine deaminase (pFH45 and pFH92) based base editors (Table S5).

EvolvR CRISPR-guided error-prone DNA polymerases have recently been shown to be able to introduce random point mutations at a targeted genomic locus (Halperin et al., 2018). Based on the Halperin et al. (2018) manuscript, we have generated a level 0 module with the wheat codon optimised version of enCas9–PolI3M–TBD (pFH77, Table S5) that could prove to be a useful tool for reverse genetics in monocot plants.

We used the above mentioned CRISPR/Cas nuclease level 0 modules to assemble a number of expression units inserted into level 1 GG vectors to be applied in monocot and dicot plant species (Table S5).

Guide RNA modules

As CRISPR/Cas is an RNA-guided nuclease, guide RNA is its essential component that must be co-expressed with the nuclease in order to achieve on-target DNA cutting. Guide RNAs are usually expressed under Pol III promoters, such as U3p or U6p, that have a defined transcription start nucleotide ('A' and 'G', respectively). A number of genomic loci can be targeted simultaneously by CRISPR/Cas by co-expressing multiple guide RNAs and the modular cloning system is highly suitable for assembling constructs carrying multiple expression units, such as the CRISPR/Cas nuclease and guide RNAs.

As part of this study, we have generated a number of level 0 Pol III promoter modules (TaU3p, OsU3p, OsU6-2p and AtU6-26p; Table S5). In addition, we have produced a number of guide RNA backbone level 0 constructs that can be used to assemble single or multiple guide RNA expression units without a PCR amplification step involved (Table S5). The cloning system we present allows guide RNAs to be expressed either under individual Pol III promoters (Figures 1-3; Supplementary methods, section 2) or from a polycistronic tRNA-sgRNA construct, which includes sgRNAs interspaced with tRNA scaffolds (Xie et al., 2015) (Figures 4 and 5; Supplementary Methods, section 3). The method with guide RNAs expressed under individual promoters enables expression of sgRNAs of the Cas9 family of CRISPR/Cas nucleases, which carry the guide sequence at the 5' end of sgRNA (Figure 2), as well as crRNAs of Cas12a (Cpf1) nucleases and related Cms1 nucleases, which have the guide sequence at the 3' end of crRNA (Figure 3). Up to four guide RNAs under individual Pol III promoters can be assembled in using the former cloning procedure (Figure 1B) and up to six sgRNAs per polycistronic construct – using the latter one (Figure 4B). It must be noted that the number of guide RNAs under individual promoters could be increased to five, if no selectable marker is needed (Figure 1B), or many more if level M/ level P vectors are used (Engler et al., 2014; Werner et al., 2012). As to tRNA-sgRNA polycistronic constructs, the total number of sgRNAs assembled into a level 2 destination vector could be up to twenty four (six per level 1), with a selectable marker, and up to thirty, if no selectable marker is used. Again, it is possible to add more than thirty sgRNAs by using level M/ level P vectors (Engler et al., 2014; Werner et al., 2012). Our GG toolkit enables the user to build such complex constructs within a week (Figure S1).

Testing of the tRNA-sgRNA CRISPR/Cas constructs in wheat protoplasts

The GG toolkit therefore allows rapid parallel assembly of constructs by streamlining the cloning process. Since building multiple CRISPR constructs using GG is a straightforward procedure, it becomes reasonable to compare the activity of several experimental CRISPR setups in a transient expression system, such as protoplast, before proceeding with stable plant transformation, which could be a highly laborious and time consuming process. Our CRISPR toolkit includes three wheat codon optimised SpCas9 versions (level 0 constructs pFH13, pFH24

and pFH25; Table S5) and their respective level 1 transcription units (pFH23, pFH66 and pFH67; Table S5). These SpCas9 variants differ by e.g. nuclear localisation signal (NLS) versions or affinity tags. We have therefore compared the activity of the three Cas9 variants in wheat protoplasts by cotransforming each of the level 1 constructs (pFH23, pFH66 and pFH67) with the level 1 plasmid containing the six sgRNAs (Figure 6A) assembled into a tRNA-sgRNA array. This has allowed us to target three different wheat genes at once (Figure 6A). We have targeted each gene by at least two sgRNAs with large deletions between Cas9 cut sites expected to be detectable by PCR due to DNA band shifts as previously described (Nekrasov et al., 2017). PCR amplification of the target genes has revealed clear additional bands corresponding to alleles carrying large CRISPR/Cas-induced deletions in protoplasts transformed with pFH66. In contrast, application of the other two Cas9 versions (pFH23 and pFH67) resulted in very faint bands of the size corresponding to amplicons carrying the deletions (Figure 6B). Our results therefore suggest a significantly higher activity of the pFH66-encoded SpCas9, as compared to the other two Cas9 variants, in wheat protoplasts.

Discussion

The modular cloning kit presented in the study enables quick and facile assembly of DNA constructs for genome editing applications in plants and is an addition to previously published collections of compatible GG modules (Castel et al., 2019; Engler et al., 2014; Raitskin et al., 2019; Weber et al., 2011). The kit includes modules encoding a number of CRISPR/Cas nucleases (SaCas9, StCas9, LbCas12a etc.) that could be used as an alternative to the most commonly utilised SpCas9. SaCas9, for instance, has proven to be an efficient tool for generating gene knockouts in a number of plant species (Kaya et al., 2016; Schmidt et al., 2019; Steinert et al., 2015; Wolter et al., 2018) and, in addition, has been shown to increase HDR efficiencies in plants (Wolter et al., 2018). Due to SaCas9 and StCas9 having longer than SpCas9 PAM motifs they are also likely to be more specific when it comes to DNA target recognition.

Cas12a (Cpf1) generates a staggered cut in DNA (Zetsche et al., 2015), while Cas9 – a blunt cut (Jinek et al., 2012). Due to this reason, Cas12a (Cpf1) could be a preferred choice of a CRISPR/Cas nuclease when it comes to HDR-based genome editing applications, such as targeted gene insertion (Begemann et al., 2017a; Wolter and Puchta, 2019). It is noteworthy that the modular cloning system is highly suitable for HDR-based applications as the DNA repair template could easily be cloned as a level 1 module into a level 2 destination vector. Also, since Cas12a (Cpf1) has a T-rich PAM motif ('TTTN' for LbCas12a (LbCpf1) and 'TTN' for

FnCas12a (FnCpf1)), it could be a better choice for targeted mutagenesis in plant species with AT-rich genomes as compared to SpCas9 ('NGG' PAM).

As part of our study, we have generated a number of guide RNA backbone level 0 modules, which are compatible with respective Pol III promoters and CRISPR/Cas nucleases (Table S1). The guide RNA backbone modules can be used for PCR-free assembly of guide RNA expression units by cloning in an annealed pair of complimentary oligos encoding the guide sequence (Figures 2 and 3). These guide RNA backbones are to be used when one wishes to express guide RNAs under individual promoters and up to four guide RNAs can be assembled into a level 2 vector together with level 1 modules encoding a CRISPR/Cas nuclease and a selectable marker (e.g. BAR, NPTII etc.; Figure 1, Tables S5 and S6).

In addition to expressing each guide RNA under its own promoter, we have generated modules that allow assembly of polycistronic tRNA-sgRNA constructs with up to six sgRNAs expressed using a single Pol III promoter (Figure 4). The tRNA-sgRNA system, originally described by Xie et al. (2015) in rice, was later successfully applied in wheat (Wang et al., 2018) and Arabidopsis (Zhang et al., 2018). Nevertheless, the previously reported tRNA-sgRNA system relies on a rather cumbersome DNA construct assembly process as it involves PCR amplification of DNA fragments carrying repeats. The level 0 GG modules we have generated (Figure 5 and Table S3) enable straightforward and efficient assembly of tRNA-sgRNA arrays without a PCR step involved. The system offers a choice of five monocot and dicot Pol III promoters, with TaU6p being a published module (Lawrenson et al., 2015), and two different SpCas9 sgRNA backbones (classic and improved; Table S3).

As stable transformation continues to be a major bottleneck for genome editing applications in many plants, including a major crop like wheat (Sparks and Jones, 2014), it is advantageous to be able to test CRISPR/Cas constructs for activity in a transient expression system, such as protoplasts, before initiating an often lengthy and labour intensive stable transformation procedure. Using the wheat protoplast system, we have compared activity of three different wheat codon optimised SpCas9 variants (Figure 6), which mostly differ in their C-terminal NLSs. The fact that one of the constructs (pFH66) performed better than the other two (Figure 6B) could be due to the histone H2B NLS, located at the C-terminus of the pFH66 variant, being more efficient at importing Cas9 into the nucleus as compared to the SV40 or nucleoplasmin NLS present in the other two constructs. A possible link between different NLS versions and Cas9 activity was previously reported in Arabidopsis (Castel et al., 2019).

The tRNA-gRNA system for Cas9 multiplexing has proven to work in monocots (Wang et al., 2018; Xie et al., 2015) as well as in dicots (Zhang et al., 2018). In dicots, it was shown that fusing tRNAs with the optimised sgRNA backbone (Dang et al., 2015) increased editing efficiencies (Zhang et al., 2018). In monocots however, only the classic sgRNA backbone (Mali

et al., 2013) was so far used in tRNA-sgRNA arrays. The wheat protoplast assay has allowed us to verify the functionality of the tRNA-sgRNA array, carrying the optimised sgRNA backbone, in a monocot species. Our results suggest that the tRNA-sgRNA array, assembled using the optimised sgRNA backbone, could also result in higher CRISPR/Cas efficiencies in stably transformed monocot plants, particularly wheat.

We believe the presented modular cloning kit will become a valuable addition to the range of already available GG modules (Castel et al., 2019; Raitskin et al., 2019; Weber et al., 2011) and expect that plant researchers, working with both monocots and dicots, will find the presented molecular tools useful for various genome editing applications. We also believe our study will contribute towards wider adoption of the GG modular cloning system by plant researchers and consequently facilitate exchange of standardised molecular cloning parts across the research community.

Author contributions

VN conceived the idea of the manuscript; FH designed the pFH-constructs and performed the experiments; AK designed the pAK-constructs and assisted with cloning; LSL performed protoplast experiments together with FH and assisted with cloning. VN and FH wrote the manuscript.

Acknowledgments

We thank Lucy Hyde (University of Bristol) for her contribution of managing the vector database and in her help at establishing the protoplast protocol. We thank Robert Hoffie (IPK Gatersleben) for the protoplast DNA extraction protocol and helpful discussions.

Funding

Vladimir Nekrasov (Rothamsted Research) receives grant-aided support from the Biotechnology and Biological Sciences Research Council (BBSRC) Designing Future Wheat project [BB/P016855/1] and Newton Fund.

Conflict of interest statement

FH, AK, LSL and VN declare no conflict of interest.

Figure legends

Figure 1 Cartoons of level 2 CRISPR/Cas constructs for targeted mutagenesis using guide RNAs expressed under individual Pol III promoters. (A) represents a construct with one and (B) – with four guide RNAs.

Figure 2 Assembly of level 1 sgRNA expression modules to be used with respective Cas9 nucleases. During the first step (A) annealed complementary oligos encoding the guide are inserted into a level 0 acceptor with the sgRNA backbone using BpiI. During the second step (B), sgRNA is fused with the respective Pol III promoter using BsaI.

Figure 3 Assembly of level 1 crRNA expression modules to be used with respective Cas12a (Cpf1) and Cms1 nucleases. During the first step (A) annealed complementary oligos encoding the guide are inserted into a level 0 acceptor carrying the sgRNA backbone using BpiI. During the second step (B), sgRNA is fused with the respective Pol III promoter using BsaI.

Figure 4 Cartoons of level 2 CRISPR/Cas constructs for targeted mutagenesis using the polycistronic SpCas9 tRNA-sgRNA system. (A) represents a construct with one and (B) – with six sgRNAs.

Figure 5 Assembly of level 1 polycistronic tRNA-sgRNA expression modules to be used with SpCas9. During the first step (A) annealed complementary oligos encoding the guide are inserted into a level 0 acceptor with the tRNA-sgRNA backbone using BpiI. During the second step (B), tRNA-sgRNA modules are fused with the respective Pol III promoter using BsaI. Assembly of 1 to 5 tRNA-sgRNA modules into a level 1 vector requires respective endlinkers (C). pFH and pAK constructs shown in black font carry the improved sgRNA backbone (Dang et al., 2015), while the ones shown in blue font – the classic sgRNA backbone (Mali et al., 2013). *This is a published module (Lawrenson et al., 2015).

Figure 6 Testing of tRNA-sgRNA CRISPR/Cas constructs in wheat protoplasts. Three wheat genes (A) were targeted: Target gene 1 (homoeologues *TraesCS1A02G338200*, *TraesCS1B02G350600* and *TraesCS1D02G340400*), Target gene 2 (homoeologues *TraesCS3A02G289300*, *TraesCS3B02G323900* and *TraesCS3D02G289100*) and Target gene 3 (homoeologues *TraesCS5A02G116500*, *TraesCS5B02G117800* and *TraesCS5D02G129600*). Red arrows illustrate positions of the 20 bp sgRNA target sites. Dashed lines illustrate expected CRISPR/Cas-induced deletions. (B) Wheat protoplasts were co-transformed with a level 1 construct carrying one of the three SpCas9 variants (pFH23, pFH66 or pFH67) and the level 1 construct (pFH94) carrying the six sgRNAs (A) in a tRNA-sgRNA array. Genotyping by PCR has revealed DNA shifted bands (marked by red asterisks) corresponding to amplicons carrying CRISPR/Cas-induced deletions of expected sizes. 'M' is the DNA marker; 'NTC' is the no template control.

References

Begemann, M. B., Gray, B. N., January, E., Gordon, G. C., He, Y., Liu, H., et al. (2017a). Precise insertion and guided editing of higher plant genomes using Cpf1 CRISPR nucleases. *Scientific Reports* 7, 11606. doi:10.1038/s41598-017-11760-6.

Begemann, M. B., Gray, B. N., January, E., Singer, A., Kesler, D. C., He, Y., et al. (2017b). Characterization and Validation of a Novel Group of Type V, Class 2 Nucleases for in vivo Genome Editing. *bioRxiv*, 192799. doi:10.1101/192799.

Belhaj, K., Chaparro-Garcia, A., Kamoun, S., Patron, N. J., and Nekrasov, V. (2015). Editing plant genomes with CRISPR/Cas9. *Current Opinion in Biotechnology* 32, 76–84. doi:http://dx.doi.org/10.1016/j.copbio.2014.11.007.

Brooks, C., Nekrasov, V., Lippman, Z. B., and Van Eck, J. (2014). Efficient Gene Editing in Tomato in the First Generation Using the Clustered Regularly Interspaced Short Palindromic Repeats/CRISPR-Associated9 System. *Plant physiology* 166, 1292–1297. doi:10.1104/pp.114.247577.

Castel, B., Tomlinson, L., Locci, F., Yang, Y., and Jones, J. D. G. (2019). Optimization of T-DNA architecture for Cas9-mediated mutagenesis in Arabidopsis. *PLOS ONE* 14, e0204778. doi:10.1371/journal.pone.0204778.

Chatterjee, P., Jakimo, N., and Jacobson, J. M. (2018). Minimal PAM specificity of a highly similar SpCas9 ortholog. *Science Advances* 4, eaau0766. doi:10.1126/sciadv.aau0766.

Dang, Y., Jia, G., Choi, J., Ma, H., Anaya, E., Ye, C., et al. (2015). Optimizing sgRNA structure to improve CRISPR-Cas9 knockout efficiency. *Genome Biology* 16, 280. doi:10.1186/s13059-015-0846-3.

Endo, A., Masafumi, M., Kaya, H., and Toki, S. (2016). Efficient targeted mutagenesis of rice and tobacco genomes using Cpf1 from *Francisella novicida*. *Scientific Reports* 6, srep38169. doi:10.1038/srep38169.

Engler, C., Youles, M., Gruetzner, R., Ehnert, T.-M., Werner, S., Jones, J. D. G., et al. (2014). A Golden Gate Modular Cloning Toolbox for Plants. *ACS Synth. Biol.* 3, 839–843. doi:10.1021/sb4001504.

Gaudelli, N. M., Komor, A. C., Rees, H. A., Packer, M. S., Badran, A. H., Bryson, D. I., et al. (2017). Programmable base editing of A•T to G•C in genomic DNA without DNA cleavage. *Nature* 551, 464. doi:10.1038/nature24644.

Hahn, F., Eisenhut, M., Mantegazza, O., and Weber, A. P. M. (2018). Homology-Directed Repair of a Defective Glabrous Gene in Arabidopsis With Cas9-Based Gene Targeting. *Front. Plant Sci.* 9. doi:10.3389/fpls.2018.00424.

Hahn, F., Mantegazza, O., Greiner, A., Hegemann, P., Eisenhut, M., and Weber, A. P. M. (2017). An Efficient Visual Screen for CRISPR/Cas9 Activity in Arabidopsis thaliana. *Front. Plant Sci.* 8. doi:10.3389/fpls.2017.00039.

Halperin, S. O., Tou, C. J., Wong, E. B., Modavi, C., Schaffer, D. V., and Dueber, J. E. (2018). CRISPR-guided DNA polymerases enable diversification of all nucleotides in a tunable window. *Nature* 560, 248–252. doi:10.1038/s41586-018-0384-8.

Hu, J. H., Miller, S. M., Geurts, M. H., Tang, W., Chen, L., Sun, N., et al. (2018). Evolved Cas9 variants with broad PAM compatibility and high DNA specificity. *Nature* 556, 57–63. doi:10.1038/nature26155.

Hua, K., Tao, X., and Zhu, J.-K. (2019). Expanding the base editing scope in rice by using Cas9 variants. *Plant Biotechnology Journal* 17, 499–504. doi:10.1111/pbi.12993.

Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J. A., and Charpentier, E. (2012). A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* 337, 816–21. doi:10.1126/science.1225829.

Kaya, H., Mikami, M., Endo, A., Endo, M., and Toki, S. (2016). Highly specific targeted mutagenesis in plants using *Staphylococcus aureus* Cas9. *Scientific Reports* 6, srep26871. doi:10.1038/srep26871.

Komor, A. C., Kim, Y. B., Packer, M. S., Zuris, J. A., and Liu, D. R. (2016). Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage. *Nature* 533, nature17946. doi:10.1038/nature17946.

Lawrenson, T., Shorinola, O., Stacey, N., Li, C., Østergaard, L., Patron, N., et al. (2015). Induction of targeted, heritable mutations in barley and Brassica oleracea using RNA-guided Cas9 nuclease. *Genome Biology* 16, 258. doi:10.1186/s13059-015-0826-7.

Li, C., Zong, Y., Wang, Y., Jin, S., Zhang, D., Song, Q., et al. (2018). Expanded base editing in rice and wheat using a Cas9-adenosine deaminase fusion. *Genome Biology* 19, 59. doi:10.1186/s13059-018-1443-z.

Mali, P., Yang, L., Esvelt, K. M., Aach, J., Guell, M., DiCarlo, J. E., et al. (2013). RNA-guided human genome engineering via Cas9. *Science* 339, 823–6. doi:10.1126/science.1232033.

Nekrasov, V., Staskawicz, B., Weigel, D., Jones, J. D., and Kamoun, S. (2013). Targeted mutagenesis in the model plant Nicotiana benthamiana using Cas9 RNA-guided endonuclease. *Nat Biotechnol* 31, 691–3. doi:10.1038/nbt.2655.

Nekrasov, V., Wang, C., Win, J., Lanz, C., Weigel, D., and Kamoun, S. (2017). Rapid generation of a transgene-free powdery mildew resistant tomato by genome deletion. *Scientific Reports* 7, 482. doi:10.1038/s41598-017-00578-x.

Nishimasu, H., Shi, X., Ishiguro, S., Gao, L., Hirano, S., Okazaki, S., et al. (2018). Engineered CRISPR-Cas9 nuclease with expanded targeting space. *Science*, eaas9129. doi:10.1126/science.aas9129.

Raitskin, O., Schudoma, C., West, A., and Patron, N. J. (2019). Comparison of efficiency and specificity of CRISPR-associated (Cas) nucleases in plants: An expanded toolkit for precision genome engineering. *PLOS ONE* 14, e0211598. doi:10.1371/journal.pone.0211598.

Schmidt, C., Pacher, M., and Puchta, H. (2019). Efficient induction of heritable inversions in plant genomes using the CRISPR/Cas system. *The Plant Journal* 98, 577–589. doi:10.1111/tpj.14322.

Shan, Q., Wang, Y., Li, J., and Gao, C. (2014). Genome editing in rice and wheat using the CRISPR/Cas system. *Nature protocols* 9, 2395–2410. doi:10.1038/nprot.2014.157.

Shimatani, Z., Kashojiya, S., Takayama, M., Terada, R., Arazoe, T., Ishii, H., et al. (2017). Targeted base editing in rice and tomato using a CRISPR-Cas9 cytidine deaminase fusion. *Nat Biotech* 35, 441–443. doi:10.1038/nbt.3833.

Sparks, C. A., and Jones, H. D. (2014). "Genetic Transformation of Wheat via Particle Bombardment," in *Cereal Genomics: Methods and Protocols* Methods in Molecular Biology., eds. R. J. Henry and A. Furtado (Totowa, NJ: Humana Press), 201–218. doi:10.1007/978-1-62703-715-0_17.

Steinert, J., Schiml, S., Fauser, F., and Puchta, H. (2015). Highly efficient heritable plant genome engineering using Cas9 orthologues from Streptococcus thermophilus and Staphylococcus aureus. *Plant J* 84, 1295–1305. doi:10.1111/tpj.13078.

Tang, X., Lowder, L. G., Zhang, T., Malzahn, A. A., Zheng, X., Voytas, D. F., et al. (2017). A CRISPR–Cpf1 system for efficient genome editing and transcriptional repression in plants. *Nature Plants* 3, nplants201718. doi:10.1038/nplants.2017.18.

Wang, W., Pan, Q., He, F., Akhunova, A., Chao, S., Trick, H., et al. (2018). Transgenerational CRISPR-Cas9 Activity Facilitates Multiplex Gene Editing in Allopolyploid Wheat. *The CRISPR Journal* 1, 65–74. doi:10.1089/crispr.2017.0010.

Weber, E., Engler, C., Gruetzner, R., Werner, S., and Marillonnet, S. (2011). A modular cloning system for standardized assembly of multigene constructs. *PLoS One* 6, e16765. doi:10.1371/journal.pone.0016765.

Werner, S., Engler, C., Weber, E., Gruetzner, R., and Marillonnet, S. (2012). Fast track assembly of multigene constructs using Golden Gate cloning and the MoClo system. *Bioengineered* 3, 38–43. doi:10.4161/bbug.3.1.18223.

Wolter, F., Klemm, J., and Puchta, H. (2018). Efficient in planta gene targeting in Arabidopsis using egg cell-specific expression of the Cas9 nuclease of Staphylococcus aureus. *The Plant Journal* 94, 735–746. doi:10.1111/tpj.13893.

Wolter, F., and Puchta, H. (2019). In planta gene targeting can be enhanced by the use of CRISPR/Cas12a. *The Plant Journal*. doi:10.1111/tpj.14488.

Xie, K., Minkenberg, B., and Yang, Y. (2015). Boosting CRISPR/Cas9 multiplex editing capability with the endogenous tRNA-processing system. *Proc Natl Acad Sci U S A* 112, 3570–5. doi:10.1073/pnas.1420294112.

Zetsche, B., Gootenberg, J. S., Abudayyeh, O. O., Slaymaker, I. M., Makarova, K. S., Essletzbichler, P., et al. (2015). Cpf1 Is a Single RNA-Guided Endonuclease of a Class 2 CRISPR-Cas System. *Cell* 163, 759–771. doi:10.1016/j.cell.2015.09.038.

Zhang, Q., Xing, H.-L., Wang, Z.-P., Zhang, H.-Y., Yang, F., Wang, X.-C., et al. (2018). Potential high-frequency off-target mutagenesis induced by CRISPR/Cas9 in *Arabidopsis* and its prevention. *Plant Mol Biol* 96, 445–456. doi:10.1007/s11103-018-0709-x.

Zong, Y., Song, Q., Li, C., Jin, S., Zhang, D., Wang, Y., et al. (2018). Efficient C-to-T base editing in plants using a fusion of nCas9 and human APOBEC3A. *Nature Biotechnology* 36, 950–953. doi:10.1038/nbt.4261.

Zong, Y., Wang, Y., Li, C., Zhang, R., Chen, K., Ran, Y., et al. (2017). Precise base editing in rice, wheat and maize with a Cas9-cytidine deaminase fusion. *Nat Biotech* 35, 438–440. doi:10.1038/nbt.3811.

A

Level 2 construct

B

Level 2 construct

Figure 1

A

Target specific annealed primers can be integrated into a level 0 acceptor vectors via BpiI

Level 0 module

B

The gRNA module can be assembled with a Pol III promoter in a level 1 acceptor via BsaI

Level 1 module

Figure 2

A

Target specific annealed primers can be integrated into level 0 acceptor vectors via BpiI

Level 0 module

B

The crRNA module can be assembled with a Pol III promoter in a level 1 vector via BsaI

Level 1 module

Figure 3

A

Level 2 construct

B

Level 2 construct

Figure 4

A **Target specific annealed primers can be integrated into a level 0 acceptor vectors via BpiI**

B **Up to six sgRNA modules can be assembled into a level 1 acceptor vector via BsaI**

C **Assembly of 1 to 5 tRNA-sgRNA modules into a level 1 vector requires endlinkers**

Figure 5

Figure 6