Paternal aging causes atypical development of vocal

communication in neonatal mice

Lingling Mai [1], Ryuichi Kimura[1], Hitoshi Inada[1], Kouta Kanno[2], Takeru Matsuda[3,4],

Ryosuke O. Tachibana[5], Kaichi Yoshizaki[6], Fumiyasu Komaki[3,4], Noriko Osumi[1]

[1]Department of Developmental Neuroscience, Graduate School of Medicine, Tohoku

University, Sendai, Japan

[2]Faculty of Law, Economic and Humanities, Kagoshima University, Kagoshima,

Japan

[3]Department of Mathematical Informatics, Graduate School of Information Science

and Technology, The University of Tokyo, Tokyo, Japan

[4]Mathematical Informatics Collaboration Unit, RIKEN Center for Brian Science,

Wako, Japan

[5]Department of Life Science, Graduate School of Arts and Sciences, The University

of Tokyo, Tokyo, Japan

[6]Department of Disease Model, Institute for Developmental Research, Aichi

Developmental Disability Center, Aichi, Japan

Abstract

Autism Spectrum Disorder (ASD) is one of the neurodevelopmental disorders and characterized with persistent impairments in social communication (including verbal and nonverbal communication) and repetitive behavior together with various comorbid symptoms. Epidemiological studies suggest a significant association between advanced paternal age and incidence of ASD in offspring, which has been modeled in rodents. However, how paternal aging makes an impact on the offspring's early communicative behavior, especially at the individual level, has not been addressed. Here we systematically analyzed postnatal development of vocal communication of pups by measuring ultrasonic vocalization (USV) induced by maternal separation that is considered to correspond to baby's cry, an earliest verbal communication in human. Maternal separation-induced USV of each offspring derived from young (3 months) or aged (>12 months) father was individually measured for 5 minutes at postnatal day 3 (P3), P6, P9, and P12. We classified USV syllables into twelve types according to Scattoni's classification with minor modifications, and analyzed duration, maximum frequency, maximum amplitude and interval of each syllable. Compared between the two groups, the offspring derived from aged fathers emitted the syllables with less number, shorter duration, different syllable components and less diversity. From an individual perspective, offspring derived from young father showed a developmental convergence in vocal communication, while those from aged fathers exhibited atypical developmental patterns. Taken together, we showed for the first time a significant influence of paternal aging on early vocal development with qualitative and quantitative aspects in individual mice.

Introduction

Autism Spectrum Disorder (ASD) is a neurodevelopmental disorder characterized by

two core symptoms; i.e., social interaction impairment (including verbal and

nonverbal communication deficits) and repetitive behavior[1-3]. In most cases, the

exact etiology of ASD remains unclear, although researchers believe that genetic,

environmental and epigenetic factors are involved in the neuropathology of ASD[4-6].

In recent years, epidemiological studies repeatedly suggest a significant association

between paternal aging and a risk of ASD in offspring. Compared with the children of

young fathers, the children were more likely to be diagnosed ASD if their fathers

were older[7-10]. Thus, paternal aging can be worth to be focused as one of the

non-genetic mechanisms leading to ASD.

In the new criteria of DSM-5, the communication and social interaction parts are

combined into one, i.e., "Social/Communication Deficits". This modification

emphasizes the importance of the social communication domain in ASD in early

infancy. Human infant crying is an innate social communication[11, 12], which has a

natural peak in frequency of approximately 2.5 hours of crying per day at around 6-8

weeks of age[13]. The infant crying can attract attention from caregivers[14] and

influences adult cognitive control[15]. It is reported that different crying patterns such

as higher frequency and shorter duration are observed in infants with a high risk of

ASD[16, 17]. Interpretation of autistic infant crying is, however, may have limitations in

separating possible etiological factors as well as in the sample size.

Rodents have been serving as suitable models to better understand genetical and

environmental factors of ASD[18, 19]. To elucidate mechanisms underlying

communication impairment in early postnatal stages, ultrasonic vocalization (USV) of

pups induced by maternal separation has recently been paid more attention. The

separation of a pup from its mother and littermates induces the pup to emit USV calls,

which triggers maternal approach and retrieval behavior[20, 21]. The patterns of USV

calls (i.e., here termed "syllables") can be seen as a form of social communication and

corresponding to the infant crying[22]. During normal postnatal development, murine

pups' USV gradually changes in acoustic features and syllable components[23, 24]. In

mouse models for ASD, researchers observe many variations of USV parameters such

as fewer calls, higher or lower frequency and shorter call duration [24-26]. However,

how paternal aging impacts early developmental patterns of detailed USV, especially

from an individual level, has not been addressed yet.

Here we measured the maternal separation-induced USV of individual mouse pups

during postnatal development at four stages. Not only comparing the developmental

patterns of USV emitted from offspring derived from young and aged father, but also

comprehensive analyses were performed by paying careful attention on individual

diversity in their syllables. We found that paternal aging influenced the trajectory of

syllable development in both quantitative and qualitative aspects, inducing more

atypical individuals derived from aged fathers.

Results

Offspring were obtained by mating young wild type female mice together with either

young (3 months) or aged (>12 months) wild type male mice (YFO and AFO,

respectively, Fig 1A). The average litter size and sample size were not significantly

different between YFO and AFO; i.e., 7.00 ± 0.35 (n = 32) and 6.6 ± 0.97 (n = 29) in

YFO and AFO, respectively (*t*-test for litter size, $p = 0.68$). Therefore, we assume that

our experimental setup was suitable to evaluate paternal aging effects on the

offspring's vocal communication.

1. USV measurement

To examine the effects of paternal aging on pup's USV, a large number of syllables

were collected from YFO and AFO at postnatal day 3 (P3), P6, P9, and P12 (Table 1).

To highlight differences between the groups of YFO and AFO clearly, the acoustic

features of USV were transformed to Z score (i.e., calculated by (Score – Mean) /

Standard Deviation), and shown in Fig 2 (raw data are shown in Supplementary Fig 1).

The syllables emitted from AFO exhibited the reduced number, shorter duration and

different trajectory of the interval between two adjacent syllables (Fig. 2A, 2B and

2E). The number of syllables emitted from all offspring reached a peak at P9 and was

affected significantly by main effects of paternal aging [father's age: $F (1, 236) =$

$12.79, p = 0.00127$] and postnatal day [day: $F (3, 236) = 4.54, p = 0.006$], but not by

interaction of paternal aging $\times$ postnatal day effect (Fig. 2A). The syllable duration

altered during postnatal development and showed a similar downward trend curve in

both groups, showing the lowest point at P6 (Fig. 2B). The duration of overall

syllables was influenced significantly by main effects of paternal aging [father's age:

$F (1, 236) = 27.16, p < 0.001$] and postnatal day [day: $F (3, 236) = 10.83, p < 0.001$],

but not by interaction of paternal aging × postnatal day effect. In our observation, no

significant paternal age effect was detected in the syllable maximum frequency and

amplitude (Fig. 2C, D). During development, the interval between two adjacent

syllables kept decreasing in the YFO, while in the AFO, the interval peaked at P6,

then decrease from P6 to P12 (Fig. 2E). A significant interaction between the main

effects of the paternal aging and postnatal day was observed [father's age × day: F (3,

236) = 10.18, $p < 0.001$]. Taken together, USV from AFO showed significantly

different developmental patterns in the overall syllable number, duration and interval

between two adjacent syllables, but not in maximum frequency and maximum

amplitude across development. These results indicate that paternal aging may alter the

developmental trajectory of overall syllables in quantitative and qualitative aspects.

2. Detailed analyses of syllable types

For deeper understanding of the paternal aging effects on the syllable patterns across

the early postnatal period, all USV syllables induced by maternal separation were

classified into twelve types (Fig. 1B) based on the shapes of spectrograms according

to a previous report [24]. The USV features of twelve types' syllables were transformed

into Z score and showed in Fig. 3. In general, a total of nine types of syllables

(upward, downward, short, chevron, wave, complex, one jump, more jump and more

jump + harmonics) showed significant alterations in the AFO. Among the altered

syllables, six types of syllables (upward, chevron, wave, complex, more jump and

more jump + harmonics) exhibited the reduced number only; the alterations of two

features were detected in the downward syllable (shorter duration and different

development trajectory in interval) and short syllable (less number and lower

maximum frequency); meanwhile, one jump syllable exhibited alterations in four

features (less number, shorter duration, lower and different trajectory of maximum

frequency, and different trajectory of interval). Therefore, paternal aging influences

distinct syllables at different levels.

2.1. The Number of individual types of syllables

After classification of the syllable types, we found that the main effect of paternal

aging significantly decreased the number of eight types of the syllables such as

"upward", "short", "chevron" "wave", "complex", "one jump", "more jump", and

"more jump + harmonics" (Fig. 3A, 3D-I and 3L, Supplementary Fig. 2A, 2D-I, 2L,

and Table 2). We did not detect "wave" syllables in the AFO at P3. These eight types

of syllables may contribute to the reduced number of the overall syllables emitted by

the AFO.

2.2. Duration of individual types of syllables

We found that the main effect of paternal aging significantly affected duration of

"downward" [father's age: $F(1, 234) = 22.74$, $p < 0.001$] and "one jump" [father's

age: $F(1, 215) = 7.99$, $p = 0.008$] syllables during development (Fig. 3B and 3H,

Supplementary Fig. 3B and 3H). Moreover, the significant interaction of two main

effects of the paternal aging and postnatal day was exhibited in the duration of "one

jump" syllables [father's age × day: $F(3, 215) = 2.97$, $p = 0.033$]. The shorter

duration of overall syllables in the AFO may attribute to the changes of "downward"

and "one jump" syllables.

2.3 Maximum frequency of individual types of syllables

Although we did not find a significant alteration of maximum frequency in the overall

syllables emitted from the AFO, detailed analyses indicated that "short" [father's age:

$F_{(1, 219)} = 11.66$, $p = 0.002$] and "one jump" [father's age: $F_{(1, 215)} = 7.49$, $p =$

0.02] syllables were influenced significantly by the main effect of paternal aging (Fig.

3D and 3H, Supplementary Fig. 4D and 4H); they showed lower maximum frequency

in the AFO. Moreover, two main effects of paternal aging and postnatal day were

interacted significantly in "one jump" syllables [father's age $\times$ day: $F_{(3, 215)} = 3.03$,

$p = 0.046$]. Paternal aging thus affected the maximum frequency at the level of two

syllables.

2.4 Maximum amplitude of individual types of syllables

After the classification of all syllables, there were again no significant main effects of

paternal aging and postnatal day in maximum amplitude (Fig. 3 and Supplementary

Fig. 5). The alteration of syllable maximum amplitude by paternal aging was not

observed after detailed analyses.

2.5 Interval of individual types of syllables

We further analyzed the interval data in all of 12 different syllable types to identify

any alteration in the interval of specific syllables. Significant interaction between the

two main effects of paternal aging and postnatal day was detected in the interval of

"downward" syllable [father's age $\times$ day: $F_{(3, 223)} = 7.22$, $p < 0.001$] and "one jump"

syllable [father's age $\times$ day: $F_{(3, 197)} = 3.15$, $p = 0.039$] (Fig. 3B and 3H,

Supplementary Fig. 6B and 6H). These data imply that relatively simple syllables

such as "downward" and "one jump" may contribute to the altered trajectory of the

syllable interval in overall syllables.

2.6 The percentage composition of the twelve types syllables

Next, we analyzed the composition of the twelve syllable types induced by maternal

separation (Fig. 4A and Table 3). MANOVA was performed to detect the difference

of syllable percentage composition between the two groups in each postnatal day we

observed. At P3, no statistical difference was detected. From P6 to P12, the AFO

emitted USV calls with a significantly different syllable component (Table 3).

Compared with the syllables of YFO in each postnatal day, AFO emitted higher

percentages of "downward" syllable at P6, P9 and P12, along with "flat", "short" and

"harmonics" syllables at P9 (by *t*-test). By contrast, other minor syllables such as

"chevron", "wave", "complex", "one jump", "more jump", "one jump + harmonics"

and "more jump + harmonics" showed lower percentage in the AFO during postnatal

development (Table 3). Therefore, the syllables emitted from the AFO had a

significantly different composition of the distinct syllables across the four postnatal

days.

2.7 Variation of syllables in offspring groups

Furthermore, we focused on the variation of syllables. Our results showed that YFO

developed to emit more types of syllables along with postnatal days. In contrast, the

AFO expanded the variation of the syllable types from P3 to P9, which became

narrower from P9 to P12 (Fig. 4B). Likewise, significantly fewer syllable types were

produced from the AFO from P3 to P12. The syllable types were affected

significantly by two main effects of paternal aging [father's age: $F (1, 236) = 58.32$, p

< 0.001] and postnatal day [day: $F (3, 236) = 13.64$, $p < 0.001$], but not by interaction

of paternal aging $\times$ postnatal day effect. To evaluate the different diversity of the

syllable types between the two groups, we calculated the entropy scores (see Methods

for the formula) as an indicator of production uniformity across the syllable types in

individual offspring (Fig. 4C). As we expected, entropy scores were also significantly

influenced by two main effects of paternal aging [father's age: $F_{(1, 224)} = 65.22$, $p <$

0.001] and postnatal day [day: $F_{(3, 224)} = 40.55$, $p < 0.001$], but not by interaction of

paternal aging $\times$ postnatal day effect. During development, a continuous rise of the

entropy scores was observed in all the offspring, but the AFO always showed lower

entropy scores compared with those from YFO across all postnatal stages. The data

reflected that the AFO exhibited fewer types and diversity of syllables, meaning that

they had a narrower vocal repertoire.

3. Difference in individual developmental trajectories of vocal communication

In this study, we addressed the "individual" differences among pups belonging to

YFO and AFO because ASD children exhibit "atypical" development. In order to

understand the longitudinal syllable development in individuals, clustering analyses

with Gaussian mixture models were applied first. Based on the Akaike Information

Criterion (AIC), the cluster number was determined objectively (see Methods for

details).

3.1 Clusters of the syllable number and duration

Because the number and duration of the overall syllables showed the positive

correlation (Pearson correlation coefficient 0.519, $p < 0.001$: YFO; 0.511, $p < 0.001$:

AFO), and the syllables emitted from AFO significantly decreased their number and

duration, we first separated the offspring into different clusters based on the syllable

number and duration. AIC implied that the choice of five clusters is optimal (Fig. 5A).

The developmental patterns of the syllable number and duration in YFO were

distributed to five clusters and concentrated on the fourth cluster, whereas the patterns

in AFO only dispersed among four clusters and focused on the third cluster (Fig.

5B-5G). Chi-square independence test revealed that the clustering patterns, i.e., the

proportion of individuals in each cluster, were significantly different between the

YFO and AFO ($p = 0.02$). These results exhibited that the dominant developmental

patterns of the syllable number and duration were different between YFO and AFO.

Repeatedly, AFO showed less variation in the syllables.

3.2 Cluster of the diversity in the syllable types

For the identification of individual development, we next classified the syllable types

into four clusters based on AIC (Fig. 6A). Interestingly, the syllables emitted from

YFO were classified into three clusters, while those from AFO were classified into

four clusters. Most of the syllable emitted from the YFO belonged to the second

cluster, whereas most of those from the AFO belonged to the first cluster (Fig. 6B-6F).

The cluster distribution was significantly different between the two groups

(chi-squared test, $p = 0.003$). Furthermore, we clustered the normalized entropy to

clarify individual development in syllable diversity. The number of clusters was

selected to five based on AIC. The cluster distribution was significantly different

between the YFO and AFO (chi-squared test, $p < 0.001$). The YFO occupied four

clusters and dominated the fourth cluster, while the AFO occupied five clusters and

dominated the first cluster (Fig 7A-7G). Data showed here indicated that the

developmental patterns of syllable diversity between the YFO, which differs from

those of AFO.

Through the clustering analyses, individual differences of the developmental patterns

in the syllable number, duration, types and diversity were clearly revealed. Therefore,

the longitudinal development patterns of offspring were found to be significantly

influenced by paternal aging.

3.3. Principal Component Analysis (PCA) of overall syllables

We further analyzed the vocal communication of offspring in regard with

developmental diversity. We extracted principal components (PC) contributing to the

individual character of overall syllables to pick up different features of individual

YFO and AFO. The PC1 and PC2 that summarized more than 80% of the variability

of the original data in overall syllables (Table 4) were plotted in Fig. 8. At P3, the

circles including 90% of the offspring's data were large, indicating that each offspring

exhibits a wide variety of individual difference in vocal communication regardless of

the paternal age. It is of note that the circle areas were almost overlapping between the

two groups of YFO (gray) and AFO (red) at P3. From P6 to P12, however, YFO

showed a developmental convergence in vocal communication patterns, while the

variability regions of AFO were kept wider, suggesting that the difference of

individual difference in the AFO was still large. In other words, there were more

individuals with "atypical development" within the AFO group. Through the

horizontal comparison of each postnatal day, the present data clearly indicated that

paternal aging resulted in greater individual diversity in regard with the syllable

patterns.

4. Correlations between body weight and USV

We wondered above described difference in vocal communication could secondarily

affect offspring's development. Therefore, we measured the body weight of offspring

after the USV recording on P3, P6, P9 and P12. Statistical analyses clearly showed

that offspring's body weight was affected significantly by two main effects of paternal

aging [father's age: $F (1, 236) = 35.84$, $p < 0.001$] and postnatal day [day: $F (3, 236) =$

874.91, $p < 0.001$], but not by interaction of paternal aging $\times$ postnatal day effect

(Supplementary Fig. 7). AFO exhibited a lower body weight gain throughout the four

stages from P3 to P12. This result suggested that paternal aging, in general, induced

lower body weight development in offspring during early postnatal development.

To clarify the correlations between body weight and USV, we first draw scatter plot

graphs to visualize the data (Fig. 9). In general, the circles involving 90% of the

offspring's data of correlations between body weight and USV were almost

overlapping between the two groups at P3 (Fig. 9A-9F). From P6 to P12, however,

the correlation between body weight and the number of syllable types gradually

became diverse between the YFO and AFO; again, AFO displayed the wider variation

than the YFO along with development (Fig. 9A). Furthermore, the correlation

coefficients and p-value were calculated by the Pearson correlation coefficient (Table

5). The body weight of YFO had positive correlations with the number of syllables,

the number of syllable types, and syllable duration at P6, as well as with syllable

frequency at P12. By contrast, the body weight of AFO had a positive correlation with

the number of syllables at P3, as well as with the number of syllable types at P6 and

P9. The strongest positive correlation (correlation coefficient $\geq |0.5|$) we found was

the correlation between body weight and the number of syllables in YFO at P6,

whereas other correlation coefficient scores were not obvious; we could not find a

strong positive correlation between the syllables and body weight in AFO. For deeper

analyses, the correlations between distinct types of syllables and body weight were

examined at four developmental stages (Supplementary Table 1). In general, all types

of syllables showed some correlations (positive or negative) with body weight at

different stages, while more strong correlations (correlation coefficient $\geq |0.5|$) were

found in YFO. These data revealed again that AFO included more diverse and

atypical individuals and abnormal developmental patterns.

Discussion

Our present study is the first report to analyze detailed vocal communication of early

postnatal stages in a mouse model of paternal aging. We revealed qualitative and

quantitative alterations of syllable development not only by comparison between YFO

and AFO groups but also at the individual perspective, across the first two postnatal

weeks. We applied comprehensive mathematical analyses because we can collect big

data by recording USV (e.g. hundreds of syllables per one pup in five minutes). This

has a great advantage compared with other conventional behavior tests that can collect

only a small number of scores. Considering epidemiological observation in human

studies suggesting paternal aging as one of the risk factors associated with autism in

offspring[7-10, 27], our study in mice indeed suggests that paternal aging actually plays

an important role in the alterations of communicative behavior in infant mice, a

consistent result with alterations of social behaviors in adult offspring in mice derived

from aged fathers[28-30].

Previous studies have shown that pup's USV can be analogous to human baby's cry

and thus used as one of a few tools to understand behavioral development during the

early postnatal period[31, 32]. We reported here that the USV features of AFO were

generally consistent with other ASD models with the decreased vocal numbers[26, 33, 34]

and shorter syllable durations[34, 35]. In our study, YFO emitted the diverse types of

syllables during the early postnatal periods, which is consistent with a previous

study[23]. However, compared with YFO, the AFO emitted a narrower spectrum of the

syllable types. The restricted syllable diversity has also been described in genetic

ASD models, i.e., *Cd157* KO mice[36] and *Tbx1*-deficient heterozygous mice[37].

Additionally, we observed the different composition of the syllables in AFO, which

has previously been reported in another genetic ASD mouse models such as *Reelin*

mutant[38], *fmr1* knockout[39] and *ScSn-Dmd^{mdx}/J* mutant[40]. It is thus reasonable to

assume that paternal aging leads to impairment in syllable development, diversity and

composition as shown in other genetic ASD models.

We noticed that the AFO exhibited lower body weight during the postnatal two weeks.

The difference was not observed when they became adults, according to our

unpublished data from different experiments used for behavior analyses in adult mice.

In human studies, lower birth weight is reported to be associated with several

neurodevelopmental difficulties such as learning disabilities[41], speech and language

problems[42], and social problems[43]. Indeed, paternal aging-associated low birth

weight has been considered as a high-risk factor of ASD[27, 44]. Therefore, it is

assumed that paternal aging might cause offspring developmental delay at the

physical level related to lower body weight, which could impact on early

communicative problems in the offspring.

Another possibility that can explain the difference in body weight might be influenced

by altered maternal care due to difference in vocal communication of offspring. It is

known that pup's USV has communicative significance[45-49], which is crucial for

maternal behaviors and preserves the social bonds between mother and infant that are

essential for the healthy development of offspring[50, 51]. In our study, even though

there were no strong correlations (correlation coefficient $\geq$ |0.5|) between the body

weight and overall syllables, some strong correlations were found between distinct

types of syllables and body weight. Expectedly, more correlations existed in YFO,

meaning "typical" developmental trajectories. It is, however, difficult to understand at

this moment what kind of information was transmitted by individual syllables or

syllable sequences and how these syllables impact maternal behaviors. In the future,

we need a more advanced method for syllable analyses to explore the deeper

biological meaning of pup's vocalization and examine the possibility of whether

maternal care may change due to altered USV of pups. So far, one of the available

methods to investigate the behaviors of maternal care and mother-infant interaction is

the playback and foster mother experiments which may help to understand whether

the abnormal syllables of AFO have an actual influence on their mothers. A previous

USV playback experiment indicated that the altered syllables in *Tbx1*-deficient

heterozygous mice led to fewer maternal responses to pups' vocalization[37]. Thus, the

alteration of USV development in AFO could be affected by not only the primary

factor of paternal aging, but also the secondary factor coming from maternal care.

Interestingly, we have previously noticed that pup's USV is a potent predictor for

sociability in adulthood; the number of USV at P6 was positively correlated with

sociability (in three-chamber social interaction test) and negatively with spatial

memory (in Morris water maze)[52]. In human infant's cry, wider variation of the

fundamental frequency is significantly reported to associate with better language and

cognitive outcomes[53]. Because of no detailed syllable classification to understand the

infant's cry, the variation of the fundamental frequency is considered as the

complexity of vocal control. In this case, since the AFO model exhibited poorer vocal

abilities than the YFO, the AFO might develop abnormal sociability (including vocal

behavior) in adulthood. We did not include adult behavior analyses in this study

because we noticed that repeated USV recording (including maternal separation)

could affect behavior in later stages (unpublished results). Actually, infancy is a

sensitive and crucial period to understand the behavioral and neuronal complications

at the beginning of life. The childhood emotional and behavioral problems often

correspond with early abnormal development in infants[54]. For instance, the infant

nonoptimal neuromotor development might predict the emotional problems in

childhood[55]; the problems of infant crying and feeding are prognostic to poor social

skills [56, 57] and adverse cognitive development[58] at the age of preschool. Therefore,

the infant's cry could be developed as an effective and noninvasive tool to evaluate

and diagnose neurodevelopmental disorders such as ASD in the future.

The highlight of this study is that we successfully modeled the "atypical" USV

development in the individual mouse by dimensional comparisons. Compared with

typically behavioral development in normal children (so called "neurotypical"), ASD

children show the variety of "atypical" behavioral phenotypes[1, 59, 60]; they do not

uniformly exhibit impaired scores in various criteria. In regard to our mouse model

during horizontal observation, the cluster preference of USV development expressed

unique patterns in each individual mouse pup and also "atypical" patterns caused by

paternal aging. In humans, some ASD-specific behaviors (e.g. impairment in social

cognition, eye contact, language abilities) are not obvious under 6 months, but

become gradually clear from the latter part of the first year and second years[54, 61, 62].

Through the vertical comparison at each postnatal day, we also found that the USV

patterns of PCA analyses were not the same in individuals but exhibited a gradual

convergence during development in the YFO. However, more diversities were

observed and remained in the AFO. At P3 (early infancy), each offspring emitted

USV calls in different variations, but after P6, the cohort of YFO gradually obtained

"typical" variations in their communication patterns by P12 (later infancy). By

contrast, the AFO still showed great variations at P12. For early intervening ASD, to

identify the "atypical" behaviors and elucidate the underlying neural basis, especially

at the early infancy (under 6 months in human and under P3 in mouse), would be the

next challenge.

The current study clearly demonstrates that detailed qualitative and quantitative

syllable analyses can be a useful tool for communicative phenotypes in a sensitive

period during postnatal stages of mice. We notice that complex syllables seemed to be

more affected in the AFO. The cerebellum is not fully developed during the first week,

and in another study of ours[30], we observed impairments in the structure of the motor

cortex and neuronal activity indicated with c-fos expression in the paraventricular

thalamus related to anxiety. Therefore, we should pay more attention to various brain

areas, e.g., including the brainstem that is important for motor control to emit USV.

Because the early postnatal period is critical for the development of the central as well

as peripheral neural systems in regard with communication behaviors, studies on

infant crying and rodent USV could synergistically contribute to understand the

neural basis for typical and atypical development, which could provide a cue for early

diagnosis and interventions for ASD.

Materials and methods

1. Animals and ethic statement

All experimental procedures were approved by the Ethics Committee for Animal

Experiments of Tohoku University Graduate School of Medicine (#2014-112) and

animals were treated according to the National Institutes of Health guidance of the

care and use of laboratory animals. Three-month-old (young) or >12-month-old (aged)

male C57BL/6J mice were crossed with 10-week-old (young) virgin female

C57BL/6J mice. After mating, each female mouse was separated from male mouse

and isolated alone to minimize possible confounding factors against the behavior of

offspring (Fig. 1A). In this study, 32 offspring were obtained from 5 young fathers

and 29 offspring from 5 aged fathers. Offspring that died during experiment periods

were excluded from analyses (mortality ratio, 5.7%: YFO vs 12.1%: AFO). At

postnatal day 3 (P3), each offspring was tattooed with an Aramis Animal Microtattoo

System (Natsume Co., Ltd., Tokyo, Japan) for individual recognition after the USV

test (described below). All animals were housed in standard cages in a temperature

and humidity-controlled room with a 12-hour light/dark cycle (light on at 8:00) and

had free access to standard lab chow and tap water.

2. USV measurement

According to previously described protocols[32, 52, 63], each offspring separated from its

mother and littermates one by one and placed on a transparent plastic dish with wood

chip bedding, and accessed within the sound-attenuating chamber for USV test on P3,

P6, P9 and P12. An ultrasound microphone (Avisoft-Bioacoustics CM16/CMPA) was

placed through a hole in the middle of the cover of the chamber, about 10 cm above

the offspring in its dish to record their vocalizations. The recorded vocalizations were

transferred to the UltraSound Gate 416H detector set (Avisoft Bioacoustics,

Germany). After a 5-min recording session, offspring were measured their body

weight and returned to the nest. This procedure was repeated in sequence until all

offspring had completed the recording phase. Both male and female pups were

analyzed. Room temperature was maintained at 22˚C.

(3) Syllable segmentation and classification

Acoustic waveforms were processed using a GUI-based MATLAB script ("usvseg")

originally developed for segmenting rodents' ultrasonic vocalizations[64]. Briefly, the

script computed the spectrograms from each waveform (60 seconds/block), put a

threshold to eliminate the noise component of the signal and detected syllables within

a frequency range of typical mice USVs (60-120 kHz). A criterion of 10-ms minimum

gap was used to separate two syllables and 2-ms as the minimum duration of a

syllable. The duration, inter-syllable interval, maximum frequency (peak frequency at

maximum amplitude) and maximum amplitude of each syllable were calculated

automatically by the program script. The syllable intervals of distinct types were

identified as the intervals between the specific type of syllable and the following

syllable. If the inter-syllable interval is wider than 250 ms, this interval will be

identified as a silence gap. Segmented syllables were manually classified into twelve

categories of syllable types by visual inspection of enhanced spectrograms which

were generated by the MATLAB program script. Ten of the syllable types (#1-10

below) were similar to those previously described by Scattoni *et al.*[24]. Noise sounds

that were mistakenly segmented by the program (e.g. scratching noise) were manually

identified and eliminated from further analyses. Definitions of syllable categories are

below (see also Fig 1B):

1) Upward syllables were upwardly modulated with a terminal frequency change ≥

6.25 kHz than the beginning of the syllable.

2) Downward syllables were downwardly modulated with a terminal frequency

change ≥ 6.25 kHz than the beginning of the syllable.

3) Flat syllables were continuous with a frequency modification ≤ 3 kHz.

4) Short syllables were displayed as a dot and shorter than or equal to 5 ms.

5) Chevron syllables were formed like a U or a reversed U.

6) Wave syllables were regulated with two directional changes in frequency > 6.25

kHz.

7) Complex syllables were regulated with three or more directional changes in

frequency > 6.25 kHz.

8) One jump syllables contained two components, in which the second component

was changed ≥10 kHz frequency than the first component and there was no time

interval between the two components.

9) More jump syllables contained three or more than three components, in which the

second component was changed ≥10 kHz frequency than the first and third

components respectively. There was no time interval between adjacent

components.

10) Harmonics syllables were displayed as one main component stacking with other

harmonically components of different frequencies.

11) One jump + harmonics syllables were contained one jump syllable and

harmonics syllable together and there was no time interval between each other.

12) More jump + harmonics syllables were contained more jump syllable and

518  harmonics syllable together and there was no time interval between each other.

520 **3. Statistical analysis**

521 For better visualization, the acoustic features of syllables are presented as Z score in

522 Fig 2 -3 (Z score = (score – Mean) / Standard Deviation). A two-way analysis of

523 variance (ANOVA) with False Discovery Rate (FDR) correction (0.05) was used to

524 investigate the statistical significance of syllables data which includes Z score and

525 original data of number, duration, maximum frequency, maximum amplitude and

526 interval of overall and distinct syllables, as well as body weight between the YFO and

527 AFO across P3, P6, P9 and P12. Two main effects (i.e., father's age and postnatal day

528 effect) and the interaction (i.e., father's age $\times$ postnatal day effect) were examined by

529 ANOVA. A multivariate analysis of variance (MANOVA) was performed to detect

530 the difference of syllable percentage component between the YFO and AFO in each

531 postnatal day with the independent variables of father's age and dependent variables

532 of the percentage of twelve syllable types. *Post-hoc* comparisons were performed

533 using two tailed *t*-test with Benjamini-Hochberg correction (BH correction) to detect

534 the difference between two groups in each postnatal day when ANOVA revealed the

535 significant interaction (paternal aging $\times$ postnatal day effect) for Supplementary Fig.

536 1-6. The correlation between body weight and USV parameters was detected by the

537 Pearson correlation coefficient.

539 To detect the diversity of syllable types, we used the information entropy as a

540 measure of uniformity in production rates across syllable types for each offspring.

541 The entropy score was ranged between 0 and 1. The score gets close to 1 when the

542 animal produced all the syllable types evenly (or diversely), while it becomes closer

543   to 0 if the animal preferred to produce fewer specific syllables types (less diversely).

544   We obtained this entropy score by the following calculation:

$$\text{entropy} = \frac{-\sum_{t=1}^{T} p_t \log_2 p_t}{\log_2 T}$$

where, T indicates the number of syllable types, and p_t means the production rate

of a specific syllable type t. Note that we excluded several offspring (i.e., 2 of YFO

and 4 of AFO at P3; 2 of AFO at P6; 2 of AFO at P9; 2 of AFO at P12) from this

analysis since the total number of syllables in 5 minutes were insufficient (less than

10) to analyze their entropies. The entropy scores were compared between the YFO

and AFO across different postnatal days by using two-way ANOVA. To understand

the individual development, clustering analysis with Gaussian mixture models

(GMMs) was applied, where the data dimension is eight corresponding to the number

and duration of syllables at four time points. We fit GMMs with diagonal covariance

Gaussian components by the MATLAB function *fitgmdist*. The number of clusters

was selected by minimizing the Akaike Information Criterion (AIC) [65-66]. Based on

the fitted GMM, we classified each individual mouse pup into the cluster with

maximum posterior probability. Then the chi-square independence test was applied to

determine whether the cluster distribution was significantly different between the two

groups. PCA was performed to objectively characterize the typical syllable patterns of

individual offspring. In the present study, the syllable data including syllable number,

number of types, duration, maximum frequency and maximum amplitude were used

as input data for the PCA to generate principal components. The first principal

component (PC1) alone is able to explain more than 50% of the variability of all input

values, and PC1 plus second principal component (PC2) can illustrate more than 80%

of the variability.

For all comparisons, significance was set at $p = 0.05$. JMP V13 Pro software (SAS

Institute, Cary, NC, USA) was used for statistical analyses. Values are shown as mean

± standard error of the mean (S.E.M.) for each group.

Acknowledgements

572 The authors thank Ms. Sayaka Makino for animal care. The authors also appreciate all

573 members of their laboratory for contributive discussions. This work was supported by

574 KAKENHI in the Innovative Areas (Grant Number 16H06530) from MEXT.

References

1.   American Psychiatric Association. Diagnostic and statistical manual of mental

disorders: DSM-5 (5th ed.). Washington, DC: American Psychiatric Association;

2013.

2.   Folstein SE, Piven J. Etiology of autism: genetic influences. Pediatrics. 1991;87(5

Pt 2):767-73. PubMed PMID: 1708492.

3.   Lotspeich LJ, Ciaranello RD. The neurobiology and genetics of infantile autism.

Int Rev Neurobiol. 1993;35:87-129. PubMed PMID: 8463065.

4.   Schaefer GB, Mendelsohn NJ. Genetics evaluation for the etiologic diagnosis of

autism   spectrum   disorders.   Genet   Med.   2008;10(1):4-12.   doi:

10.1097/GIM.0b013e31815efdd7. PubMed PMID: 18197051.

5.   Tordjman S, Somogyi E, Coulon N, Kermarrec S, Cohen D, Bronsard G, et al.

Gene x environment interactions in autism spectrum disorders: role of epigenetic

mechanisms. Frontiers in Psychiatry. 2014;5. doi: 10.3389/fpsyt.2014.00053. PubMed

PMID: 25136320

6.   Nardone S, Sams DS, Zito A, Reuveni E, Elliott E. Dysregulation of cortical

neuron DNA methylation profile in autism spectrum disorder. Cereb Cortex.

2017;27(12):5739-54. doi: 10.1093/cercor/bhx250. PubMed PMID: 29028941.

7.   Reichenberg A, Gross R, Weiser M, Bresnahan M, Silverman J, Harlap S, et al.

Advancing paternal age and autism. Arch Gen Psychiatry. 2006;63(9):1026-32. doi:

10.1001/archpsyc.63.9.1026. PubMed PMID: 16953005.

8.   Hultman CM, Sandin S, Levine SZ, Lichtenstein P, Reichenberg A. Advancing

paternal age and risk of autism: new evidence from a population-based study and a

meta-analysis of epidemiological studies. Mol Psychiatry. 2011;16(12):1203-12. doi:

10.1038/mp.2010.121. PubMed PMID: 21116277.

9.   Lundstrom S, Haworth CM, Carlstrom E, Gillberg C, Mill J, Rastam M, et al.

Trajectories leading to autism spectrum disorders are affected by paternal age:

findings from two nationally representative twin studies. J Child Psychol Psychiatry.

2010;51(7):850-6.   doi: 10.1111/j.1469-7610.2010.02223.x.   PubMed   PMID:

20214699.

10. Lampi KM, Hinkka-Yli-Salomaki S, Lehti V, Helenius H, Gissler M, Brown AS,

et al. Parental age and risk of autism spectrum disorders in a Finnish national birth

cohort. J Autism Dev Disord. 2013;43(11):2526-35. doi: 10.1007/s10803-013-1801-3.

PubMed PMID: 23479075.

11. Acebo C, Thoman EB. Crying as social behavior. Infant Mental Health Journal.

1992;13(1):67-82.

12. Acebo C, Thoman EB. Role of infant crying in the early mother-infant dialogue.

Physiology and Behavior. 1995;57(3):541-7.

13. St James-Roberts I, Halil T. Infant crying patterns in the first year: normal

community and clinical findings. J Child Psychol Psychiatry. 1991;32(6):951-68.

PubMed PMID: 1744198.

14. Soltis J. The signal functions of early infant crying. Behav Brain Sci.

2004;27:443-90. doi: 10.1017/s0140525x0400010x.

15. Dudek J, Faress A, Bornstein MH, Haley DW. Infant cries rattle adult cognition.

PLoS One. 2016;11(5):e0154283. doi: 10.1371/journal.pone.0154283. PubMed PMID:

27191845; PubMed Central PMCID: PMCPMC4871531.

16. Sheinkopf SJ, Iverson JM, Rinaldi ML, Lester BM. Atypical cry acoustics in

6-month-old infants at risk for autism spectrum disorder. Autism Res. 2012;5(5):331-9.

doi: 10.1002/aur.1244. PubMed PMID: 22890558; PubMed Central PMCID:

PMCPMC3517274.

17. Unwin LM, Bruz I, Maybery MT, Reynolds V, Ciccone N, Dissanayake C, et al.

Acoustic properties of cries in 12-Month old infants at high-risk of autism spectrum

disorder. J Autism Dev Disord. 2017;47(7):2108-19. doi: 10.1007/s10803-017-3119-z.

PubMed PMID: 28425020.

18. Crawley JN. Designing mouse behavioral tasks relevant to autistic-like behaviors.

Ment Retard Dev Disabil Res Rev. 2004;10(4):248-58. doi: 10.1002/mrdd.20039.

PubMed PMID: 15666335.

19. Kazdoba TM, Leach PT, Yang M, Silverman JL, Solomon M, Crawley JN.

Translational mouse models of autism: advancing toward pharmacological

therapeutics. Curr Top Behav Neurosci. 2016;28:1-52. doi: 10.1007/7854_2015_5003.

PubMed PMID: 27305922; PubMed Central PMCID: PMCPMC5116923.

20. D'Amato F, Scalera E, Sarli C, Moles A. Pups call, mothers rush: does maternal

responsiveness affect the amount of ultrasonic vocalizations in mouse pups? . Behav

Genet. 2005 Jan;35 (1):103-12. doi: 10.1007/s10519-004-0860-9. PubMed PMID:

15674537.

21. Hahn ME, Lavooy MJ. A Review of the methods of studies on infant ultrasound

production and maternal retrieval in small rodents. Behav Genet. 2005;35 (1):31-52.

doi: 10.1007/s10519-004-0854-7. PubMed PMID: 15674531.

22. Esposito G, Hiroi N, Scattoni ML. Cry, baby, cry: expression of distress as a

biomarker and modulator in autism spectrum disorder. International Journal of

Neuropsychopharmacology. 2017;20(6):498-503. doi: 10.1093/ijnp/pyx014.

23. Grimsley JM, Monaghan JJ, Wenstrup JJ. Development of social vocalizations in

mice. PLoS One. 2011;6(3):e17460. doi: 10.1371/journal.pone.0017460.

24. Scattoni ML, Gandhy SU, Ricceri L, Crawley JN. Unusual repertoire of

vocalizations in the BTBR T+tf/J mouse model of autism. PLoS One.

2008;3(8):e3067. doi: 10.1371/journal.pone.0003067. PubMed PMID: 18728777;

PubMed Central PMCID: PMCPMC2516927.

25. Wohr M. Ultrasonic vocalizations in Shank mouse models for autism spectrum

disorders: detailed spectrographic analyses and developmental profiles. Neurosci

Biobehav Rev. 2014;43:199-212. doi: 10.1016/j.neubiorev.2014.03.021. PubMed

PMID: 24726578.

26. Dougherty JD, Maloney SE, Wozniak DF, Rieger MA, Sonnenblick L, Coppola G,

et al. The disruption of Celf6, a gene identified by translational profiling of

serotonergic neurons, results in autism-related behaviors. J Neurosci.

2013;33(7):2732-53. doi: 10.1523/JNEUROSCI.4762-12.2013. PubMed PMID:

23407934; PubMed Central PMCID: PMCPMC3711589.

27. Itzchak BE, Lahat E, Zachor DA. Advanced parental ages and low birth weight in

autism spectrum disorders--rates and effect on functioning. Res Dev Disabil.

2011;32(5):1776-81. Epub 2011 Apr 17. doi: 10.1016/j.ridd.2011.03.004. PubMed

PMID: 21498045.

28. Foldi CJ, Eyles DW, McGrath JJ, Burne TH. Advanced paternal age is associated

with alterations in discrete behavioural domains and cortical neuroanatomy of

C57BL/6J   mice.   Eur   J   Neurosci.   2010;31(3):556-64.   doi:

10.1111/j.1460-9568.2010.07074.x. PubMed PMID: 20105239.

29. Janecka M, Manduca A, Servadio M, Trezza V, Smith R, Mill J, et al. Effects of

advanced paternal age on trajectories of social behavior in offspring. Genes Brain

Behav. 2015;14(6):443-53. doi: 10.1111/gbb.12227. PubMed PMID: 26096767.

30. Yoshizaki K, Koike T, Kimura R, Kikkawa T, Oki S, Koike K, et al. Paternal age

affects offspring's behavior prossibly via an epigenetic machanism recruiting a

transciptional repressor REST. bioRxiv. 2019. doi: 10.1101/550095.

31. Branchi I, Santucci D, Alleva E. Ultrasonic vocalisation emitted by infant rodents:

a tool for assessment of neurobehavioural development. Behav Brain Res.

2001;125(1-2):49-56.   doi: 10.1016/S0166-4328(01)00277-7.   PubMed   PMID:

11682093.

32. Scattoni ML, Crawley J, Ricceri L. Ultrasonic vocalizations: a tool for

behavioural phenotyping of mouse models of neurodevelopmental disorders. Neurosci

Biobehav Rev. 2009;33(4):508-15. doi: 10.1016/j.neubiorev.2008.08.003. PubMed

PMID: 18771687; PubMed Central PMCID: PMCPMC2688771.

33. Wohr M, Roullet FI, Hung AY, Sheng M, Crawley JN. Communication

impairments in mice lacking Shank1: reduced levels of ultrasonic vocalizations and

scent   marking   behavior.   PLoS   One.   2011;6(6):e20631.   doi:

10.1371/journal.pone.0020631. PubMed PMID: 21695253; PubMed Central PMCID:

PMCPMC3111434.

34. Yang L, Faraone SV, Zhang-James Y. Autism spectrum disorder traits in Slc9a9

knock-out mice. Am J Med Genet B Neuropsychiatr Genet. 2016;171B(3):363-76. doi:

10.1002/ajmg.b.32415. PubMed PMID: 26755066.

35. Gaub S, Groszer M, Fisher SE, Ehret G. The structure of innate vocalizations in

Foxp2-deficient mouse pups. Genes Brain Behav. 2010;9(4):390-401. doi:

10.1111/j.1601-183X.2010.00570.x. PubMed PMID: 20132318; PubMed Central

PMCID: PMCPMC2895353.

36. Lopatina OL, Furuhara K, Ishihara K, Salmina AB, Higashida H. Communication

Impairment in Ultrasonic Vocal Repertoire during the Suckling Period of Cd157

Knockout Mice: Transient Improvement by Oxytocin. Frontiers in Neuroscience.

2017;11. doi: 10.3389/fnins.2017.00266.

37. Takahashi T, Okabe S, Broin PO, Nishi A, Ye K, Beckert MV, et al. Structure and

function of neonatal social communication in a genetic mouse model of autism. Mol

Psychiatry. 2016;21(9):1208-14. doi: 10.1038/mp.2015.190. PubMed PMID:

26666205; PubMed Central PMCID: PMCPMC4909589.

38. Romano E, Michetti C, Caruso A, Laviola G, Scattoni ML. Characterization of

neonatal vocal and motor repertoire of reelin mutant mice. PLoS One.

2013;8(5):e64407. doi: 10.1371/journal.pone.0064407. PubMed PMID: 23700474;

PubMed Central PMCID: PMCPMC3660261.

39. Lai JK, Sobala-Drozdowski M, Zhou L, Doering LC, Faure PA, Foster JA.

Temporal and spectral differences in the ultrasonic vocalizations of fragile X knock

out mice during postnatal development. Behav Brain Res. 2014;259:119-30. doi:

10.1016/j.bbr.2013.10.049. PubMed PMID: 24211451.

40. Miranda R, Nagapin F, Bozon B, Laroche S, Aubin T, Vaillend C. Altered social

behavior and ultrasonic communication in the dystrophin-deficient mdx mouse model

of Duchenne muscular dystrophy. Mol Autism. 2015;6:60. doi:

10.1186/s13229-015-0053-9. PubMed PMID: 26527530; PubMed Central PMCID:

716 PMCPMC4627616.

717 41. Johnson EO, Breslau N. Increased risk of learning disabilities in low birth weight

718 boys at age 11 years. Biol Psychiatry. 2000;47(6):490-500. doi:

719 10.1016/S0006-3223(99)00223-1.

720 42. Aram DM, Hack M, Hawkins S, Weissman BM, Borawski-Clark E.

721 Very-low-birthweight children and speech and languange development. Journal of

722 Speech, Language, and Hearing Research. 1991 34:1169-79. doi:

723 10.1044/jshr.3405.1169. PubMed PMID: 1749247.

724 43. Hack M, Taylor HG, Klein N, Eiben R, Schatschneider C, Mercuri-Minich N.

725 School-age outcomes in children with birth weights under 750g. N Engl J Med.

726 1994;331:753-9. doi: 10.1056/NEJM199409223311201. PubMed PMID: 7520533.

727 44. Kolevzon A, Gross R, Reichenberg A. Prenatal and perinatal risk factors for

728 autism: a review and integration of findings. Arch Pediatr Adolesc Med.

729 2007;161(4):326-33. doi: 10.1001/archpedi.161.4.326. PubMed PMID: 17404128.

730 45. Hashimoto H, Saito TR, Furudate S, Takahashi K. Prolactin levels and maternal

731 behavior induced by ultrasonic vocalizations of the rat pup. Exp Anim.

732 2001;50(4):307-12. doi: 10.1538/expanim.50.307. PubMed PMID: 11515093.

733 46. Liu RC, Miller KD, Merzenich MM, Schreiner CE. Acoustic variability and

734 distinguishability among mouse ultrasound vocalizations. J Acoust Soc Am.

735 2003;114(6):3412-22. doi: 10.1121/1.1623787. PubMed PMID: 14714820.

736 47. Liu RC, Linden JF, Schreiner CE. Improved cortical entrainment to infant

737 communication calls in mothers compared with virgin mice. Eur J Neurosci.

738 2006;23(11):3087-97. doi: 10.1111/j.1460-9568.2006.04840.x. PubMed PMID:

739 16819999.

740 48. Ehret G, Bernecker C. Low-frequency sound communication by mouse pups

(Mus musculus): wriggling calls release maternal behaviour. Animal Behaviour. 1986;34(3):821-30. doi: 10.1016/S0003-3472(86)80067-7.

49. Kikusui T, Hiroi N. A Self-Generated Environmental Factor as a Potential Contributor to Atypical Early Social Communication in Autism. Neuropsychopharmacology. 2017;42(1):378. doi: 10.1038/npp.2016.225. PubMed PMID: 27909329; PubMed Central PMCID: PMCPMC5143512.

50. Mogi K, Takakuda A, Tsukamoto C, Ooyama R, Okabe S, Koshida N, et al. Mutual mother-infant recognition in mice: The role of pup ultrasonic vocalizations. Behav Brain Res. 2017;325(Pt B):138-46. doi: 10.1016/j.bbr.2016.08.044. PubMed PMID: 27567527.

51. Mogi K, Nagasawa M, Kikusui T. Developmental consequences and biological significance of mother-infant bonding. Prog Neuropsychopharmacol Biol Psychiatry. 2011;35(5):1232-41. doi: 10.1016/j.pnpbp.2010.08.024. PubMed PMID: 20817069.

52. Yoshizaki K, Koike K, Kimura R, Osumi N. Early postnatal vocalizations predict sociability and spatial memory in C57BL/6J mice: Individual differences in behavioral traits emerge early in development. PLoS One. 2017;12(11):e0186798. doi: 10.1371/journal.pone.0186798. PubMed PMID: 29091920; PubMed Central PMCID: PMCPMC5665508.

53. Shinya Y, Kawai M, Niwa F, Imafuku M, Myowa M. Fundamental Frequency Variation of Neonatal Spontaneous Crying Predicts Language Acquisition in Preterm and Term Infants. Front Psychol. 2017;8. PubMed PMID: doi:10.3389/fpsyg.2017.02195; PubMed Central PMCID: PMCPMC5744644.

54. Zwaigenbaum L, Bryson S, Rogers T, Roberts W, Brian J, Szatmari P. Behavioral manifestations of autism in the first year of life. Int J Dev Neurosci. 2005;23(2-3):143-52. doi: 10.1016/j.ijdevneu.2004.05.001. PubMed PMID:

15749241.

55. Serdarevic F, Ghassabian A, van Batenburg-Eddes T, Tahirovic E, White T,

Jaddoe VWV, et al. Infant Neuromotor Development and Childhood Problem

Behavior. Pediatrics. 2017;140(6). doi: 10.1542/peds.2017-0884. PubMed PMID:

29138362.

56. Schmid G, Schreier A, Meyer R, Wolke D. A prospective study on the persistence

of infant crying, sleeping and feeding problems and preschool behaviour. Acta

Paediatr. 2010;99(2):286-90. doi: 10.1111/j.1651-2227.2009.01572.x. PubMed PMID:

19886897.

57. Winsper N, Wolke D. Infant and Toddler Crying, Sleeping and Feeding Problems

and Trajectories of Dysregulated Behavior Across Childhood. Journal of Abnormal

Child Psychology. 2014;42(5):831-43. doi: 10.1007/s10802-013-9813-1).

58. Wolke D, Schmid G, Schreier A, Meyer R. Crying and feeding problem in

Infancy and cognitive outcome in preschool children born at risk: a prospective

population   study.   J   Dev   Behav   Pediatr.   2009;30(3):226-38.   doi:

10.1097/DBP.0b013e3181a85973. PubMed PMID: 19433987.

59. Grzadzinski R, Huerta M, Lord C. DSM-5 and autism spectrum disorders (ASDs):

an opportunity for identifying ASD subtypes. Mol Autism. 2013;4(12). doi:

10.1186/2040-2392-4-12. PubMed PMID: 2375638.

60. Wozniak RH, Leezenbaum NB, Northrup JB, West KL, Iverson JM. The

development of autism spectrum disorders: variability and causal complexity. Wiley

Interdiscip Rev Cogn Sci. 2017;8(1-2):e1426. doi: 10.1002/wcs.1426. PubMed PMID:

27906524.

61. Piven J, Elison JT, Zylka MJ. Toward a conceptual framework for early brain and

behavior development in autism. Mol Psychiatry. 2017;22(10):1385-94. doi:

10.1038/mp.2017.131. PubMed PMID: 28937691; PubMed Central PMCID:

PMCPMC5621737.

62. Ozonoff S, Losif AM, Baguio F, Cook IC, Hill MM, Hutman T, et al. A

Prospective study of the emergence of early behavioral signs of autism. J Am Acad

Child   Adolesc Psychiary. 2010;49(3):256-66. PubMed PMID: 20410715.

63. Yoshizaki K, Furuse T, Kimura R, Tucci V, Kaneda H, Wakana S, et al. Paternal

Aging Affects Behavior in Pax6 Mutant Mice: A Gene/Environment Interaction in

Understanding Neurodevelopmental Disorders. PLoS One. 2016;11(11):e0166665.

doi: 10.1371/journal.pone.0166665. PubMed PMID: 27855195; PubMed Central

PMCID: PMCPMC5113965.

64. Tachibana RO, Kanno K, Okabe S, Kobayasi KI, Okanoya K. Usvseg: A robust

segmentation of rodents' ultrasonic vocalization. bioRxiv. 2019. doi: 10.1101/572743.

65. Konishi S, Kitagawa G. Information Criteria and Statistical Modeling.

Springer-Verlag New York Press; 2008.

66. Mclachlan G, Peel D. Finite Mixture Models. 1st ed. Wiley-Interscience Press;

2000.

Figure Legends

Fig. 1. Experiment design and syllable types.

(A) Three-month-old (young) or >12-month-old (aged) male C57BL/6J mice were

crossed with 10-week-old (young) virgin female C57BL/6J mice to obtain offspring.

On P3, P6, P9 and P12, each offspring was separated from the mother to record the

USV for 5 minutes. After data collection, detailed analyses were performed. (B)

Typical sonograms of USV are classified into twelve types of syllables such as

downward, one jump, harmonics, upward, chevron, wave, complex, more jump +

harmonics, short, one jump + harmonics, more jump, and flat. Scale bar = 50 kHz, 20

ms. The colors of magenta, blue and grey spectra indicate the syllables with increased,

decreased and non-specifically changed percentages, respectively, during the

development of offspring derived from young father.

Fig. 2. The Z score trajectories of USV features in overall syllable.

Z scores of syllable number (A), duration (B), maximum frequency (C), maximum

amplitude (D) and interval (E) were obtained from USV data of offspring derived

from young (YFO) and aged (AFO) fathers. Two-way ANOVA was used to compare

the differences between the two groups. Data are presented as means ± SEM. $^{\dagger\dagger}p<$

0.01, $^{\dagger\dagger\dagger}p < 0.001$ indicates a significant main effect of father's age, $^{\S\S\S}p < 0.001$

indicates a significant interaction of father's age $\times$ day effect.

Fig. 3. The Z score trajectories of USV features in individual type of syllables.

The syllable number, duration, maximum frequency, maximum amplitude and

interval of individual type of syllables from offspring derived from young (YFO) and

aged (AFO) fathers are transformed to Z score. (A)-(L) show the developmental

trajectories of five USV features in the individual type of syllables. Two-way

ANOVA was used to compare the differences between the two groups. Data are

presented as means ± SEM. $^{\dagger}p < 0.05$, $^{\dagger\dagger}p < 0.01$, $^{\dagger\dagger\dagger}p < 0.001$ indicates a significant

main effect of father's age. $^{\S}p < 0.05$, $^{\S\S}p < 0.01$, $^{\S\S\S}p < 0.001$ indicates a significant

interaction of father's age $\times$ day effect.

Fig. 4. The composition of the twelve types syllables

(A) Pie graphs exhibited the percentage composition of the twelve types of syllables

from P3 to P12 in offspring derived from young (YFO) and aged (AFO) fathers. Data

are presented as means. The colors of magenta, blue and grey spectra indicate the

syllables with increased, decreased and non-specifically changed percentages,

respectively, during the development of the YFO. (B) The developmental trajectories

in regard to the variation of the syllable types in YFO and AFO. Data are presented as

means ± SEM. ††† $p < 0.001$ indicates a significant main effect of the father's age. (C)

Normalized entropy scores of syllables emitted from YFO and AFO. Two-way

ANOVA was used to compare the differences between the groups across development.

Data are presented as means ± SEM. ††† $p < 0.001$ indicates a significant main effect

of the father's age.

Fig. 5. Separation of individual patterns according to the syllable number and

duration

(A) The cluster number of syllable number and duration were chosen as five by using

the Akaike Information Criterion (AIC) as the minimum number (shown on X-axis)

with AIC values (Y-axis). (B)-(F): Clustering the syllable number and duration. The

clustering analyses with Gaussian mixture models (GMMs) were applied and

separated the individual offspring into five clusters according to the syllable number

and duration across four postnatal days. Offspring derived from young (YFO) and

aged (AFO) fathers were arranged to each cluster with significantly different

populations. (G) The different percentages of offspring in each cluster. The heat map

summarized the percentages of YFO and AFO that were distributed to each cluster.

Fig. 6. Separation of individual patterns according to syllable types

(A) The cluster number of syllable types was chosen as four by using AIC as the

minimum number (shown on X-axis) with AIC values (Y-axis). (B)-(E) Clustering the

syllable types. The clustering analyses with Gaussian mixture models (GMMs) were

applied and separated the offspring into four clusters according to the syllable types

across four postnatal stages. The young father-derived offspring (YFO) and aged

father-derived offspring (AFO) were arranged to each cluster with significantly

different populations. (F) The percentages of offspring in each cluster. The heat map

summarized the percentages of YFO and AFO that were distributed in each cluster.

Fig. 7. Separation of individual patterns according to normalized entropy

(A) The cluster number of syllable types was chosen as five by using AIC as the

minimum number (shown on X-axis) with AIC values (Y-axis). (B)-(F) Clustering the

normalized entropy. The clustering analyses with Gaussian mixture models (GMMs)

were applied and separated the offspring into five clusters according to the normalized

entropy across four postnatal days. The young father-derived offspring (YFO) and

aged father-derived offspring (AFO) were arranged to each cluster with significantly

different populations. (G) The percentages of offspring in each cluster. The heat map

summarized the percentages of the YFO and AFO that were distributed to each

cluster.

Fig. 8. Principal component analysis of overall syllables.

PCA plots showing individual differences of syllable development among the

individual offspring derived from young (YFO) and aged (AFO) fathers. At P3, PCA

of syllable patterns, where the Y-axis (PC1) explained 55.6% variance, whereas the

X-axis (PC2) explained a 25.6% variance of the data. At P6, PCA of syllable patterns,

where the Y-axis (PC1) explained 62.3% variance, whereas the X-axis (PC2)

explained a 21.2% variance of the data. At P9, PCA of syllable patterns, where the

Y-axis (PC1) explained 62.6% variance, whereas the X-axis (PC2) explained 20.3%

variance of the data. At P12, PCA of syllable patterns, where the Y-axis (PC1)

explained 59.2% variance, whereas the X-axis (PC2) explained 23.4% variance of the

data. The circles in gray or red included a 90% population of YFO and AFO.

Fig. 9. Correlation between body weight and USV

The correlation between body weight and USV parameters was detected by the

Pearson correlation coefficient. The correlation coefficient and *p*-value were showed

in Table 5. Scatter plot showing the correlation between body weight and USV in

individuals from P3 to P12 among offspring derived from young (YFO) and aged

(AFO) fathers. Correlation between body weight and the number of syllable types (A),

the number of syllables (B), syllable duration (C), syllable maximum frequency (D),

syllable maximum amplitude (E), and syllable interval (F), respectively. The circles in

gray or red included a 90% population of YFO or AFO.

Supplementary Fig. 1. The developmental trajectories of USV features in overall

syllable.

The numbers (A), durations (B), maximum frequency (C), maximum amplitude (D),

and interval (E) of syllables emitted from offspring derived from young (YFO) and

aged (AFO) fathers. Two-way ANOVA followed by *post-hoc* comparisons (*t*-test

with BH correction) was used to compare the differences between the two groups.

Data are presented as means ± SEM. $^{\dagger\dagger}p < 0.01$, $^{\dagger\dagger\dagger}p < 0.001$ indicates a significant

main effect of father's age, $^{\S\S\S}p < 0.001$ indicates a significant interaction of father's

age × day effect. ** $p < 0.01$ indicates a significant decrease in AFO; * $p < 0.05$

indicates a significant increase in AFO.

Supplementary Fig. 2. The Number of individual type of syllables

(A)-(L) The developmental trajectory in regard to the number of the twelve types of

syllables. Two-way ANOVA followed by *post-hoc* comparisons *t*-test with BH

correction was used to compare the syllable number between the offspring derived

from young (YFO) and aged (AFO) fathers at each postnatal day. Data are presented

as means ± SEM. $^{†}p < 0.05$, $^{††}p < 0.01$, $^{†††}p < 0.001$ indicates a significant main

effect of father's age. $^{§}p < 0.05$, $^{§§}p < 0.01$, $^{§§§}p < 0.001$ indicates significant

interaction of father's age × day effect. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

indicates a significant decrease of the syllable number in the AFO at individual

postnatal day (*t*-test with BH correction).

Supplementary Fig. 3: Duration of individual types of syllables

(A)-(L) The developmental trajectory in regard to the duration of the twelve types of

syllables. Two-way ANOVA followed by *post-hoc* comparisons *t*-test with BH

correction was used to compare the syllable duration between the offspring derived

from young (YFO) and aged (AFO) fathers at each postnatal day. Data are presented

as means ± SEM. $^{†}p < 0.05$, $^{††}p < 0.01$, $^{†††}p < 0.001$ indicates a significant main

effect of father's age. $^{§}p < 0.05$, $^{§§}p < 0.01$, $^{§§§}p < 0.001$ indicates a significant

interaction of father's age × day effect. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

indicates a significant decrease of the syllable number in the AFO at individual

postnatal day (*t*-test with BH correction).

Supplementary Fig. 4. Maximum frequency of individual types of syllables

(A)-(L) The developmental trajectory in regard to the maximum frequency of the

twelve types of syllables. Two-way ANOVA followed by *post-hoc* comparisons *t*-test

with BH correction was used to compare the syllable number between the offspring

derived from young (YFO) and aged (AFO) fathers. Data are presented as means $\pm$

SEM. $^{\dagger}p<0.05$, $^{\dagger\dagger}p<0.01$, $^{\dagger\dagger\dagger}p<0.001$ indicates a significant main effect of father's

age. $^{\S}p<0.05$, $^{\S\S}p<0.01$, $^{\S\S\S}p<0.001$ indicates a significant interaction of father's age

$\times$ day effect. *$p<0.05$, **$p<0.01$, ***$p<0.001$ indicates a significant decrease of the

syllable number in the AFO at individual postnatal day (*t*-test with BH correction).

Supplementary Fig. 5. Maximum amplitude of individual types of syllables

(A)-(L) The developmental trajectory in regard to the maximum amplitude of the

twelve types of syllables. Two-way ANOVA was used to compare the syllable

number between the offspring derived from young (YFO) and aged (AFO) fathers.

Data are presented as means $\pm$ SEM. No significant differences were detected.

Supplementary Fig. 6: Interval of individual types of syllables

(A)-(L) The developmental trajectory in regard to the interval of the twelve types of

syllables. Two-way ANOVA followed by *post-hoc* comparisons *t*-test with BH

correction was used to compare the syllable number between the offspring derived

from young (YFO) and aged (AFO) fathers at each postnatal day. Data are presented

as means $\pm$ SEM. $^{\S}p < 0.05$, $^{\S\S}p < 0.01$, $^{\S\S\S}p < 0.001$ indicates a significant

interaction of father's age $\times$ day effect. *$p < 0.05$ indicates a significant increase of

the syllable interval in the AFO at individual postnatal day (*t*-test with BH correction).

Supplementary Fig. 7. Body weight development.

Body weight of offspring derived from young (YFO) and aged (AFO) fathers.

Two-way ANOVA was used to compare the body weight between the groups. Data

are presented as means ± SEM. ††† $p < 0.001$ indicates a significant main effect of the

father's age.

Table 1. The number of syllables was collected from the offspring at each postnatal day.

	P3	P6	P9	P12
YFO	4212	6041	7122	7039
AFO	2990	3276	5454	3296

YFO: young father-derived offspring; AFO: aged father-derived offspring.

Table 2. Eight types of syllables with decreased numbers in the AFO from P3 to P12.

Syllable types	Results of ANOVA
Upward	father's age: $F (1, 236) = 6.78$, $p = 0.015$
Short	father's age: $F (1, 236) = 30.50$, $p < 0.001$
Chevron	father's age: $F (1, 236) = 28.91$, $p < 0.001$
Wave	No wave syllable at P3 in the AFO, from P6 to P12:
	father's age: $F (1, 177) = 30.76$, $p < 0.001$
	father's age × day: $F (2, 177) = 6.69$, $p = 0.002$
Complex	father's age: $F (1, 236) = 17.64$, $p < 0.001$
	father's age × day: $F (3, 236) = 2.77$, $p = 0.042$
One jump	father's age: $F (1, 236) = 7.67$, $p = 0.018$
More jump	father's age: $F (1, 236) = 16.03$, $p < 0.001$
More jump + harmonics	father's age: $F (1, 236) = 10.47$, $p = 0.002$

Table 3. The percentage composition of syllable types was significantly different between the YFO and AFO.

Postnatal day	Results of MANOVA	Syllable types	Results of *t*-test
P3	F (11, 49) = 1.65, p = 0.114	Chevron	8.88 ± 1.34%: YFO vs 4.33 ± 0.88%: AFO, p = 0.006
		Wave	0.26 ± 0.11%: YFO vs 0%: AFO, p = 0.026
P6	F (11, 49) = 2.92, p = 0.005	Downward	46.76 ± 2.27%: YFO vs 56.29 ± 4.21%: AFO, p = 0.046
		Chevron	6.01 ± 0.63%: YFO vs 3.32 ± 0.57%: AFO, p = 0.003
		Complex	0.70 ± 0.27%: YFO vs 0.08 ± 0.043%: AFO, p = 0.037
		One jump	16.27 ± 2.01%: YFO vs 8.03 ± 2.01%: AFO, p = 0.006
		More jump	2.87 ± 0.74%: YFO vs 0.71 ± 0.28%: AFO, p = 0.011
		One jump + harmonics	3.05 ± 0.5%: YFO vs 1.35 ± 0.34%: AFO, p = 0.008
P9	F (11, 49) = 5.37, p < 0.001	Upward	15.54 ± 1.57%: YFO vs 10.63 ± 1.74%: AFO, p = 0.039
		Downward	36.54 ± 1.49%: YFO vs 46.03 ± 2.34%: AFO, p < 0.001
		Flat	1.86 ± 0.27%: YFO vs 4.78 ± 1.36%: AFO, p = 0.031
		Short	5.6 ± 0.67%: YFO vs 9.76 ± 2.05%: AFO, p = 0.050
		Chevron	11.46 ± 0.78%: YFO vs 5.77 ± 0.87%: AFO, p < 0.001
		Wave	3.08 ± 0.43%: YFO vs 0.62 ± 0.16%: AFO, p < 0.001
		Complex	2.08 ± 0.4%: YFO vs 0.29 ± 0.1%: AFO, p < 0.001
		More jump	2.22 ± 0.47%: YFO vs 0.87 ± 0.22%:AFO, p = 0.015
		Harmonics	2.15 ± 0.46%: YFO vs 4.39 ± 0.99%: AFO, p = 0.038
		More jump + harmonics	0.86 ± 0.2%: YFO vs 0.2 ± 0.07%: AFO, p = 0.004
P12	F (11, 49) = 2.73, p = 0.008	Downward	33.14 ± 1.53%: YFO vs 40.57 ± 2.52%: AFO, p = 0.013
		Wave	5.51 ± 0.71%: YFO vs 2.00 ± 0.47%: AFO, p < 0.001
		Complex	4.41 ± 0.64%: YFO vs 2.38 ± 0.66%: AFO, p = 0.030
		More jump	2.88 ± 0.4%: YFO vs 1.31 ± 0.47%: AFO, p = 0.012
		Harmonics	1.98 ± 0.36%: YFO vs 1.05 ± 0.93: AFO, p = 0.047
		More jump + harmonics	1.45 ± 0.31%: YFO vs 0.31 ± 0.14%: AFO, p = 0.002

YFO: young father-derived offspring; AFO: aged father-derived offspring.

Table 4. The variabilities of PC1 and PC2 from P3 to P12.

	P3	P6	P9	P12
PC1	55.6%	62.3%	62.6%	59.2%
PC2	25.6%	21.1%	20.3%	23.4%
PC1 + PC2	81.2%	83.4%	82.9%	82.6%

Table 5. The correlation coefficients and *p*-value between the body weight and USV.

		YFO	AFO
		Body weight	Body weight
P3	Number of types	0.03 (0.890)	0.32 (0.095)
	Number of syllables	0.23 (0.211)	**0.39 (0.037)**
	Duration	-0.14 (0.453)	-0.00 (0.999)
	Max frequency	0.27 (0.143)	-0.04 (0.848)
	Max amplitude	-0.12 (0.531)	0.06 (0.757)
	Interval	0.14 (0.500)	0.18 (0.384)
P6	Number of types	**0.42 (0.016)**	**0.40 (0.032)**
	Number of syllables	**0.50 (0.004)**	0.37 (0.051)
	Duration	**0.41 (0.019)**	0.13 (0.517)
	Max frequency	0.04 (0.838)	0.22 (0.249)
	Max amplitude	0.16 (0.373)	0.05 (0.807)
	Interval	0.22 (0.227)	-0.35 (0.107)
P9	Number of types	-0.15 (0.427)	**0.40 (0.031)**
	Number of syllables	0.01 (0.971)	0.31 (0.101)
	Duration	0.08 (0.666)	0.32 (0.090)
	Max frequency	0.05 (0.770)	0.07 (0.725)
	Max amplitude	-0.10 (0.601)	0.27 (0.158)
	Interval	-0.18 (0.317)	-0.24 (0.219)
P12	Number of types	-0.26 (0.151)	0.14 (0.456)
	Number of syllables	-0.31 (0.086)	0.23 (0.222)
	Duration	-0.15 (0.416)	0.36 (0.057)
	Max frequency	**0.44 (0.012)**	-0.15 (0.433)
	Max amplitude	-0.31 (0.080)	0.30 (0.111)
	Interval	-0.01 (0.951)	-0.25 (0.221)

Correlation coefficients (p-value)

The bold values are statistically significant (p < 0.05).

Fig 1

A

B

- downward
- one jump
- harmonics
- upward
- chevron
- wave
- complex
- more jump + harmonics
- short
- one jump + harmonics
- more jump
- flat

Fig 2

Fig 3

Fig 4

A

P3   P6   P9   P12

YFO

AFO

- ● downward   ● one jump   ● harmonics
- ● upward   ● chevron   ● wave   ● complex   ● more jump + harmonics
- ● short   ● one jump + harmonics   ● more jump   ● flat

B

number of syllable types

P3   P6   P9   P12

†††

C

entropy score

1
0.8
0.6
0.4
0.2
0

P3   P6   P9   P12

†††

— YFO
— AFO

Fig 5

Fig 6

YFO
AFO

A

B
cluster 1

C
cluster 2

D
cluster 3

E
cluster 4

F

Fig 7

A

B

cluster 1

C cluster 2   D cluster 3   E cluster 4

F cluster 5   G

Fig 8

Fig 9