Systemic Commitments of the Context-Dependency; Basics of Elementary Coordination in

Circadian Cycles

Authors: Chulwook Park[1]

[1] International Institute for Applied Systems Analysis (IIASA), A-2361 Laxenburg, Austria.

Correspondence should be addressed to Chulwook Park; parkc@iiasa.ac.at

Abstract

The present study was attempted to measure whether the dynamics of bimanual coordination is influenced by an overarching temporal structure that is embedded in circadian rhythms (part 1) as well as the systemic proof associated with the intelligent capabilities (part 2). For part 1, evidence of entrainment or any influence of the embedding rhythm (a fixed-phase shift, variability, and entropy) were examined on the stability or attractor location. The estimations from the dynamics of the relative phase between the two oscillations show that while (i) circadian effects under the artificially perturbed temperature manipulation were not straightforward along the day-night temperature cycle, (ii) the circadian effect divided by the ordinary circadian seems to be constant along the day-night temperature cycle. Corresponding to this evidence related to performance consequences depending on the organism and environmental interaction, the part 2 determined the impact of circadian (mis)alignment on biological functions and raised the possibility that the disruption of circadian systems may contribute to physical complications. The observations entail rules that self-attunement of current performance may develop not at a single component but across many nested, inter-connected scales. These inter-dependencies from different object phase may allow a potential context-dependent explanation for goal-oriented movements and the emergent assumption of a principle of organisms embedded into their environmental contexts.

Keywords: Physical intelligence, bi-manual coordination, circadian rhythm, thermodynamics, context dependency.

Acknowledgments

This research was supported by an NSF Grant BCS-1344725 (INSPIRE Track 1), and the National Research Foundation of Korea (2016K2A9A1A02952017). All of them provided written informed consent to the study approved by the local ethics committee (SNUIRB No.1509/002-002) and conformed to the ethical standards of the 1964 Declaration of Helsinki (Collaborative Institutional Training Initiative Program, report ID 20481572).

Systemic Commitments of the Context-Dependency; Basics of Elementary Coordination in

Circadian Cycles

Men, earthworms, single-cell organisms, and even plants are aware of their

surroundings and organize actions according to their circumstances (Cariani, 1993; Ford,

2008; Strong & Ray, 1975). Exhibits of autonomy and control of functions have been the

goal in contemporary research to explain agency more scientifically (Shaw & Kinsella-Shaw,

1988). To understand this directed behavior as a call for the organism-environment system

rather than as merely pertaining to the organism (Turvey & Carello, 2012), we seek to expose

the laws that underlie intelligent capabilities.

One useful strategy looks for cycles at all time scales and aims to show how

interacting cyclic processes account for the emergence of new entities, many of which are

similarly cyclic (Iberall, 2001). The central idea is that the earth's cycles—the geophysical,

hydrological, meteorological, geochemical, and biochemical—have interacted to create self-

replicating living systems abiding particular cyclicities (Soodak & Iberall, 1978). We

inquired into whether something akin to attunement to the environmental 24 day/night

temperature cycles (Maury, Ramsey, Bass, 2010) may be apparent in an experimental setting

of bimanual coordination, a setting that has been used to examine self-organization in

biological systems (Kugler & Turvey, 1987). We asked whether unintentional coordination

with an environmental rhythm might raise by assessing whether the dynamics of bimanual

coordination is influenced by an overarching temporal structure that is irrelevant to the task

(see Part 1).

We should embrace, or at least acknowledge, methodological and theoretical diversity given any observations. As one of the prototypes, circadian systems can be considered to be a comparatively simple frame for dealing with complicated physical phenomena. Presenting strategy that includes very many processes and parameters qualitative understanding (Stepp & Turvey, 2009). We take it that to be proof for the possibility as a way of mechanistic principle in living systems in order to address its pervasiveness. This makes us to its mechanistic identifiable in that how components of certain structures or elementary interact dynamically for that phenomena (Bechtel, 2009a,b). The circadian oscillation considered must be incorporated as an endogenous system that represents of the primary clock in both biology and environment, it is worthwhile to be contacted with appropriate methodological basic (Stepp, Chemero & Turvey, 2011). Contributing from this assumption may open a hypothesis of coupling of organism and environment, as an alternative to understanding systems dynamics (Warren, 2006) (see Part 2).

Part 1: Embedded Motor Synchrony in Circadian Rhythm

The present study of this section does so in two principal ways in order to determine the physical characteristics in an effort to find that approximations under certain conditions serve these self-potentials. The first involves an increase in the capability to self-generate forces along the lines of the roles of the fundamental dimensions of environments (temperature embedding in light-dark cycles). The second tied to observe availability of an internally based source (coordination) or sources of force (stability and entropy) within dynamical boundaries in systematic ways.

Method of the first experimental design

Experiments 1 embedded a bimanual coordination task in an ordinary 24-hour day-night cycle (5:00, 12:00, 17:00, and 24:00). Data sets were subjected to an analysis of variance. The setting asks "Is our system influenced by an ecological feature?" A metronome was used to impose a rhythm reflecting the natural period of the pendulum system. The ordinary temperature condition served as a replication of standard in-phase bimanual coordination tasks. The participant was seated with his or her arms supported on armrests and a pendulum in each hand held firmly to prohibit within-grasp motions. Gaze was elevated to prevent viewing the pendulum oscillations which arose from motions strictly about the wrist. Experiment 1 (n = 8) used a setting that minimized variability: in-phase coordination of the two pendulums (relative phase $\phi = 0°$) with no detuning (i.e., the two pendulums had the same eigenfrequency). In the experiments, bimanual coordination was performed at a metronome beat (1.21 s) — this period was chosen because it corresponded to the natural period of the pendulum system: Amazeen, Amazeen, Trffner, & Turvey, 1997—without concern over amplitude or frequency (see Table 1).

Condition	Participants (N)	Circadian points	Trials at each circadian	Task/rest (min)
Normal	8	5:00 12:00 17:00 00:00	6	1/5

Table 1.1: Data collection for the experiment 1. 8 participants, 4 circadian points, 6 trials at each circadian point. Note: Participants were asked to swing in-phase of their limbs in different anatomy points [192 data set (3-level: wrist, elbow, and shoulder)], but used only wrist joint data (64 set) for analysis. Duration of each trial is 1 minute and 5 minutes rest interval between trials.

Results of the first experimental design

A variety of measures (e.g., phase shift, variability, entropy) were examined for evidence of entrainment or any influence of the embedding rhythm on stability or attractor location. Absolute differences in parameters (fixed point shift, variability as a function of frequency competition, and entropy production) were found, especially, at circadian 17:00 point. The parameters changed in the opposite direction from the core body temperature. While the core body temperature rhythm shows a minimum at 05:00 h but has a maximum at about 17:00 h, behavioural performance (entropy; Shannon, 1948) shows a maximum at 05:00 h but has a more clearly defined minimum at about 17:00 h in day-night cycle (see Figure 1).

	Circadian 5:00		Circadian 12:00		Circadian 17:00		Circadian 00:00	
	ϕ_{ave}	$SD\phi$	ϕ_{ave}	$SD\phi$	ϕ_{ave}	$SD\phi$	ϕ_{ave}	$SD\phi$
P1	1.1	6.0	0.8	5.2	1.5	6.7	6.0	43.1
P2	4.0	43.1	2.8	10.8	5.0	6.2	3.8	16.0
P3	11.2	6.7	4.0	6.0	1.5	5.4	3.9	7.7
P4	2.67	77.3	1.7	83.7	6.6	10.1	7.6	6.7
P5	6.18	40.7	3.0	13.6	2.7	11.3	4.6	20.6
P6	1.95	38.0	5.6	59.7	5.8	33.3	4.62	41.1
P7	3.52	40.1	4.4	20.9	0.8	6.1	3.8	55.2
P8	4.25	17.3	5.5	40.0	1.9	35.9	7.1	44.1
T(C°)	36.6		36.8		37.0		36.6	

Table 1.2: Experimental result 1. Each type of raw value for the normal day-night temperature effects. P denotes the participants, numbered 1 ~ 8, ϕ_{ave} = fixed point shift, $SD\phi$ = variability as a function of the frequency competition. T = core body temperature (Celsius).

Figure 1.1. General tendencies in the normal condition. Normalized = standard score (Z calculation), a.u. = arbitrary unit, blue line = fixed point shift, red line = variability as a function of the frequency competition, Temp = temperature (Celsius), 5:00, 12:00, 17:00, and 00:00.

Figure 1.2. Uncertainty characteristics in the normal condition. The figures denote the estimated entropy states according to the time series. Left = 5:00, middle left = 12:00, middle right 17:00, right = 00:00. Note: for this realization, participant 3's data was used which could be represented as the most similar scores with the arbitrarily normalized scores from the averaged all participants' scores as a function of the frequency competition [$SD\phi$ (rad)].

Method of the second experimental design

Experiments 2 and 3 asked, "How does our system adapt to regular or irregular thermal structures?" Normal and abnormal day-night circadian temperature effects were compared at dawn (5 a.m., approximately when core temperature reaches its minimum) and dusk (5 p.m., approximately when core temperature reaches its maximum; Aschoff, 1983). In-phase coordination without detuning was performed at dawn and dusk. A metronome was used to impose a rhythm reflecting the natural period of the pendulum system. In addition, a short-term, thermodynamic manipulation was introduced. Prior to half of the sessions, participants (n = 8) donned a heated vest for 30 min that increased their core temperature by certain (within 0.5˚, *SD* 0.2˚) degrees (Exp 2). And participants (n = 8) donned an ice vest for 30 min that reduced their core temperature by certain (within 0.4˚, *SD* 0.15˚) degrees (Exp 3). The expectation was that the thermal (increasing or decreasing) manipulation's influence on coordination would interact with a time of day (see Table 2, 3).

Condition	Participants (N)	Circadian points	Trials at each circadian	Task/rest (min)
Normal	8	5:00 17:00	6	1/5
Abnormal (heat based)	8	5:00 17:00	6	1/5

Table 2: Data collection for the experiment 2. 2 conditions, 8 participants, 2 circadian points, 6 trials at each circadian point. Note: Participants were asked to swing in-phase of their limbs in different anatomy points [192 data set (3-level: wrist, elbow, and shoulder)], but used only wrist joint data (64 set) for analysis. Duration of each trial is 1 minute and 5 minutes rest interval between trials.

Condition	Participants (N)	Circadian points	Trials at each circadian	Task/rest (min)
Normal	8	5:00 17:00	6	1/5
Abnormal (ice based)	8	5:00 17:00	6	1/5

Table 3: Data collection for the experiment 3. 2 conditions, 8participants, 2 circadian points, 6 trials at each circadian point. Note: Participants were asked to swing in-phase of their limbs in different anatomy points [192 data set (3-level: wrist, elbow, and shoulder)], but used only wrist joint data (64 set) for analysis. Duration of each trial is 1 minute and 5 minutes rest interval between trials.

Results of the second experimental design

Systems state was estimated from the dynamics of the relative phase between the two limbs oscillating at the wrists. Stability of the performance was affected by the temporal locus during the circadian cycle, as well as the introduction of the heated vest (Exp 2) and the ice vest (Exp 3); the influence of the thermal manipulation was not identical. Even if the same external temperature perturbations were given, the influence of the vest was negatively exaggerated (increasing entropy) at dawn, but the influence of the vest was positively exaggerated (decreasing entropy) in the evening (see Figure 2, 3).

Figure 2. Uncertainty characteristics in the normal vs. abnormal (heat based) conditions. The figures denote the estimated entropy states according to the time series between normal (5:00 and 17:00), vs. abnormal (5:00 and 17:00). N = normal, Ab = abnormal in terms of heat_based experimental design. Note: for this realization, participant 2's data was used which could be represented as the most similar scores with the arbitrarily normalized scores from the averaged all participants' scores as a function of the frequency competition [$SD\phi$ (rad)].

Figure 3. Uncertainty characteristics in the normal vs. abnormal (ice based) conditions. The figures denote the estimated entropy states according to the time series between normal (5:00 and 17:00), vs. abnormal (5:00 and 17:00). N = normal, Ab = abnormal in terms of ice_based experimental design. Note: for this realization, participant 5's data was used which could be represented as the most similar scores with the arbitrarily normalized scores from the averaged all participants' scores as a function of the frequency competition [$SD\phi$ (rad)].

Estimated dynamics from the relative phase between the two limbs, oscillatory bimanual coordination was affected by the temporal locus during the circadian cycle. Results at this biological scale correspond to a theoretical study which has shown that the rate of entropy production is changed when a new energy source is accessed via a nonequilibrium phase transition process (Frank, 2011).

Figure 4. Entropy maximizations for the all uncertainty characteristics. Each figure denotes the estimated entropy forces according to the time series for each experimental design. The figure on the left denotes the entropy forces of the normal condition (red line = 5:00, green line = 12:00, black line = 17:00), which data was from the Figure 1. The figure in the middle denotes the entropy forces of the heat-based normal (red line = 5:00, black line = 17:00) vs. abnormal (green line = 5:00, yellow line = 17:00) conditions, which data was from the Figure 2. The figure on the right denotes the entropy forces of the ice-based normal (red line = 5:00, black line = 17:00) vs. abnormal (green line = 5:00, yellow line = 17:00) conditions, which data was from the Figure 3.

Remarks of the results (theoretical implications 1)

The biology may be a complex sensor of its environment that can very effectively adapt itself in a broad range of different circumstances. Organism may convert internal energy of itself so efficiently that if it were to produce what is physically (thermodynamically) possible (England, 2013). These results from Experiment 1, 2, and 3 reflect that accessing a new energy source differs as a function of the circadian cycle, access that can be manipulated by a temporary thermal manipulation. A generalization of this model may not distinguish organism (x) and environment (y), but points to relevant characteristics that may include the context or circumstances in which physical dynamics takes place as a mutually constraining factor. Because, while (a) circadian effect divided by the artificially perturbed temperature manipulation is not constant according to different day-night

temperature cycle, (b) circadian effect divided by the ordinary temperature seems may be constant according to the different day-night temperature cycle.

Part 2: Systemic Proof of the Context-Dependency

Pervasive interconnectedness—everything is connected with some other thing or things (Mahner & Bunge, 1997) —suggests that behavior is adapted to perceiving both the nested environmental properties and one's own nested behaviors, a union that organizes actions on surrounding circumstances (Reed, 1996). The discovery of direct and robust relation between biological aspects (body temperature and motor synchrony) an environmental process (circadian temperature cycle) may echo adaptation of our biological system to the environment.

These relations of both inner (bi-manual coordination) and outer (circadian temperature cycles) sources will provide a sign post with address intelligence as a natural consequence of the principles pertaining to the emergence of functional systems. Thus, physical principles might be methodologically reducible to any level of things (Iberall, 1977), and in this sense, it will motivate methodological and measurement advances which today constitute a strategy for examining self-organization in biological systems. The effect must extend the reach of thermodynamic theory and perhaps identify general principles that apply more broadly to complex systems.

Conditioning main variables with regards to the components' relationships

A core cycles of our system were influenced by temperature embedding on 24 hour light-dark oscillation (called circadian rhythm) according to the observation (Part 1); biochemical, physiological, or behavioral processes that persist under constant conditions with a period length of ~24 hours (Maury, Ramsey, Bass, 2010). We think that it is worth taking engaged variables when it comes to the experimental measurement and defined relatively independent processes: "circadian" and "temperature." Simultaneously, these patterns or characteristics were constrained by the dynamical criteria of stability pattern when it comes to such systems possess a static equilibrium point, which can be called a point attractor (Kugler, Kelso, & Turvey, 1980).

As discussed above definition about the ϕ (state of the coupling strength given time), we say that these two objects are entangled which means knowing something about one causes we to know something about the other. It means the source of a relative phase is energetically emitting two correlated states in opposite directions. Suppose we take a particle (ϕ) in the state ($\dot{\phi}$) and subjected to experiment with two possible outcome (different directions) states.

Figure 5. Schematic drawing of the measurement system. x_axis means time and y_axis represents space of the two entangled particles (ϕ). The particle departs from a source and move apart in opposite directions ($\pm$). It is detected by a correlated two space with two possible outcome states ($\dot{\phi}$).

The emission of one state is followed quickly by the second, and they share the same plane of stability, one up and one down. The measurement calls these parameters (and label the outcomes) as the biological coupling strength ($\dot{\phi}$), positive (+ = 1) and negative (- = 0). It assumed that a pair of coupling one-half states formed somehow in the singlet coupling state, moving freely in opposite directions. As it is, in classical mechanics, if the measurement of the component is one vector (>), the value of one side must yield the value of the other, opposite, side.

$$\chi \rightarrow \vec{f} \rightarrow f(x), \quad f(x) = \begin{cases} x + 1 \ \ if \ x = 0 \\ x - 1 \ \ if \ x = 1 \end{cases} \tag{1}$$

$$0\ 0\ 1\ 1 \xrightarrow{f(x)} 1\ 1\ 0\ 0 \tag{2}$$

$$2\ binary\ objects = \psi_1(a|0> \ + b|1>)\ \psi_2(c|0> \ - d|1>) \tag{3}$$

$$\therefore \psi = ac|00> \ - ad|01> \ + bc|10> \ -bd|11> \tag{4}$$

This must be state 0 plus 1 for the first particle (ψ_1), with the second particle (ψ_2) being in state 0 minus 1. This is the term 0, 0 for ac, plus the term 0,1 for ad with a minus ad, plus the term 1,0 for bc, and finally plus the term 1,1 for bd.

Because this logic can predict in advance the result of measuring any chosen component of $\dot{\phi}(\alpha, >)$ and $\dot{\phi}(\beta, >)$, by previously measuring the same component of $\dot{\phi}(\alpha, >)$ and $\dot{\phi}(\beta, >)$, it follows that the result of any such measurement must actually be predetermined. The interpretation devises mechanical correlations such that each vector variable ($\vec{x}$) is a single continuous parameter, but the measurements of these parameters made

in state (Ω) have no influence on those in other states. The expected value of the components is then determined as follows;

$$\vec{X} \in \Omega \implies \int P(\vec{x})d\vec{x} = 1 \tag{5}$$

Here, $\vec{X}$ equals the parameter in the state space (Ω). $\int P(\vec{x})d\vec{x}$ is the normalized distribution of these parameters, stating that the system is in a certain interval ($\chi, \chi + d\chi$). However, the present experimental design has another vector apart from the "$\vec{\alpha}$ = circadian" and "$\vec{\beta}$ = temperature" factors, which are always embedded in the system's state as an interconnected component known as "homeostasis = $\vec{\gamma}$." According to the circadian temperature correlates of performance, a circadian temperature component and a homeostatic process are assumed to be mutually interdependent of both processes (Carrier & Monk, 2000). They are relatively independent but always correlated each other in that they are always communicating with one another regardless of how far apart they are according to the same processes. As it is, with normalization [$\int P(\vec{x})d\vec{x} = 1$] mapping onto the three components of β, α, and the value of γ, with the measurement variable $\dot{\phi}$, the results found different component conditions of biological states (entropy) have significant context dependency on each variable combination (see the results of the Part 1).

To describe the collections of the three macroscopic processes results, the interpretation of this finding undertakes a putative approximation. If a measurement considers the symmetric coupling strength ($\dot{\phi}$) as positive (+) and negative (-) with a value of 1 and 0 (to simplify a two-by-two matrix), respectively, for the state of the particle, the measurement can reach a possible outcome function having three components (sine waves during the day).

$$\dot\phi(1) \qquad\qquad \dot\phi(1) \qquad\qquad \dot\phi(1)$$

$$\phi \qquad\qquad\qquad \phi \qquad\qquad\qquad \phi$$

$$\dot\phi(0) \qquad\qquad \dot\phi(0) \qquad\qquad \dot\phi(0)$$

α = Circadian   β = Temperature   γ = Homeostasis

Figure 6. Illustration of a three double-valued parameters specification of a physical property. α means circadian component. β denotes temperature component. γ represents homeostatic component. 1 is outcome of the $\dot\phi$ change, while 0 is not change.

Thus, the following binary objects function as a unit vector (λ), meaning that basic logic of this measurement takes the state of 0, resulting in 1. On the other hand, it takes 1 and gives 0.

$$Unitary\ vector\ (\lambda) = |0> \to |1>, or\ |1> \to |0> \qquad (6)$$

Considering the possible outcomes of these components underlying the unitary vector operation, there must be eight possibilities (see Table 4; parameter). Let this measurement considers a situation where ϕ faces the process of the first component with a change but faces the next process with no change, labelling this as $\vec{\alpha}\,\vec{\beta}'$ (the circadian changes and not the temperature); that is, α equals 1 and β equals 0. There are two rows in a table for which α equals 1 and β equals 0. Next, if considering the situation where ϕ passes process $\vec{\beta}\,\vec{\gamma}'$ (a temperature changes and no homeostasis), this logic has two rows where β equals 1 and γ equals 0. Finally, consider the passing of $\vec{\alpha}\,\vec{\gamma}'$ (a circadian change and no homeostasis), the logic also has two rows where α equals 1 and γ equals 0.

		1	2	3	4	5	6	7	8
Parameters	$\vec{\alpha}$ = Circadian	0	0	0	0	1	1	1	1
	$\vec{\beta}$ = Temperature	0	0	1	1	0	0	1	1
	$\vec{\gamma}$ = Homeostasis	0	1	0	1	0	1	0	1
Conditions	$\vec{\alpha}\,\vec{\beta}'$					∨	∨		
	$\vec{\beta}\,\vec{\gamma}'$				∨			∨	
	$\vec{\alpha}\,\vec{\gamma}'$						∨		∨

Table 4: Visualizing the number of objects with combination outcomes. Notice: as indicated by these lines, whenever $\vec{\alpha}\,\vec{\gamma}'$ occurs either $\vec{\alpha}\,\vec{\beta}'$ or $\vec{\beta}\,\vec{\gamma}'$ must also happen, conversely, table for which $\vec{\alpha}\,\vec{\gamma}'$ happens but either $\vec{\alpha}\,\vec{\beta}'$ or $\vec{\beta}\,\vec{\gamma}'$ does not happen.

Proving components' relationship by way of inequality

Assuming these combinations, the interpretation can write the formula considering the probability distribution of the three unit vectors, as follows;

$$\lambda \in \Omega = \phi\left(\vec{\alpha}\vec{\beta}'\right) + \phi\left(\vec{\beta}\vec{\gamma}'\right) \geq \phi\left(\vec{\alpha}\vec{\gamma}'\right) \tag{7}$$

Because the statistical predictions of this correlation cannot be arbitrarily equal to the vectors combinations (Bell, 1964), the formal proof of this can be surmised as shown below.

Figure 7. Representing the number of objects with properties. α = circadian component. β = temperature component. γ = homeostatic component.

Figure 7 shows some of the interconnected relationship of the objects. The parameters α, β, and γ refer to measurements of the 1:1 frequency-locked synchronization phase ($\dot{\phi}$). If any property enters another property first, the component can subsequently include the others and change simultaneously. The measurement has calculations which keep track of the area of the circles $\vec{\alpha}\,\vec{\beta}'$, $\vec{\beta}\,\vec{\gamma}'$, and $\vec{\alpha}\,\vec{\gamma}'$. These relationships can be indicated as the given proportion (A; circle area).

$$A\left(\vec{\alpha}\vec{\beta}'\right) \quad + \quad A\left(\vec{\beta}\vec{\gamma}'\right) \quad \geq \quad A\left(\vec{\alpha}\vec{\gamma}'\right)$$

Figure 8. Mapping of the objects with the area of the circles. α = circadian component. β = temperature component. γ = homeostatic component.

The area of the circle of the $\vec{\alpha}\,\vec{\beta}'$ proportion plus the area of the circle of the $\vec{\beta}\,\vec{\gamma}'$ proportion must be greater or equal to the area of the circle of the $\vec{\alpha}\,\vec{\gamma}'$ proportion. As indicated by the combination table, this is because the last combination ($\vec{\alpha}\,\vec{\gamma}'$) cannot occur without one and the other two combinations ($\vec{\alpha}\,\vec{\beta}'$, $\vec{\beta}\,\vec{\gamma}'$) also transpiring. However, it is possible for one of the first two combinations ($\vec{\alpha}\,\vec{\beta}'$, $\vec{\beta}\,\vec{\gamma}'$) to occur without the third combination ($\vec{\alpha}\,\vec{\gamma}'$) transpiring. Their two frequencies (rows) in the table $\vec{\alpha}\,\vec{\beta}'$ or $\vec{\beta}\,\vec{\gamma}'$ are captured, while $\alpha\vec{\gamma}$ is not. Therefore, the measurement has an inequality on the left side [$A\left(\vec{\alpha}\,\vec{\beta}'\right) + A\left(\vec{\beta}\,\vec{\gamma}'\right)$] which is greater than or equal to that on the right side [$A\left(\vec{\alpha}\,\vec{\gamma}'\right)$].

Because this assumption is true for any collection of relationships, this logic assumes a type of realism in the proof that these parameters have properties regardless of whether they

are measured or specified or not; the logic rewrites these proportions in terms of the sector

area ($\hat{A}$) of each circle.

$$\lambda \in \Omega = \hat{A}\left(\vec{\alpha}\,\vec{\beta}'\right) + \hat{A}\left(\vec{\beta}\,\vec{\gamma}'\right) \geq \hat{A}\left(\vec{\alpha}\,\vec{\gamma}'\right) \tag{8}$$

Based on the above logic, this representation simulates the sector area from each

circle for which two lines have identical radiuses. These two radiuses from the center of the

circle create the area of the circle. It is necessary to measure the proportion of the large area

or the sector of the circle because, as shown in the diagram (Figure 8), the proportion of the

$\vec{\alpha}\,\vec{\beta}'$ area is indicated as the large part of the circle. Next, the representation gives the radius

tentatively as a value of two inches and give the large area of the circle as ¾ (the small area is

¼). The area of the circle is pi (π) multiplied by the radius squared ($A = \pi r^2$). Given that the

overall angle of the circle α is 360 degrees, the $\vec{\alpha\beta}$ portion of the circle as referred to here is

270 degrees (small area: 90 degree). If the measurement takes this fraction and multiplies it

by the area of the entire circle, the result is as follows;

$$\hat{A} = \frac{\vec{\alpha}\,\vec{\beta}'^{\,\circ}}{\vec{\alpha}^{\,\circ}}(\pi r^2) \tag{9}$$

Here, $\hat{A}$ denotes the sector area of component α. $\vec{\alpha\beta}^{\circ}$ is a large portion of θ°(270

degree), and α° is the entire portion of θ°(360 degree). The equation for $\vec{\alpha\beta}^{\circ}$ over α° requires

the multiplication of the area of the entire circle, which is (πr^2). If the measurement

simplifies both the numerator and denominator of this fraction, the result is ¾, after which the

logic multiplies this value by π multiplied by the radius (2) squared, giving a solution to the

proportions of the large sector area of these three components [3/4(π4) = 9. 42 sq.in.]. These

complex portions are represented via the Cartesian system for a linear relationship (complex system for a nonlinear), as shown below;

$$\hat{A}\left(\vec{\alpha}\,\vec{\beta}'\right) + \hat{A}\left(\vec{\beta}\,\vec{\gamma}'\right) \qquad\qquad \geq \qquad\qquad \hat{A}\left(\vec{\alpha}\,\vec{\gamma}'\right)$$

Figure 9. representing the probability of objects with Cartesian coordinate system. Left side of the upper plot = combined proportion of $\vec{\alpha}\,\vec{\beta}'$ and $\vec{\beta}\,\vec{\gamma}'$; [relative portion: $3.5/4(\pi 4) = 10.99$ sq.in.]. Right side of the upper plot = proportion of $\vec{\alpha}\,\vec{\gamma}'$; [relative portion: $3/4(\pi 4) = 9.42$ sq.in.]. Bottom shows their inequalities of the system consideration of the complex behavior [Initial value of x is 0.l, iteration frequency 20, and X_{n+1} is $r\chi_n(1 - X_n)$]. Left plot is showing when parameter value of r equals 3.5 (combined proportion of $\alpha\vec{\beta}$ and $\beta\vec{\gamma}$; 3.5/4), while right is showing the r is 3 (proportion of $\alpha\vec{\gamma}$; 3/4).

The fact that all objects each have all properties including the others corresponds to the changes. The main principle of this equation is the simultaneous measure, for instance, property "$\vec{\alpha}$" and property "$\vec{\alpha}\,\vec{\beta}'$" of the same property. These interpretations are taken as

proof that our experimental feature by which the principles of the state of action and the antecedence do apply to the system. Regardless of how many trials are run, because we have three parameters, it will always be a case where left combinations ($\vec{\alpha}$, $\vec{\beta}$, and $\vec{\gamma}$) must greater than or equal to right combinations ($\vec{\alpha}$ and $\vec{\gamma}$), such as our perturbed morning entropy being greater than normal condition.

However, here is the experimental complex which must be proved regarding the temperature perturbed pm condition: the measurement assumes the experimental set again with wave function $[g(x)]$ similar to when object (ϕ) faces the parameters with the states of their angles up ($\uparrow$) or down ($\downarrow$).

$$f(x) \rightarrow \vec{g} \rightarrow g[f(x)], \quad g[f(x)] = g(x) = \begin{cases} up\ (\uparrow)\ if\ f(x) = 1 \\ down\ (\downarrow)\ if\ f(x) = 0 \end{cases} \tag{10}$$

The first parameter ($\vec{\alpha}$) changes the state to up or down at 0, and ($\vec{\beta'}$) changes the state up or down at θ. The second parameter ($\vec{\beta}$) then changes the state up or down at θ, while ($\vec{\gamma'}$) changes the state up and down at 2θ. The third parameter ($\vec{\alpha}$) changes the state up or down at 0, and ($\vec{\gamma'}$) changes the state up and down at 2θ.

Angle / Process	0_ϕ $g(x)_\alpha$	θ_ϕ $g(x)_\beta$	$2\theta_\phi$ $g(x)_\gamma$
First	($\uparrow$)	($\downarrow$)	
Second		($\uparrow$)	($\downarrow$)
Third	($\uparrow$)		($\downarrow$)

Table 5: Representing the process (first, second, and third) with properties. α = circadian process. β = temperature process. γ = homeostatic process. ($\uparrow$), ($\downarrow$) object's (ϕ) state up or down based on $g(x)$ function.

This function can simplify the inequality considering the experimental relative phase change at each parameter set as follows;

$$\lambda \in \Omega = \phi\left(\vec{\alpha}\,\vec{\beta}'\right) + \phi\left(\vec{\beta}\,\vec{\gamma}'\right) \geq \phi\left(\vec{\alpha}\,\vec{\gamma}'\right) \tag{11}$$

$$\psi = \dot{\phi}_1(\uparrow_0, \downarrow_\theta) + \dot{\phi}_2(\uparrow_\theta, \downarrow_{2\theta}) \geq \dot{\phi}_3(\uparrow_0, \downarrow_{2\theta}) \tag{12}$$

The expressions above assume that with $\dot{\phi}_1$, if the measurement of one side (the entangled relative phase) is up or down at 0, the opposite side of the state must be up or down at θ because they are entangled with a unitary vector operator. For $\dot{\phi}_2$, the next state of one side is then going to be up or down at θ, and the opposite side of the state must then be up or down at 2θ. For $\dot{\phi}_3$, one side is up or down at 0, and the state of the opposite side must then be up or down at 2θ. This inequality can be computed with the common angle scales shown below,

$$0 = Cos\left(\frac{\pi}{2}\right) = \downarrow_0 \tag{13}$$

$$1 = Sin\left(\frac{\pi}{2}\right) = \uparrow_0 \tag{14}$$

When the measurement refers to angles, the two common "scales" are degrees and radians. In degrees, the measurement uses the π notation. Understand that π (radians) is equivalent to 180 degrees, meaning that $\pi/2$ equals 90 degrees. In this case, the sine of $\pi/2$ equals 1/1, and when the sine of $\pi/2$ is 1, cos $\pi/2$ becomes 0. However, to prove this, because

this logic cannot be defined by the relative phase calculation $[\phi = \Delta\omega(t) + \phi(0)]$ simply as

a π value ($\phi \neq \pi$), the change in the relative phase angle is replaced by a putative degree of

theta (θ) after which each state is expressed as follows;

$$\downarrow_\theta = Cos\left(\frac{\theta}{2}\right)\downarrow_0 + i\,Sin\left(\frac{\theta}{2}\right)\uparrow_0 \tag{15}$$

In this equation, if the phase angle is down at theta $[Cos\left(\frac{\theta}{2}\right)\downarrow_0]$, the probability that

the phase being up at 0 is $[i\,Sin\left(\frac{\theta}{2}\right) = \uparrow_0]$. It should be $\downarrow_\theta$ because $Cos\left(\frac{\theta}{2}\right) + i\,Sin\left(\frac{\theta}{2}\right)$ equals

1 (i is the imaginary unit). Thus, the probability of the first state phase being up or down at 0

is $[Sin^2\left(\frac{\theta}{2}\right)]$. Likewise, the next probability that the phase is up or down at theta ($\uparrow_\theta$) and up

or down at 2 theta ($\downarrow_{2\theta}$) must be identical, as the rotation of theta gives us an identical value.

$$\dot{\phi}_1(\uparrow_0, \downarrow_\theta) = Sin^2(x) \times \frac{\theta}{2} = \frac{1}{2}\theta Sin^2(x) = Sin^2\left(\frac{\theta}{2}\right) \tag{16}$$

$$\dot{\phi}_2(\uparrow_\theta, \downarrow_{2\theta}) = Sin^2(x) \times \frac{\theta}{2} = \frac{1}{2}\theta Sin^2(x) = Sin^2\left(\frac{\theta}{2}\right) \tag{17}$$

However, the probability that the phase is up at 0 ($\uparrow_0$) and down at 2 theta ($\downarrow_{2\theta}$) is the

sine squared of theta, as this state only takes a factor of 2 for theta.

$$\dot{\phi}_3(\uparrow_0, \downarrow_{2\theta}) = Sin^2(x)\theta = Sin^2(\theta) \tag{18}$$

Based on these wave functions, the terms can be redefined as follows;

$$Sin^2\left(\frac{\theta}{2}\right) + Sin^2\left(\frac{\theta}{2}\right) \quad ? \quad Sin^2(\theta) \tag{19}$$

Hence, the sine squared theta is less than 1 ($\theta < 1$), and this function can be simplified on the left side as follows;

$$\dot{\phi}_1(\uparrow_0, \downarrow_\theta) = Sin^2\left(\frac{\theta}{2}\right) = \left(\frac{\theta}{2}\right)^2 \tag{20}$$

$$\dot{\phi}_2(\uparrow_\theta, \downarrow_{2\theta}) = Sin^2\left(\frac{\theta}{2}\right) = \left(\frac{\theta}{2}\right)^2 \tag{21}$$

$$\dot{\phi}_1(\uparrow_0, \downarrow_\theta) + \dot{\phi}_2(\uparrow_\theta, \downarrow_{2\theta}) = \left(\frac{\theta}{2}\right)^2 + \left(\frac{\theta}{2}\right)^2 = \frac{\theta^2}{4} + \frac{\theta^2}{4} = \frac{2\theta^2}{4} = \frac{\theta^2}{2} \tag{22}$$

This can also be done on the right side, as follows;

$$\dot{\phi}_3(\uparrow_0, \downarrow_{2\theta}) = Sin^2(\theta) = \theta^2 \tag{23}$$

The measurement can finally be written in the terms below, to reevaluate the inequality.

$$\psi = \frac{\theta^2}{2} < \theta^2 \tag{24}$$

As shown above, the left-hand side $[\phi\left(\vec{\alpha}\,\vec{\beta}'\right) + \phi\left(\vec{\beta}\,\vec{\gamma}'\right)]$ cannot be greater than or equal to the right-hand side $[\phi\left(\vec{\alpha}\,\vec{\gamma}'\right)]$.

$$\left\{\lambda \in \Omega | - \dot{\phi}_1\left(\vec{\alpha}\vec{\beta}'\right) + \dot{\phi}_2\left(\vec{\beta}\vec{\gamma}'\right) < -\dot{\phi}_3\left(\vec{\alpha}\vec{\gamma}'\right)\right\} \tag{25}$$

Using these observations, the measurement finds an explicit violation of a previous inequality (Harrison, 1982) and suggest this as empirical proof underlying the probability of temperature-perturbed pm events. This set solution can also be represented by the Cartesian system for linear relationships (complex system for a nonlinear), as follows;

Figure 10. Representing the probability of objects with Cartesian coordinate system(Upper). Upper left = combined proportion of $\vec{\alpha}\,\vec{\beta}'$ and $\vec{\beta}\,\vec{\gamma}'$; [relative portion: 1.5/4(π4) = 4. 71 sq.in.]. Upper right = proportion of $\vec{\alpha}\,\vec{\gamma}'$; [relative portion: 3/4(π4) = 9. 42 sq.in.]. Bottom plots illustrate the complex behavior [Initial value of x is 0.l, iteration frequency 20, and X_{n+1} is $r\chi_n(1 - X_n)$]. Left plot is showing when parameter value of r equals 1.5 (combined proportion of $\alpha\vec{\beta}$ and $\beta\vec{\gamma}$; 1.5/4), while right is showing the r is 3 (proportion of $\alpha\vec{\gamma}$; 3/4).

The result shows that the predictions of outcomes for the states by the system are inconsistent (Hensen, Bernien, Dréau, Reiserer, Kalb, Blok, ... & Amaya, 2015; Wiseman, 2015) with the parameters (different combinations) and measurements (different calculations). In the context of interaction between all parameters and measurements, the deviation of phase inequality asserts a fundamental limit on the precision of certain pairs of physical position and momentum properties simultaneously. The remarkable possibility, this interpretation can estimate, is that whereas the relationship between objects' energy and reaction rates obtains only under certain constraints, having derived inequality here indicates

that the entangled objects' states applies in cases where many degrees of freedom in the system states out of thermal equilibrium (England, 2013). These entangled features prove the heterogeneity of the system in an ecologically dependent process context.

Remarks of the results (theoretical implications 2)

As evident from the related biological scale, we encourage the belief that there is no common prediction of bias for uncertainty toward the system of ecology. The space between antecedents and consequences is the heart of the process, and we must look at how the variables change. It is recursive in that each variable and process can affect each other depending on where in the flow of behavior one begins. A few scholars in science have investigated experimental models that guide the circadian process. Differential time-of-day variations for different tasks were observed under a normal day-night condition. No attempt was made to distinguish variations in performance due to endogenous circadian factors relative to those linked to the amount of time since awaking. Perhaps the main conclusion to be drawn from studies of the effects of the time of day on performance is that the best time to perform a particular task depends on the nature of that task (Folkard, 1983).

One study showed an early morning peak of mental arithmetic performance in children (Rutenfranz & Helbruegge 1957), while another study found an evening peak for this type of performance in highly practiced young adults (Blake, 1967). With a low working memory load, performance was correlated positively with the circadian rhythm of the body temperature (Folkard, Knauth, Monk, & Rutenfranz, 1967). The majority of the performance-related components (flexibility, muscle strength, short-term memory) appear to vary with the time of day. In particular, contemporary models of subjective alertness and performance

efficiency view these variables as being determined both by a homeostatic process (number of hours since waking) and by input from the circadian timing system (Monk, Weitzman, Fookson, Moline, Kronauer, & Gander, 1983; Johnson et al., 1992). There is still much work to do before one can understand which performance tasks will show different time-of-day effects and which will define the mechanisms that underlie these differences.

Frequency locked 1:1 coupling embedded on a bi-manual pendulum ($\dot{\phi} = \theta_{Left} - \theta_{Right}$) and the homeostatic process (temperature) embedded on the circadian ($\dot{\phi} = \theta_{Circadian\ cycle} - \theta_{homeostatic\ process}$) can be illuminated by a similar relative phase dynamic function. The result of the part 1 shows that the system's stability decreases more at the circadian AM time when the system is perturbed as compared to the normal circadian AM condition. However, the stability of the system increases more at the circadian PM time in spite of being perturbed as compared to normal circadian PM condition. What relations must hold for the formulae to be an interpretation of a biological state with respect to an environmental process in the system? Assumption of this interpretation does not intend to have any precise, predicative knowledge corresponding to the systemic proof in part 2. For a biological state to be related to the environmental process represented by the system, the system must be considered in conjunction with its basis. This is simply the relationship between one and another with infinite distinct representations of the system's productivities.

Conclusion

Inquiry into the possibility of relating perception-action to dynamics began in the 1970s with the problem of coordination: could a principled dynamical account be given of fundamental rhythmic capabilities involving multiple joints, scores of muscles, and millions

of cells? Efforts to address this question invoked the concepts and tools of nonlinear

dynamics (e.g., Kugler & Turvey, 1987). One useful approach to the relationship between

dynamics and self-organization is provided by *Homeokinetics* (e.g., Soodak & Iberall, 1978),

a strategy that looks for cycles at all time scales and aims to show how interacting cyclic

processes account for the emergence of new entities, many of which are similarly cyclic. The

central idea is that the cycles have interacted to create self-replicating living systems abiding

particular cyclicities (Iberall, 2001). Attunement to these cycles is integral to life. Circadian

rhythms are cycles of particular prominence in contemporary research on living things.

Rather than crediting the rhythms to "clock genes," the dynamic approach considers them as

an emergent property of the system.

They are found in most living things – especially in humans. These rhythms are not to

be confused with the biological system. However, they are related in that the circadian system

drives the biological system. The circadian rhythms that control biological rhythms are

grouped interacting molecules in cells throughout the body that act in sync with each other.

This has been being a major issue, and it is hugely influential on us because we are in effect

physical, emotional and performance systems in our bodies (Monk et al., 1983).

Contemporary models of subjective alertness and performance efficiency view these

variables as being determined both by a homeostatic process (amount of hours since waking)

and by input from the circadian timing system (Johnson et al., 1992). However, there is still

much work to do before one can understand which performance tasks will show different

time-of-day effects with various environmental variables and which mechanisms underlie

these differences. Corresponding to a theoretical study about the rate of the stability of a

system and how it can vary when a new energy source is accessed via a nonequilibrium phase

transition process (Frank, 2011), the results from this experiment appear to reconfirm that

access differs as a function of thermodynamic variables (circadian temperature), which means access that can be manipulated by temporary thermal manipulation methods.

One must know the larger system to characterize the smaller system, but one cannot know the larger system in the absence of the characterization of the smaller system. Such that may be called this the problem of impoverished entailment (Shaw, 2001). This does not address directly, however, the question as to why the baseline of the different behavioral modes differ as they do. The ultimate significance of a recurrence of behavioral modes is that it allows the animal to balance the entropic (order-reducing) degradations associated with the processes, thereby ensuring persistence of its characteristic forms and functions (Iberall & Soodak, 1987). The biology may be a complex sensor of its environment that can very effectively adapt itself in a broad range of different circumstances. Organism may convert internal energy of itself so efficiently that if it were to produce what is thermodynamically possible (England, 2013). As it is, generalization may not distinguish organism (x) from environment (y), or larger (biology) factors from smaller (anatomy); such systems may include the context or contexts in which physical dynamics can exist as a mutual factor. This suggestion would not response the question of where or how the system emergent, but provides a coherent account between biology and environment.

Everything is connected with some other thing or things (Mahner & Bunge, 1997), and adapt their behavior perceiving both the nested environmental properties and their own nested behaviors as a union (Turvey et al., 1981) in order to organize actions according to their surrounded circumstance (Cariani, 1993; Ford, 2008; Reed, 1996; Strong & Ray, 1975). Which may reflect that the mechanism of the system's state that is not a specific component of special properties but a general co-activity encompassing all components (Gottlieb, 2000). It will be a reduction of physical principles (Iberall, 1977) classifying them as having a

nervous system is not necessary condition. Spontaneously evolved engineering of this internal characteristics embedding in external context awaits future investigation so that we labeled these sorts of things to be deserved calling as physical intelligent (Turvey & Carello, 2012).

Reference

Amazeen, E. L., Amazeen, P. G., Treffner, P. J., & Turvey, M. T. (1997). Attention and handedness in bimanual coordination dynamics. *Journal of Experimental Psychology: Human Perception and Performance*, *23*(5), 1552.

Aschoff, J. (1983). Circadian control of body temperature. *Journal of thermal Biology*, *8*(1), 143-147.

Bell, J. S. (1964). On the einstein-podolsky-rosen paradox. *Physics*, *1*(3), 195-200.

Blake, M. J. F. (1967). Time of day effects on performance in a range of tasks. *Psychonomic science*, *9*(6), 349-350.

Cariani, P. (1993). To evolve an ear. Epistemological implications of gordon pask's electrochemical devices. *Systems research*, *10*(3), 19-33.

Carrier, J., & Monk, T. H. (2000). Circadian rhythms of performance: new trends. *Chronobiology international*, *17*(6), 719-732.

England, J. L. (2013). Statistical physics of self-replication. *The Journal of chemical physics*, *139*(12), 121923.

Folkard, S. (1983). Diurnal variation. *Stress and fatigue in human performance*, 245-272.

Folkard, S., Knauth, P., Monk, T. H., & Rutenfranz, J. (1976). The Effect of Memory Load on the Circadian Variation in Performance Efficiency Under a Rapidly Rotating Shift System*. *Ergonomics*, *19*(4), 479-488.

Ford, B. J. (2008). Microscopical substantiation of intelligence in living cells. *Infocus Magazine*, (12), 4.

Frank, T. D. (2011). Collective behavior of biophysical systems with thermodynamic feedback loops: a case study for a nonlinear Markov model, the Takatsuji system. *Modern Physics Letters, B, 25*, 551-568.

Gottlieb, G. (2000). Environmental and behavioral influences on gene activity. *Current directions in psychological science*, *9*(3), 93-97.

Harrison, D. (1982). Bell's inequality and quantum correlations. *American Journal of Physics*, *50*(9), 811-816.

Hensen, B., Bernien, H., Dréau, A. E., Reiserer, A., Kalb, N., Blok, M. S., ... & Amaya, W. (2015). Loophole-free Bell inequality violation using electron spins separated by 1.3 kilometres. *Nature*, *526*(7575), 682-686.

Iberall, A. S. (1977). A field and circuit thermodynamics for integrative physiology. I. Introduction to the general notions. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, *233*(5), R171-R180.

Iberall, A. S. (2001). The physics, chemical physics, and biological physics of the origin of life on earth. *Ecological Psychology, 13,* 315-327.

Iberall, A. S., & Soodak, H. (1987). A physics for complex systems. In *Self-organizing systems* (pp. 499-520). Springer US.

Johnson, M. P., Duffy, J. F., Dijk, D. J., Ronda, J. M., Dyal, C. M., & Czeisler, C. A. (1992). Short-term memory, alertness and performance: a reappraisal of their relationship to body temperature. *Journal of sleep research*, *1*(1), 24-29.

Kräuchi, K., & Wirz-Justice, A. (2001). Circadian clues to sleep onset mechanisms. *Neuropsychopharmacology*, *25*, S92-S96.

Kugler, P. N., Kelso, J. S., & Turvey, M. T. (1980). On the concept of coordinative structures as dissipative structures: I. Theoretical lines of convergence. *Tutorials in motor behavior*, *3*, 3-47.

Kugler, P. N., Kelso, J. S., & Turvey, M. T. (1982). On the control and coordination of naturally developing systems. *The development of movement control and coordination*, 5-78.

Kugler, P. N., & Turvey, M. T. (1987). *Information, natural law and the self-assembly of rhythmic movement*. Hillsdale, NJ: Erlbaum.

Mahner, M., & Bunge, M. (1997). *Foundations of biophilosophy*. Springer Science & Business Media.

Maury, E., Ramsey, K. M., & Bass, J. (2010). Circadian rhythms and metabolic syndrome from experimental genetics to human disease. *Circulation research*, *106*(3), 447-462.

Monk, T. H., Weitzman, E. D., Fookson, J. E., Moline, M. L., Kronauer, R. E., & Gander, P. H. (1983). Task variables determine which biological clock controls circadian rhythms in human performance.

Reed, E. S. (1996). *Encountering the world: Toward an ecological psychology* (Vol. 1). New York: Oxford University Press.

Rutenfranz, J., & Hellbrügge, T. (1957). über Tagesschwankungen der Rechengeschwindigkeit bei 11jÄhrigen Kindern. *Zeitschrift für Kinderheilkunde*, *80*(1), 65-82.

Shannon, C. E. (1948). A note on the concept of entropy. *Bell System Tech. J*, *27*, 379-423.

Shaw, R. E. (2001). Processes, acts, and experiences: Three stances on the problem of intentionality. *Ecological Psychology*, *13*(4), 275-314.

Shaw, R. E., & Kinsella-Shaw, J. (1988). Ecological mechanics: A physical geometry for intentional constraints. *Human movement science*, *7*(2), 155-200.

Soodak, H., & Iberall, A. (1978). Homeokinetics. *Science, 201,* 579-582.

Stepp, N., Chemero, A., & Turvey, M. T. (2011). Philosophy for the rest of cognitive science. *Topics in Cognitive Science*, *3*(2), 425-437.

Stepp, N., & Turvey, M. T. (2010). On strong anticipation. *Cognitive systems research*, *11*(2), 148-164.

Strong Jr, D. R., & Ray Jr, T. S. (1975). Host Tree Location Behavior of a Tropical Vine (M on sters gi gont ea) by S kototropism.

Turvey, M. T., & Carello, C. (2012). On intelligence from first principles: Guidelines for inquiry into the hypothesis of physical intelligence (PI). *Ecological Psychology*, *24*(1), 3-32.

Warren, W. H. (2006). The dynamics of perception and action. *Psychological review*, *113*(2), 358.

Wiseman, H. (2015). Quantum physics: Death by experiment for local realism. *Nature*, *526*(7575), 649-650.