Chromosome-level assemblies of multiple Arabidopsis genomes reveal hotspots of rearrangements with altered evolutionary dynamics

Wen-Biao Jiao[1] and Korbinian Schneeberger[1,2,*]

[1]Max Planck Institute for Plant Breeding Research, Department for Plant Developmental Biology, Carl-von-Linné-Weg 10, 50829 Cologne, Germany

[2]Faculty of Biology, LMU Munich, Großhaderner Str. 2, 82152 Planegg-Martinsried, Germany

*Author for correspondence: schneeberger@mpipz.mpg.de

Abstract

We report chromosome-level, reference-quality assemblies of seven *Arabidopsis thaliana* accessions selected across the global range of this predominately ruderal plant. Each genome revealed between 13-17 Mb rearranged and 5-6 Mb novel sequence introducing copy-number changes in ~5,000 genes, including ~1,900 genes which are not part of the current reference annotation. Analyzing the collinearity between the genomes revealed ~350 regions (4.1% of the euchromatin) where accession-specific tandem duplications destroyed the syntenic gene order between the genomes. These *hotspots of rearrangements* were characterized by the loss of meiotic recombination in hybrids within these regions and the enrichment of genes implicated in biotic stress response. Together this suggests that hotspots of rearrangements are governed by altered evolutionary dynamics as compared to the rest of the genome, which are based on new mutations and not on the recombination of existing variation, and thereby enable a quick response to the ever-evolving challenges of biotic stress.

The individual genomes of sexually reproducing species are typically highly collinear to

enable physical exchange of alleles (chromosome arms) during meiosis. This exchange

ensures the generation of diversity and the removal of deleterious alleles[1–3] and at the time

protects the offspring from major mutations changing the karyotype of a genome[4,5]. Despite

the obvious importance to preserve a common karyotype, the presence of genomic

rearrangements suggests that the genomes are in fact not entirely collinear. Genomic

rearrangements (and the resulting lack of allelic exchange) have been shown to contribute

to population diversification including the evolution of different sexes[6,7] or life-history traits[8].

But even though the absence of collinearity can have drastic effects, there is hardly

anything known about the actual degree of collinearity within populations as most of the

current genome studies are not based on chromosome-level assemblies. The first complete

assembly of a plant genome was the reference sequence of *A. thaliana* (Col-0), which was

based on a minimal tiling path of BACs sequenced with Sanger technology[9]. Since then

multiple hundred Arabidopsis genomes have been studied, however, most of these studies

relied on short-read based resequencing or reference-guided assembly, where the

identification of genomic rearrangements remained challenging[10–16]. In contrast, reference-

independent, chromosome-level assemblies with almost complete reconstruction of the

nucleotide sequence enable accurate identification of all sequence differences and would

therefore reveal the degree of synteny across the genome[17]. So-far, however, there are only

few whole-genome *de-novo* assemblies for *A. thaliana* available and the assemblies have

not been thoroughly compared to each other[18–21].

Using deep PacBio (45-71x) and Illumina (56-78x) whole-genome shotgun sequencing,

we assembled the genomes of seven accessions from geographically diverse regions

including An-1 (Antwerpen, Belgium), C24 (Coimbra, Portugal), Cvi-0 (Cape Verde Islands),

Eri-1 (Eringsboda, Sweden), Kyo (Kyoto, Japan), L*er* (Gorzów Wielkopolski, Poland) and

Sha (Shahdara, Tadjikistan) (Supplementary Table S1) (see Methods). The assembly of L*er*

was already described in the context of the development of a whole-genome comparison

tool used in this study[17]. The seven accessions (together with the reference accession Col-0)

were initially used as the founder lines of Arabidopsis Multi-parent Recombination

Inbreeding Lines (AMPRIL)[22] population and were selected to maximize the genetic

diversity in this set. The contig assemblies featured N50 values from 4.8 - 11.2 Mb and

chromosome-normalized L50 (CL50)[23] values of 1 or 2 indicating that nearly all

chromosome arms were assembled into a few contigs only (Fig. 1, Table 1 and

Supplementary Table 2). We arranged 43 - 73 contigs per genome to chromosome-level

assemblies based on homology to the reference sequence. Even though these assemblies

do rely on the reference sequence, we would like to point out that the sequence assembly

itself was independent of the reference sequence, and that the contigs were large in general,

implying that it is unlikely that we misplaced any of the contigs. To confirm this, we
compared two of the chromosome-level assemblies with three different genetic maps,
where we did not find even a single mis-placed contig (Supplementary Table 3). The seven
chromosome-level assemblies reached a total length of 117.7 - 118.8 Mb, which is very
similar to the 119.1 Mb of the reference sequence (Table 1) and even included parts of the
highly complex regions of centromeres, telomeres and rDNA clusters (Supplementary Table
4 and 5). The remaining unanchored contigs had a total length of 1.5 - 3.3 Mb and consisted
almost entirely of repeats, which agrees with gaps between the contigs, which were mostly
introduced due to repetitive regions (Supplementary Table 6). Overall, we annotated 27,098
to 27,574 protein-coding genes in each of the assemblies, which is similar to the 27,445
genes annotated in the reference sequence[24] (Table 1 and Supplementary Table 7-9) (see
Methods).

By comparing each of the new assemblies against the reference sequence using the
whole-genome comparison tool *SyRI* (V1.1)[17], we found 102.2 - 106.6 Mb of collinear
regions and 12.6 - 17.0 Mb of rearranged regions in each of the genomes (Fig. 2a). The
rearrangements included 1.5 - 4.2 Mb (33 - 46) inversions, 1.8 – 2.9 Mb (729 - 1,192)
translocations, and - most abundant - polymorphic duplications, which comprised 7.2 – 8.7
Mb (4,288 – 5,150) within each of the individual genomes (Supplementary Table 10).
Similar to small-scale sequence variation[25], rearrangements were not evenly distributed
along the chromosomes, but were enriched in pericentromeres (Supplementary Table 11).
Their lengths ranged from a few dozen bp to hundreds of kb and even Mb scale (Fig. 2b),
including a 2.48 Mb inversion specific to chromosome 3 of Sha (Supplementary Fig. 1 and
Supplementary Table 12) which explains the suppression of meiotic recombination in this
region in hybrids including the Sha haplotype[26,27]. Sequence divergence in rearranged
regions was generally higher as compared to collinear regions mostly due an excess of
local copy-gain and copy-loss variation in rearranged regions (Fig. 2a, Supplementary Fig. 2
and Supplementary Table 13).

Genomic rearrangements have the potential to delete, create or duplicate genes
resulting in gene copy number variation (CNV). Based on the clustering of orthologous
genes across all eight accessions[28] we found 22,040 gene families with conserved copy
number, while 4,957 gene families showed differences in gene copy numbers in at least one
accession (Fig. 2c and Supplementary Table 14). Almost 99% of these copy-variable gene
families had a maximum copy number of 5 or less, while only less than 10% of them
showed more than two different copy numbers across the eight accessions (Supplementary
Fig. 3). Among the copy-variable genes we found 1,941 non-reference gene families
including 891 gene families present in at least two of the other accessions (Fig. 2c). Around
23% of the non-reference gene families featured orthologs in the closely related genome of

Arabidopsis lyrata and, according to RNA-seq read mapping, 26%-40% of them showed

evidence of expression (Supplementary Table 15). The remaining 1,050 non-reference

(accessions-specific) gene families were evenly distributed across the accessions (Fig. 2c),

with the exceptions of Cvi-0, where we found nearly twice as many (214) accession-specific

genes which is in agreement with the divergent ancestry of this relict accession[12,29].

Based on all possible pairwise genome comparisons, we identified 5.1 - 6.5 Mb

accession-specific sequence and used this to estimate a pan-genome size of ~135 Mb

including ~30,000 genes and a core-genome size of ~105 Mb with ~24,000 genes (Fig.

2d)[30,31] illustrating that one reference genome is not sufficient to capture the entire

sequence diversity within *A. thaliana*.

As most of the *A. thaliana* genomes have been analyzed with short read sequencing

hardly anything is known about their collinearity. In contract, chromosome-level assemblies

allow for a first analysis of the conservation of the contiguity between multiple individuals.

To quantify the collinearity we developed a new parameter called *Synteny Diversity* π_{syn},

which is similar to *Nucleotide Diversity*[32], however, instead of measuring average sequence

differences it measures the degree of collinearity between the genomes of a population (see

Methods). π_{syn} values can range from 0 to 1, where 1 refers to the complete absence of

collinearity between any of the genomes and 0 to regions where all genomes are collinear.

π_{syn} can be calculated in any given region, however, the annotation of collinearity still needs

to be established within the context of the whole genomes to avoid false assignments of

homologous but non-allelic regions.

We calculated π_{syn} in 5-kb sliding windows across the genome using pair-wise

comparisons of all eight accessions (Fig. 3a). As expected, π_{syn} was generally high in

pericentromeric regions and low in chromosome arms. Overall, this revealed around 90 Mb

(76% of the genome) where all genomes were collinear to each other, while for the

remaining 29 Mb (24%) the collinearity between the genomes was not conserved. This, for

example, included a region on chromosome 3 (ranging from Mb ~2.8 - 5.3), where π_{syn} was

increased to ~0.25 (i.e. one genome is not collinear to all other seven genomes) due to the

2.48 Mb inversion in the Sha genome (Fig. 3a, arrow labelled with (A)).

Unexpectedly, however, some regions featured π_{syn} values even larger than 0.5. This

implied that not only two, but multiple independent, non-collinear haplotypes segregate in

these regions. In turn, this suggest that these regions are more likely to undergo or

conserve complex mutations as compared to the rest of the genomes and thereby create

hotspots of rearrangements (HOT regions) where multiple accessions independently

evolved diverse haplotypes. Overall, we found 576 of such HOT regions with a total size of

10.2 Mb including 351 HOT regions in the gene-rich chromosome arms with a total length of

4.1 Mb (or 4% of euchromatic genome) (Supplementary Table 16).

Even though HOT regions in euchromatic regions included more transposable elements

and less genes as compared to the collinear regions, they still contained significant

numbers of genes, many of which occurred at high and variable copy-number between the

accessions (Fig. 3b,c). For example, a HOT region on chromosome 4, which overlapped

with the *RPP4/RPP5* R gene cluster[33,34], displayed 5-15 intact or truncated copies of the

RPP5 gene across the eight genomes (Fig. 3d and Supplementary Table 17). The different

gene copies were primarily introduced by an accumulation of forward tandem duplications

and large indels (Fig. 3e).

This remarkable pattern of forward tandem duplications and large indels was shared by

many of the HOT regions (Fig. 3c and Supplementary Fig. 4). The clear pattern of almost

exclusively forward tandem duplications suggested higher mutation (duplication) rates,

which are specific to these regions in each of the accessions. In contrast, the borders of the

HOT regions were surprisingly well conserved across the accessions (Fig. 3f). This

suggested that either different selection regimes introduced clear-cut borders between the

HOT regions and their vicinity, or that HOT regions are specific targets of increased tandem

duplication rates. Such a local increase of mutation rates could potentially be mediated by

non-allelic homologous recombination which could be triggered by the high number of local

repeats in these regions[35,36]. Fig. 4 shows two more examples of these complex regions.

In contrast, meiotic recombination in Arabidopsis was shown to be supressed by

structural diversity[37]. To test if HRs are indeed depleted for meiotic recombination, we

overlapped rearranged regions with 15,683 crossover (CO) sites previously identified within

Col-0/Ler F2 progenies[37,38]. Only 64 of them partially overlapped with non-syntenic regions

while all other COs were found in syntenic regions (Fig. 5a), suggesting that HOT regions

are almost completely silenced for COs (P-value < 2e-16). In consequence, this would imply

that HOT regions are segregating as large non-recombining regions. To test this, we

analysed the linkage disequilibrium (LD) within 1,135 genomes of the 1001 Genomes

Project[12] around and across the HOT regions. LD increased in the vicinity of the HOT

regions, with increasing LD close to the HOT regions implying reduced recombination in the

regions surrounding the HOT regions. Likewise, LD was also high within HOT regions

corroborating the recombination suppression in HOT regions. However, when calculated

across the border of these regions, LD was significantly lower supporting the idea that HOT

regions are not strongly linked to the surrounding haplotypes and that they hardly exchange

alleles (Fig. 5b).

Reduced meiotic recombination has been linked to the accumulation of new (deleterious)

mutations[39]. In agreement with this, HOT regions showed an accumulation of SNPs with low

allele frequencies and potentially deleterious variation as compared to other regions in the

genome (Fig. 5c,d and Supplementary Fig. 5). Moreover, reduced recombination combined
with geographic isolation can provide the basis for the development of alleles, which are
incompatible with distantly related haplotypes leading to intra-species incompatibilities[40]. To
test this, we searched the location of nine recently reported genetic incompatible loci[41]
(*DM1-9*) and found that all except of one overlapped with HOT regions, while *DM3*, the
locus which did not overlap with a HOT region, was closely flanked by two HOT regions
(Supplementary Fig. 6-12). In addition, we also checked the locus of a recently published
single-locus genetic incompatibility[42] and found that it was also residing in a HOT region
(Supplementary Fig. 13).

The high structural diversity of the HOT regions was reminiscent of the patterns that
have been described for R gene clusters[43–50]. In fact, the 808 reference genes in HOT
regions were significantly enriched for genes involved in defense response, signal
transduction and secondary metabolite biosynthesis (Fig. 5e) suggesting a reoccurring role
of HOT regions in the adaptation to biotic stress.

As biotic stress is an evolving environmental challenge, the Red Queen hypothesis
suggests that the genomes of *A. thaliana* are in the constant need to diversify their
offspring[51]. It has been proposed that in response to this, meiotic recombination might
increase and thereby diversified offspring is generated[52]. However, exclusively shuffling
existing variation might not be sufficient to respond to the evolution of pathogens. Instead, it
has been proposed that the accumulation of new gene duplicates could enable a rapid
genomic response of plants against pathogens[36,53–55]. The hotspots of rearrangements have
the potential to build the basis for such a response, as frequent gene duplications could
build the basis for an evolutionary playground to evolve a quick response to the challenges
of biotic stress and overcome fitness valleys during the evolution of more complex function.
This, in turn, comes at the costs of loss of synteny and the loss of meiotic recombination
between distant haplotypes. Though it still needs to be analyzed whether local populations
show the same level of diversity or if their haplotypes in HOT regions are more similar and
still exchange alleles, we have observed the negative consequences of reduced meiotic
recombination in this small world-wide population including the accumulation of deleterious
alleles and incompatible epistatic effects between distant genotypes.

Taken together, using chromosome-level genome assemblies of a small, highly diverse
population of *A. thaliana*, we have identified regions where genome collinearity (a
prerequisite for the allelic exchange in populations) was lost due to genome-specific
accumulation of rearrangements. We propose that these regions, which we called *hotspots*
of rearrangements or HOT regions, facilitate evolutionary responses to rapidly changing
environmental challenges through the accumulation of mutations and not by shuffling alleles.
These regions are thus undergoing different evolutionary dynamics as compared to the rest

of the genome, where each region segregates with only few haplotypes. Future genome-
wide screens for selection patterns might need to take the nature of such regions into
account.

Methods

Plant material and whole genome sequencing

We received the seeds of all seven accessions from Maarten Koornneef (MPI for Plant Breeding Research), and grew them under normal greenhouse conditions. DNA preparation and next generation sequencing was performed by the Max Planck Genome center. The DNA samples were prepared as previously described for the L*er*[17], and sequenced with a PacBio Sequel system. For each accession, data from two SMRT cells were generated. Besides, Illumina paired-end libraries were prepared and sequenced on the Illumina HiSeq system.

Genome assembly

PacBio reads were filtered for short (<50bp) or low quality (QV<80) reads using SMRTLink5 package. *De novo* assembly of each genome was initially performed using three different assembly tools including Falcon[21], Canu[56] and MECAT[57]. The resulting assemblies were polished with Arrow from the SMRTLink5 package and then further corrected with mapping of Illumina short reads using BWA[58] to remove small-scale assembly errors which were identified with SAMTools[59]. For each genome, the final assembly was based on the Falcon assembly as these assemblies always showed highest assembly contiguity. A few contigs were further connected or extended based on whole genome alignments between Falcon and Canu or MECAT assemblies. Contigs were labelled as organellar contigs if they showed alignment identity and coverage both larger than 95% when aligned against the mitochondrial or chloroplast reference sequences. A few of contigs aligned to multiple chromosomes and were split if no Illumina short read alignments supported the conflicting regions. Assembly contigs larger than 20kb were combined to pseudo-chromosomes according to their alignment positions when aligned against the reference sequence using MUMmer4[60]. Contigs with consecutive alignments were concatenated with a stretch of 500 Ns. To note, the assembly of the L*er* accession was already described in a recent study[17].

Assembly evaluation

We evaluated the assembly completeness by aligning the reference genes against each of the seven genomes using Blastn[61]. Reference genes which were not aligned or only partially aligned might reveal genes which were missed during the assembly. To examine whether they were really missed, we mapped Illumina short reads from each genome against the reference genome using the BWA[58] and checked the mapping coverage of these genes. The genes, which were missing in the assembly, should show fully alignment

coverage (Supplementary Table 7).

Centromeric and telomeric tandem repeats were annotated by searching for the 178 bp

tandem repeat unit[62] and the 7 bp tandem repeat unit of TTTAGGG[63]. rDNA clusters were

annotated with Infernal version 1.1[64].

The assembly contiguity of Cvi-0 and L*er* were further tested using three previously

published genetic maps[65–67] (Supplementary Table 3). For this we aligned the marker

sequences against the chromosome-level assemblies and checked the order of the markers

in the assembly versus their order in the genetic map. The ordering of contigs to

chromosomes was perfectly supported by all three maps. Overall, only six (out of 1,156)

markers showed conflicts between the genetic and physical map. In all six cases we found

evidence that the conflict was likely caused by structural differences between the parental

genomes.

Gene annotation

Protein-coding genes were annotated based on *ab initio* gene predications, protein

sequence alignments and RNA-seq data. Three *ab initio* gene predication tools were used

including Augustus[68], GlimmerHMM[69] and SNAP[70]. The reference protein sequences from

the Araport 11[24] annotation were aligned to each genome assembly using exonerate[71] with

the parameter setting "--percent 70 --minintron 10 --maxintron 60000". For five accessions

(An-1, C24, Cvi-0, Ler-0, and Sha) we downloaded a total of 155 RNA-seq data sets from

the NCBI SRA database (Supplementary Table 8). RNA-seq reads were mapped to the

corresponding genome using HISAT2[72] and then assembled into transcripts using

StringTie[73] (both with default parameters). All different evidences were integrated into

consensus gene models using Evidence Modeler[74].

The resulting gene models were further evaluated and updated using the Araport 11[24]

annotation. Firstly, for each of the seven genomes, the predicted gene and protein

sequences were aligned to the reference sequence, while all reference gene and protein

sequences were aligned to each of the other seven genomes using Blast[61]. Then,

potentially mis-annotated genes including mis-merged (two or more genes are annotated as

a single gene), mis-split (one gene is annotated as two or more genes) and unannotated

genes were identified based on the alignments using in-house python scripts. Mis-annotated

or unannotated genes were corrected or added by incorporating the open reading frames

generated by *ab initio* predications or protein sequence alignment using Scipio[75].

Noncoding genes were annotated by searching the Rfam database[76] using Infernal

version 1.1[64]. Transposon elements were annotated with RepeatMasker

(http://www.repeatmasker.org). Disease resistance genes were annotated using RGAugury

[77]. NB-LRR R gene clusters were defined based on the annotation from a previous study[78].

Pan-genome analysis

Pan-genome analysis

Pan-genome analyses were performed at both sequence and gene level. To construct a

pan-genome of sequences, we generated pair-wise whole genome sequence alignments of

all possible pairs of the eight genomes using the nucmer in the software package

MUMmer4[60]. A pan-genome was initiated by choosing one of the genomes, followed by

iteratively adding the non-aligned sequence of one of the remaining genomes. Here, non-

aligned sequences were required to be longer than 100bp without alignment with an identity

of more than 90%. The core genome was defined as the sequence space shared by all

sampled genomes. Like the pan-genome, the core-genome analysis was initiated with one

genome. Then all other genomes were iteratively added, while excluding all those regions

which were not aligned against each of the other genomes. The pan- and core-genome of

genes was built in a similar way. The pan-genome of genes was constructed by selecting

the whole protein coding gene set of one of the accessions followed by iteratively adding the

genes of one of the remaining accessions. Likewise, the core-genome of genes was defined

as the genes shared in all sampled genomes.

For each pan or core genomes analysis, all possible $(\sum_{n=1}^{8} \left(\frac{8!}{n!(8-n)} \right))$ combinations of

integrating the eight genomes (or a subset of them) were evaluated. The exponential

regression model y = A e^{Bx} + C to was then used to model the pan-genome/core-genomes

by fitting medians using the least square method implemented in the nls function of R.

Analysis of structural rearrangements and gene CNV

All assemblies were aligned to the reference sequence using nucmer from the

MUMmer4[60] toolbox with parameter setting "-max -l 40 -g 90 -b 100 -c 200". The resulting

alignments were further filtered for alignment length (>100) and identity (>90). Structural

rearrangements and local variations were identified using SyRI[17]. The functional effects of

sequence variation were annotated with snpEff[79]. The gene CNV were identified according

to the gene family clustering using the tool OrthoFinder[28] based on all protein sequences

from the eight accessions.

Synteny diversity, hotspots of rearrangements and diversity estimates

Synteny Diversity was defined as the average fraction of non-syntenic sites found within

all pairwise genome comparisons within a given population. Here we denote *Synteny*

Diversity as

317
$$\pi_{syn} = \sum_{ij} x_i x_j \pi_{ij},$$

where x_i and x_j refer to the frequencies of sequence i and j and π_{ij} to the average

probability of a position in i to be non-syntenic. Note, π_{syn} can be calculated in a given

region or for the entire genome. However even when calculated for small regions the

annotation of synteny still needs to established within the context of the whole genomes to

avoid false assignments of homologous but non-allelic sequence. Here we used the

annotation of *SyRI* to define syntenic regions. π_{syn} values can range from 0 to 1, with higher

values referring to a higher average degree of non-syntenic regions between the genomes.

For the analyses, we calculated π_{syn} in 5-kb sliding windows with 1kb step-size across

the entire genome. HOT regions were defined as regions with π_{syn} larger than 0.5.

Neighboring regions were merged into one HOT region if their distance was shorter than

2kb.

The nucleotide and haplotype diversity were calculated with the R package

PopGenome[80] using SNP markers (with MAF > 0.05) from 1001 Genomes Project[12]. LD

were calculated as correlation coefficients r^2 using SNP markers with MAF > 0.05. GO

enrichment analysis was performed using the webtool DAVID[81,82]

References

1. Barton, N. H. & Charlesworth, B. Why Sex and Recombination. *Science (80-.).* **281**, 1986 LP – 1990 (1998).

2. Rice, W. R. & Chippindale, A. K. Sexual recombination and the power of natural selection. *Science (80-.).* **294**, 555–559 (2001).

3. McDonald, M. J., Rice, D. P. & Desai, M. M. Sex speeds adaptation by altering the dynamics of molecular evolution. *Nature* (2016).

4. Heng, H. H. Q. Elimination of altered karyotypes by sexual reproduction preserves species identity. *Genome* (2007).

5. Gorelick, R. & Heng, H. H. Q. Sex reduces genetic variation: A multidisciplinary review. *Evolution (N. Y).* **65**–**4**, 1088–1098 (2011).

6. Küpper, C. *et al.* A supergene determines highly divergent male reproductive morphs in the ruff. *Nat. Genet.* **48**, 79–83 (2015).

7. Lamichhaney, S. *et al.* Structural genomic changes underlie alternative reproductive strategies in the ruff (Philomachus pugnax). *Nat. Genet.* **48**, 84–88 (2015).

8. Lowry, D. B. & Willis, J. H. A widespread chromosomal inversion polymorphism contributes to a major life-history transition, local adaptation, and reproductive isolation. *PLoS Biol.* **8**, (2010).

9. The Arabidopsis Genome Initiative. Analysis of the genome sequence of the flowering plant Arabidopsis thaliana. *Nature* **408**, 796–815 (2000).

10. Cao, J. *et al.* Whole-genome sequencing of multiple Arabidopsis thaliana populations. *Nat. Genet.* **43**, 956–965 (2011).

11. Long, Q. *et al.* Massive genomic variation and strong selection in Arabidopsis thaliana lines from Sweden. *Nat. Genet.* **45**, 884–890 (2013).

12. Alonso-Blanco, C. *et al.* 1,135 Genomes Reveal the Global Pattern of Polymorphism in Arabidopsis thaliana. *Cell* **166**, 481–491 (2016).

13. Schneeberger, K. *et al.* Reference-guided assembly of four diverse Arabidopsis thaliana genomes. *PNAS* **108**, 10249–10254 (2011).

14. Gan, X. *et al.* Multiple reference genomes and transcriptomes for Arabidopsis thaliana. *Nature* **477**, 419–423 (2011).

15. Schneeberger, K. *et al.* Simultaneous alignment of short reads against multiple genomes. *Genome Biol.* (2009). doi:10.1186/gb-2009-10-9-r98

16. Schmitz, R. J. *et al.* Patterns of population epigenomic diversity. *Nature* **495**, 193–198 (2013).

17. Goel, M., Sun, H., Jiao, W.-B. & Schneeberger, K. SyRI: identification of syntenic and rearranged regions from whole-genome assemblies. *bioRxiv* (2019).

doi:10.1101/546622

18. Zapata, L. *et al.* Chromosome-level assembly of Arabidopsis thaliana Ler reveals the

extent of translocation and inversion polymorphisms. *Proc. Natl. Acad. Sci.* **113**,

E4052–E4060 (2016).

19. Michael, T. P. *et al.* High contiguity Arabidopsis thaliana genome assembly with a

single nanopore flow cell. *Nat. Commun.* **9**, 1–8 (2018).

20. Pucker, B. *et al.* A chromosome-level sequence assembly reveals the structure of the

Arabidopsis thaliana Nd-1 genome and its gene set. *PLoS One* (2019).

doi:10.1371/journal.pone.0216233

21. Chin, C.-S. *et al.* Phased diploid genome assembly with single-molecule real-time

sequencing. *Nat Methods* **13**, 1050–1054 (2016).

22. Huang, X. *et al.* Analysis of natural allelic variation in *Arabidopsis* using a multiparent

recombinant inbred line population. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 4488–4493

(2011).

23. Jiao, W.-B. *et al.* Improving and correcting the contiguity of long-read genome

assemblies of three plant species using optical mapping and chromosome

conformation capture data. *Genome Res.* **27**, 778–786 (2017).

24. Cheng, C. Y. *et al.* Araport11: a complete reannotation of the Arabidopsis thaliana

reference genome. *Plant J.* (2017). doi:10.1111/tpj.13415

25. Clark, R. M. *et al.* Common sequence polymorphisms shaping genetic diversity in

Arabidopsis thaliana. *Science (80-.).* **317**, 338–342 (2007).

26. Loudet, O., Chaillou, S., Camilleri, C., Bouchez, D. & Daniel-Vedele, F. Bay-0 x

Shahdara recombinant inbred line population: A powerful tool for the genetic

dissection of complex traits in Arabidopsis. *Theor. Appl. Genet.* **104**, 1173–1184

(2002).

27. Simon, M. *et al.* Quantitative trait loci mapping in five new large recombinant inbred

line populations of Arabidopsis thaliana genotyped with consensus single-nucleotide

polymorphism markers. *Genetics* **178**, 2253–2264 (2008).

28. Emms, D. M. & Kelly, S. OrthoFinder: solving fundamental biases in whole genome

comparisons dramatically improves orthogroup inference accuracy. *Genome Biol.*

(2015). doi:10.1186/s13059-015-0721-2

29. Durvasula, A. *et al.* African genomes illuminate the early history and transition to

selfing in *Arabidopsis thaliana. Proc. Natl. Acad. Sci.* **114**, 5213–5218 (2017).

30. Tettelin, H. *et al.* Genome analysis of multiple pathogenic isolates of Streptococcus

agalactiae: Implications for the microbial 'pan-genome'. *Proc. Natl. Acad. Sci.* **102**,

13950–13955 (2005).

31. Medini, D., Donati, C., Tettelin, H., Masignani, V. & Rappuoli, R. The microbial pan-

genome. *Current Opinion in Genetics and Development* (2005).

32.   Nei, M. & Li, W. H. Mathematical model for studying genetic variation in terms of

restriction endonucleases. *Proc. Natl. Acad. Sci.* (1979).

33.   Parker, J. E. The Arabidopsis Downy Mildew Resistance Gene RPP5 Shares

Similarity to the Toll and Interleukin-1 Receptors with N and L6. *PLANT CELL*

ONLINE (1997).

34.   Van Der Biezen, E. A., Freddie, C. T., Kahn, K., Parker, J. E. & Jones, J. D. G.

Arabidopsis RPP4 is a member of the RPP5 multigene family of TIR-NB-LRR genes

and confers downy mildew resistance through multiple signalling components. *Plant J.*

29, 439–451 (2002).

35.   Wicker, T., Yahiaoui, N. & Keller, B. Illegitimate recombination is a major evolutionary

mechanism for initiating size variation in plant resistance genes. *Plant J.* (2007).

36.   Leister, D. Tandem and segmental gene duplication and recombination in the

evolution of plant disease resistance genes. *Trends in Genetics* (2004).

37.   Rowan, B. A. *et al.* An ultra high-density Arabidopsis thaliana crossover map that

refines the influences of structural variation and epigenetic features. *bioRxiv* (2019).

38.   Serra, H. *et al.* Massive crossover elevation via combination of HEI10 and recq4a

recq4b during Arabidopsis meiosis . *Proc. Natl. Acad. Sci.* (2018).

39.   Kondrashov, A. S. Deleterious mutations and the evolution of sexual reproduction.

Nature (1988).

40.   Bomblies, K. & Weigel, D. Hybrid necrosis: Autoimmunity as a potential gene-flow

barrier in plant species. *Nat. Rev. Genet.* **8**, 382–393 (2007).

41.   Chae, E. *et al.* Species-wide genetic incompatibility analysis identifies immune genes

as hot spots of deleterious epistasis. *Cell* **159**, 1341–1351 (2014).

42.   Smith, L. M., Bomblies, K. & Weigel, D. Complex evolutionary events at a tandem

cluster of Arabidopsis thaliana genes resulting in a single-locus genetic

incompatibility. *PLoS Genet.* **7**, (2011).

43.   Michelmore, R. W. & Meyers, B. C. Clusters of resistance genes in plants evolve by

divergent selection and a birth-and-death process. *Genome Research* (1998).

44.   Meyers, B. C., Shen, K. A., Rohani, P., Gaut, B. S. & Michelmore, R. W. Receptor-

like genes in the major resistance locus of lettuce are subject to divergent selection.

Plant Cell (1998).

45.   Noel, L. Pronounced Intraspecific Haplotype Divergence at the RPP5 Complex

Disease Resistance Locus of Arabidopsis. *Plant Cell Online* **11**, 2099–2112 (1999).

46.   McDowell, J. M. *et al.* Intragenic recombination and diversifying selection contribute

to the evolution of downy mildew resistance at the RPP8 locus of arabidopsis. *Plant*

Cell (1998).

47. Botella, M. A. *et al.* Three genes of the arabidopsis RPP1 complex resistance locus
recognize distinct Peronospora parasitica avirulence determinants. *Plant Cell* (1998).

48. Barragan, C. A. *et al.* RPW8/HR repeats control NLR activation in Arabidopsis
thaliana. *PLOS Genet.* (2019).

49. Bakker, E. G. A Genome-Wide Survey of R Gene Polymorphisms in Arabidopsis.
Plant Cell Online **18**, 1803–1818 (2006).

50. Guo, Y.-L. *et al.* Genome-Wide Comparison of Nucleotide-Binding Site-Leucine-Rich
Repeat-Encoding Genes in Arabidopsis. *Plant Physiol.* **157**, 757–769 (2011).

51. Wilson, D. S., Gleeson, S. K. & Bell, G. *The Masterpiece of Nature: The Evolution*
and Genetics of Sexuality. Evolution (1983). doi:10.2307/2408354

52. Singh, N. D. *et al.* Fruit flies diversify their offspring in response to parasite infection.
Science (80-.). (2015).

53. Dangl, J. L. & Jones, J. D. G. Plant pathogens and integrated defence responses to
infection. *Nature* (2001).

54. Boller, T. & He, S. Y. Innate immunity in plants: An arms race between pattern
recognition receptors in plants and effectors in microbial pathogens. *Science* (2009).

55. Kondrashov, F. A. Gene duplication as a mechanism of genomic adaptation to a
changing environment. *Proceedings of the Royal Society B: Biological Sciences*
(2012).

56. Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer
weighting and repeat separation. *Genome Res.* (2017).

57. Xiao, C. Le *et al.* MECAT: Fast mapping, error correction, and de novo assembly for
single-molecule sequencing reads. *Nat. Methods* **14**, 1072–1074 (2017).

58. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754–1760 (2009).

59. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078–2079 (2009).

60. Marçais, G. *et al.* MUMmer4: A fast and versatile genome alignment system. *PLoS*
Comput. Biol. (2018).

61. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local
alignment search tool. *J. Mol. Biol.* (1990).

62. Heslop-Harrison, J. S., Murata, M., Ogura, Y., Schwarzacher, T. & Motoyoshi, F.
Polymorphisms and Genomic Organization of Repetitive DNA from Centromeric
Regions of Arabidopsis Chromosomes. *Plant Cell* **11**, 31 LP – 42 (1999).

63. Richards, E. J. & Ausubel, F. M. Isolation of a higher eukaryotic telomere from
Arabidopsis thaliana. *Cell* (1988).

64. Nawrocki, E. P. & Eddy, S. R. Infernal 1.1: 100-fold faster RNA homology searches.

Bioinformatics (2013).

65.   Simon, M. *et al.* Quantitative trait loci mapping in five new large recombinant inbred

line populations of *Arabidopsis thaliana* genotyped with consensus single-nucleotide

polymorphism markers. *Genetics* (2008).

66.   Singer, T. *et al.* A high-resolution map of Arabidopsis recombinant inbred lines by

whole-genome exon array hybridization. *PLoS Genet.* (2006).

67.   Giraut, L. *et al.* Genome-wide crossover distribution in Arabidopsis thaliana meiosis

reveals sex-specific patterns along chromosomes. *PLoS Genet.* (2011).

68.   Stanke, M. & Waack, S. Gene prediction with a hidden Markov model and a new

intron submodel. *Bioinformatics* **19**, (2003).

69.   Majoros, W. H., Pertea, M. & Salzberg, S. L. TigrScan and GlimmerHMM: Two open

source ab initio eukaryotic gene-finders. *Bioinformatics* (2004).

70.   Korf, I. Gene finding in novel genomes. *BMC Bioinformatics* (2004).

71.   Slater, G. S. C. & Birney, E. Automated generation of heuristics for biological

sequence comparison. *BMC Bioinformatics* **6**, 31 (2005).

72.   Kim, D., Langmead, B. & Salzberg, S. L. HISAT: A fast spliced aligner with low

memory requirements. *Nat. Methods* (2015).

73.   Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from

RNA-seq reads. *Nat. Biotechnol.* (2015).

74.   Haas, B. J. *et al.* Automated eukaryotic gene structure annotation using

EVidenceModeler and the Program to Assemble Spliced Alignments. *Genome Biol.* **9**,

R7 (2008).

75.   Keller, O., Odronitz, F., Stanke, M., Kollmar, M. & Waack, S. Scipio: Using protein

sequences to determine the precise exon/intron structures of genes and their

orthologs in closely related species. *BMC Bioinformatics* **9**, 1–12 (2008).

76.   Kalvari, I. *et al.* Rfam 13.0: Shifting to a genome-centric resource for non-coding RNA

families. *Nucleic Acids Res.* (2018). doi:10.1093/nar/gkx1038

77.   Li, P. *et al.* RGAugury: A pipeline for genome-wide prediction of resistance gene

analogs (RGAs) in plants. *BMC Genomics* (2016). doi:10.1186/s12864-016-3197-x

78.   Choi, K. *et al.* Recombination Rate Heterogeneity within Arabidopsis Disease

Resistance Genes. *PLoS Genet.* **12**, 1–30 (2016).

79.   Cingolani, P. *et al.* A program for annotating and predicting the effects of single

nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster

strain w1118; iso-2; iso-3. *Fly (Austin).* (2012).

80.   Pfeifer, B., Wittelsbürger, U., Ramos-Onsins, S. E. & Lercher, M. J. PopGenome: An

efficient swiss army knife for population genomic analyses in R. *Mol. Biol. Evol.* **31**,

1929–1936 (2014).

81.   Huang, D. W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis
of large gene lists using DAVID bioinformatics resources. *Nat. Protoc.* (2009).
82.   Huang, D. W., Sherman, B. T. & Lempicki, R. A. Bioinformatics enrichment tools:
Paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids*
Res. (2009).
523

Data availability

Raw sequencing data, assemblies and annotations can be accessed in the European Nucleotide Archive under the project accession number PRJEB31147. Assemblies, annotation, variation and orthologs can be found on the *1001 Arabidopsis thaliana Genomes* webpage (www.1001genomes.org).

Code availability

Custom code used in this study can be found online at *https://github.com/schneebergerlab/AMPRIL-genomes*.

Supplementary Information

Tables S1 to S17

Figs. S1 to S15

Acknowledgements.

The authors would like to thank Beth R. Rowan (UC Davis) for providing the CO breakpoint list prior to publication, Bruno Hüttel (Max Planck Genome center) for support in genome sequencing, Sigi Effgen and Maarten Koornneef (Max Planck Institute for Plant Breeding Research) for providing seeds, Onur Dogan (Max Planck Institute for Plant Breeding Research) for help in the greenhouse, Angela M. Hancock (Max Planck Institute for Plant Breeding Research) for helpful discussions, and Raphael Mercier and Padraic J. Flood (Max Planck Institute for Plant Breeding Research) for helpful comments on the manuscript and the interpretation of HOT regions. **Funding.** K.S. gratefully acknowledges support from European Research Council (ERC) Grant "INTERACT" (802629). **Authors contributions.** W.-B.J. and K.S. designed the study. W.-B.J. performed all analysis. K.S. supervised the study. W.-B.J. and K.S. wrote the manuscript. All authors read and approved the final manuscript.

Authors contributions

W.-B.J. and K.S. designed the study. W.-B.J. performed all analysis. K.S. supervised the study. W.-B.J. and K.S. wrote the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare no competing interests.

555 **Tables**

556 **Table 1. Genome assembly and annotation of seven *A. thaliana* accessions**

	Col-0[*]	An-1	C24	Cvi-0	Eri-1	Kyo	L*er*	Sha
Contigs	-	151	167	140	200	230	149	143
Pseudo-molecules	5	5	5	5	5	5	5	5
Contig N50 (Mbp)	-	8.2	4.8	7.4	4.8	9.1	11.2	7.0
Contig CL50[**]	-	2	2	2	2	2	1	1
Chr. length (Mbp)	119.1	118.4	117.7	118.3	117.7	118.8	118.5	118.4
Genes	27,445	27,342	27,214	27,098	27,285	27,574	27,376	27,293

558 [*] Reference sequence

559 [**] Chromosome number normalized L50 [23]

Figure legends

Fig. 1. Chromosome-level genome assemblies of seven *A. thaliana* accessions
The light grey bars outline each of the chromosomes, whereas the dark gray inlays show the extend of each of the pericentromeric regions. The contig arrangements of the chromosome assemblies is shown in green for contigs > 1 Mb and dark grey for contigs < 1 Mb. The location of centromeric tandem repeat arrays and rDNA clusters within the assemblies are marked by yellow and blue boxes above each of the chromosomes.

Fig. 2. Structural and sequence differences between the genomes
(a) Schematic of the structural differences (upper panel) and sequence variation (lower panel) that can be identified between chromosome-level assemblies. Note, local sequence variation can reside in syntenic as well as in rearranged regions. The barplots on the right upper side show the total span of syntenic and rearranged regions between the reference and each of other accessions (colors match the schematic on the left): The left barplots shows the sequence span in respect to the reference sequence, while the right plot shows the sequence space, which is specific to each of the accessions. The barplots on the right lower side show local sequence variation (per kb) in syntenic (left) and rearranged (right) regions between the reference and each of other accessions (again colors match the schematic on the left). **(b)** Size distributions of different types of structural and sequence variation. **(c)** Gene copy number variations between the reference and each of the accessions. The left barplots shows the fraction of (reference) genes which are in gene families with conserved or variable copy numbers. The right barplots shows the number of non-reference genes found in at least two accessions, or found to be specific to an accession genome. **(d)** Pan-genome and core-genome estimations for sequence (upper plot) and gene space (lower plot) were based on all pairwise whole-genome and gene set comparisons across all eight accessions. Each black point corresponds to a pan- or core-genome size estimated with a particular combination of genomes. Pan-genome (blue) and core-genome (red) estimations were fitted using an exponential model.

Fig. 3. Quantitative analysis of synteny reveals hotspots of rearrangements
(a) *Synteny Diversity* along each chromosome: (100kb sliding windows with a step-size of 50kb in blue; 5kb sliding windows with a step-size of 1kb in grey). The red bars under the x-axes indicate the location of R gene clusters. Gray rectangles indicate the location of centromeric regions. The dashed green and red lines indicate thresholds for Synteny Diversity values of 0.25 and 0.50 indicative for the segregation of two (0.25) or three (0.50) non-syntenic haplotypes. The arrow labelled with "(A)" indicates a 2.48 Mb inversion in the

Sha genome. The arrow labelled with "(B)" indicates the location of the example show in (b).

(b) Gene and TE densities in syntenic (SYN) and hotspots of rearrangements (HOT)

regions. **(c)** The number of variable copy-number alleles in syntenic (SYN) and

rearrangements (HOT) regions**. (d)** An example of a HOT region which includes the

RPP4/RPP5 R gene cluster. The upper panel shows the distribution of *Synteny Diversity*

(blue curve) nucleotide diversity (gray background) and haplotype diversity (pink

background) in a 5kb sliding window with a step-size of 1kb. Both the nucleotide diversity

and the haplotype diversity were calculated based on informative markers (MAF >= 0.05,

missing rate < 0.2) from the 1001 Genomes Project [12]. The marker density is shown as the

heatmap on top. The green and red dashed lines indicate the value 0.25 and 0.50 of

synteny diversity, respectively. The schematic in the lower part shows the protein-coding

genes (colored rectangles) annotated in each of the eight genomes. In blue, genes without

function implicated in disease resistance. Rectangles with other colors represent resistance

genes, where genes with the same color belong to the same gene family. The gray links

between the rectangles indicate the homologous relationships between non-resistance

genes. The red lines indicate the positions of HOT regions**. (e)** A dot plot of Col-0 and C24

sequence from the HOT region shown in (D). Red lines indicate regions with homology

between the two genomes. **(f)** The distribution of *Synteny Diversity* values in 1kb sliding

windows around and in HOT regions.

Fig. 4. Two examples for hotspots of rearrangements

Visualization of **(a)** the *DM6* locus (*RPP7*) and **(b)** an unnamed R gene cluster on

chromosome 5. Descriptions for the plots can be found in the legend of Fig. 3d.

Fig. 5. The causes and consequences of hotspots of rearrangements

(a) Crossover (CO) breakpoints [37,38] identified in Col-0 x L*er* hybrids were checked for their

overlaps in syntenic or rearranged regions. Only unique CO intervals smaller than 5kb were

used. Chi-square test was applied. **(b)** Linkage disequilibrium (LD) calculated in 4kb

windows in and around each of the HOT regions as shown in the lower part. LD was

calculated as the correlation coefficient (r^2) based on informative SNP markers (MAF > 0.05,

missing rate < 0.2) selected from the 1001 Genomes Project data [12]. **(c)** Minor allele

frequency of SNP markers in syntenic, partially syntenic, and HOT regions. The SNP

markers (MAF >0.005, missing rate < 0.2) from 1001 Genomes Project were used. **(d)**

Frequency of deleterious mutations in syntenic regions and HOT regions. Deleterious

mutations include SNPs and small indels that introduce premature stop codons, loss of start

or stop codons, frameshifts, splicing sites mutations or deletions of exons. (**e**) GO term

enrichment analysis of protein-coding genes in HOT regions (P-values cut-off = 0.05).

a

Structural differences

Syntenic | Inversion | Translocation | Duplication loss

Reference

Accession

Genome-specific

Duplication gain

Matching the reference

Sha
Ler
Kyo
Eri-1
Cvi-0
C24
An-1

0   100   110   120

Accession specific

0   4   8   12
Mb

Sequence differences

SNP

Deletion | Highly divergent region | Copy loss

Insertion | Copy gain

Syntenic regions

Sha
Ler
Kyo
Eri-1
Cvi-0
C24
An-1

0   40

Rearranged regions

0   40   80   120   160
bp / kb

b

Inversion
Transloc.
Duplication
Deletion
Insertion

100   1,000   10,000   100,000   1,000,000

c

Reference genes

Sha
Ler
Kyo
Eri-1
Cvi-0
C24
An-1

0   24,000   25,000   26,000   27,000

Non-ref. genes

0   200   400   600

Conserved copy number
Copy-number decrease
Copy-number increase
Reference-specific

Shared by at least 2 accessions
Accession-specific

d

Pan-genome
Core-genome

130 Mb
120 Mb
110 Mb
100 Mb

Genome size (Mb)

Pan-genome size (Gene number)

30000
28000
26000
24000

0   2   4   6   8   10   12
Number of genomes