Heart Rate n-Variability (HRnV) and Its Application to Risk Stratification of Chest Pain Patients in the Emergency Department

Nan Liu, PhD[1,2]*

Dagang Guo, PhD[3]

Zhi Xiong Koh, BEng[4]

Andrew Fu Wah Ho, MBBS[1,3,5]

Feng Xie, BSc[1]

Takashi Tagami, MD, PhD[6]

Jeffrey Tadashi Sakamoto, MD[7]

Pin Pin Pek, MPH[1,4]

Bibhas Chakraborty, PhD[1]

Swee Han Lim, MBBS[4]

Jack Wei Chieh Tan, MBBS, MMed[8]

Marcus Eng Hock Ong, MBBS, MPH[1,2,4]

[1] Duke-NUS Medical School, National University of Singapore, Singapore

[2] Health Services Research Centre, Singapore Health Services, Singapore

[3] SingHealth Duke-NUS Emergency Medicine Academic Clinical Programme, Singapore

[4] Department of Emergency Medicine, Singapore General Hospital, Singapore

[5] National Heart Research Institute Singapore, National Heart Centre, Singapore

[6] Department of Emergency and Critical Care Medicine, Nippon Medical School Musashikosugi Hospital, Tokyo, Japan

[7] School of Medicine, Stanford University, Stanford, CA, USA

[8] National Heart Centre Singapore, Singapore

* **Corresponding Author**

Nan Liu

Centre for Quantitative Medicine and Programme in Health Services and Systems Research, Duke-NUS Medical School, Academia, 20 College Road, Singapore 169856

Email: liu.nan@duke-nus.edu.sg

Abstract

Background: Chest pain is one of the most common complaints among patients presenting to the emergency department (ED). Causes of chest pain can be benign or life threatening, making accurate risk stratification a critical issue in the ED. In addition to the use of established clinical scores, prior studies have attempted to create predictive models with heart rate variability (HRV). In this study, we proposed heart rate n-variability (HRnV), an alternative representation of beat-to-beat variation in electrocardiogram (ECG) and investigated its association with major adverse cardiac events (MACE) for ED patients with chest pain.

Methods: We conducted a retrospective analysis of data collected from the ED of a tertiary hospital in Singapore between September 2010 and July 2015. Patients >20 years old who presented to the ED with chief complaint of chest pain were conveniently recruited. Five to six-minute single-lead ECGs, demographics, medical history, troponin, and other required variables were collected. We developed the HRnV-Calc software to calculate HRnV parameters. The primary outcome was 30-day MACE, which included all-cause death, acute myocardial infarction, and revascularization. Univariable and multivariable logistic regression analyses were conducted to investigate the association between individual risk factors and the outcome. Receiver operating characteristic (ROC) analysis was performed to compare the HRnV model (based on leave-one-out cross-validation) against other clinical scores in predicting 30-day MACE.

Results: A total of 795 patients were included in the analysis, of which 247 (31%) had MACE within 30 days. The MACE group was older and had a higher proportion of male patients. Twenty-one conventional HRV and 115 HRnV parameters were calculated. In univariable analysis, eleven HRV parameters and 48 HRnV parameters were significantly associated with 30-day MACE. The multivariable stepwise logistic regression identified 16 predictors that were strongly associated with the MACE outcome; these predictors consisted of one HRV, seven HRnV parameters, troponin, ST segment changes, and several other factors. The HRnV model outperformed several clinical scores in the ROC analysis.

Conclusions: The novel HRnV representation demonstrated its value of augmenting HRV and traditional risk factors in designing a robust risk stratification tool for patients with chest pain at the ED.

Keywords: Heart rate variability (HRV), heart rate n-variability (HRnV), electrocardiogram, chest pain, risk stratification, emergency department.

Background

Chest pain is one of the most common presenting complaints in the emergency department (ED) [1, 2], which may be due to life-threatening myocardial infarction (MI) or benign musculoskeletal pain [3]. Majority of chest pain patients are subjected to extensive diagnostic tests to rule out acute coronary syndrome (ACS), resulting in oftentimes, unnecessary prolonged and costly ED admission, since only a small proportion of these patients will eventually receive a diagnosis of ACS [3]. Hence, early identification of chest pain patients at high-risk of developing adverse cardiac events has been a pressing issue to contend with in the ED. Several established clinical scores have been used for risk stratifying chest pain patients in the ED [4, 5], including the History, ECG, Age, Risk factors and Troponin (HEART) [6], the Thrombolysis in Myocardial Infarction (TIMI) [7], and the Global Registry of Acute Coronary Events (GRACE) [8] scores. Of the available clinical scores, the HEART score is the most accurate and widely used [5, 9-12]. Further research has focused on developing risk score-based clinical pathways for safe discharge of low-risk patients [1, 3, 13, 14].

In a recent review of clinical scores for ED patients with chest pain [5], heart rate variability (HRV) provided an alternative approach to build predictive models for accurate risk stratification [15-17]. HRV is a widely adopted tool for evaluating changes in cardiac autonomic regulation, and has been shown to be strongly associated with the autonomic nervous system (ANS) [18-20]. HRV analysis characterizes the beat-to-beat variation in an electrocardiogram (ECG) by utilizing time domain, frequency domain, and nonlinear analyses [19]. Reduced HRV has been found to be a significant predictor of adverse cardiac outcomes [21]. Given the complexity of quantifying HRV representation, several tools such as the PhysioNet Cardiovascular Signal Toolbox [22] and Kubios HRV [23] have been developed to standardize HRV analyses.

Based on the principle of parameter calculation on normal R-R intervals (RRIs; in this paper, RRIs are equivalent to normal-to-normal [NN] intervals, in which abnormal beats have been

removed), HRV analysis generates only one set of parameters from a fixed length of ECG record. This limits the amount of information that can be extracted from raw ECG signals. In this paper, we proposed a novel representation of beat-to-beat variation, named as heart rate n-variability (HRnV) [24] to characterize RRIs from a different perspective. With the use of HRnV measures, multiple sets of parameters are calculated from the same ECG record, which significantly increases the amount of extracted information. Our study is the first clinical application of the HRnV representation, in which the value of this novel measure is evaluated in risk stratification of chest pain patients in the ED. We hypothesize that HRnV is closely related to conventional HRV while providing supplementary information that is associated with adverse cardiac events. Further, we will investigate the potential use of HRnV parameters to develop risk prediction tools.

Methods

Study Design and Setting

We conducted a retrospective analysis of data collected in our previous study on risk stratification of chest pain patients in the ED [9]. A convenience sample of patients was recruited at the ED of Singapore General Hospital between September 2010 and July 2015. This is a tertiary hospital with around-the-clock primary percutaneous coronary intervention capabilities and a median door-to-balloon time of 101 minutes [25]. At ED triage, patients are assigned a Patient Acuity Category Scale (PACS) score ranging from 1 to 4 to indicate the urgency for consultation; smaller number means higher acuity. In this study, patients >20 years old who presented to the ED with chief complaint of chest pain and were triaged to PACS 1 or 2 were included. Patients were excluded if they had ST-elevation myocardial infarction (STEMI) or an obvious non-cardiac etiology of chest pain diagnosed by the primary emergency physician. Patients were also excluded if their ECGs had high percentage of noise or they were in non-sinus rhythm; these criteria were applied to ensure the quality of HRV and HRnV analyses. The ethical approval was obtained from the Centralized Institutional Review Board (CIRB, Ref: 2014/584/C) of SingHealth, the largest public healthcare system in Singapore that includes the Singapore General Hospital as a key partner. In the ethical approval, patient consent was waived.

Data Collection

During the data collection period, five to six-minute single-lead (lead II) ECG tracings were retrieved from the X-Series Monitor (ZOLL Medical Corporation, Chelmsford, MA). The first set of vital signs and troponin values from the recruited patients were extracted from the hospital's electronic health records (EHR). In this study, high-sensitivity troponin-T was used, and an abnormal value was defined as >0.03 ng/mL [26]; it was further stratified into three groups and coded as 0 if the value was ≤0.03 ng/mL, 1 if the value was between 1 and 3 times the normal limit, and 2 if the value was >3 times the normal limit. Additionally, the patients' first 12-lead ECGs were interpreted by two independent clinical reviewers. Pathologic ST-elevation, ST-depression, T-wave inversions, and Q-waves were recorded. Patient demographics, medical history, and information required for computing the HEART, TIMI, and GRACE scores were retrospectively reviewed and obtained from EHR.

Proposed HRnV Representation of Beat-to-Beat Variation in ECG

<u>HR_nV: A Novel Measure with Non-Overlapped RRIs</u>
Prior to introducing the new HR_nV measure, we define a new type of RRI called RR_nI, where $1 \leq n \leq N$, and $N \ll \hat{N}$. $\hat{N}$ is the total number of RRIs. The definition of RR_nI is illustrated in Figure 1a. When $n = 1$, RR_nI becomes conventional RRI, that is, RR_1I is equal to RRI. When $n > 1$, every n adjacent RRI is connected to form a new sequence of RR_nIs. By using this strategy, we can create a maximum number of $(N - 1)$ new RR_nI sequences from conventional single RRI sequence. With these newly generated RR_nI sequences, the calculation of HR_nV parameters is straightforward and can be accomplished by applying established quantitative methods including time domain analysis, frequency domain analysis, and nonlinear analysis [18, 19]. In describing this new measure, we use the term "HR_nV" prior to parameter names to indicate that these parameters are calculated from RR_nI sequences. As noted in the above, HR_nV is a novel measure based on newly generated, non-overlapped RR_nIs. The computed HR_nV parameters include but are not limited to the following: the average of RR_nIs (HR_nV mean NN), standard deviation of RR_nIs (HR_nV SDNN), square root of the mean squared differences between RR_nIs (HR_nV RMSSD), the number of times that the absolute difference between two successive RR_nIs exceeds 50 ms (HR_nV NN50), HR_nV NN50 divided by the total number of RR_nIs (HR_nV pNN50), the integral of the RR_nI histogram divided by the height of the histogram (HR_nV triangular index), low frequency power (HR_nV LF power), high frequency power (HR_nV HF power),

approximate entropy (HR$_n$V ApEn), sample entropy (HR$_n$V SampEn), and detrended fluctuation analysis (HR$_n$V DFA), among others. Notably, two new parameters NN50n and pNN50n are created, where 50×n ms is set as the threshold to assess the difference between pairs of consecutive RR$_n$Is.

<u>HR$_n$V$_m$: A Novel Measure with Overlapped RRIs</u>

Like RR$_n$I that is used in HR$_n$V, to define the HR$_n$V$_m$ measure we introduce another type of RRI called RR$_n$I$_m$, where $1 \leq n \leq N$, $1 \leq m \leq N - 1$, and $N \ll \widehat{N}$. In the RR$_n$I$_m$ sequence, m is used to indicate the level of overlap between consecutive RR$_n$I$_m$ sequences. As illustrated in Figure 1b, $(n - m)$ RRIs form the overlapped portions. When $m = n$, RR$_n$I$_m$ becomes RR$_n$I; therefore, the upper limit of m is $N - 1$. By controlling the overlap among these newly generated RR$_n$I$_m$ sequences, we can create a maximum number of $(N \times (N - 1)/2)$ RR$_n$I$_m$ sequences (excluding the RR$_n$I sequence) from conventional single RRI sequence. For each of the newly created RR$_n$I$_m$ sequences, we apply time domain analysis, frequency domain analysis, and nonlinear analysis to calculate HR$_n$V$_m$ parameters. We add the term "HR$_n$V$_m$" prior to the parameters to denote that they are computed from RR$_n$I$_m$ sequences. For example, the average RR$_n$I$_m$ intervals and the sample entropy are written as HR$_n$V$_m$ mean NN and HR$_n$V$_m$ SampEn, respectively. The HR$_n$V$_m$ measure extracts additional information than HR$_n$V does, by adopting a strategy of controlling sequence overlap.

HRnV Analysis and Parameter Calculation

We developed the HRnV-Calc software suite (https://github.com/HRnV) to calculate HRnV parameters. The HRnV-Calc software integrates functions from the PhysioNet Cardiovascular Signal Toolbox [22] to perform standardized ECG signal processing and QRS complex detection. Given the short ECG records in this study, the upper limit of n was set as three; thus, six sets of parameters were calculated, namely HRV, HR$_2$V, HR$_2$V$_1$, HR$_3$V, HR$_3$V$_1$, and HR$_3$V$_2$.

Clinical Outcomes

The primary endpoint in this study was a composite outcome of major adverse cardiac events (MACE) [27], including all-cause death, acute myocardial infarction (AMI), and revascularization (coronary artery bypass graft [CABG] or percutaneous coronary intervention [PCI]) within 30 days of ED presentation.

Statistical Analysis

Continuous variables were presented in terms of mean and standard deviation and compared between the two categories of the primary outcome (MACE) using two-sample t-test. Categorical variables were presented in terms of frequency and percentage and compared between the two categories of the primary outcome (MACE) using chi-square test. A statistically significant difference was defined as $p<0.05$. To evaluate the HRnV parameters and other risk factors, we conducted univariable and multivariable analyses and subsequently developed simple prediction models using traditional logistic regression. In building the HRnV prediction model, we selected candidate variables with $p<0.2$ in the univariable analysis and fed them into multivariable stepwise logistic regression. To evaluate the predictive performance, we used leave-one-out cross-validation (LOOCV) to conduct the analysis.

The receiver operating characteristic (ROC) analysis [28] was performed to compare prediction performances among the HRnV model, HEART, TIMI and GRACE scores. The area under the ROC curve (AUC), sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV) were reported as the predictive measures. Data preparation, descriptive analysis, and predictive model development were performed in R version 3.6.0 (R Foundation, Vienna, Austria), and the ROC analysis was conducted in MATLAB R2019a (MathWorks, Natick, MA).

Results

A total of 795 patients were selected from the originally recruited 922 patients [9]. 28 were excluded for ECG recording issues, four were excluded for clearly non-cardiac chest pain, and 95 were excluded for irregular rhythm/artifacts. Among the included 795 patients, 247 (31%) met the primary outcome of 30-day MACE. Table 1 shows patient baseline characteristics. Patients with the primary outcome were older (mean age 61 years vs. 59 years, p=0.035) and had a higher proportion of males (76.1% vs. 64.6%, p=0.002). There were no statistically significant differences between MACE and non-MACE groups in terms of patient ethnicity. Factors such as history of diabetes and current smoking status were significantly more prevalent in the group with MACE.

Descriptive analyses of HRV and HRnV parameters are tabulated in Table 2. In this clinical case study, N was set as 3, thus HR_2V, HR_2V_1, HR_3V, HR_3V_1 and HR_3V_2 parameters were calculated. Among time domain parameters such as mean NN, SDNN and RMSSD, the HR_nV and HR_nV_m values were generally incremental with an increase in n. Notably, HR_2V NN50 and HR_3V NN50 were much lower than conventional HRV NN50. Moreover, NN50n and pNN50n are parameters specifically applicable to the HRnV representation. Like time domain parameters, the same trend of changes in frequency domain parameters were observed. The magnitude of increment in VLF power and LF power was larger than that of HF power with increasing n. However, one exception was the normalized HF power, whose HR_nV and HR_nV_m parameters were smaller than that of HRV. In nonlinear analysis, the differences in Poincaré SD2 values were obvious between HRV and HRnV parameters. HR_2V SampEn and HR_3V SampEn were remarkably larger compared to SampEn parameters of HRV, HR_2V_1, HR_3V_1, and HR_3V_2, as the confidence interval of SampEn was wide when data points were less than 200 [19], since our ECG recordings were only five to six-minute long. HR_2V_1, HR_3V_1 and HR_3V_2 were free from this issue as they were calculated from overlapping RR_nI_m sequences where more than 200 data points were available.

Table 3 presents the results of univariable analyses of HR_nV and HR_nV_m parameters. Eleven out of 21 conventional HRV parameters were statistically significant. Additionally, 13 HR_2V, six HR_3V, 11 HR_2V_1, seven HR_3V_1 and 11 HR_3V_2 parameters were also significant. Overall, additional 115 HRnV parameters were derived, among which 48 showed statistical significance between patients who had 30-day MACE and who did not. Among all HRV and HRnV parameters, mean NN, SDNN, RMSSD, NN50, pNN50, HF power, Poincaré SD1 and SD2 appeared statistically significant in at least five out of six measures (i.e., HRV, HR_2V, HR_2V_1, HR_3V, HR_3V_1, and HR_3V_2). Furthermore, skewness, LF power, SampEn, and ApEn, which did not demonstrate statistical significance in conventional HRV analysis, became statistically significant in HRnV representation. Table 4 presents the results of multivariable analyses on HR_nV and HR_nV_m parameters by adjusting for age and sex. Compared with the results in Table 3, several parameters became statistically insignificant, e.g. NN50 of HR_3V and HR_3V_2, and triangular index of HRV, HR_2V, and HR_3V_2. On the other hand, ApEn of HR_2V, HR_2V_1, and HR_3V_2 became statistically significant after the adjustment.

Table 5 lists the 16 variables that were selected through multivariable stepwise logistic regression, among which there were one conventional HRV parameter and seven HRnV parameters. In addition to traditional predictors of adverse cardiac outcomes such as ST segment changes and troponin, HR_2V ApEn (OR=0.095; 95% CI 0.014-0.628), HR_2V_1 ApEn (OR=19.700; 95% CI 2.942-131.900) and HR_3V skewness (1.560; 95% CI 1.116-2.181) also demonstrated strong predictive power in assessing the risk of 30-day MACE. Table 6 presents the results of ROC analysis in evaluating the predictive performance by the HRnV model (based on LOOCV), HEART, TIMI, and GRACE scores. The HRnV model achieved the highest AUC value and outperformed HEART, TIMI, and GRACE scores in terms of specificity, PPV, and NPV at the optimal cut-off scores, defined as the points nearest to the upper-left corner of the ROC curves.

Discussion

HRV has been well investigated in the past decades [18, 19, 29]. While the majority of efforts in HRV research were focused on development of advanced nonlinear techniques to derive novel parameters [30, 31], few investigated alternative approaches to analyze RRIs. Vollmer [32] used relative RRIs to describe the relative variation of consecutive RRIs, with which HRV was analyzed. Likewise, we proposed a novel HRnV representation, providing additional HRnV parameters than conventional HRV analysis. In this paper, we introduced two measures of HRnV, namely HR_nV and HR_nV_m. HR_nV is calculated based on non-overlapped RR_nI sequences, while HR_nV_m is computed from overlapped RR_nI_m sequences. HRnV is not developed to replace the conventional HRV; instead, this representation is a natural extension of HRV. It enables the creation of additional parameters from raw ECGs, and thus empowers the extraction of supplementary information.

In our clinical case study, we investigated the predictive value of HRnV parameters to assess the risk of 30-day MACE for chest pain patients in the ED. In addition to 21 HRV parameters, 115 HRnV parameters were derived, of which 48 were found to be statistically significant in their association with the primary outcome. Notably, even with a small n (three in our study), newly generated HRnV parameters have greatly boosted the number of candidate predictors. When longer ECG records are available, more HRnV parameters can be calculated. With HRnV parameters, HRV parameters, vital signs, and several established risk factors, we

conducted multivariable logistic regression analysis and selected age, diastolic BP, pain score, ST-elevation, ST-depression, Q wave, cardiac history, troponin, HRV NN50, and seven HRnV parameters. In addition to traditional risk factors like ST segment changes, HR_2V ApEn, HR_2V_1 ApEn, and HR_3V skewness were found as strong predictors for 30-day MACE. Comparing with the HEART, TIMI, and GRACE scores, the HRnV model achieved the highest AUC, specificity, PPV, and NPV values at the optimal cut-off points in ROC analysis. Therefore, HRnV may be clinically useful in determining the risk of 30-day MACE for ED patients with chest pain.

Due to the wide differential diagnosis for chest pain, accurate stratification is vital, particularly for preventing low-risk patients from obtaining expensive and unnecessary medical testing and intervention [3]. The TIMI and GRACE scores have been validated for risk prediction of patients with chest pain in the ED [4, 33, 34]. However, because they were originally developed for post-acute myocardial infarction, certain criteria within the TIMI and GRACE scores may be inappropriate for the undifferentiated chest pain cohorts in the ED [1]. In comparison, the HEART score was derived from a population of ED patients with chest pain, and has been extensively validated worldwide [10, 13, 27, 35]. It has proven its applicability in identifying both low-risk patients for possible early discharge and high-risk patients for urgent intervention. Built upon established scores, many chest pain pathways [14, 36-38] have been implemented and tested, particularly for the management of low-risk patients. Than et al. [38] evaluated a TIMI score-based accelerated diagnostic protocol (ADP) with a reported sensitivity of 99.3% and NPV of 99.1%. Similarly, a systematic review by Laureano-Phillips et al. [39] reported that the HEART score achieved both sensitivity and NPV of 100% in several validation studies. Furthermore, a cost-effectiveness study conducted in Brisbane, Australia reported economic benefits by adopting an ADP in the ED, with its associated reduction in expected cost and length of stay amongst patients with chest pain [40].

Most established clinical scores use conventional risk factors such as biomarkers, medical history, and presenting vital signs. However, patient history can sometimes be subjective and blood tests, such as troponin, take time to result. HRV, as a noninvasive measure, can be easily calculated from ECGs; it is an objective tool to assess the activities of the ANS [19]. It also has the advantage of requiring only several minutes to acquire (five to six minutes in our protocol), which is much faster than serum biomarkers. Over the past decades, HRV has been

widely investigated in a broad range of clinical applications, particularly in cardiovascular research. HRV was found to be closely associated with sudden cardiac death [18]. It also showed significant correlations with adverse clinical outcomes in prehospital setting [41] and with MACE outcomes in ED patients with chest pain [17]. HRV parameters have been integrated with other risk factors into machine learning algorithms to predict adverse outcomes [42, 43]. These promising results motivated the use of HRV to develop objective and computerized risk stratification tools for chest pain patients [44, 45]. In an updated review of clinical scores for chest pain, Liu et al. [5] summarized several studies which aimed to develop alternative techniques for risk stratification.

This study aimed to present the novel HRnV representation and its measures and investigate their association with clinical outcomes. Although the HRnV parameters showed promising performance in identifying high-risk chest pain patients, this study was not intended to create a ready-to-use clinical tool. Instead, we have demonstrated the feasibility of utilizing HRnV parameters to augment conventional HRV and risk factors in designing a prediction tool/score. These HRnV parameters can be readily calculated without the collection of supplementary data. In this study, with five to six-minute ECG and $n = 3$, five-fold more HRnV parameters were calculated compared to HRV alone. When longer ECGs are available and parameter n is set as a larger number, more HRnV parameters can be derived. To build a HRnV-based risk stratification tool, a systematic approach is needed to derive a point-based, consistent score to ease its clinical application and practical implementation.

As a natural extension of conventional HRV, HRnV representation creates the opportunity to generate additional parameters. This representation could also serve as a smoother for RRIs, making them less sensitive to sudden changes caused by abnormal heart beats (e.g., very short or very long RRI). However, since HRnV is a novel representation of beat-to-beat variations in ECG, many technical issues need to be addressed in future research. For instance, as shown in Table 2, SampEn became larger when the available number of data points was less than 200 [19], suggesting that additional research is required to investigate its applicability to short ECG records. Moreover, parameters NN50n and pNN50n are newly introduced in HRnV representation only. They characterize the number of times that the absolute difference between two successive RR$_n$I sequences exceeds 50×n ms, by assuming that the absolute difference may be magnified when the corresponding RR$_n$I is n times longer

than RRI. Thus, in-depth investigations are required in the selection of appropriate thresholds. More importantly, physiological interpretations of the HRnV parameters and their normal values [29] need to be determined through numerous research. One example is the identification of frequency bands that correlate with certain physiological phenomenon. In the current analysis, the conventional cut-off values were adopted (i.e., ≤0.04 Hz as very low frequency range; 0.04–0.15 Hz as low frequency range; 0.15–0.4 Hz as high frequency range). With the increase in n, frequency domain analysis may need to be changed accordingly.

Beyond its use in risk stratification of ED patients with chest pain, HRnV foresees potentials in many clinical domains where conventional HRV has been extensively investigated [46-49]. With the augmented RR_nI and RR_nI_m sequences, HRnV could possibly capture more dynamic changes in cardiac rhythms than HRV. This capability enables the extraction of extra information from limited raw ECGs. This study utilized HRnV parameters as independent risk factors and analyzed them with traditional biostatistical methods. There are multiple ways to use HRnV parameters, e.g., each set of HRnV parameters can be analyzed individually and are subsequently combined with an ensemble learning [50] (a special type of machine learning algorithm) architecture to reach a decision. However, artificial intelligence and machine learning methods generally create black-box predictive models, making interpretation a challenge [51].

Limitations

This study has several limitations. First, no scoring tool was developed for practical clinical use. The primary aim of this study was to demonstrate the feasibility of using HRnV parameters and common risk factors to build predictive models. Second, the HRnV model was evaluated with LOOCV strategy due to small sample size. Ideally, separate patient cohorts are needed to train and test prediction models. When a new scoring tool is developed, it is necessary to conduct external validations on cohorts with diverse patient characteristics. Furthermore, properly designed clinical pathways are needed as well. Third, the patients included in this study were mainly from the high acuity group, resulting in a higher 30-day MACE rate (i.e., 31%) compared to other similar studies [10, 39]. As a result, the generalizability of the HRnV model developed in this study may be uncertain in other patient cohorts. Fourth, the calculated HRnV and HRV parameters depended on the choice of tools and methods for ECG signal analysis. Thus, the values of these parameters may vary across studies. Lastly, the physiology of HRnV and interpretations of its measures are mostly

unknown; calculation of some parameters also needs to be standardized. All these require future collaborative efforts between clinicians and scientists to address.

Conclusions

In this study, we proposed a novel HRnV representation and investigated the use of HRnV and established risk factors to develop a predictive model for risk stratification of patients with chest pain in the ED. Multiple HRnV parameters were found as statistically significant predictors, which effectively augmented conventional HRV, vital signs, troponin, and cardiac risk factors in building an effective model with good discrimination performance. The HRnV model outperformed the HEART, TIMI, and GRACE scores in the ROC analysis. It also demonstrated its capability in identifying low-risk patients, which may enable its use in building a new clinical pathway. Moving forward, we suggest further development of a point-based, ready-to-use HRnV risk stratification tool. Although some issues remain to be addressed, we hope to stimulate a new stream of research on HRnV. We believe that future endeavors in this field will lead to the possibility of in-depth evaluation of the associations between HRnV measures and various human diseases.

List of abbreviations

ACS, acute coronary syndrome

ADP, accelerated diagnostic protocol

AMI, acute myocardial infarction

ANS, autonomic nervous system

AUC, area under the curve

ApEn, approximate entropy

CABG, coronary artery bypass graft

DFA, detrended fluctuation analysis

ED, emergency department

EHR, electronic health records

GRACE, Global Registry of Acute Coronary Events

HEART, History, ECG, Age, Risk factors and Troponin

HF, high frequency

HRnV, Heart Rate n-Variability

HRV, heart rate variability

LF, low frequency

LOOCV, leave-one-out cross-validation

MACE, major adverse cardiac events

Mean NN, average of R-R intervals

MI, myocardial infarction

NN50, the number of times that the absolute difference between 2 successive R-R intervals exceeds 50 ms

NN50n, the number of times that the absolute difference between 2 successive RRnI/RRnIm sequences exceeds 50×n ms

NPV, negative predictive value

PACS, Patient Acuity Category Scale

PCI, percutaneous coronary intervention

pNN50, NN50 divided by the total number of R-R intervals

pNN50n, NN50n divided by the total number of RRnI/RRnIm sequences

PPV, positive predictive value

RMSSD, square root of the mean squared differences between R-R intervals

ROC, receiver operating characteristic

RRI, R-R interval

SampEn, sample entropy

SD, standard deviation

SDNN, standard deviation of R-R intervals

STEMI, ST-elevation myocardial infarction

TIMI, Thrombolysis in Myocardial Infarction

VLF, very low frequency

Declarations

Ethics approval and consent to participate

The ethical approval was obtained from the Centralized Institutional Review Board (CIRB, Ref: 2014/584/C) of SingHealth, in which patient consent was waived.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

NL and MEHO hold patents related to using heart rate variability and artificial intelligence for medical monitoring. NL, DG, ZXK, and MEHO are currently advisers to TIIM SG. The other authors report no conflicts.

Funding

This work was supported by the SHF Foundation Research Grant (SHF/FG652P/2017).

Authors' contributions

NL invented the HRnV representation, conceived the study, supervised the project, and wrote the first draft of the manuscript. NL, DG, ZXK, and FX performed the analyses. All authors contributed to evaluation of the HRnV measures, interpretation of the results, and revision of the manuscript. All authors approved the final manuscript.

Acknowledgements

We would like to thank and acknowledge the contributions of doctors, nurses, and clinical research coordinators from the Department of Emergency Medicine, Singapore General Hospital.

References

1.   Long B, Koyfman A: **Best Clinical Practice: Current Controversies in the Evaluation of Low-Risk Chest Pain with Risk Stratification Aids. Part 2**. *The Journal of emergency medicine* 2017, **52**(1):43-51.

2.   Long B, Koyfman A: **Best Clinical Practice: Current Controversies in Evaluation of Low-Risk Chest Pain-Part 1**. *The Journal of emergency medicine* 2016, **51**(6):668-676.

3.   Hollander JE, Than M, Mueller C: **State-of-the-Art Evaluation of Emergency Department Patients Presenting With Potential Acute Coronary Syndromes**. *Circulation* 2016, **134**(7):547-564.

4.   Backus BE, Six AJ, Kelder JH, Gibler WB, Moll FL, Doevendans PA: **Risk scores for patients with chest pain: evaluation in the emergency department**. *Curr Cardiol Rev* 2011, **7**(1):2-8.

5.   Liu N, Ng JCJ, Ting CE, Sakamoto JT, Ho AFW, Koh ZX, Pek PP, Lim SH, Ong MEH: **Clinical scores for risk stratification of chest pain patients in the emergency department: an updated systematic review**. *Journal of Emergency and Critical Care Medicine* 2018, **2**(2).

6.   Six AJ, Cullen L, Backus BE, Greenslade J, Parsonage W, Aldous S, Doevendans PA, Than M: **The HEART score for the assessment of patients with chest pain in the emergency department: a multinational validation study**. *Crit Pathw Cardiol* 2013, **12**(3):121-126.

7.   Antman EM, Cohen M, Bernink PJ, McCabe CH, Horacek T, Papuchis G, Mautner B, Corbalan R, Radley D, Braunwald E: **The TIMI risk score for unstable angina/non-ST elevation MI: A method for prognostication and therapeutic decision making**. *Jama* 2000, **284**(7):835-842.

8.   Granger CB, Goldberg RJ, Dabbous O, Pieper KS, Eagle KA, Cannon CP, Van De Werf F, Avezum A, Goodman SG, Flather MD *et al*: **Predictors of hospital mortality in the global registry of acute coronary events**. *Arch Intern Med* 2003, **163**(19):2345-2353.

9.   Sakamoto JT, Liu N, Koh ZX, Guo D, Heldeweg MLA, Ng JCJ, Ong MEH: **Integrating heart rate variability, vital signs, electrocardiogram, and troponin to triage chest pain patients in the ED**. *Am J Emerg Med* 2017.

10.   Poldervaart JM, Langedijk M, Backus BE, Dekker IMC, Six AJ, Doevendans PA, Hoes AW, Reitsma JB: **Comparison of the GRACE, HEART and TIMI score to predict major adverse cardiac events in chest pain patients at the emergency department**. *International journal of cardiology* 2017, **227**:656-661.

11.   Fernando SM, Tran A, Cheng W, Rochwerg B, Taljaard M, Thiruganasambandamoorthy V, Kyeremanteng K, Perry JJ: **Prognostic Accuracy of the HEART Score for Prediction of Major Adverse Cardiac Events in Patients Presenting With Chest Pain: A Systematic Review and Meta-analysis**. *Academic*

emergency medicine : official journal of the Society for Academic Emergency Medicine 2019, **26**(2):140-151.

12.   Reaney PDW, Elliott HI, Noman A, Cooper JG: **Risk stratifying chest pain patients in the emergency department using HEART, GRACE and TIMI scores, with a single contemporary troponin result, to predict major adverse cardiac events**. *Emergency medicine journal : EMJ* 2018, **35**(7):420-427.

13.   Van Den Berg P, Body R: **The HEART score for early rule out of acute coronary syndromes in the emergency department: a systematic review and meta-analysis**. *European heart journal Acute cardiovascular care* 2018, **7**(2):111-119.

14.   Mahler SA, Riley RF, Hiestand BC, Russell GB, Hoekstra JW, Lefebvre CW, Nicks BA, Cline DM, Askew KL, Elliott SB *et al*: **The HEART Pathway randomized trial: identifying emergency department patients with acute chest pain for early discharge**. *Circ Cardiovasc Qual Outcomes* 2015, **8**(2):195-203.

15.   Liu N, Lin Z, Cao J, Koh ZX, Zhang T, Huang G-B, Ser W, Ong MEH: **An intelligent scoring system and its application to cardiac arrest prediction**. *IEEE Trans Inf Technol Biomed* 2012, **16**(6):1324-1331.

16.   Liu N, Koh ZX, Chua ECP, Tan LML, Lin Z, Mirza B, Ong MEH: **Risk scoring for prediction of acute cardiac complications from imbalanced clinical data**. *IEEE J Biomed Health Inform* 2014, **18**(6):1894-1902.

17.   Ong MEH, Goh K, Fook-Chong S, Haaland B, Wai KL, Koh ZX, Shahidah N, Lin Z: **Heart rate variability risk score for prediction of acute cardiac complications in ED patients with chest pain**. *Am J Emerg Med* 2013, **31**(8):1201-1207.

18.   Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology: **Heart rate variability: standards of measurement, physiological interpretation and clinical use**. *Circulation* 1996, **93**(5):1043-1065.

19.   Rajendra Acharya U, Paul Joseph K, Kannathal N, Lim CM, Suri JS: **Heart rate variability: a review**. *Med Biol Eng Comput* 2006, **44**(12):1031-1051.

20.   Billman GE: **Heart rate variability - a historical perspective**. *Front Physiol* 2011, **2**:86.

21.   Carpeggiani C, L'Abbate A, Landi P, Michelassi C, Raciti M, Macerata A, Emdin M: **Early assessment of heart rate variability is predictive of in-hospital death and major complications after acute myocardial infarction**. *International journal of cardiology* 2004, **96**(3):361-368.

22. Vest AN, Da Poian G, Li Q, Liu C, Nemati S, Shah AJ, Clifford GD: **An open source benchmarked toolbox for cardiovascular waveform and interval analysis**. *Physiological measurement* 2018, **39**(10):105004.

23. Tarvainen MP, Niskanen JP, Lipponen JA, Ranta-Aho PO, Karjalainen PA: **Kubios HRV–heart rate variability analysis software**. *Comput Methods Programs Biomed* 2014, **113**.

24. Liu N, Guo D, Koh ZX, Ho AFW, Ong MEH: **Heart rate n-variability (HRnV): A novel representation of beat-to-beat variation in electrocardiogram**. *bioRxiv* 2018:449504.

25. Ho AF, Fook-Chong S, Pek PP, Ng YY, Wong AS, Ong ME: **Prehospital presentation of patients with ST-segment elevation myocardial infarction in Singapore**. *International journal of cardiology* 2013, **168**(4):4273-4276.

26. Januzzi JL, Jr., Bamberg F, Lee H, Truong QA, Nichols JH, Karakas M, Mohammed AA, Schlett CL, Nagurney JT, Hoffmann U *et al*: **High-sensitivity troponin T concentrations in acute chest pain patients evaluated with cardiac computed tomography**. *Circulation* 2010, **121**(10):1227-1234.

27. Sakamoto JT, Liu N, Koh ZX, Fung NX, Heldeweg ML, Ng JC, Ong ME: **Comparing HEART, TIMI, and GRACE scores for prediction of 30-day major adverse cardiac events in high acuity chest pain patients in the emergency department**. *International journal of cardiology* 2016, **221**:759-764.

28. Fawcett T: **An introduction to ROC analysis**. *Pattern Recogn Lett* 2006, **27**(8):861-874.

29. Shaffer F, Ginsberg JP: **An Overview of Heart Rate Variability Metrics and Norms**. *Frontiers in public health* 2017, **5**:258.

30. Peng CK, Havlin S, Stanley HE, Goldberger AL: **Quantification of scaling exponents and crossover phenomena in nonstationary heartbeat time series**. *Chaos* 1995, **5**(1):82-87.

31. Barbieri R, Matten EC, Alabi AA, Brown EN: **A point-process model of human heartbeat intervals: new definitions of heart rate and heart rate variability**. *American journal of physiology Heart and circulatory physiology* 2005, **288**(1):H424-435.

32. Vollmer M: **A robust, simple and reliable measure of heart rate variability using relative RR intervals**. In: *Computing in Cardiology Conference (CinC): 6-9 Sept. 2015 2015*; 2015: 609-612.

33.   Chase M, Robey JL, Zogby KE, Sease KL, Shofer FS, Hollander JE: **Prospective validation of the Thrombolysis in Myocardial Infarction Risk Score in the emergency department chest pain population**. *Annals of emergency medicine* 2006, **48**(3):252-259.

34.   Pollack CV, Sites FD, Shofer FS, Sease KL, Hollander JE: **Application of the TIMI risk score for unstable angina and non-ST elevation acute coronary syndrome to an unselected emergency department chest pain population**. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine* 2006, **13**(1):13-18.

35.   Marcoon S, Chang AM, Lee B, Salhi R, Hollander JE: **HEART score to further risk stratify patients with low TIMI scores**. *Crit Pathw Cardiol* 2013, **12**(1):1-5.

36.   Long B, Oliver J, Streitz M, Koyfman A: **An end-user's guide to the HEART score and pathway**. *Am J Emerg Med* 2017, **35**(9):1350-1355.

37.   Than MP, Pickering JW, Aldous SJ, Cullen L, Frampton CM, Peacock WF, Jaffe AS, Goodacre SW, Richards AM, Ardagh MW *et al*: **Effectiveness of EDACS Versus ADAPT Accelerated Diagnostic Pathways for Chest Pain: A Pragmatic Randomized Controlled Trial Embedded Within Practice**. *Annals of emergency medicine* 2016, **68**(1):93-102 e101.

38.   Than M, Cullen L, Reid CM, Lim SH, Aldous S, Ardagh MW, Peacock WF, Parsonage WA, Ho HF, Ko HF *et al*: **A 2-h diagnostic protocol to assess patients with chest pain symptoms in the Asia-Pacific region (ASPECT): a prospective observational validation study**. *Lancet (London, England)* 2011, **377**(9771):1077-1084.

39.   Laureano-Phillips J, Robinson RD, Aryal S, Blair S, Wilson D, Boyd K, Schrader CD, Zenarosa NR, Wang H: **HEART Score Risk Stratification of Low-Risk Chest Pain Patients in the Emergency Department: A Systematic Review and Meta-Analysis**. *Annals of emergency medicine* 2019, **74**(2):187-203.

40.   Cheng Q, Greenslade JH, Parsonage WA, Barnett AG, Merollini K, Graves N, Peacock WF, Cullen L: **Change to costs and lengths of stay in the emergency department and the Brisbane protocol: an observational study**. *BMJ Open* 2016, **6**(2):e009746.

41.   Ong ME, Padmanabhan P, Chan YH, Lin Z, Overton J, Ward KR, Fei DY: **An observational, prospective study exploring the use of heart rate variability as a**

predictor of clinical outcomes in pre-hospital ambulance patients. *Resuscitation* 2008, **78**(3):289-297.

42. Liu N, Zhang Z, Ho AFW, Ong MEH: **Artificial intelligence in emergency medicine**. *Journal of Emergency and Critical Care Medicine* 2018, **2**:82.

43. Chiew CJ, Liu N, Tagami T, Wong TH, Koh ZX, Ong MEH: **Heart rate variability based machine learning models for risk prediction of suspected sepsis patients in the emergency department**. *Medicine* 2019, **98**(6):e14197.

44. Liu N, Sakamoto JT, Cao J, Koh ZX, Ho AFW, Lin Z, Ong MEH: **Ensemble-Based Risk Scoring with Extreme Learning Machine for Prediction of Adverse Cardiac Events**. *Cognitive Computation* 2017, **9**(4):545-554.

45. Liu N, Koh ZX, Goh J, Lin Z, Haaland B, Ting BP, Ong MEH: **Prediction of adverse cardiac events in emergency department patients with chest pain using machine learning for variable selection**. *BMC Med Inform Decis Mak* 2014, **14**:75.

46. Quintana DS, Alvares GA, Heathers JA: **Guidelines for Reporting Articles on Psychiatry and Heart rate variability (GRAPH): recommendations to advance research communication**. *Transl Psychiatry* 2016, **6**:e803.

47. Kloter E, Barrueto K, Klein SD, Scholkmann F, Wolf U: **Heart Rate Variability as a Prognostic Factor for Cancer Survival - A Systematic Review**. *Front Physiol* 2018, **9**:623.

48. Schroeder EB, Chambless LE, Liao D, Prineas RJ, Evans GW, Rosamond WD, Heiss G: **Diabetes, glucose, insulin, and heart rate variability: the Atherosclerosis Risk in Communities (ARIC) study**. *Diabetes Care* 2005, **28**(3):668-674.

49. Buchman TG, Stein PK, Goldstein B: **Heart rate variability in critical illness and critical care**. *Curr Opin Crit Care* 2002, **8**(4):311-315.

50. Polikar R: **Ensemble based systems in decision making**. *IEEE Circuits Syst Mag* 2006, **6**(3):21-44.

51. Lundberg SM, Nair B, Vavilala MS, Horibe M, Eisses MJ, Adams T, Liston DE, King-Wai Low D, Newman SF, Kim J *et al*: **Explainable machine-learning predictions for the prevention of hypoxaemia during surgery**. *Nature biomedical engineering* 2018, **2**(10):749-760.

Table 1: Patient baseline characteristics.

	Total (n=795)	MACE (n=247)	Non-MACE (n=548)	p-value
Age, mean (SD)	59.63 (12.88)	61.06 (11.38)	58.99 (13.47)	0.035
Male gender, n (%)	542 (68.2)	188 (76.1)	354 (64.6)	0.002
Race, n (%)				0.623
Chinese	492 (61.9)	159 (64.4)	333 (60.8)	
Indian	129 (16.2)	34 (13.8)	95 (17.3)	
Malay	150 (18.9)	46 (18.6)	104 (19.0)	
Other	24 (3.0)	8 (3.2)	16 (2.9)	
Medical history, n (%)				
Ischemic heart disease	343 (43.1)	115 (46.6)	228 (41.6)	0.22
Diabetes	278 (35.0)	106 (42.9)	172 (31.4)	0.002
Hypertension	509 (64.0)	161 (65.2)	348 (63.5)	0.707
Hypercholesterolemia	476 (59.9)	151 (61.1)	325 (59.3)	0.683
Stroke	58 (7.3)	15 (6.1)	43 (7.8)	0.458
Cancer	29 (3.6)	7 (2.8)	22 (4.0)	0.537
Respiratory disease	31 (3.9)	5 (2.0)	26 (4.7)	0.102
Chronic kidney disease	87 (10.9)	26 (10.5)	61 (11.1)	0.32
Congestive heart failure	38 (4.8)	9 (3.6)	29 (5.3)	0.407
History of PCI	199 (25.0)	68 (27.5)	131 (23.9)	0.316
History of CABG	71 (8.9)	26 (10.5)	45 (8.2)	0.355
History of AMI	133 (16.7)	48 (19.4)	85 (15.5)	0.288
Active smoker	197 (24.8)	73 (29.6)	124 (22.6)	0.003

MACE, major adverse cardiac events; SD, standard deviation; PCI, percutaneous coronary intervention; CABG, coronary artery bypass graft; AMI, acute myocardial infarction.

Table 2: Descriptive analyses of heart rate variability (HRV) and heart rate n-variability (HRnV) parameters.

	HRV	HR$_2$V	HR$_2$V$_1$	HR$_3$V	HR$_3$V$_1$	HR$_3$V$_2$
Mean NN (s)	829.40 (169.49)	1656.65 (339.85)	1658.81 (338.99)	2484.80 (509.33)	2488.22 (508.50)	2485.02 (509.84)
SDNN (s)	38.16 (25.49)	62.28 (45.45)	68.81 (47.00)	82.06 (62.47)	97.79 (67.46)	87.77 (64.52)
RMSSD (s)	30.04 (23.07)	32.61 (26.68)	33.79 (25.67)	34.83 (28.86)	36.27 (26.50)	34.98 (27.43)
Skewness	-0.65 (2.34)	-0.41 (1.66)	-0.59 (1.95)	-0.29 (1.29)	-0.55 (1.69)	-0.38 (1.42)
Kurtosis	14.59 (26.83)	7.33 (13.58)	10.17 (17.90)	5.15 (8.13)	8.06 (12.92)	5.98 (9.75)
Triangular index	7.68 (4.19)	10.38 (5.10)	12.60 (6.45)	11.47 (5.29)	16.25 (7.94)	13.06 (6.04)
NN50 (count)	21.08 (33.98)	14.46 (20.35)	29.35 (40.03)	11.57 (15.05)	35.29 (44.34)	17.41 (22.51)
pNN50 (%)	6.31 (11.08)	8.66 (13.18)	8.75 (12.97)	10.31 (14.27)	10.38 (13.95)	10.28 (14.20)
NN50n (count)	-	4.16 (9.72)	8.45 (18.76)	1.37 (3.72)	4.37 (10.72)	2.08 (5.48)
pNN50n (%)	-	2.60 (6.67)	2.64 (6.47)	1.32 (3.95)	1.39 (3.86)	1.33 (3.87)
Total power (ms^2)	2518.30 (4797.05)	7797.46 (16947.44)	9156.26 (17970.75)	13904.78 (37182.24)	18714.67 (37620.26)	15706.11 (34845.52)
VLF power (ms^2)	985.18 (1991.52)	3401.42 (6569.37)	3922.74 (7987.46)	6503.53 (14205.11)	8772.26 (17986.63)	7567.79 (14666.32)
LF power (ms^2)	732.36 (1841.88)	2626.83 (7593.16)	2782.48 (7212.62)	5091.49 (18402.20)	5740.99 (15243.38)	5397.76 (16001.18)
HF power (ms^2)	527.27 (1232.69)	1328.86 (4033.96)	1361.53 (3433.55)	1661.69 (7237.55)	1762.45 (4851.11)	1761.05 (6477.63)
LF power norm (nu)	56.76 (19.20)	66.82 (18.17)	66.42 (17.35)	76.53 (15.32)	77.65 (14.55)	77.93 (14.95)
HF power norm (nu)	43.24 (19.20)	33.18 (18.17)	33.58 (17.35)	23.47 (15.32)	22.35 (14.55)	22.07 (14.95)
LF/HF	1.99 (1.93)	3.24 (2.95)	3.04 (2.73)	5.60 (5.21)	5.79 (4.99)	6.06 (5.18)
Poincaré SD1 (ms)	21.27 (16.34)	23.12 (18.93)	23.92 (18.18)	24.72 (20.50)	25.68 (18.77)	24.80 (19.46)
Poincaré SD2 (ms)	48.82 (33.29)	84.47 (62.15)	93.88 (64.58)	112.87 (86.62)	135.55 (94.02)	121.20 (89.72)
SampEn	1.57 (0.51)	83.84 (2324.24)	1.33 (0.48)	248.48 (4020.64)	1.06 (0.41)	1.14 (0.45)
ApEn	0.99 (0.20)	0.72 (0.18)	0.91 (0.17)	0.60 (0.15)	0.84 (0.17)	0.70 (0.15)
DFA, α1	0.99 (0.31)	1.24 (0.29)	1.23 (0.27)	1.41 (0.27)	1.42 (0.23)	1.42 (0.25)
DFA, α2	0.95 (0.22)	0.98 (0.35)	0.98 (0.22)	0.86 (0.65)	1.01 (0.22)	1.02 (0.36)

HRV, heart rate variability; mean NN, average of R-R intervals; SDNN, standard deviation of R-R intervals; RMSSD, square root of the mean squared differences between R-R intervals; NN50, the number of times that the absolute difference between 2 successive R-R intervals exceeds 50 ms; pNN50, NN50 divided by the total number of R-R intervals; NN50n, the number of times that the absolute difference between 2 successive RR$_n$I/RR$_n$I$_m$ sequences exceeds 50×n ms; pNN50n, NN50n divided by the total number of RR$_n$I/RR$_n$I$_m$ sequences; VLF, very low frequency; LF, low frequency; HF, high frequency; SD, standard deviation; SampEn, sample entropy; ApEn, approximate entropy; DFA, detrended fluctuation analysis.

Table 3: Univariable analysis of HR_nV and HR_nV_m parameters.

	HRV		HR₂V		HR₃V	
	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Mean NN	0.999 (0.998-1.000)	0.023*	0.999 (0.999-1.000)	0.023*	1.000 (0.999-1.000)	0.023*
SDNN	0.992 (0.986-0.999)	0.023*	0.996 (0.992-1.000)	0.028*	0.997 (0.995-1.000)	0.060
RMSSD	0.990 (0.982-0.998)	0.010*	0.992 (0.985-0.998)	0.011*	0.994 (0.988-0.999)	0.030*
Skewness	1.059 (0.991-1.132)	0.088	1.079 (0.981-1.186)	0.118	1.139 (1.006-1.290)	0.040*
Kurtosis	1.006 (1.000-1.011)	0.038*	1.009 (0.998-1.019)	0.113	1.011 (0.993-1.029)	0.242
Triangular index	0.961 (0.925-0.998)	0.039*	0.967 (0.938-0.997)	0.032*	0.978 (0.950-1.007)	0.133
NN50	0.993 (0.987-0.998)	0.008*	0.989 (0.981-0.998)	0.012*	0.988 (0.977-0.999)	0.031*
pNN50	0.978 (0.962-0.995)	0.009*	0.984 (0.971-0.997)	0.014*	0.987 (0.976-0.999)	0.027*
NN50n	-	-	0.982 (0.964-1.001)	0.065	0.952 (0.905-1.002)	0.059
pNN50n	-	-	0.974 (0.946-1.002)	0.069	0.951 (0.903-1.001)	0.054
Total power	1.000 (1.000-1.000)	0.031*	1.000 (1.000-1.000)	0.021*	1.000 (1.000-1.000)	0.072
VLF power	1.000 (1.000-1.000)	0.132	1.000 (1.000-1.000)	0.070	1.000 (1.000-1.000)	0.133
LF power	1.000 (1.000-1.000)	0.077	1.000 (1.000-1.000)	0.023*	1.000 (1.000-1.000)	0.063
HF power	1.000 (0.999-1.000)	0.002*	1.000 (1.000-1.000)	0.014*	1.000 (1.000-1.000)	0.074
LF power norm	1.001 (0.994-1.009)	0.738	0.999 (0.99-1.007)	0.733	0.994 (0.985-1.004)	0.248
HF power norm	0.999 (0.991-1.007)	0.738	1.001 (0.993-1.01)	0.733	1.006 (0.996-1.015)	0.248
LF/HF	1.034 (0.959-1.116)	0.381	1.014 (0.964-1.066)	0.592	1.001 (0.973-1.031)	0.923
Poincaré SD1	0.986 (0.975-0.997)	0.010*	0.988 (0.979-0.997)	0.011*	0.991 (0.983-0.999)	0.029*
Poincaré SD2	0.995 (0.990-1.000)	0.032*	0.997 (0.994-1.000)	0.032*	0.998 (0.996-1.000)	0.063
SampEn	0.813 (0.604-1.095)	0.173	0.730 (0.545-0.977)	0.035*	1.000 (1.000-1.000)	0.932
ApEn	1.645 (0.752-3.598)	0.213	2.319 (1.003-5.357)	0.049*	1.241 (0.463-3.327)	0.667
DFA, α1	0.953 (0.585-1.552)	0.846	1.031 (0.611-1.741)	0.908	0.968 (0.560-1.672)	0.907
DFA, α2	1.532 (0.773-3.034)	0.221	1.202 (0.782-1.848)	0.401	1.184 (0.934-1.500)	0.163
	HR₂V₁		HR₃V₁		HR₃V₂	
	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Mean NN	0.999 (0.999-1.000)	0.023*	1.000 (0.999-1.000)	0.023*	1.000 (0.999-1.000)	0.023*
SDNN	0.996 (0.993-1.000)	0.034*	0.997 (0.995-1.000)	0.042*	0.997 (0.995-1.000)	0.034*
RMSSD	0.991 (0.984-0.998)	0.010*	0.992 (0.986-0.999)	0.016*	0.993 (0.986-0.999)	0.016*
Skewness	1.061 (0.980-1.149)	0.144	1.072 (0.978-1.176)	0.139	1.098 (0.982-1.227)	0.100
Kurtosis	1.007 (0.999-1.015)	0.082	1.006 (0.994-1.017)	0.333	1.010 (0.995-1.025)	0.195
Triangular index	0.981 (0.958-1.005)	0.119	0.982 (0.963-1.001)	0.065	0.974 (0.949-0.999)	0.040*
NN50	0.995 (0.991-0.999)	0.018*	0.996 (0.993-1.000)	0.052	0.992 (0.985-0.999)	0.035*
pNN50	0.984 (0.972-0.997)	0.020*	0.988 (0.977-1.000)	0.049*	0.988 (0.976-0.999)	0.035*
NN50n	0.989 (0.979-1.000)	0.043*	0.982 (0.964-1.000)	0.054	0.974 (0.943-1.007)	0.118
pNN50n	0.969 (0.939-0.999)	0.046*	0.947 (0.895-1.002)	0.058	0.960 (0.914-1.009)	0.109
Total power	1.000 (1.000-1.000)	0.048*	1.000 (1.000-1.000)	0.072	1.000 (1.000-1.000)	0.029*
VLF power	1.000 (1.000-1.000)	0.139	1.000 (1.000-1.000)	0.145	1.000 (1.000-1.000)	0.074
LF power	1.000 (1.000-1.000)	0.084	1.000 (1.000-1.000)	0.092	1.000 (1.000-1.000)	0.027*
HF power	1.000 (1.000-1.000)	0.005*	1.000 (1.000-1.000)	0.010*	1.000 (1.000-1.000)	0.022*
LF power norm	1.000 (0.991-1.008)	0.937	0.995 (0.985-1.006)	0.382	0.995 (0.986-1.005)	0.356

HF power norm	1.000 (0.992-1.009)	0.937	1.005 (0.994-1.015)	0.382	1.005 (0.995-1.015)	0.356
LF/HF	1.024 (0.970-1.080)	0.387	1.003 (0.973-1.033)	0.863	0.999 (0.971-1.029)	0.966
Poincaré SD1	0.987 (0.978-0.997)	0.010*	0.989 (0.980-0.998)	0.016*	0.989 (0.981-0.998)	0.016*
Poincaré SD2	0.997 (0.995-1.000)	0.039*	0.998 (0.996-1.000)	0.045*	0.998 (0.996-1.000)	0.037*
SampEn	0.854 (0.623-1.171)	0.328	0.802 (0.553-1.161)	0.242	0.709 (0.500-1.005)	0.053
ApEn	2.065 (0.842-5.064)	0.113	1.207 (0.499-2.922)	0.677	2.558 (0.906-7.222)	0.076
DFA, α1	0.888 (0.514-1.537)	0.672	1.039 (0.547-1.971)	0.907	1.004 (0.549-1.835)	0.991
DFA, α2	1.557 (0.782-3.098)	0.208	1.554 (0.780-3.093)	0.210	1.169 (0.764-1.789)	0.472

HRV, heart rate variability; OR, odds ratio; CI, confidence interval; mean NN, average of R-R intervals; SDNN, standard deviation of R-R intervals; RMSSD, square root of the mean squared differences between R-R intervals; NN50, the number of times that the absolute difference between 2 successive R-R intervals exceeds 50 ms; pNN50, NN50 divided by the total number of R-R intervals; NN50n, the number of times that the absolute difference between 2 successive RR_nI/RR_nI_m sequences exceeds 50×n ms; pNN50n, NN50n divided by the total number of RR_nI/RR_nI_m sequences; VLF, very low frequency; LF, low frequency; HF, high frequency; SD, standard deviation; SampEn, sample entropy; ApEn, approximate entropy; DFA, detrended fluctuation analysis.

Table 4: Multivariable analysis of HR_nV and HR_nV_m parameters by adjusting for age and sex.

	HRV		HR_2V		HR_3V	
	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Mean NN	0.999 (0.998-1)	0.005*	0.999 (0.999-1.000)	0.005*	1.000 (0.999-1.000)	0.005*
SDNN	0.993 (0.986-0.999)	0.035*	0.996 (0.992-1.000)	0.040*	0.998 (0.995-1.000)	0.093
RMSSD	0.990 (0.982-0.998)	0.011*	0.992 (0.985-0.999)	0.016*	0.994 (0.988-1.000)	0.047*
Skewness	1.064 (0.995-1.138)	0.068	1.082 (0.983-1.191)	0.109	1.140 (1.005-1.293)	0.042*
Kurtosis	1.005 (1.000-1.011)	0.047*	1.008 (0.997-1.019)	0.139	1.011 (0.993-1.030)	0.238
Triangular index	0.967 (0.93-1.006)	0.093	0.971 (0.940-1.002)	0.070	0.982 (0.953-1.013)	0.256
NN50	0.993 (0.988-0.999)	0.013*	0.991 (0.982-0.999)	0.030*	0.990 (0.979-1.001)	0.078
pNN50	0.979 (0.963-0.996)	0.015*	0.986 (0.972-0.999)	0.033*	0.989 (0.977-1.001)	0.063
NN50n	-	-	0.983 (0.964-1.002)	0.081	0.954 (0.906-1.005)	0.077
pNN50n	-	-	0.975 (0.947-1.004)	0.086	0.952 (0.903-1.004)	0.069
Total power	1.000 (1.000-1.000)	0.042*	1.000 (1.000-1.000)	0.026*	1.000 (1.000-1.000)	0.104
VLF power	1.000 (1.000-1.000)	0.167	1.000 (1.000-1.000)	0.082	1.000 (1.000-1.000)	0.152
LF power	1.000 (1.000-1.000)	0.093	1.000 (1.000-1.000)	0.033*	1.000 (1.000-1.000)	0.105
HF power	1.000 (0.999-1.000)	0.003*	1.000 (1.000-1.000)	0.016*	1.000 (1.000-1.000)	0.101
LF power norm	1.002 (0.994-1.011)	0.589	0.999 (0.990-1.007)	0.769	0.994 (0.984-1.003)	0.202
HF power norm	0.998 (0.989-1.006)	0.589	1.001 (0.993-1.010)	0.769	1.006 (0.997-1.016)	0.202
LF/HF	1.039 (0.961-1.124)	0.336	1.013 (0.962-1.066)	0.620	0.999 (0.970-1.028)	0.928
Poincaré SD1	0.986 (0.975-0.997)	0.011*	0.989 (0.980-0.998)	0.016*	0.992 (0.983-1.000)	0.047*
Poincaré SD2	0.995 (0.990-1.000)	0.050*	0.997 (0.994-1.000)	0.046*	0.998 (0.996-1.000)	0.098
SampEn	0.852 (0.630-1.152)	0.297	0.752 (0.559-1.010)	0.058	1.000 (1.000-1.000)	0.956
ApEn	1.669 (0.754-3.693)	0.207	2.668 (1.139-6.246)	0.024*	1.507 (0.555-4.096)	0.421
DFA, α1	0.991 (0.593-1.654)	0.971	1.072 (0.622-1.848)	0.802	0.962 (0.550-1.682)	0.891
DFA, α2	1.499 (0.750-2.993)	0.252	1.204 (0.782-1.853)	0.400	1.193 (0.941-1.512)	0.146
	HR_2V_1		HR_3V_1		HR_3V_2	
	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Mean NN	0.999 (0.999-1.000)	0.005*	1.000 (0.999-1.000)	0.005*	1.000 (0.999-1.000)	0.005*
SDNN	0.996 (0.993-1.000)	0.052	0.998 (0.995-1.000)	0.064	0.997 (0.995-1.000)	0.049*
RMSSD	0.992 (0.985-0.998)	0.015*	0.993 (0.986-0.999)	0.023*	0.993 (0.987-0.999)	0.023*
Skewness	1.066 (0.984-1.156)	0.118	1.079 (0.983-1.185)	0.108	1.099 (0.982-1.229)	0.099
Kurtosis	1.007 (0.999-1.015)	0.096	1.005 (0.994-1.017)	0.377	1.010 (0.994-1.025)	0.218
Triangular index	0.985 (0.960-1.010)	0.234	0.985 (0.965-1.005)	0.137	0.977 (0.951-1.003)	0.088
NN50	0.996 (0.991-1.000)	0.047*	0.997 (0.993-1.001)	0.130	0.993 (0.986-1.001)	0.084
pNN50	0.986 (0.973-1.000)	0.046*	0.990 (0.979-1.002)	0.111	0.989 (0.978-1.001)	0.076
NN50n	0.990 (0.980-1.000)	0.059*	0.982 (0.963-1.001)	0.064	0.975 (0.943-1.008)	0.142
pNN50n	0.971 (0.941-1.002)	0.063	0.947 (0.893-1.004)	0.067	0.962 (0.915-1.012)	0.131
Total power	1.000 (1.000-1.000)	0.064	1.000 (1.000-1.000)	0.096	1.000 (1.000-1.000)	0.035*
VLF power	1.000 (1.000-1.000)	0.173	1.000 (1.000-1.000)	0.180	1.000 (1.000-1.000)	0.086
LF power	1.000 (1.000-1.000)	0.100	1.000 (1.000-1.000)	0.108	1.000 (1.000-1.000)	0.037*
HF power	1.000 (1.000-1.000)	0.006*	1.000 (1.000-1.000)	0.014*	1.000 (1.000-1.000)	0.025*
LF power norm	1.000 (0.991-1.009)	0.960	0.995 (0.984-1.005)	0.324	0.995 (0.985-1.005)	0.329

HF power norm	1.000 (0.991-1.009)	0.960	1.005 (0.995-1.016)	0.324	1.005 (0.995-1.015)	0.329
LF/HF	1.023 (0.968-1.081)	0.428	0.999 (0.969-1.030)	0.940	0.996 (0.967-1.026)	0.786
Poincaré SD1	0.988 (0.979-0.998)	0.015*	0.990 (0.981-0.999)	0.023*	0.990 (0.981-0.999)	0.023*
Poincaré SD2	0.997 (0.995-1.000)	0.059	0.998 (0.997-1.000)	0.068	0.998 (0.996-1.000)	0.052
SampEn	0.870 (0.632-1.197)	0.393	0.842 (0.578-1.227)	0.371	0.716 (0.504-1.019)	0.064
ApEn	2.520 (1.009-6.298)	0.048*	1.413 (0.575-3.471)	0.451	3.461 (1.201-9.971)	0.021*
DFA, α1	0.898 (0.508-1.587)	0.710	1.068 (0.555-2.058)	0.843	1.005 (0.543-1.838)	0.988
DFA, α2	1.507 (0.751-3.025)	0.249	1.500 (0.746-3.014)	0.255	1.172 (0.764-1.798)	0.467

HRV, heart rate variability; OR, odds ratio; CI, confidence interval; mean NN, average of R-R intervals; SDNN, standard deviation of R-R intervals; RMSSD, square root of the mean squared differences between R-R intervals; NN50, the number of times that the absolute difference between 2 successive R-R intervals exceeds 50 ms; pNN50, NN50 divided by the total number of R-R intervals; NN50n, the number of times that the absolute difference between 2 successive RR_nI/RR_nI_m sequences exceeds $50 \times n$ ms; pNN50n, NN50n divided by the total number of RR_nI/RR_nI_m sequences; VLF, very low frequency; LF, low frequency; HF, high frequency; SD: standard deviation; SampEn, sample entropy; ApEn, approximate entropy; DFA: detrended fluctuation analysis.

Table 5: Multivariable analysis with logistic regression on all variables.

Variable	Adjusted OR	95% CI
Age	1.021	1.002-1.041
Diastolic BP	1.018	1.003-1.034
Pain score	1.082	1.003-1.168
ST-elevation	6.449	2.762-15.059
ST-depression	4.827	2.511-9.277
Q wave	3.383	1.668-6.860
Cardiac history†	7.838	5.192-11.832
Troponin	4.406	3.218-6.033
HRV NN50	0.981	0.970-0.991
HR_2V skewness	0.806	0.622-1.045
HR_2V SampEn	0.600	0.348-1.035
HR_2V ApEn	0.095	0.014-0.628
HR_2V_1 ApEn	19.700	2.942-131.900
HR_3V RMSSD	1.024	1.008-1.040
HR_3V skewness	1.560	1.116-2.181
HR_3V_2 HF power	1.000	1.000-1.000

BP, blood pressure; HRV, heart rate variability; OR, odds ratio; CI, confidence interval; mean NN, average of R-R intervals; RMSSD, square root of the mean squared differences between R-R intervals; NN50, the number of times that the absolute difference between 2 successive R-R intervals exceeds 50 ms; LF, low frequency; HF, high frequency; SampEn, sample entropy; ApEn, approximate entropy.

† Cardiac history was a numeric value that was derived from the narrative in the hospital charts. Its value was zero if the patient history contained characteristics of atypical cardiac chest pain; Its value was two if the history contained characteristics of typical cardiac chest pain; Its value was one if the history contained characteristics of both atypical and typical cardiac chest pain.

Table 6: Comparison of performance of the HRnV model (based on leave-one-out cross-validation), HEART, TIMI, and GRACE scores in predicting 30-day major adverse cardiac events (MACE).

	AUC (95% CI)	Cut-off	Sensitivity (95% CI)	Specificity (95% CI)	PPV (95% CI)	NPV (95% CI)
HRnV Model	0.888 (0.860-0.917)	0.3611†	77.3% (72.1% - 82.5%)	81.8% (78.5% - 85.0%)	65.6% (60.2% - 71.1%)	88.9% (86.1% - 91.6%)
	-	0.0352	99.2% (98.1% - 100.0%)	39.6% (35.5% - 43.7%)	42.5% (38.5% - 46.6%)	99.1% (97.8% - 100.0%)
HEART	0.841 (0.808-0.874)	5†	78.9% (73.9% - 84.0%)	72.8% (69.1% - 76.5%)	56.7% (51.4% - 61.9%)	88.5% (85.5% - 91.4%)
	-	3	99.6% (98.8% - 100.0%)	35.8% (31.8% - 39.8%)	41.1% (37.2% - 45.1%)	99.5% (98.5% - 100.0%)
TIMI	0.681 (0.639-0.723)	2†	63.6% (57.6% - 69.6%)	58.4% (54.3% - 62.5%)	40.8% (35.9% - 45.7%)	78.0% (74.0% - 82.1%)
	-	0	98.4% (96.8% - 100.0%)	19.3% (16.0% - 22.7%)	35.5% (31.9% - 39.1%)	96.4% (92.9% - 99.9%)
GRACE	0.665 (0.623-0.707)	107†	64.0% (58.0% - 70.0%)	60.8% (56.7% - 64.9%)	42.4% (37.3% - 47.4%)	78.9% (75.0% - 82.8%)
	-	60	98.8% (97.4% - 100.0%)	8.0% (5.8% - 10.3%)	32.6% (29.3% - 36.0%)	93.6% (86.6% - 100.0%)

AUC, area under the curve; CI, confidence interval; PPV, positive predictive value; NPV, negative predictive value; HEART, History, ECG, Age, Risk factors and Troponin; TIMI, Thrombolysis in Myocardial Infarction; GRACE, Global Registry of Acute Coronary Events.
† Optimal cut-off values, defined as the points nearest to the upper-left corner on the ROC curves.

Figure Legends

Figure 1: **(a)** Illustration of R-R intervals (RRIs) and the definition of RR_nI where $1 \leq n \leq N$ and $N \ll \widehat{N}$. $\widehat{N}$ is the total number of RRIs; **(b)** Illustration of RRIs and the definition of RR_nI_m where $1 \leq n \leq N$, $1 \leq m \leq N-1$, and $N \ll \widehat{N}$. $\widehat{N}$ is the total number of RRIs and m indicates the non-overlapped portion between two consecutive RR_nI_m sequences.

Figure 2: The receiver operating characteristic (ROC) curves produced by the heart rate n-variability (HRnV) model (performance was based on leave-one-out cross-validation), the History, ECG, Age, Risk factors and Troponin (HEART) score, the Thrombolysis in Myocardial Infarction (TIMI) score, and the Global Registry of Acute Coronary Events (GRACE) score.

(a)

(b)

Figure 1: **(a)** Illustration of R-R intervals (RRIs) and the definition of RR_nI where $1 \leq n \leq N$ and $N \ll \widehat{N}$. $\widehat{N}$ is the total number of RRIs; **(b)** Illustration of RRIs and the definition of RR_nI_m where $1 \leq n \leq N$, $1 \leq m \leq N-1$, and $N \ll \widehat{N}$. $\widehat{N}$ is the total number of RRIs and m indicates the non-overlapped portion between two consecutive RR_nI_m sequences.

Figure 2: The receiver operating characteristic (ROC) curves produced by the heart rate n-variability (HRnV) model (performance was based on leave-one-out cross-validation), the History, ECG, Age, Risk factors and Troponin (HEART) score, the Thrombolysis in Myocardial Infarction (TIMI) score, and the Global Registry of Acute Coronary Events (GRACE) score.