# Rats Sniff off Toxic Air

Haoxuan Chen[#], Xinyue Li[#] and Maosheng Yao*

State Key Joint Laboratory of Environmental Simulation and Pollution Control,

College of Environmental Sciences and Engineering, Peking University, Beijing

100871, China

**Corresponding author:**

Maosheng Yao, PhD

Boya Distinguished Professor

State Key Joint Laboratory of Environmental Simulation and Pollution Control,

College of Environmental Sciences and Engineering, Peking University, Beijing

100871, China

Email: yao@pku.edu.cn, Ph: +86 01062767282

# Both authors contributed equally

Beijing, China

January 2020

**Toxic Air**

**Distinctive Breath-borne VOCs**

VOC Sensor

Rats Sniff Off Toxic Air
*(RSTair)*

32
33
34
35

**Abstract**

Breathing air is a fundamental human need, yet its safety, when challenged by

various harmful or lethal substances, is often not properly guarded. For example, air

toxicity is currently monitored only for single or limited number of known toxicants,

thus failing to fully warn against possible hazardous air. Here, we discovered that

within minutes living rats emitted distinctive profiles of volatile organic compounds

(VOCs) via breath when exposed to various airborne toxicants such as endotoxin, $O_3$,

ricin, and $CO_2$. Compared to background indoor air, when exposed to ricin or

endotoxin aerosols breath-borne VOC levels, especially that of carbon disulfide, were

shown to decrease; while their elevated levels were observed for $O_3$ and $CO_2$

exposures. A clear contrast in breath-borne VOCs profiles of rats between different

toxicant exposures was observed with a statistical significance. Differences in

MicroRNA regulations such as miR-33, miR-146a and miR-155 from rats' blood

samples revealed different mechanisms used by the rats in combating different air

toxicant challenges. Similar to dogs, rats were found here to be able to sniff against

toxic air by releasing a specific breath-borne VOC profile. The discovered science

opens a new arena for online monitoring air toxicity and health effects of pollutants.

**Keywords**: Online Monitoring Toxic Air, Rat, Volatile Organic Compound,

microRNA,

## 1. Introduction

Polluted air is a complex mixture of various pollutants, including particulate matter (PM), biologicals, and also gaseous substances such as $O_3$ and NOx. Inhaling these pollutants can cause a variety of serious health problems: respiratory and cardiovascular diseases, and even death. [1-2] For example, PM alone was shown to have resulted in 4.2 million deaths in 2015. [2-3] Exhibiting a positive correlation with daily mean mortality, ground ozone exposure resulted in decreased lung function and airway inflammation. [4-5] In the meantime, exposure to pathogenic bioaerosols including bacteria, fungi, virus, et al in the air can cause respiratory infections with grave human and economic costs. [6-9] Apart from these, there is a growing risk of terrorist attacks by intentionally releasing biological and chemical agents into the air to cause substantial civilian casualties. [10-12] Apparently, inhaling unsafe air has become an increasing health concern. Yet, in many high profile events, in addition to the public sectors, the air being inhaled is not readily protected or properly guarded. Real-time monitoring of air toxicity is of great importance, which however is a long-standing significant challenge in the field.

For monitoring hazardous substances in the air, a variety of real-time online monitoring methods have been previously developed or tested for individual pollutants such as the PM and other chemicals. [13-16] While for bioaerosols, the adenosine triphosphate (ATP) bioluminescence technology, surface-enhanced Raman spectroscopy (SERS), bioaerosol mass spectrometry (BAMS), ultraviolet aerodynamic particle sizer (UVAPS) as well as silicon nanowire (SiNW) biosensor were investigated and attempted over the years. [17-19] It is well known that these existing or developed technologies are mostly restricted to either single agent or

overall microbial concentration levels without identifying species. In addition,

airborne pollutants and toxicity could vary greatly from one location to another [20-21],

thus presenting location-specific air toxicity and health effects. Current

epidemiological or toxicological methods involving data analysis or animal and cells

experiments cannot provide in situ air toxicity information, accordingly failing to

represent the response at the time of exposure since biomarker levels evolve over time.

[22] In addition, under certain scenarios (high profile meetings or locations) a rapid

response to air toxicity needs to be in place in order to protect the interests. However,

the response time is very demanding for an immediate effective counter-measure, for

example, usually several minutes can be tolerated. [10, 23] In many air environments,

multiple hazardous pollutants could also co-exist even with unknown ones at a

particular time, which makes protecting the air rather difficult, if not impossible,

using current technologies of species level detection and warning.

Previously, olfactory receptors of mouse cells for odors [24], immune B cells [25] for

pathogen detections, and silicon nanowire sensor arrays for explosives[26] were studied.

Recently, a breath-borne biomarker detection system (dLABer) integrating rat's breath

sampling, microfluidics and a silicon nanowire field effect transistor (SiNW FET)

device has been developed for real-time tracking biological molecules in the breath of

rats exposed to particulate matter (PM).[27] The dLABer system was shown to be able

to online detect interleukins-6 (IL-6) level in rat's breath, and capable of

differentiating between different PM toxicities from different cities using the

biomarker level. However, as observed in the study the production of protein

biomarkers could significantly lag behind the pollutant exposure, thus falling short of

providing a timely warning against toxic air. Nonetheless, exhaled breath is

increasingly being used in biomarker analysis in both medical and environmental

health studies. [28-30] In addition to protein biomarkers, a large number of volatile

compounds (e.g., nitric oxide, carbon monoxide, hydrocarbons) have been also

studied to assess health status and even developed for clinical diagnosis.[31-32] For

example, ethane and n-pentane detected in the breath were linked to the *in vivo* level

of lipid peroxidation and oxidative stress [33]; breath-borne acetone was shown to

correlate with the metabolic state of diabetic patients.[34] In addition, exhaled VOCs

have been used for the diagnosis of asthma, lung cancer and other diseases.[35-37]

Undoubtedly, breath-borne VOC has emerged as a promising biomarker for health or

environmental exposure monitoring.

Inspired by the dog sniff for explosive, the work here was conducted to

investigate if we can use breath-borne VOCs from living rats to real-time monitor

toxic air. Particularly, we wanted to investigate: 1) When rats are exposed to air

toxicants, whether the VOC species and concentration in the exhaled breath change?

If yes, how long does the change need to occur? 2) Are there any specific exhaled

VOCs in response to different toxicants exposure including both chemical and

biological agents? 3) To develop an online air toxicity analyzing system via real-time

monitoring of exhaled VOCs of rats. The work here has demonstrated a great promise

for online air toxicity monitoring using the method developed, and opened a new

arena for studying health effects of air pollution.

**2. Materials and methods**

**2.1 *Rat breeding***

The Jugular Vascular Catheterizations (JVC) rat model described in our previous

study was used in this work.[38] Weighing 200–240 g at an age of 10 weeks, a total of

20 male Wistar rats with jugular vein catheterization operation and a 100 mL/min

inhalation rate were purchased from a local provider (Beijing Vital River Laboratory

Animal Technology Co., Ltd.). With about 1 centimeter out of the skin, a flexible

sterile catheter was embedded into the jugular vein and fixed on the back with staples.

Under a 12 h light/12 h dark cycle, all the rats were raised in an animal care facility

naturally with a normal chow diet. After 1 week of acclimation, the rats were

randomly divided into 5 groups (4 rats in each group) for the exposure of different air

toxicants. All animal experiments were approved by the Institutional Review Board of

Peking University and relevant experiments were performed in accordance with the

ethical standards (approval # LA2019294).

*2.2 Preparation of toxic air*

In this work, four different exposure toxicants (ricin, endotoxin, ozone and

carbon dioxide) and indoor air (as a background control) (5 groups) for rats were

prepared for the exposure experiments. Ricin was extracted from the seeds of castor

produced in Xinjiang, China, and prepared by Institute of Microbiology and

Epidemiology, Academy of Military Medical Sciences in Beijing. The endotoxin was

purchased from Associates of Cape Cod, Inc., USA. The ricin and endotoxin

suspensions were prepared by vigorously vortexing 40 μg of ricin or 50 ng endotoxin

per ml deionized (DI) water for 20 min at a vortex rate of 3200 rpm (Vortex Genie-2,

Scientific Industries Co., Ltd., USA). Detailed information about ricin preparation and

exposure can be also found in another work.[39] Here, ozone was generated by an ozone

generator (Guangzhou Environmental Protection Electric Co., Ltd., China) using

corona discharge method. The ozone was further diluted with indoor air for rat

exposure experiments, and the ozone concentration in the exposed chamber was

approximately 5 ppm. Carbon dioxide was purchased from Beijing Haike Yuanchang

Utility Gas Co., Ltd., and diluted 20 times with indoor air to a concentration of about

5% ($5 \times 10^4$ ppm). The major objective of this work was to study whether there will be

specific VOC release by rats when challenged with different toxicants, not specific

dose-response for each toxicant. Nonetheless, the selection of specific exposure levels

for different toxicants was provided and discussed in Supporting Information.

**2.3 *Rats sniff off toxic air (RST$_{air}$) system and experimental procedure***

To investigate whether we can use breath-borne VOCs from living systems to

real-time monitor toxic air, we have developed the system named as *RSTair* (Rats

Sniff Off Toxic Air). As shown in Figure 1 and Figure S1 (Supporting Information),

the system is composed of four major parts: toxicant generator, exposure chamber,

exhaled breath sampling and online VOC analysis. Indoor air was used as carrier gas

for generating toxicant aerosols (ricin and endotoxin) using a Collison Nebulizer (BGI,

Inc., USA) or diluting toxicants gas (ozone and carbon dioxide). The toxicant aerosol

or toxicant gas was introduced together with indoor air into the exposure chamber at a

flow rate of 1 L/min. As shown in Figure S1, a cage was used as the exposure

chamber which can allow rat's feces and urine to fall from below quickly so as to

reduce their influences on VOC analysis. In addition, teflon tubes and vales were also

applied to reducing adsorption loss of VOCs. For ricin and endotoxin, they are small

molecules and relevant particle loss on the aerosolization tubing could be negligible.

When performing the experiments, the rats were first placed in the exposure chamber.

Indoor air was continuously introduced into the chamber at a flow rate of 1 L/min for

10 minutes, then followed by each of tested toxicants together with indoor air at the

same flow rate for about 10 minutes to conduct exposure tests.

For each toxicant experiment, the TVOC was first measured for the control (rat

with indoor air at a flow rate of 1 L/min), then followed by the toxicant aerosol/gas

exposure (rat + toxic air) at the same flow rate for 10 min. Here, the photoionization

detector (PID) (MOCON, Inc., USA) coupled with an air pump was used to real-time

monitor TVOC at a flow rate of 0.6 L/min. After the exposure, the TVOC was

continuously measured again using the PID sensor, reaching a relatively stable level

approximately within 7-8 min. In the meantime, VOC samples were also collected

both before (control) and after the toxicant exposure. A 3.2 L Silonite canister (Entech

Instruments, Inc., USA) was used to collect VOC samples at the flow rate of 0.8

L/min at least 10 min after the exposure test, and the VOCs species were further

analyzed using a gas chromatograph-mass spectrometry/flame ionization detection

(GC–MS/FID) system (Agilent Technologies, Inc., USA). A total of 117 VOC species

were screened and quantified for all exposure tests using the GC-MS system. During

the VOC sample collection, the PID sensor was switched to measure the indoor air

TVOC levels. The detailed descriptions of the working principles of PID and VOC

species analysis by GC–MS/FID are provided in Supporting Information. In this work,

both TVOC and VOC species in the exposure chamber were analyzed for all the

experiments: 1) when the rats were not in the chamber (background air); 2) rats in the

chamber (before toxicant exposure) and rats in the chamber (after toxicant exposure).

The air flows in and out of the exposure chamber were the same both for TVOC

measurements and VOC collection. Based on the dimensions of the exposure chamber,

the residence time of pollutants in the chamber was about 5 min under the

experimental conditions, e.g., the air flow rate was about 1 L/min. We collected VOC

samples at least 10 min after the toxicant exposure, therefore remaining secondary

pollutants produced, if any, during the exposure tests such as $O_3$ will be flushed out by

the indoor air. For each toxicant, the above experiments were repeated four times with

a different rat each time. Each rat (total four) either for toxicant or control exposure

group experienced the exposure only once, not repeated exposures. For each of four

tested toxicants, the same experiments were repeated. All tests were performed inside

a Class 2 Type A2 Biological Safety Cabinet (NuAire, Plymouth, MN), and all

exposure toxicants were ventilated out via the built-in and lab ventilation system.

.

**Figure 1.** Rats sniff off toxic air (*RST*air) system: different toxicants were introduced

or aerosolized into the chamber for exposure. The VOCs from the rat placed in the

chamber before and after the toxicant exposure for 10 min were analyzed by the PID

directly and also by GC–MS/FID method coupled with a VOC sampler silonite

canister. During the VOC collection and measurements by the PID, the toxicant

exposure was terminated. Each time only one rat was placed into the chamber. For

each toxicant, the experiment was independently repeated four times with a different

rat each time.

***2.4 Blood microRNA detection***

Right before and 20 min after each 10-min toxicant exposure, 0.75 mL blood

samples from rats were taken through the catheter using sterile syringes with 23G

flat-end needles and kept at -20 $^{o}$C for microRNA analysis. The blood microRNAs in

the blood samples such as miR-125b, miR-155, miR-146a, miR-21, miR-20b,

miR-210, miR-122, and miR-33 were analyzed using a RT-qPCR array (Wcgene

Biotech, Inc., China). Total RNAs in the blood samples, including microRNAs, were

extracted using a Trizol reagent (Sigma Aldrich, Inc., USA) according to the

manufacturer's instructions. Subsequently, the purified RNAs were polyadenylated

through a poly(A) polymerase reaction and was then reversed-transcribed into

complementary DNA (cDNA). TIANGEN® miRcute Plus miRNA First-Strand cDNA

Kit (Code No. KR211) was used in the reverse transcriptional reaction system of total

10 μL, including 5 μL 2x miRNA RT Reaction Buffer, 1 μL miRNA RT Enzyme

Mix and 4 μL RNA sample. The reaction conditions are 40 $^{o}$C for 60 mins and 95 $^{o}$C

for 3 mins. The cDNA was quantified in real-time SYBR Green RT-qPCR reactions

with the specific microRNA qPCR Assay Primers. TIANGEN® miRcute Plus miRNA

qPCR Kit (SYBR Green) (Code No. FP411) was used in the qPCR reaction system of

total 10 μL, including 5 μL 2x miRcute Plus miRNA PreMix (SYBR&ROX),

0.2 μL Forward Primer, 0.2 μL Reverse Primer, 1 μL 50X ROX Reference Dye,

1 μL DNA Sample and 2.6 μL ddH$_2$O. The cycling conditions were 95 $^{o}$C for 15

min, followed by 40 cycles at 94 $^{o}$C for 20s, 60 $^{o}$C for 15s and 72$^{o}$C for 30s. The

primers used for qPCR are presented in Table S1 (Supporting Information).

**3. Statistical analysis**

In this study, the TVOC levels for all samples detected by the PID sensor were not normally distributed, while their TVOC change rates were. The TVOC change rate was calculated by dividing the TVOC level after exposure using the TVOC level before exposure. Therefore, the Mann-Whitney rank sum test was used to analyze the differences in TVOC levels before and after each toxicant exposure, and t-test was used to analyze the differences in TVOC change rates between each toxicant exposure group and the control group (indoor air). For individual VOC concentrations by GC-MS/FID, the paired t-test was used to analyze differences for each VOC species before and after the exposure. The software Canoco 4.5 was used to visualize the VOC profile distance and relatedness between the samples of different groups using the principal component analysis (PCA). Besides, the concentrations of micro-RNAs in blood samples from different rat groups were determined by RT-qPCR. For the group exposed to carbon dioxide, blood samples were only taken from two rats (before and after the 10-min exposure) because of the catheter blockage (unable to draw bloods) for the other two. For the other three groups, blood samples were obtained for all four rats. The differences between micro-RNA levels in blood samples before and after the exposure in one group were analyzed using a paired t-test (data exhibited a normal distribution) or Wilcoxon signed rank test (data did not follow a normal distribution). The outliers for normally distributed data were examined and eliminated by a Grubbs test. All the statistical tests were performed via the statistical component of SigmaPlot 12.5 (Systat Software, Inc., USA), and a *p-value* of less than 0.05 indicated a statistically significant difference at a confidence level of 95%.

**4. Results and discussion**

*4.1 TVOC monitoring for the toxicants exposure*

As described in the experimental section, four toxicants (ricin, endotoxin, ozone and carbon dioxide) and indoor air (as a background control) were used for inhalation exposure in rats. For each group, the TVOC levels of only 3 rats (PID instrument/software failure for one out of 4 rats) were shown in Figure 2. The indoor air background TVOC was found to be less than 0.04±0.02 ppm. After one rat was placed in the exposure chamber, the TVOC level in the cage was shown to first gradually increase, then reach a relatively stable level after about 500 seconds. The TVOC level before the exposure (rat + indoor air) when one rat was in the chamber was about 2 ppm, except for the $CO_2$ group it was about 0.5-0.8 ppm (These differences, if any, applied to both before and after exposure in one test, thus presenting no influences on the same experiments). The differences in TVOC levels for indoor air exposures (different times: "before" and "after", but the same indoor air) were small (the average change rate was about -4%±1.4% (95% confidence interval)), although the Mann-Whitney Rank Sum Test showed that for each of the rats, the difference (over some time for the indoor air) was significant (*p-value*<0.001). The background indoor air TVOC levels were 1.22-5.1% (detected) of the chamber TVOC with rats together with indoor air or other toxicants). The fluctuations were taken into account for each toxicant exposure test. Nonetheless, the fluctuations, if any, from background indoor air had minor impacts on the TVOC levels measured for rats' exposure tests. The change rates (n=3) of the control group (indoor air) then served as the reference for other toxicant exposures in the statistical analysis. During the indoor air experiment, the rats were seen to carry out normal life activities in the exposure

chamber, and correspondingly the TVOCs in the chamber were shown to remain

relatively stable.

**Figure 2.** Real-time continuous measurements of exhaled TVOC levels in the

chamber when rats were exposed to different toxicants via inhalation for 10 mins:

Indoor air, ricin, endotoxin, $O_3$ and $CO_2$. During the exposure processes, the PID

sensor was turned off. Data lines (measurement time was 1000 s) represent results

from three individual rats (#1, #2, #3) before or after exposure to each of the air

toxicants (aerosolized amounts described in the experimental section) tested. Each

exposure test was independently repeated with four rats from the same group (PID

instrument/software failure for one rat).

In contrast, the TVOC levels were shown to vary greatly with different toxicant

exposures as shown in Figure 2. For example, when rats were exposed to the

aerosolized ricin, the TVOC level was observed to exhibit an average change rate of

-3%±1.6. Compared to the control group (indoor air + rat before the exposure) shown

in Figure 2, the difference of the TVOC change rate was not significant for the ricin

exposure (t-test, *p-value*=0.426). For ricin exposure, its concentration (40 μg/mL

suspension aerosolized) might be too low in aerosol state after the aerosolization from

the liquid to produce a detectable response from the rats. It was previously reported

that 0.1 mg/mL ricin solution was used for aerosolization and subsequent exposure to

mice (Ref # 5, Supporting Information). After 30 minutes ricin aerosol exposure, the

exposed mice became poisoned. The concentration of ricin used in this experiment

was lower and the exposure time was shorter, therefore the toxicity reaction may be

mild. In contrast to the ricin exposure, however we observed a different phenomenon

for the endotoxin (50 ng/mL aerosolized) tests as shown in Figure 2. Upon the

endotoxin exposure, the TVOC level was observed to first increase, and then

decreased to a level that was about 21-46% below the pre-exposure level after four

minutes (Mann-Whitney Rank Sum Test, all *p-values*<0.001). Compared to the

control group (indoor air + rat), the difference of the TVOC change rate was

statistically significant for endotoxin (t-test, *p-value*=0.0147). The observed

differences from the ricin and endotoxin exposures could be due to different

mechanisms initiated by different substances involved. Ricin is derived from plant,

while endotoxin is from Gram-negative bacterial membrane. They could interact

differently with relevant human respiratory or other body cells. Nonetheless, for both

ricin and endotoxin, they were probably causing health effects by immuno-toxicity,

while $O_3$ and $CO_2$ both induce harm first by chemical manners.

After exposure to gaseous toxicants such as ozone and carbon dioxide, the levels

of TVOCs in the exposure chamber with rats were observed to have increased

significantly, as observed in Figure 2. As can be seen from the figure, the TVOC

levels has increased for about 44-110% for ozone and about 109-265% for carbon

dioxide exposure (Mann-Whitney Rank Sum Test, all *p-values*<0.001). The t-test

showed that differences of the TVOC change rates of both ozone and $CO_2$ exposures

compared to the control group (indoor air) were statistically significant

(*p-value*=0.0219 and 0.0296, respectively). These data indicated that rat exposure to

both ozone and $CO_2$ has resulted in significant elevations of TVOCs, suggesting rats

were actively responding to the exposure challenges. The behavior observation from a

video also indicated that rats after the exposure to $O_3$ seemed to be suffering from the

challenge (Video S1, Supporting Information).

*4.2 Changes in exhaled VOCs after exposure to different toxicants*

Using GC-MS/FID method, a total of 31 different VOCs out of 117 screened

were detected and shown in Figure S2 (Supporting Information) for background

indoor air with and without rats. Among detected VOCs as shown in Figure S2

(Supporting Information), the VOCs with the highest concentrations in indoor air

were n-hexane, ethyl acetate and acetone. When rats were placed in the exposure

chamber (one rat at each time), the most abundant VOC species was detected to be

acetone, which was about 4 times more than that of the indoor air background.

Statistical tests found that the concentrations of ethylene and ethane in the chamber

containing one rat were significantly lower than those of the background (paired t-test,

*p-value*<0.05), which in part could be due to the air dilution by the rat's breath.

Namely, when rats' breath with specific higher or lower VOC species replaced

equivalent indoor air inside the chamber, diluting effects for higher indoor VOC

species and enhancing effects for lower indoor VOC species could take place.

**Figure 3.** Differentiations of VOC species from indoor air and those from the rats'

exhaled breath under different air toxicity with exposure to ricin, $O_3$, endotoxin and

$CO_2$. The red lines show that the average change ratios of every toxicant calculated by

the level after the exposure divided by the level before exposure (right axis). The

dotted line is the baseline with a change ratio of 1. Percentages refer to specific VOC

percentage change before and after the exposure. Values represent averages and

standard deviations from three different rats. "*" indicates a significant difference at

*p-value*<0.05.

Differentiations of VOC species from the rat's exhaled breath under different

toxicants exposures were also shown in Figure 3. There were no significant

differences in the concentrations of any VOCs before and after the exposure for the

control group, i.e., indoor air (t-test, *p-value*=0.05). This suggests that indoor air is

relatively less toxic at a level that is unable to detect a VOC change. In contrast,

specific VOC species had experienced significant changes when rats were exposed to

ricin, endotoxin, $O_3$ and $CO_2$ as observed from Figure 3. For example, exposure to

ricin caused significant higher concentration of ethyl acetate (183% higher), while

lower concentration of carbon disulfide (22% lower). As shown in Figure 3, after the

endotoxin exposure process, concentrations of five VOC species: ethane, acetone,

cyclopentane, carbon disulfide and methylcyclopentane were shown to be

significantly different with those of before the exposure (t-test, all *p-values*<0.05). As

can be seen from the results of the ozone exposure group in Figure 3, the

concentrations of propionaldehyde, pentane, 2-butanone, hexane and 2-methylpentane

exhibited significant differences before and after exposure (t-test, all *p-values*<0.05),

in which all the VOCs except 2-methylpentane were elevated. In comparison, rat

exposure to $CO_2$ resulted in acetone level increase by 34% (t-test, *p-value*=0.0016).

These data suggest that exposure to different toxicants had led to production of

different profiles of VOC species in addition to their level changes.

***4.3 Detection of micro RNAs in Blood Samples***

To further explore the VOC response mechanism of rats to toxicants exposure,

microRNAs (miRNA) in the blood samples were examined by an RT-qPCR assay.

Fold-changes in microRNA levels after toxicants exposure were shown in Table S2

(Supporting Information). The level of miR-33 in the blood of rats was shown to be

significantly lower than that before ricin exposure (*p-value*<0.05); after exposure to

ozone, miR-146a level in the blood samples of rats was significantly higher than those

before the exposure (*p-value* <0.05), while miR-155 was significantly lower than that

before the exposure (*p-value* <0.05). For other microRNAs as listed in Table S2, the

changes seemed to be insignificant (t-test, *p-values*>0.05).

As observed from Figure 4, PCA results revealed a clear contrast in breath-borne

VOC profiles of rats between different toxicants exposures. The VOC profiles of the

ozone exposure group was very different from that of the control group (indoor air)

and the VOCs profile of the ricin exposure group was the closest to that of the control

group, which agreed with TVOC level and VOC species profiles obtained above.

Overall, the experiments showed that rats responded differently to different toxicants

by releasing different VOC species owing to different mechanisms of toxicity: ozone

caused significant increases in various breath-borne VOCs; while endotoxin exposure

generally decreased the releasing of VOCs; and ricin and carbon dioxide exposure

resulted in one or two significant VOC species changes. In general, the results of

qualitative and quantitative analysis by the GC-MS/FID method agreed with the

TVOC level monitored by the PID sensor.

**Figure 4.** PCA ordinations of exhaled breath-borne VOCs profiles under exposures to different toxicants: ricin, endotoxin, $O_3$, $CO_2$ and control (indoor air). PC1 and PC2 are the first and second principal components. The VOCs species involved in the PCA analysis were the 12 species which were shown to have undergone changes after each toxicant exposure. Data presented in the figure were from three independent rats exposed to each toxicant.

As observed from Figure 3, exposure to ricin caused 183±143 % higher concentration of ethyl acetate, while 22±8 % lower concentration of carbon disulfide. It was previously reported the concentration of ethyl acetate was significantly higher in exhaled breath from people with cancer compared to the healthy group.[40,41] In addition, *in vitro* experiments have shown the human umbilical vein endothelial cells (HUVEC) can produce ethyl acetate, which is presumably generated by a reaction of ethanol with acetic acid.[42] It was demonstrated that ricin is not only responsible for the ricin intoxication through ribosomal inactivation and subsequent inhibition of protein synthesis and cell death, but also shows endothelial toxicity by acting as a natural disintegrin binding to and damaging human endothelial cells.[43] Therefore, the

toxicity of ricin on the endothelial cells might be the source of the higher

concentration of ethyl acetate observed in this work. As a disease biomarker, carbon

disulfide was observed in the exhaled breath.[44-45] Recently, it was suggested that the

carbon disulfide may be generated endogenously and play a role as a bioregulator. [46]

Here, we observed that exposure to both ricin and endotoxin resulted in lower levels

of breath-borne carbon disulfide compared to the control.

For $CO_2$ and endotoxin exposure, the observation for acetone was the opposite as

shown in Figure 3. Acetone in exhaled breath was widely investigated in many studies

as an important biomarker related to blood glucose and diabetes.[34,47] Acetone is

produced in the fatty acids metabolism by hepatocytes via decarboxylation of excess

acetyl coenzyme A (Acetyl–CoA), and then oxidized via the Krebs cycle in peripheral

tissue.[48] As shown in Figure 3, the acetone level increased by 34±9% as a result of

$CO_2$ exposure, suggesting $CO_2$ caused hypoxia in rats, and led to increased respiration

from rats. These increases in acetone level corresponded to TVOC level increase as

determined by the PID sensor after the exposure to $CO_2$. However, when exposed to

endotoxin, the acetone level in the exposure chamber decreased by about 10±6%,

indicating that the respiration of the rats may be attenuated by the exposure of

endotoxin. Clearly, the involved mechanisms by which endotoxin and $CO_2$ cause

health effects to rats could be very different.

As observed in Figure 3, the increase of ethane level by endotoxin exposure

suggested that lipid damage was induced by oxidative stress in the rat's body since

ethane is acknowledged as a marker of lipid peroxidation and described to be

generated by peroxidation of ω-3 polyunsaturated fatty acids.[33] In addition to ethane,

the increase of methylcyclopentane level as shown in Figure 3 might also be the result

of the endotoxin exposure. Endotoxin has been shown to trigger inflammation through

its interaction with the TLR4/CD14/MD2 receptor and then initiates a signal cascade.

This reaction correspondingly results in the activation of transcription factor such as

NF-κB leading to the production of pro-inflammatory cytokines and type 1 interferons

(IFNs), and finally results in systemic inflammatory response syndrome.[49] In general,

in terms with the average fold changes, the concentration of total VOCs in the exhaled

breath was relatively lower after the endotoxin exposure, which agreed with the

results of TVOC obtained by the PID sensors.

In addition to these biologicals, we have also shown that exposure to chemicals

such as ozone and $CO_2$ also resulted in in-vivo changes in VOC levels. From the fold

changes of various VOC species such as propionaldehyde, N-pentane, 2-Butanone,

and Hexane, the ozone exposure has resulted in an overall increase of VOCs in rats'

exhaled breath, which agreed with the TOVC monitoring shown in Figure 2 by the

PID sensors. It was previously indicated that increase in propionaldehydes, further

products of lipid peroxidation, indicated more severe oxidative damage in rats

following exposure to ozone.[50] Here, we also showed that after exposure to ozone

changes in specific breath-borne VOCs and also certain miRNA regulations were

detected. Ozone was described as a strong oxidizing agent, and can cause intracellular

oxidative stress through ozonide and hydroperoxide formation.[51] The mechanism of

ozone oxidative damage involves the activation of Nrf2, heat shock protein 70, and

NF-κB, thus increasing expression of a range of proinflammatory cytokines such as

TNFα and interleukin 1β, and chemokines such as interleukin 8.[52] The results above

show that regardless of toxicant types breath-borne VOCs from rats experienced *in*

*vivo* changes.

Results of miRNAs from rats' blood as shown in Table S4 revealed different

mechanisms by rats when exposed to various types of toxicants. miRNAs are short

non-coding RNA sequences that regulate gene expression at the posttranscriptional

level; and many miRNAs have already been identified to influence physiological

processes such as immune reaction, adaptation to stress, and widely investigated in

environmental exposure studies.[53] Among these microRNAs, miR-125b, miR-155,

miR-146a, and miR-21 are mainly shown to regulate oxidative stress and

inflammatory processes *in vivo*, and widely investigated in air pollution related

studies.[53-54] For example, among them, miR-155 has a positive regulation function,

and the other three are negative regulation.[55-56] However, in this study, the ozone

exposure resulted in decreased levels of miR-155 (t-test, *p-value*<0.05), which are

different from previous reports in the literature. [55] The difference could be due to

different exposure toxicant, i.e., ozone used here, leading to acute lung damages

compared to mild airway inflammation or asthma problems in other studies. Previous

studies have shown that miR-20b and miR-210 are hypoxia regulators in animals, and

miR-122 and miR-33 are mainly responsible for regulating lipid metabolism and

glucose metabolism in the body.[57-58] However, these microRNAs, except for the

decreased level of miR-33 and increased level of miR-146a caused by ricin and ozone

exposure, respectively, were shown to have not undergone significant changes in this

study after the exposures to four different toxicants (*p-values*=0.05). The possible

reason may be that microRNAs act as post-transcriptional regulators by degrading

mRNA or inhibiting their translation, thus failing to respond in a timely manner

during short-term exposure (blood samples taken 20 min after the 10-min exposure ).

To further understand the problem, a yeast model is currently being used to fully

investigate the mechanisms underlying the miRNA regulation when exposed to the

toxicants used here. Nonetheless, these results here reflected that exposure to the

toxicants led to specific miRNAs either up- or down-regulated. On the other hand, the

results here also suggest, especially for those no changes observed for the miRNAs,

higher toxicant level or specific longer time might be needed to allow miRNA

regulation change to occur.

To address the major objective of this work, we repeated a total of 20 times using

5 different exposure agents such as ozone, $CO_2$, ricin, endotoxin together with the

background indoor air as a control, and 20 different rats. Clearly, above results

indicate that when rats are exposed to toxic substances their certain metabolic

activities are immediately affected, i.e., these exposures promoted or inhibited

specific VOC productions. Based on the results we obtained from this work, the

following VOC emission mechanisms of rats when exposed to different toxicants are

proposed and illustrated in Figure 5. Previously, it was suggested that VOCs are

produced during the normal metabolism in the body; while pathological processes,

such as metabolic disorders, can also produce new species of VOCs or alter the levels

of existing VOCs.[59] Therefore, cell or tissue injuries caused by external toxicants

exposure also can alter the exhaled VOCs profile by disturbing the normal process.[60]

The exact toxic effect mechanism as observed from this work could vary from one

toxicant to another. For some pollutants such as endotoxin and ricin, there are specific

receptors to recognize them and then start the chain of responses or reactions.[49,61]

Among these various mechanisms, the ROS (reactive oxygen species) and oxidative

stress are recognized to be the central and the common mechanism in various forms of

pathophysiology, as well as the health effect of various air pollutants including

ambient particulate matter (PM). [1, 62-63] Oxidative stress is essentially a compensatory

state of the body and can trigger redox-sensitive pathways leading to different

biological processes such as inflammation and cell death.[51,64] For example, the strong

oxidants such as ozone might cause oxidative stress through direct effects on lipids

and protein[65-66], which mostly caused the generation and release of hydrocarbons and

aldehydes, such as ethane, ethylene, and propionaldehyde.[50,67] While carbon dioxide

tends to make the redox balance tilted toward the reduction side by reducing oxygen

supply and thus influencing the energy metabolism in cells.[68-70] For ricin and

endotoxin exposure, the underlying mechanisms seem to be different from ozone and

$CO_2$, and they could cause oxidative stress indirectly through the activation of

intracellular oxidant pathways. Nonetheless, all toxicants share a common effect of

disrupting the redox balance, and thus interfere with normal biochemical reactions or

cause material damages in cells, accordingly changing the VOC profile released into

the breath. As discussed above, in this work, the VOCs profile of rats changed

significantly after exposure to different toxicants. Therefore, regardless of toxicant

types, breath-borne VOCs from the rats seem to be capable of serving as a proxy for

real-time monitoring air toxicity. It was reported about 100 years ago that three mice

were also carried on all British submarines for sensing small leakage of gasoline as

the rats could squeak to notify the crew (Figure S3, Supporting Information). Some

previously detected VOCs such as hexane, pentane, acetaldehyde, butanone, and

acetone from rat's exhaled breath [71, 72] were also detected in this work. Nonetheless,

due to variations in rat breed, experimental conditions, VOC sampling and analysis

methods, it is rather difficult to exactly compare the VOCs from rat's exhaled breath

across different studies. There might be some breath-borne VOC species not identified

yet for the rats used here.   Interestingly, rat L6 skeletal muscle cells cultivated with

α-MEM containing 10% FCS were shown to release 7 VOC species and uptake 16

VOC species.[73] Similarly, the increased VOCs detected here from rat's exhaled breath

might be released from lung cells or other impacted ones by the exposure. On another

front, previous work showed that changes were also detected in breath-borne VOCS

from people with upper respiratory infections compared to the health individuals (Ref

#7, Supporting Information). All these studies and data support the results and validity

of our work from various aspects. For improving $RST_{air}$ performance, GC-MS can be

also replaced using the Proton Transfer Reaction-Mass Spectrometry (PTR-MS) for

fast online VOC species analysis. By using this discovered fundamental science, the

invented *RSTair* system here showed its great promise of revolutionizing the air

toxicity monitoring, and providing significant technological advances for air security

in related fields such as military defense, customs, counter-terrorism and security

assurances for important events or special locations.

**Figure 5.** Proposed mechanisms of toxic effects and VOCs releasing in rats when exposed to the environmental toxicants via inhalation. The black arrows represent the toxic effects of different toxicants and the possible pathways of VOCs generation. The blue arrows stand for the principle and working process of the invented *RST*air system for real-time air toxicity monitoring. The corresponding references cited are: ① [59]; ② [66]; ③ [63]; ④ [48]; ⑤ [42]; ⑥ [48, 70].

**Supporting Information**

Measurements of exhaled VOCs by PID and GC-MS/FID, Selection of exposure levels for different toxicant, experimental setup for *RSTair* system, indoor air background VOCs, Photo of UK postcard for three mice carried on British submarines, Primers used for RT-qPCR analysis of microRNA and mi-RNA expression level changes after exposure, video of rat after exposed to ozone were

provided as Supporting Information.

**Acknowledgements**

This study was supported by the NSFC Distinguished Young Scholars Fund

Awarded to M. Yao (21725701), and the Ministry of Science and Technology

(2016YFC0207102).

**Conflict of interests**

The authors declare no competing financial interest.

**References**

1.   Brook, R. D.; Rajagopalan, S.; Pope, C. A., III.; Brook, J. R.; Bhatnagar, A.;

Diez-Roux, A., V.; Holguin, F.; Hong Y.; Luepker, R., V.; Mittleman, M. A.; Peters, A.;

Siscovick, D.; Smith, S. C., Jr.; Whitsel, L.; Kaufman, J. D. , Particulate matter air

pollution and cardiovascular disease an update to the scientific statement from the

American Heart Association. *Circulation* **2010,** *121* (21), 2331-2378.

2.   Cohen, A. J.; Brauer, M.; Burnett, R.; Anderson, H.  R.; Frostad, J.; Estep, K.;

Balakrishnan, K.; Brunekreef, B.; Dandona, L. Dandona, R.; Feigin, V.; Freedman, G.;

Hubbell, B.; Jobling, A.; Kan, H.; Knibbs, L.; Liu, Y.; Martin, R.; Morawska, L.; C

Pope, C. A., III.; Shin, H.; Straif, K.; Shaddick, G.; Thomas, M.; Dingenen, R. v.;

Donkelaar, A. v.; Vos, T.; Murray, C. J. L.; Forouzanfar, M. H.  Estimates and

25-year trends of the global burden of disease attributable to ambient air pollution: an

analysis of data from the Global Burden of Diseases Study 2015. *Lancet* **2017,** *389*

(10082), 1907.

3.  GBD 2015 Risk Factors Collaborators.  Global, regional, and national

comparative risk assessment of 79 behavioural, environmental and occupational, and

metabolic risks or clusters of risks, 1990-2015: a systematic analysis for the Global

Burden of Disease Study 2015. *Lancet* **2016,** *388* (10053), 1659-1724.

4.   EPA, U. S., Integrated Science Assessment (ISA) of Ozone and Related

Photochemical Oxidants (Final Report, Feb 2013). Agency, U. S. E. P., Ed.

Washington, DC, 2013; Vol. EPA/600/R-10/076F.

5.   Jerrett, M.; Burnett, R. T.; Pope, C. A. III.; Ito, K.; Thurston, G.; Krewski,  D.;

Shi, Y.; Calle, E.; Thun, M. Long-term ozone exposure and mortality. *N. Engl. J. Med.*

**2009,** *360* (11), 1085-1095.

6.   Thompson, W. W.; Shay, D. K.; Weintraub, E.;Brammer, L.; Cox, N.; Anderson,

L. J.; Fukuda, K. Mortality associated with influenza and respiratory syncytial virus in the United States. *JAMA* **2003,** *289* (2), 179-186.

7.   Jones, R. M.; Brosseau, L. M. Aerosol transmission of infectious disease. *J. Occup. Environ. Med.* **2015,** *57* (5), 501-508.

8.   Fernstrom, A.; Goldblatt, M. Aerobiology and its role in the transmission of infectious diseases. *J Pathog* **2013,** Article ID: 493960, doi: http://dx.doi.org/10.1155/2013/493960.

9.   Yu, I. T. S.; Li, Y.; Wong, T. W.; Tam, W.; Chan, A. T.; Lee, J. H.W.; Leung, D. Y. C.; Ho, T. Evidence of airborne transmission of the Severe Acute Respiratory Syndrome Virus. *N. Engl. J. Med.* **2004,** *350* (17), 1731-1739.

10. Green, M. S.; LeDuc, J.; Cohen, D.; Franz, D. R. Confronting the threat of bioterrorism: realities, challenges, and defensive strategies. *Lancet Infect Dis* **2019,** *19* (1), e2-e13.

11. Okumura, T.; Takasu, N.; Ishimatsu, S. Miyanoki, S.; Mitsuhashi, A.; Kumada. K.; Tanaka, K.; Hinohara, S. Report on 640 Victims of the Tokyo subway Sarin attack. *Ann. Emerg. Med.* **1996,** *28* (2), 129-135.

12. Jernigan, D. B.; Raghunathan, P. L.; Bell, B. P.; Brechner, R.; Bresnitz, E. A.; Butler, J. C.; Cetron, M.; Cohen, M.; Doyle, T.; Fischer, M.; Greene, C.; Griffith, K. S.; Guarner, J.; Hadler, J. L.; Hayslett, J. A.; Meyer, R.; Petersen, L. R.; Phillips, M.; Pinner, R.; Popovic, T.; Quinn, C. P.; Reefhuis, J.; Reissman, D.; Rosenstein, N.; Schuchat, A.; Shieh, W.; Siegal, L.; Swerdlow, D. L.; Tenover, F. C.; Traeger, M.; Ward, J. W. Weisfuse, I.; Wiersma, S.; Yeskey, K.; Zaki, S.; Ashford, D. A. Perkins, B. A. Ostroff, S.; Hughes, J.; Fleming, D.; Koplan, J. P.; Gerberding, J. L. and the National Anthrax Epidemiologic Investigation Team. Investigation of bioterrorism-related anthrax, United States, 2001: epidemiologic findings. *Emerg.*

*Infect. Dis.* **2002,** *8* (10), 1019.

13. Marć, M.; Tobiszewski, M.; Zabiegała, B.; Guardia, M. d. l.; Namieśnik, J. Current air quality analytics and monitoring: A review. *Anal. Chim. Acta* **2015,** *853*, 116-126.

14. Sameenoi, Y.; Koehler, K.; Shapiro, J.; Boonsong, K.; Sun, Y.; Collett, J. Jr.; Volckens, J.; Henry, C. S. Microfluidic electrochemical sensor for on-line monitoring of aerosol oxidative activity. *J. Am. Chem. Soc.* **2012,** *134* (25), 10562-10568.

15. Ishihara, S.; Azzarelli, J. M.; Krikorian, M.; Swager, T. M. Ultratrace detection of toxic chemicals: triggered disassembly of supramolecular nanotube wrappers. *J. Am. Chem. Soc.* **2016,** *138* (26), 8221-8227.

16. Azzarelli, J. M.; Mirica, K. A.; Ravnsbæk, J. B.; Swager, T. M. Wireless gas detection with a smartphone via rf communication. *Proc. Natl. Acad. Sci.* **2014,** *111* (51), 18162.

17. Jung, J. H.; Park, S. Y.; Lee, J. E.; Lee, B. U.; Bae, G. N. Distinguishing biotic and abiotic particles using an ultraviolet aerodynamic particle sizer for real-time detection of bacterial bioaerosols. *Environ. Eng. Sci.* **2012,** *29* (9), 866-874.

18. Schwarzmeier, K.; Knauer, M.; Ivleva, N. P.; Niessner, R.; Haisch, C. Bioaerosol analysis based on a label-free microarray readout method using surface-enhanced Raman scattering. *Anal. Bioanal. Chem.* **2013,** *405* (16), 5387-5392.

19. Shen, F.; Wang, J.; Xu, Z.; Wu, Y.; Chen, Q.; Li, X.; Jie, X.; Li, L.; Yao, M.; Guo, X.; Zhu, T. Rapid flu diagnosis using silicon nanowire sensor. *Nano Lett.* **2012,** *12* (7), 3722-3730.

20. Jin, L.; Xie, J.; Wong, C. K. C.; Chan, S. K.; Abbaszade, G.; Schnelle-Kreis, J.; Zimmermann R.; Li, J.; Zhang, G.; Fu, P.; Li, X. Contributions of city-specific fine particulate matter ($PM_{2.5}$) to differential in vitro oxidative stress and toxicity

implications between Beijing and Guangzhou of China. *Environ. Sci. Technol.* **2019,** *53* (5), 2881-2891.

21. Li, J.; Chen, H.; Li, X.; Wang, M.; Zhang, X.; Cao, J.; Shen, F.; Wu, Y.; Xu, S.; Fan, H.; Da, G.; Huang R.; Wang, J.; Chan, C. K.; Jesus, A. L. D.; Morawska, L.; Yao, M. Differing toxicity of ambient particulate matter (PM) in global cities. *Atmos. Environ.* **2019,** *212*, 305-315.

22. Pleil, J. D. Breath biomarkers in toxicology. *Arch. Toxicol.* **2016,** *90* (11), 2669-2682.

23. Council, N. R. *Sensor systems for biological agent attacks: protecting buildings and military bases*; 030909576X; National Academies Press: 2005.

24. Kida, H.; Fukutani, Y.; Mainland, J. D.; Claire, A.; Vihani, A.; Li, Y. R.; Chi, Q.; Toyama, A.; Liu, L.; Kameda, M.; Yohda, M.; Matsunami, H. Vapor detection and discrimination with a panel of odorant receptors. *Nat Commun* **2018,** *9* (1), 4556.

25. Rider, T. H.; Petrovick, M. S.; Nargi, F. E.; Harper, J. D.; Schwoebel, E. D.; Mathews, R. H.; Blanchard, D. J.; Bortolin, L. T.; Young, A. M.; Chen, J.; Hollis, M. A. A B cell-based sensor for rapid identification of pathogens. *Science* **2003,** *301* (5630), 213.

26. Engel, Y.; Elnathan, R.; Pevzner, A.; Davidi, G.; Flaxer, E.; Patolsky, F. Supersensitive detection of explosives by silicon nanowire arrays. *Angew. Chem. Int. Ed.* **2010,** *49* (38), 6830-6835.

27. Chen, H.; Li, J.; Zhang, X.; Li, X.; Yao, M.; Zheng, G. Automated in vivo nanosensing of breath-borne protein biomarkers. *Nano Lett.* **2018,** *18* (8), 4716-4726.

28. Vereb, H.; Dietrich, A. M.; Alfeeli, B.; Agah, M. The possibilities will take your breath away: breath analysis for assessing environmental exposure. *Environ. Sci. Technol.* **2011,** *45* (19), 8167-8175.

29. Haick, H.; Broza, Y. Y.; Mochalski, P.; Ruzsanyi, V.; Amann, A. Assessment,

origin, and implementation of breath volatile cancer markers. *Chem. Soc. Rev.* **2014,**

*43* (5), 1423-1449.

30. Li, X.; Sinues, P. M. L.; Dallmann, R.; Bregy, L.; Hollmén, M.; Proulx, S.; Brown,

A. S.; Detmar, M.; Kohler, M.; Zenobi, R. Drug pharmacokinetics determined by

real-time analysis of mouse breath. *Angew. Chem. Int. Edit.* **2015,** *54* (27), 7815-7818.

31. Broza, Y. Y.; Mochalski, P.; Ruzsanyi, V. Amann, A.; Haick, H. Hybrid

volatolomics and disease detection. *Angew. Chem. Int. Edit.* **2015,** *54* (38),

11036-11048.

32. Wang, F.; Swager, T. M. Diverse chemiresistors based upon covalently modified

multiwalled carbon nanotubes. *J. Am. Chem. Soc.* **2011,** *133* (29), 11181-11193.

33. Aghdassi, E.; Allard, J. P. Breath alkanes as a marker of oxidative stress in

different clinical conditions. *Free Radic. Biol. Med.* **2000,** *28* (6), 880-886.

34. Righettoni, M.; Tricoli, A.; Pratsinis, S. E. Si:WO3 sensors for highly selective

detection of acetone for easy diagnosis of diabetes by breath analysis. *Anal. Chem.*

**2010,** *82* (9), 3581-3587.

35. Peng, G.; Tisch, U.; Adams, O.; Hakim, M.; Shehada, N.; Broza, Y. Y.; Billan,

S.; Abdah-Bortnyak, R.; Kuten, A. Haick, H. Diagnosing lung cancer in exhaled

breath using gold nanoparticles. *Nat Nanotechnol* **2009,** *4* (10), 669.

36. Fens, N.; Zwinderman, A. H.; Schee, M. P. v. d.; Nijs, S. B. d,; Dijkers, E.;

Roldaan, A. C.; Cheung, D.; Bel, E. H.; Sterk, P. J. Exhaled breath profiling enables

discrimination of chronic obstructive pulmonary disease and asthma. *Am. J. Respir.*

*Crit. Care Med.* **2009,** *180* (11), 1076-1082.

37. Ahmed, W. M.; Lawal, O.; Nijsen, T. M.; Goodacre, R.; Fowler, S. J. Exhaled

volatile organic compounds of infection: a systematic review. *ACS Infect Dis* **2017,** *3*

(10), 695-710.

38. Zhang, X.; Kang, J.; Chen, H.; Yao, M.; Wang, J. $PM_{2.5}$ meets blood: in vivo

damages and immune defense. *Aerosol Air Qual Res* **2018,** *18*, 456-470.

39. Wei, K.; Li, W.; Gao, S.; Ji, B.; Zang, Y.; Su, B.; Wang, K.; Yao, M.; Zhang, J.;

Wang, J. Inactivation of ricin toxin by nanosecond pulsed electric fields including

evidences from cell and animal toxicity. *Sci. Rep.* **2016,** *6*, 18781.

40. Buszewski, B.; Ligor, T.; Jezierski, T.; Wenda-Piesik, A.; Walczak, M.; Rudnicka,

J. Identification of volatile lung cancer markers by gas chromatography–mass

spectrometry: comparison with discrimination by canines. *Anal. Bioanal. Chem.* **2012,**

*404* (1), 141-146.

41. Amal, H.; Leja, M.; Funka, K.; Lasina, I.; Skapars, R.; Sivins, A.; Ancans, G.;

Kikuste, I.; Vanags, A.; Tolmanis, I.; Kirsners, A.; Kupcinskas, L.; Haick, H. Breath

testing as potential colorectal cancer screening tool. *Int. J. Cancer* **2016,** *138* (1),

229-236.

42. Mochalski, P.; Theurl, M.; Sponring, A.; Unterkofler, K.; Kirchmair, R.; Amann,

A. Analysis of volatile organic compounds liberated and metabolised by human

umbilical vein endothelial cells (HUVEC) in vitro. *Cell Biochem. Biophys.* **2015,** *71*

(1), 323-329.

43. Machesky, N. J.; Rusnak, J. M.; Moore, E. H.; Dorsey, C. B.; Ward, L. A.

Development of a highly sensitive in vitro endothelial cell toxicity assay for

evaluating ricin toxin A chain-based vaccines or therapeutics. *Toxicon* **2019,** *167*,

152-161.

44. Fernández del Río, R.; O'Hara, M. E.; Holt, A.; Pemberton, P.; Shah, T.;

Whitehouse, T.; Mayhew, C. A. Volatile biomarkers in breath associated with liver

cirrhosis — comparisons of pre- and post-liver transplant breath samples.

*EBioMedicine* **2015,** *2* (9), 1243-1250.

45. Daniel, D. A. P.; Thangavel, K. Breathomics for gastric cancer classification

using back-propagation neural network. *J Med Signals Sens* **2016,** *6* (3), 172-182.

46. DeMartino, A. W.; Zigler, D. F.; Fukuto, J. M.; Ford, P. C. Carbon disulfide. Just

toxic or also bioregulatory and/or therapeutic? *Chem. Soc. Rev.* **2017,** *46* (1), 21-39.

47. Cao, W.; Duan, Y. Breath Analysis: Potential for clinical diagnosis and exposure

assessment. *Clin. Chem.* **2006,** *52* (5), 800-811.

48. Miekisch, W.; Schubert, J. K.; Noeldge-Schomburg, G. F. E. Diagnostic potential

of breath analysis—focus on volatile organic compounds. *Clin. Chim. Acta* **2004,** *347*

(1), 25-39.

49. Wang, X.; Quinn, P. J. *Endotoxins: structure, function and recognition*. Springer

Science & Business Media: 2010; Vol. 53.

50. Traxler, S.; Bischoff, A.-C.; Saß, R.; Trefz, P.; Gierschner, P.; Brock, B.;

Schwaiger, T.; Karte, C.; Blohm, U.; Schröder, C.; Miekisch, W.; Schubert, J. K. VOC

breath profile in spontaneously breathing awake swine during Influenza A infection.

*Sci. Rep.* **2018,** *8* (1), 14857-14857.

51. Lodovici, M.; Bigagli, E. Oxidative stress and air pollution exposure. *J. Toxicol.*

**2011,** *2011*, 487074-487074.

52. Nichols, B. G.; Woods, J. S.; Luchtel, D. L.; Corral, J.; Koenig, J. Q. Effects of

ozone exposure on nuclear factor-κB activation and tumor necrosis factor-α

expression in human nasal epithelial cells. *Toxicol. Sci.* **2001,** *60* (2), 356-362.

53. Vrijens, K.; Bollati, V.; Nawrot Tim, S. MicroRNAs as potential signatures of

environmental exposure or effect: a systematic review. *Environ. Health Perspect.*

**2015,** *123* (5), 399-411.

54. Wei, J.; Li, F.; Yang, J.; Liu, X.; Cho, W. C. MicroRNAs as regulators of airborne

pollution-induced lung inflammation and carcinogenesis. *Arch. Toxicol.* **2015,** *89* (5), 677-685.

55. Deshpande, D.; Dileepan, M.; Walseth, T.; Subramanian, S.; Kannan, M. S. MicroRNA regulation of airway inflammation and airway smooth muscle function: relevance to asthma. *Drug Dev. Res.* **2015,** *76* (6), 286-295.

56. Kong, A. P. S.; Xiao, K.; Choi, K. C.; Wang, G.; Chan, M. H.; Ho, C. S.; Chan, I.; Wong, C. K.; Chan, J. C.N.; Szeto, C. C. Associations between microRNA (miR-21, 126, 155 and 221), albuminuria and heavy metals in Hong Kong Chinese adolescents. *Clin. Chim. Acta* **2012,** *413* (13), 1053-1057.

57. Huang, X.; Le, Q.-T.; Giaccia, A. J. MiR-210 – micromanager of the hypoxia pathway. *Trends Mol. Med.* **2010,** *16* (5), 230-237.

58. Aranda, J. F.; Madrigal-Matute, J.; Rotllan, N.; Fernández-Hernando, C. MicroRNA modulation of lipid metabolism and oxidative stress in cardiometabolic diseases. *Free Radic. Biol. Med.* **2013,** *64*, 31-39.

59. Hakim, M.; Broza, Y. Y.; Barash, O.; Peled, N.; Phillips, M.; Amann, A.; Haick, H. Volatileorganic compounds of lung cancer and possible biochemical pathways. *Chem. Rev.* **2012,** *112* (11), 5949-5966.

60. Filipiak, W., Mochalski, P., Filipiak, A.; Ager, C.; Cumeras, R.; Davis, E. C.; Agapiou, A.; Unterkofler, K.; Troppmair, J. A compendium of volatile organic compounds (VOCs) released by human cell lines. *Curr. Med. Chem.* **2016**, 23(20), 2112-2131.

61. Lord, M. J.; Jolliffe, N. A.; Marsden, C. J.; Pateman, C. S.C.; Smith, D. C.; Spooner, R. A.; Watson, P. D.; Roberts, L. M. Ricin: mechanisms of cytotoxicity. *Toxicol. Rev.* **2003,** *22* (1), 53-64.

62. Hybertson, B. M.; Gao, B.; Bose, S. K.; McCord, J. M. Oxidative stress in health

and disease: the therapeutic potential of Nrf2 activation. *Mol. Aspects Med.* **2011,** *32*

(4), 234-246.

63. Maritim, A. C.; Sanders, R. A.; Watkins Iii, J. B. Diabetes, oxidative stress, and

antioxidants: A review. *J. Biochem. Mol. Toxicol.* **2003,** *17* (1), 24-38.

64. Birben, E.; Sahiner, U. M.; Sackesen, C.; Erzurum, S.; Kalayci, O. Oxidative

stress and antioxidant defense. *World Allergy Organ J* **2012,** *5* (1), 9-19.

65. Kosmider, B.; Loader, J. E.; Murphy, R. C.; Mason, R. J. Apoptosis induced by

ozone and oxysterols in human alveolar epithelial cells. *Free Radic. Biol. Med.* **2010,**

*48* (11), 1513-1524.

66. Filipiak, W.; Ruzsanyi, V.; Mochalski, P.; Filipiak, A.; Bajtarevic, A.; Ager, C.;

Denz, H.; Hilbe, W.; Jamnig, H.; Hackl, M.; Dzien, A. Dependence of exhaled breath

composition on exogenous factors, smoking habits and exposure to air pollutants. *J.*

*Breath Res.* **2012,** *6* (3), 21.

67. Amann, A.; Costello, B. d. L.; Miekisch, W.; Schubert, J.; Buszewski, B.; Pleil, J.;

Ratcliffe, N.; Risby, T. The human volatilome: volatile organic compounds (VOCs) in

exhaled breath, skin emanations, urine, feces and saliva. *J. Breath Res.* **2014,** *8* (3),

034001.

68. Vander Heiden, M. G.; Cantley, L. C.; Thompson, C. B. Understanding the

Warburg effect: the metabolic requirements of cell proliferation. *Science (New York,*

*N.Y.)* **2009,** *324* (5930), 1029-1033.

69. Hochachka, P. W. Defense strategies against hypoxia and hypothermia. *Science*

**1986,** *231* (4735), 234.

70. Laffel, L. Ketone bodies: a review of physiology, pathophysiology and

application of monitoring to diabetes. *Diabetes Metab. Res. Rev.* **1999,** *15* (6),

412-426.

71. Wlodzimirow, K.A., Abu-Hanna, A.; Schultz, M. J.; Maas, M. A. W.; Bos, L. D. J.;

Sterk, P. J.; Knobele, H.H.; Soers, R.J.T.; Chamuleau, R. A. F. M., Exhaled breath

analysis with electronic nose technology for detection of acute liver failure in rats.

*Biosens. Bioelectron.* **2013**, 53, 129-134.

72. Simone M., Kyu-Bok J., Barbara B.; Blake, D. R.; Vaziri, N. D., Exhaled breath

and fecal volatile organic biomarkers of chronic kidney disease. *Biochim. Biophys.*

*Acta, Gen. Subj.* **2013**, 1830(3), 2531-2537.

73. Mochalski P.; Al-Zoairy R.; Niederwanger A.; Unterkofler K.; Amann A.,

Quantitative analysis of volatile organic compounds released and consumed by rat L6

skeletal muscle cells in vitro. *J. Breath Res.* **2014**, 8(4), 046003.