Spherical deconvolution with tissue-specific response functions and multi-shell diffusion MRI to estimate multiple fiber orientation distributions (mFODs)

Alberto De Luca[1], Fenghua Guo[1], Martijn Froeling[2], Alexander Leemans[1]

[1]Image Sciences Institute, UMC Utrecht, Utrecht, the Netherlands
[2]Department of Radiology, UMC Utrecht, Utrecht, the Netherlands

Corresponding author: Alberto De Luca
Heidelberglaan 100, 3584CX
Utrecht, the Netherlands

Submitted to Neuroimage

Abstract

In diffusion MRI, spherical deconvolution approaches can estimate local white matter (WM) fiber orientation distributions (FOD) which can be used to produce fiber tractography reconstructions. The applicability of spherical deconvolution to grey matter (GM), however, is still limited, despite its critical role as start/endpoint of WM fiber pathways. The advent of multi-shell diffusion MRI data offers additional contrast to model the GM signal but, to date, only isotropic models have been applied to GM. Evidence from both histology and high-resolution diffusion MRI studies suggests a marked anisotropic character of the diffusion process in GM, which could be exploited to improve the description of the cortical organization. In this study, we investigated whether performing spherical deconvolution with tissue specific models of both WM and GM can improve the characterization of the latter while retaining state-of-the-art performances in WM. To this end, we developed a framework able to simultaneously accommodate multiple anisotropic response functions to estimate multiple, tissue-specific, fiber orientation distributions (mFODs). As proof of principle, we used the diffusion kurtosis imaging model to represent the WM signal, and the neurite orientation dispersion and density imaging (NODDI) model to represent the GM signal. The feasibility of the proposed approach is shown with numerical simulations and with data from the Human Connectome Project (HCP). The performance of our method is compared to the current state of the art, multi-shell constrained spherical deconvolution (MSCSD). The simulations show that with our new method we can accurately estimate a mixture of two FODs at SNR$\geq$50. With HCP data, the proposed method was able to reconstruct both tangentially and radially oriented FODs in GM, and performed comparably well to MSCSD in computing FODs in WM. When performing fiber tractography, the trajectories reconstructed with mFODs reached the cortex with more spatial continuity and for a longer distance as compared to MSCSD and allowed to reconstruct short trajectories tangential to the cortical folding. In conclusion, we demonstrated that our proposed method allows to perform spherical deconvolution of multiple anisotropic response functions, specifically improving the performances of spherical deconvolution in GM tissue.

Introduction

Diffusion Magnetic Resonance Imaging (dMRI) is a non-invasive technique sensitive to the microscopic motion (diffusion process) of water molecules[1]. In biologic tissues, the diffusion process is influenced by the presence of biologic membranes and macromolecules[2], which can hinder and/or restrict the molecular random walk in both isotropic and anisotropic fashions. In 1994, Basser et al.[3] proposed the diffusion tensor imaging (DTI) framework to characterize the anisotropy of hindered diffusion processes. Later, it was shown that when the diffusion process exhibits a sufficient degree of anisotropy, its principal direction can be computed[3] and eventually tracked[4], ultimately leading to what is nowadays known as fiber tractography[5,6] of the brain white matter (WM).

DTI-based fiber tractography is efficient in terms of acquisition and computation but suffers from a number of limitations. Firstly, the technique fails when multiple brain pathways coexist in the same spatial location[7], which has been estimated to be the case in over 90% of the human WM[8] and which is not likely to be solved by technological improvements as higher imaging resolutions[9]. Secondly, DTI-based fiber tractography only applies to data acquired within the Gaussian diffusion regime, and thus cannot take advantage of the strong diffusion weightings which are typical for high angular resolution diffusion imaging methods[10]. Due to these considerations, DTI is not likely the method of choice for a heterogeneous structure as grey matter (GM), which is characterized by low fractional anisotropy (FA) and, hence, high uncertainty of the principal diffusion direction.

Spherical deconvolution approaches[11,12] are the current gold standard to reconstruct fiber orientation distributions (FOD) and to perform fiber tractography in WM. However, their application to the study of GM remains limited, as FODs derived in the cortex are noisy and exhibit spurious peaks, a challenge when performing fiber tractography. Importantly, limitations in fiber tractography propagate to other analysis techniques making use of fiber reconstructions as, among others, structural connectivity approaches[13]. Indeed, fiber tracking is often seeded (or terminated) at the grey/white matter interface, but the inability to properly follow WM bundles in the cortex generates uncertainty onto the endpoints of the bundles. This has been partially addressed with geometric heuristics[14], but these oversimplify the way axons traverse the cortex[15].

The advent of specialized techniques for diffusion imaging, such as ultra-strong gradients[16] and simultaneous multi-slice[17], are rapidly increasing the imaging quality and resolution achievable in dMRI. This allows to appreciate new details of the human cortex, as recently reviewed by Assaf[18]. In ex-vivo studies[15,19,20], it has been consistently shown that the diffusion process observed in the cortex has a large anisotropic component perpendicular to the cortical folding, but also that tangential directions can be observed, for instance, in the central sulcus[21]. Such high-quality data acquisitions may soon be available for in-vivo applications, offering unprecedented research possibilities[22] in fiber tractography of GM, but a dedicated framework has not yet been investigated.

Multi-tissue spherical deconvolution approaches[23] have been suggested to further increase the accuracy of FOD reconstructions in WM by accounting for partial volume effects, e.g. in spatial locations where WM overlaps with GM or cerebrospinal fluid (CSF). Although GM and CSF are merely considered nuisance factors[24], such framework highlights the feasibility of explicitly accounting for multiple tissues in spherical deconvolution approaches. By separating multiple partial volume effects, multi-tissue approaches have shown improved FOD reconstructions in adults[23], neonates[25] and in patients with Parkinson's disease[26].

In current state-of-the-art multi-shell spherical deconvolution, only WM is assumed to be anisotropic, while GM is regarded as an isotropic tissue. To estimate the WM FOD, a WM fiber model – or response function – is estimated from voxels with a single fiber population, which are typically located in large bundles[27] such as, for example, the corpus callosum or the corticospinal tract. The response function is then assumed to be valid throughout the brain. While the validity of this representation has been questioned even in WM itself[28], it is highly unlikely that it can adequately represent the anisotropic signature of axons and dendrites in the cortex.

In this study, we investigate whether the spherical deconvolution approach can be revisited to consider the different signal characteristics of WM and GM by explicitly modelling both signal mixtures simultaneously. This novel framework allows to reconstruct multiple FODs (mFOD) and is ultimately aimed at improving the FOD estimation and fiber tractography of GM while retaining state-of-the-art performances in WM. In our proposed method, we disregard the existence of a single response function for the brain, and investigate whether it is feasible to simultaneously estimate multiple anisotropic FODs corresponding to tissue-specific response functions. We also evaluate whether this improves the fiber orientation characterization in GM, and fiber tractography in GM as compared to existing spherical deconvolution methods. Preliminary results of this work have been presented at the 2019 ISMRM meeting in Montreal, Canada[29].

Theory and methods

The following paragraphs present the mFOD framework and explain the qualitative and quantitative analyses that we performed using both simulations and in-vivo data.

Generalized Richardson Lucy (GRL)

In previous work[30,31], we introduced a multi-shell spherical deconvolution framework named Generalized Richardson Lucy (GRL) to account for multiple tissue types, such as WM, GM and cerebrospinal-fluid (CSF), as shown in Eq. 1.

$$\frac{S}{S_0} = f_{WM} \begin{bmatrix} H_{WM,1}^1 & \cdots & H_{WM,k}^1 \\ \vdots & \ddots & \vdots \\ H_{WM,1}^n & \cdots & H_{WM,k}^n \end{bmatrix} \otimes \text{FOD}_{WM} + f_{GM} \begin{bmatrix} H_{GM}^1 \\ \vdots \\ H_{GM}^1 \end{bmatrix} + f_{CSF} \begin{bmatrix} H_{CSF}^1 \\ \vdots \\ H_{CSF}^1 \end{bmatrix} \quad [1]$$

In Eq. [1], S is the signal collected at n diffusion weightings, S_0 is the non-diffusion-weighted signal, the H matrices represent the signal model associated to WM, GM and CSF, f is the associated signal fraction, and FOD_{WM} is the fiber orientation distribution associated to WM. The columns of the deconvolution matrix shown in Eq. [1] represent the possible solution of the spherical deconvolution on the unit sphere, resulting in a total number of $k+2$ columns, as GM and CSF are both considered isotropic.

Multiple fiber orientation distributions (mFOD)

In the mFOD framework, we redesigned the deconvolution matrix to allow for an arbitrary number of independent anisotropic deconvolution models. Considering solving the spherical deconvolution problem once more for WM, GM and CSF, but now accounting for anisotropic diffusion in both WM and GM, we can write:

$$\frac{S}{S_0} = \begin{bmatrix} H_{WM,1}^1 & \cdots & H_{WM,k}^1 & H_{GM,1}^1 & \cdots & H_{GM,k}^1 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ H_{WM,1}^n & \cdots & H_{WM,k}^n & H_{GM,1}^n & \cdots & H_{GM,k}^n \end{bmatrix} \otimes \begin{bmatrix} FOD_{WM} \\ FOD_{GM} \end{bmatrix} \begin{bmatrix} f_{WM} \\ f_{GM} \end{bmatrix} + f_{CSF} \begin{bmatrix} H_{CSF}^1 \\ \vdots \\ H_{CSF}^1 \end{bmatrix} \qquad [2]$$

Eq. [2] can be solved using the iterative least-squares minimizer previously introduced for GRL but replacing the Richardson-Lucy deconvolution scheme with a regularized non-negative least-squares solver[32], producing multiple independent FODs.

Signal models

To underline the model independence of the mFOD framework, the H-matrices of WM, GM and CSF, shown in Eq. 2, were evaluated with three different models. H_{CSF} was generated using the ADC model with the typical diffusion coefficient of free water at 37 degrees (3.0 x 10^{-3} mm2/s). The columns of H_{WM} and H_{GM} were generated using the DKI model[33] with isotropic kurtosis[34], and the NODDI model[35], respectively, both in simulations and with in-vivo data. The DKI model was chosen as it can represent the anisotropy of the WM signal while also considering non-Gaussian diffusion effects. The NODDI model has been shown to characterize the GM microstructure better than tensor models[36], and allows to take into account the larger presence of cell bodies in GM as compared to WM. The parameters of the models were determined as explained in the following paragraphs.

Simulations

The feasibility of simultaneously computing multiple anisotropic FODs corresponding to different models was evaluated numerically. H_{WM} was generated with the DKI model by setting the eigenvalues of the diffusion tensor as (1.7, 0.3, 0.3) x 10^{-3} mm^2/s, and isotropic kurtosis equal to 0.6. H_{GM} was generated with the NODDI model by simulating values of intra to extra-cellular ratio 0.4, parallel diffusivity 1.7x10^{-3} mm^2/s, concentration parameter of the Watson distribution equal to 1.0, and no partial volume of free-water was considered as the mFOD approach explicitly accounts for it. The parameters used for generating H_{WM} and H_{GM} matched the average values observed with in-vivo data.

Two set of simulations were generated. Simulation I aimed at evaluating the robustness to noise of the mFOD approach, whereas simulation II characterized its ability to separate partial volume effects for a given SNR level.

Simulation I

Three configurations were generated by considering partial volume effects between i) WM and CSF, ii) GM and CSF, and iii) WM and GM. The simulated partial volume ratio was 0.8/0.2, with an angle of approximately 90 degrees between the main direction of WM and the main direction of GM. The effect of noise was simulated by adding 1000 Rician realizations in correspondence of SNR levels 10, 20, 30, 40, 50, 60, 70, and 150. For each SNR level, we evaluated the error between the estimated and the simulated signal fractions, as well as the error between the simulated and the estimated main diffusion direction.

Simulation II

Three partial volume configurations were generated in analogy with simulation I, but increasing the ratio between the two simulated tissues from 0 to 1 with step 0.1. Rician noise was added by considering 1000 realizations at SNR 50. The error between the simulated and the estimated signal fractions was determined.

In-vivo data processing

Two subjects from the Human Connectome Project (HCP)[16,37] were randomly chosen and included in this study. The datasets included a T1-weighted image acquired at a resolution of 0.7 mm isotropic, and a dMRI dataset acquired at 1.25 mm isotropic. The diffusion datasets consisted of 18 b = 0 s/mm^2 volumes in addition to 270 volumes acquired by sampling diffusion weightings b = 1000, 2000, and 3000 s/mm^2 along 90 directions each. One of the two subjects was chosen to showcase the main results of the mFOD approach, using the native dMRI space (1.25 mm) as analysis space. Generalizability of the methods was then shown on the second subject, using the T1 resolution (0.7 mm) to highlight some features of the mFOD method. The T1-weighted data of the first subject was rigidly registered to the pre-processed dMRI data. Then, the tissue type segmentations of both datasets in WM, GM and CSF were derived with FSL FAST[38].

The dMRI data was fit with both the "Multi-Shell Constrained Spherical Deconvolution" (MSCSD)[23] (only data at 1.25 mm) and the mFOD approaches using custom implementations in MATLAB R2018b (The Mathworks Inc.) and functions from the ExploreDTI toolbox[39]. The code of mFOD is freely available online as part of the "MRIToolkit" toolbox for MATLAB (www.mritoolkit.org). MSCSD was initialized as previously suggested by using the tissue maps derived from the T1-weighted image. For the generation of the H-matrices used for the mFOD approach, a whole brain DKI and NODDI fit were performed on the data at 1.25 mm resolution. For the WM H-matrix, the eigenvalues of the diffusion tensor and the isotropic kurtosis were averaged within a white matter mask defined by fractional anisotropy values above 0.7. This resulted in eigenvalues [1.3, 0.5, 0.5] x 10^{-3} mm^2/s and isotropic kurtosis value 0.64. For the H-matrix of GM, the NODDI metrics were averaged within the GM segmentation obtained with the T1-weighted image, which resulted in intra to extra-cellular ratio 0.4, parallel diffusivity 1.7x10^{-3}mm^2/s and Watson's concentration parameter equal to 1.

FODs and signal fractions

The MSCSD fit produced signal fraction maps associated to WM, GM and CSF, and one FOD describing WM orientations. The fit of the mFOD approach resulted in the same number of fractional maps but two FODs, mFOD$_1$ describing WM geometry, and mFOD$_2$ describing GM geometry, respectively.

The signal fraction maps computed with MSCSD and mFOD were quantitatively compared by evaluating their Pearson correlations. To this end, f$_{WM}$ and f$_{GM}$ were evaluated, respectively, in WM and GM masks derived from the T1-weighted image. Next, the signal fraction maps from both mFOD and MSCSD were visually compared to those obtained from the T1-weighted volume. Further, the FODs computed with MSCSD and the mFOD framework were compared in selected cortical areas, to determine whether mFOD$_2$ provided additional information in GM as compared to mFOD$_1$ and MSCSD.

The FODs computed with the mFOD method can be either analyzed separately, or merged to combine accurate geometrical information in both WM and GM. As a proof of concept, we combined the two FODs by considering their normalized sum weighted by the respective signal fractions (mFOD-WS).

One of the difficulties when validating in-vivo dMRI techniques is the lack of ground truth. MSCSD can be regarded as a gold standard in WM, but debatably not in GM. In a first analysis, we investigated to which extent the principal diffusion directions of the mFOD-WS agree with those of MSCSD in both WM and GM. Per voxel, up to three peaks were identified for both FODs, then sorted per descending amplitude values.

Subsequently, the angle between the MSCSD and mFOD-WS results was determined as the minimum dyadic angle between any pair of peaks.

If discrepancies are observed in GM, it remains impossible to ultimately judge which method is the most accurate, but comparisons with a third method could provide additional interpretations. Anatomical knowledge as well as previous MRI studies[22] suggest that the dominant diffusion directions in GM should be perpendicular to the cortical folding, with exceptions such as the primary visual cortex[15]. We therefore derived the normal vectors to the cortical folding based on the structural tensor[40]. In short, a 3D volume was initialized with 1-values in correspondence of the GM/WM and GM/CSF interfaces, as derived from the T1-weighted segmentation. The linear interpolation field between the two surfaces was derived, and the corresponding vectors computed by computing the X-Y-Z first order derivatives of the volume. Subsequently, we determined the minimum dyadic angle between the normal vectors and any peak orientation determined with the MSCSD and mFOD-WS results, respectively.

Fiber tractography

Tracking parameters

We investigated fiber tractography results obtained with MSCSD, $mFOD_1$, $mFOD_2$ and mFOD-WS, with a particular focus on GM areas. Deterministic fiber tractography was performed in *ExploreDTI* by using identical parameters for all methods: FOD threshold 0.1 (default), minimum/maximum fiber length equal to 20/500mm, angle threshold 45 degrees, and step-size 0.6 mm. Fiber tractography was also performed in an axial region of interest overlapping the central sulcus in a single slice, and using FOD thresholds of 0.1 and 0.3, minimum fiber lengths of 10 and 50 mm, and step-size of 0.35 mm.

Firstly, FODs computed with the MSCSD and mFOD-WS approaches were computed in a frontal region and color encoded with a custom colormap assigning red, yellow and blue to pathways traversing WM, GM and CSF, respectively. Subsequently, whole-brain tractography of all FODs was performed by seeding all voxels with an FOD value above the default threshold. The fiber bundles were color encoded both with the conventional RGB color scheme[41] and the above-mentioned tissue-type color scheme, to convey both directional information and specificity of the FODs to the different tissue types.

Fiber pathways obtained with MSCSD and mFOD in a 3.75 mm wide sagittal slab were color encoded by normalized distance between the GM/WM interface and the WM/CSF interface – as obtained from the T1-weighted segmentation – and also directly overlaid, to appreciate differences in GM tracking distance and density between the compared methods.

Subsequently, we evaluated the number of tract pathways generated with MSCSD and mFOD-WS, which is informative given that the seed-points were kept constant and that a deterministic tracking algorithm was used. Additionally, we counted the fraction of plausible tract pathways, i.e., trajectories having at least one endpoint in grey matter, and the spatial coverage of tractography within GM when considering only start/endpoints of tracks.

To qualitatively assess whether the mFOD framework can reconstruct previously reported fibers tangential to the cortex[21], we investigated the fiber tractography results with the mFOD-WS approach. Fiber tractography was seeded in a single axial slice, then compared to the cortical surface reconstructed with the CAT12 toolbox[42] for MATLAB.

Results

Simulations

Figure 1 shows the signal fractions estimated with the mFOD approach in simulation I, where different mixtures of WM, GM and CSF were simulated in correspondence of multiple SNR Levels. At SNR level equal or greater than 20, the mFOD method disentangles mixtures of WM and CSF with signal fraction errors below 2% and angular errors below 3 degrees. The separation of GM from both WM and CSF is more challenging and requires higher SNR to achieve an error below 10%. In both cases, the signal fraction errors at SNR 50 are about 9%, 11% and 2%, for WM, GM and CSF, respectively. At the same SNR level, the angular error on the GM FOD is 8 degrees for the GM and CSF mixture. With regards to the GM and WM mixture, the angular errors of the WM and GM FODs are 4 degrees and 10 degrees at SNR 50, respectively.

Figure 1: Results of simulation I. The first row shows the $25^{th} - 75^{th}$ percentile (shaded area) and the median (solid line) of the signal fractions estimated with the mFOD approach for different SNR levels and partial volume mixtures (each column). The second row shows the angular error of the main direction of the WM and GM FODs as compared to the ground truth value. The solid line represents the median value, whereas the error bars the 25^{th} and 75^{th} percentile. The simulations were performed using anisotropic response functions for WM and GM, and an isotropic response function for CSF. To properly detect the main peak orientation associated to GM, a high SNR ($\geq$50) is required.

Figure 2 shows the signal fractions estimated in simulation II when linearly varying the simulated WM, GM and CSF fractions at SNR 50. For all the three cases, the estimated average signal fractions show excellent agreement with the simulated values. The maximum errors observed for WM, GM and CSF are 13%, 20%

and 4%, respectively. The Pearson correlation coefficient between the simulated and estimated signal fractions have values between 0.99 and 1.

Figure 2: The median value (solid line) and the 25th – 75th percentile (error bars) of the signal fractions estimated with the mFOD approach for the simulated anisotropic WM and GM response functions (columns 1-2) and the isotropic CSF response function (column 3). Different mixtures of the three tissue types were simulated with increasing partial volume and 1000 realizations of Rician noise at SNR 50. The largest errors were observed when mixing GM and CSF in correspondence of low GM signal fractions.

HCP data

Signal fractions

The WM, GM and CSF signal fractions estimated with the MSCSD and mFOD methods, as well as those derived from the segmentation of the T1-weighted image are shown for an example axial slice in Fig. 3. The maps estimated with the mFOD method exhibit excellent agreement with both MSCSD and the T1-derived segmentation, despite the numerical ill-conditioning of the mFOD approach. The Pearson

correlation coefficient between mFOD and MSCSD were 0.96 and 0.95 for f_{WM} and f_{GM}, respectively. Notice that the WM fractional maps estimated with both the MSCSD and mFOD methods have non-zero values in some GM areas, in contrast to the T1-derived segmentation.

Figure 3: An example axial slice of the signal fractions estimated with the MSCSD and mFOD methods in correspondence of different response functions as compared to the tissue segmentation obtained from the structural T1 image. Estimates with the MSCSD and mFOD approaches are remarkably similar and have high anatomical correspondence with the tissue types that they are designed to represent.

FODs

The FODs estimated with the MSCSD and mFOD methods in two cortical areas are shown in Fig. 4 and Fig. 5 for an example axial slice and an example coronal slice, respectively. The WM fiber orientations estimated with the mFOD method (mFOD$_1$) show remarkable similarity to the orientations estimated with MSCSD. Both FODs have very sharp and comparable orientations in the vicinity of WM and tend to become smaller towards deeper cortical GM. In contrast, the GM fiber orientations with the mFOD method (mFOD$_2$) show dominant and anatomically plausible peaks in most GM voxel, with orientations mostly perpendicular to that of the underlying WM. When mFOD$_1$ and mFOD$_2$ are linearly combined (mFOD-WS), a continuous vector field from WM to deep cortical GM is revealed.

Figure 4: Fiber orientation distributions (FODs) obtained with MSCSD and the mFOD method in an axial slice. The middle and the right columns show increasing zoom levels to highlight details of FODs. The $mFOD_1$ and $mFOD_2$ refer to the FODs computed when using response functions meant to represent WM and GM, respectively. The mFOD-WS is the sum of $mFOD_1$ and $mFOD_2$ weighted by their corresponding signal fractions. The $mFOD_2$ is non-zero only in GM and its proximity, and shows dominant orientations also in the deeper cortex, in contrast to both MSCSD and $mFOD_1$. The mFOD-WS combines the information content of $mFOD_1$ in WM with that of $mFOD_2$ in GM.

Figure 5: Fiber orientation distributions (FODs) obtained with MSCSD and the mFOD method in a coronal slice. The middle and the right columns show increasing zoom levels to highlight how the FODs change in vicinity of GM. The mFOD$_1$ and mFOD$_2$ refer to the FODs computed when using response functions meant to represent WM and GM, respectively. The mFOD$_2$ is non-zero only in proximity of GM, and shows dominant orientations also in the deeper cortex, in contrast to both MSCSD and mFOD$_1$. The mFOD$_1$ provides directional information in strong agreement with MSCSD but occasionally shows more peaks.

Fig. 6 shows the histogram of the peak deviations observed between MSCSD and mFOD-WS in GM and WM, as well as the angle between the FODs and the normal to the cortical folding as derived from the T1-weighted image with a structural tensor approach. When comparing the peak direction of MSCSD and mFOD-WS in WM, we observe that more than 70% of the WM voxels have an angular deviation smaller or equal than 10 degrees. In GM, the angular deviation between the two methods is larger, and the peak of the distribution is centered at about 5 degrees. When looking at the angle with the cortex normal, the histogram of both peaks is centered at 20 degrees, but the mFOD-WS method results on average in a smaller angular deviation than MSCSD.

Figure 6: The left histogram shows the normalized frequency (probability) of the angle between the main orientation computed with MSCSD and mFOD-WS in GM and WM. The right histogram shows the probability of the angle difference between the normal direction to the GM/WM surface and the main direction with MSCSD or mFOD-WS. With regards to the left plot, the two methods have very small angle differences in WM, within 10 degrees in over 70% of voxels, whereas larger differences are observed in GM. mFOD-WS resulted in lower angle deviation with the surface normal (derived with the structural tensor technique) as compared to MSCSD.

Fiber tractography

Fiber tractography results in a frontal lobe region computed with the MSCSD and mFOD-WS methods are shown in Fig. 7. The fiber pathways are color encoded according to the tissue type they are spatially traversing, and show that the mFOD-WS method generates more dense tractography results in GM. Further, the fiber bundles reconstructed with the mFOD-WS approach cover the GM more uniformly and traverse the tissue with a more plausible shape and to a greater extent.

Figure 7: Fiber pathways obtained with the MSCSD and mFOD-WS methods by seeding a frontal region (shown in yellow in the bottom left corner) with identical tracking parameters and FOD thresholds. The fiber pathways are color encoded according to the tissue type they transverse, as determined on the structural T1 image. The tracking with the mFOD-WS method resulted in a more complete tracking of the frontal cortex as compared to MSCSD, with the reconstructed trajectories covering a larger distance within the cortex.

Subsequently, we performed the whole-brain fiber tractography with the MSCSD and mFOD methods, both individually (mFOD$_1$, mFOD$_2$) and after the weighted sum procedure (mFOD-WS). The tracking with mFOD$_1$ produces a whole brain reconstruction that is largely comparable to that of MSCSD, suggesting preservation of the WM tracking performance. Conversely, the tracking of mFOD$_2$ results mostly in cortical reconstructions and short U-fibers, in-line with the GM-specific nature of this FOD. For pathways reconstructed with the mFOD-WS approach, 42.1% have their end points in GM, as compared to 34.1% when performing tractography with MSCSD. This is qualitatively visible in the tissue-type encoded reconstruction in Fig. 8, which shows that trajectories reconstructed with the mFOD-WS method cover the GM more extensively as compared to MSCSD. Note, however, that this also results in introducing extra spurious fibers.

Figure 8 – Whole-brain tracking with MSCSD and the mFOD methods. The first row shows the fiber pathways color encoded with the conventional color scheme, whereas the second row shows the same tracking color encoded by traversed tissue type (subsampled with factor 8). MSCSD and mFOD$_1$ provide similar tracking results with the latter showing better characterization of the cerebellar cortex, as shown by the more homogeneous red coloring in the tissue type encoded tractography. The tracking with mFOD$_2$ resulted exclusively in cortical and short (U-fibers) connections. With mFOD-WS, a more complete coverage of the GM is achieved as compared to both mFOD$_1$ and MSCSD.

An example sagittal slab of the trajectories reconstructed with MSCSD and the mFOD-WS approach was obtained by cropping the whole brain fiber tractography, as shown in Fig. 9. In the first column, the trajectories are color encoded according to their distance from the WM/GM interface (blue) to the GM/CSF interface (red). It can be seen that the fiber bundles reconstructed with the mFOD-WS method consistently travel a longer distance within the cortical GM than with MSCSD, in-line with observations in Fig. 7. Further, the frontal, parietal and occipital side of the cortex appear to be better reconstructed with the mFOD-WS method.

Figure 9 – A sagittal slab (red line in the bottom left picture) of the fiber tractography results obtained with MSCSD and the mFOD-WS method. The first column shows the fiber pathways color encoded according to their normalized distance from the GM/WM and the GM/CSF interfaces. The distance was determined in 3D as flow (linear interpolation field) between the two surfaces using the same method we previously used to determine the cortex normal. The central and right column show the same sagittal slab color encoded by reconstruction method, and its zoom in a cortical region. The WM-GM flow color encoding shows that a deeper cortical tracking was achieved with the mFOD-WS method as compared to MSCSD, especially in the frontal and occipital areas. When looking at the tissue type encoded images, mFOD-WS resulted in a more extensive and contiguous reconstruction of the cortical folding, although a larger number of spurious reconstructions can also be observed.

Fiber tractography of the mFOD-WS method using the HCP data is shown in Fig. 10 for different choices of tracking criteria. Good coverage of the cortical structure is observed for all settings, even when enforcing a minimum length of 50 mm, ensuring that most of the reconstructions originating in GM can reach the WM. When using an FOD threshold equal to 0.1, more spurious reconstructions are revealed as compared to a more restrictive threshold (0.3), but a denser tract reconstruction in the cortical ribbon and more extensive coverage of the contralateral hemisphere can be observed. Allowing for trajectories as short as 10 mm, results in a very dense reconstruction of the cortical gyri, showing short bundles that seem to travel tangentially to GM, as well as many short radial reconstructions.

Figure 10 – Results of fiber tractography of the mFOD-WS method with a seed ROI in an axial slice including the left central superior cortex. The tractography results are shown for two choices of FOD thresholds and minimum tract length. Reconstruction of WM bundles reaching the cortex can be observed with all tested settings. A lower FOD threshold (0.1) results in a more contiguous representation of the gyri as compared to the more restrictive setting (0.3), but also in more false positive reconstructions. Lowering the minimum tract length from 50 mm to 10 mm allows one to reconstruct short fibers tangent to the cortex (white arrows) that appear to travel within the cortical ribbon.

Discussion

In this work, we have introduced the mFOD framework to perform spherical deconvolution of multiple anisotropic response functions. With simulations and HCP data, we have shown that it is possible to reconstruct two anisotropic FODs, specific to WM and GM, while also accounting for the CSF partial volume effect. Further, we have shown that the mFOD approach improves the reconstruction of FODs in GM as compared to existing state-of-the-art methods, and that this results in improved fiber tractography reconstruction in regions near or in the GM.

Feasibility of mFOD

The reconstruction of multiple anisotropic FODs is a complicated, numerical challenge that requires the inversion of a matrix whose rank (number of independent rows/columns) is much lower than the number of possible solutions (FOD amplitudes on the unit sphere), a recurring challenge in deconvolution problems. Previous studies, as Tournier et al.[27] and Jeurissen et al.[23] used an L2 regularization term enforcing continuity (smoothness) in the FOD amplitudes. In this work, we used a non-negative least-squares solver with L2 regularization encapsulated in an iterative solver[31]. Taking into consideration that the mFOD formulation (eq. [2]) hosts multiple FODs, this might be suboptimal and introduce discontinuities in the solution. Nevertheless, our results suggest that the mFOD approach does not propagate smoothness between multiple FODs, likely due to the attenuation of small perturbations imposed by the median operator[30]. Indeed, the results of simulations, shown in Fig. 1 and Fig. 2, suggest that this approach allows to effectively disentangle the partial volume effects between mixtures of isotropic (n=1) and anisotropic (n=2) signal mixtures, if sufficient SNR is provided. In particular, the partial volume effects between WM and CSF are the easiest to disentangle, and an SNR equal or greater than 20 is sufficient to obtain accurate separation of signal fractions and minimal angular error on the WM FODs. In contrast, the separation of GM from both WM and CSF is a harder task, probably as a result of the limited anisotropy of this tissue type.

If an SNR equal or above 50 is achieved, the mFOD framework can preserve the ratios between different tissue types and can estimate their mixtures with minimal biases (Fig. 2). While an SNR of about 50 might be difficult to achieve in clinical settings, accurate pre-processing and denoising[43], or even deep-learning based reconstructions[44] might mitigate this need. Furthermore, in both simulation I and simulation II, the diffusion direction of GM was set perpendicular to that of WM, but in-vivo the two are likely to coincide. Indeed, white matter bundles travel to their cortical endpoint perpendicularly[45] to the cortical folding, with exceptions such as the primary visual cortex[15] or the U-fibers adjacent to the cortex[46]. If the direction of the axons leaving the cortex (to the WM) would not match that of intra-cortical axons, the results of the simulations suggest that the GM FOD would be estimated with an angular error between 10 (SNR 30) and 5 degrees (SNR 50), which are likely to perturbate the tracking trajectory. In practice, very high SNR is likely required to reconstruct fiber pathways tangential to the cortical folding, but less critical to detect radial trajectories. Furthermore, Fig. 3 shows that with in-vivo data the signal fraction map associated to GM is zero-valued in all non-GM voxels and has values of 1 in almost all GM voxels, which suggests excellent separation of the two tissue types.

GM modelling

While recent work has introduced the reconstruction of multiple WM FODs[25] to account for inadequacies in WM modelling in infants, the mFOD framework has the unique characteristic of reconstructing multiple FODs that are specific of remarkably different tissue types as WM and GM, as shown in Fig. 4 and Fig. 5. The figure shows that WM fiber orientations estimated with mFOD ($mFOD_1$) and MSCSD are very similar in WM, and that both exhibit very small values in GM. In contrast, the GM fiber orientations estimated with $mFOD_2$ show consistent and well-defined diffusion directions for a large extent of the cortical GM. The performance improvement over MSCSD of the mFOD framework in GM is explained by considering the improved tissue modelling. Approaches as MSCSD essentially describe the WM and GM mixture as

the sum of an anisotropic and an isotropic signal, similarly to the ball-and-stick model[47], where the anisotropic signal is modelled with properties extracted from single fiber population WM[27,48,49]. Therefore, in practice, by applying any constrained spherical deconvolution method in GM, one assumes the axonal component of GM to produce a similar signal to that of a WM bundle without crossing fibers. However, assuming the cortical axons and dendrites of the cortex to be identical to the axons in the corpus callosum or other major WM bundles seems a stretch given their different diffusion signature[50]. In this work, we represent the GM signal with a special case of the NODDI model[35], which represents a first approach to take into account additional effects as tortuosity[51] and intra-to-extra neurite signal ratios into the deconvolution problem.

How to use multiple FODs

One of the open challenges in the mFOD framework regards how to combine the multiple FODs computed by the method. In this work, we adopted a pragmatic approach and considered their weighted sum (mFOD-WS). This choice produces convincing and contiguous FODs, as shown in Fig. 4 and Fig. 5. However, this might be sub-optimal for fiber tracking approaches, as it might blur the peaks detected in individual FODs. Future work is needed to establish whether dynamically tracking the locally larger FOD, or dynamically choosing the FODs most aligned to the tracking direction might represent better alternatives. Additionally, sparsity constraints as L1 or L0 norm[52] could be taken into consideration to improve the FOD separation and minimize overfit. Nevertheless, when we compared the main peaks of mFOD-WS to those obtained with MSCSD (Fig. 6), we observed remarkable similarity between the two in WM. Further, the average angular error between the two techniques was below 10 degrees in over 70% of WM voxels, suggesting that the mFOD-WS approach does not affect the performance of spherical deconvolution in WM. In GM, the mFOD-WS approach resulted in smaller angular deviations with respect to the cortex normal as compared to MSCSD. While the cortex normals might not always be representative of the direction with which axons are organized in the cortex, they represent an accepted assumption[14], and might serve as a comparison term between the two methods. The average angular deviation between the cortex normals and mFOD-WS is 20 degrees, which is in-line with the angular error predicted in case of WM and GM mixtures at SNR 50 (Fig. 1).

Improved fiber tractography in GM

Fiber tractography is one of the main applications of spherical deconvolution techniques. Here, we used a deterministic approach to facilitate the interpretation of pathway reconstructions. As shown in Fig. 7, we obtained a higher percentage of tract pathways reaching a GM region with the mFOD-WS approach than with MSCSD given identical tracking parameters. Furthermore, the fiber pathways reconstructed with the mFOD-WS method covered the cortical structure more extensively, and their geometrical configuration appeared more similar to previous reports of high resolution cortical dMRI[22]. While a similar tractography result might also be achievable with MSCSD by tuning the FOD threshold[53], this is likely to introduce not only false positives in WM, but also to worsen the tracking in GM itself due to a larger number of spurious peaks[54]. When performing whole-brain tractography, the use of the mFOD-WS approach resulted in more tract pathways having a start or endpoint in the GM (+8%), and generally in more extensive GM coverage, as shown in Fig. 8. This was even more apparent when looking at a sagittal slab (Fig. 9), which showed how fiber bundles could be followed in GM up to the GM/CSF interface, at the price of potentially more spurious reconstructions as compared to MSCSD. In practice, mFOD can thus have a major impact on applications such as pre-surgical planning[55] or structural connectivity studies[56], where following the fiber pathways up to their cortical endpoint is crucial. Existing fiber tractography

methods mostly do not define the seed points in the cortex, but rather at the GM/WM interface or in the adjacent WM, potentially introducing a number of false positives[57]. For similar reasons, connectivity studies are often performed by first seeding WM or the WM/GM interface, then filtering the trajectories with anatomical constraints[58], and finally assigning the fiber bundles to the closest GM node after spatial dilation, potentially introducing uncertainty in the assignment[59]. The mFOD framework may address these shortcomings, allowing one to follow WM bundles with higher accuracy than existing methods from/to their cortical start/endpoint. Taking into account the results shown in Fig. 10, it seems possible to also effectively reconstruct some crossing bundles in the GM itself, where radial projections cross previously reported tangential fibers[15,19,21].

Methodological considerations

The models used here to describe WM and GM properties were chosen from current popular methods, but their combination might not be the optimal representations to estimate multiple FODs. The DKI model, here used in its simplified version with isotropic kurtosis[34], efficiently represents non-Gaussian signals with a limited number of parameters but is sub-optimal to account for direction dependent signal restrictions. Alternatively, geometric representations of the WM signal[60], as well as data driven methods[49] could also be explored. Note that the diffusion approaches suggested for WM (DKI) and GM (NODDI) could be further adjusted or even replaced by other more specific tissue models to further improve the mFOD framework.

In Fig. 6, we used the cortex normal direction as a comparison for FODs computed with both MSCSD and mFOD. While it is reasonable to assume that most axonal projections will traverse the cortex with a perpendicular trajectory, this is not always the case. Furthermore, the method employed to compute the normal trajectory is not accurate when the sulci are separated by less than 2 voxels and should therefore be interpreted cautiously. The performance of the mFOD framework has been shown for two subjects and, although promising, should be further evaluated on larger cohorts and with more clinically achievable protocols. Finally, while we have shown that the mFOD framework improves the performance of spherical deconvolution and fiber tractography in GM, it remains to be shown to which extent such improvements will benefit clinical studies and structural connectivity investigations.

In conclusion, we have presented the mFOD framework to perform tissue-specific spherical deconvolution which can simultaneously reconstruct multiple distributions of fiber orientations per voxel. By integrating specific WM and GM models, we have shown that the mFOD approach can improve the performance of spherical deconvolution in GM while retaining the performance of state-of-the-art methods designed for WM.

Acknowledgements

The research of A.D.L. is supported by the ERA-NET Neuron grant R.4195 – "Repetitive Subconcussive Head Impacts - Brain Alterations and Clinical Consequences" (REPIMPACT). The research of F.G. is funded by the Chinese Scholarship Council (CSC), No. 201306080017. The research of A.L. is supported by VIDI Grant 639.072.411 from the Netherlands Organisation for Scientific Research (NWO).

References

1.   Einstein, A. Über die von der molekularkinetischen Theorie der Wärme geforderte Bewegung von in ruhenden Flüssigkeiten suspendierten Teilchen. *Ann. Phys.* **322**, 549–560 (1905).

2.   Walter, J. A. & Hope, A. B. Proton magnetic resonance studies of water in slime mould plasmodia. *Aust. J. Biol. Sci.* **24**, 497–507 (1971).

3.   Basser, P. J., Mattiello, J. & LeBihan, D. MR diffusion tensor spectroscopy and imaging. *Biophys. J.* **66**, 259–67 (1994).

4.   Jones, D. K., Simmons, A., Williams, S. C. & Horsfield, M. A. Non-invasive assessment of axonal fiber connectivity in the human brain via diffusion tensor MRI. *Magn. Reson. Med.* **42**, 37–41 (1999).

5.   Mori, S., Crain, B. J., Chacko, V. P. & van Zijl, P. C. Three-dimensional tracking of axonal projections in the brain by magnetic resonance imaging. *Ann. Neurol.* **45**, 265–9 (1999).

6.   Mori, S., Wakana, S., van Zijl, P. C. M. & Nagae-Poetscher, L. M. *MRI Atlas of Human White Matter*. (Elsevier Science, 2005).

7.   Tuch, D. S. *et al.* High angular resolution diffusion imaging reveals intravoxel white matter fiber heterogeneity. *Magn. Reson. Med.* **48**, 577–582 (2002).

8.   Jeurissen, B., Leemans, A., Tournier, J.-D., Jones, D. K. & Sijbers, J. Investigating the prevalence of complex fiber configurations in white matter tissue with diffusion magnetic resonance imaging. *Hum. Brain Mapp.* **34**, 2747–66 (2013).

9.   Schilling, K. *et al.* Can increased spatial resolution solve the crossing fiber problem for diffusion MRI? *NMR Biomed.* **30Schillin**, 1–16 (2017).

10.   Xie, S. *et al.* How does B-value affect HARDI reconstruction using clinical diffusion MRI data? *PLoS One* **10**, 1–19 (2015).

11.   Tournier, J.-D., Calamante, F., Gadian, D. G. & Connelly, A. Direct estimation of the fiber orientation density function from diffusion-weighted MRI data using spherical deconvolution. *Neuroimage* **23**, 1176–85 (2004).

12.   Dell'acqua, F. *et al.* A Deconvolution Approach Based on Multi-Tensor Model to Solve Fiber Crossing in Diffusion-MRI. *Conf. Proc. IEEE Eng. Med. Biol. Soc.* **2**, 1415–8 (2005).

13.   Hagmann, P. *et al.* Mapping the structural core of human cerebral cortex. *PLoS Biol.* **6**, 1479–1493 (2008).

14.   Tozer, D. J. *et al.* Linking white matter tracts to associated cortical grey matter: A tract extension methodology. *Neuroimage* **59**, 3094–3102 (2012).

15.   Kleinnijenhuis, M. *et al.* Layer-specific diffusion weighted imaging in human primary visual cortex invitro. *Cortex* **49**, 2569–2582 (2013).

16.   McNab, J. A. *et al.* The Human Connectome Project and beyond: initial applications of 300 mT/m gradients. *Neuroimage* **80**, 234–45 (2013).

17.   Feinberg, D. A. & Setsompop, K. Ultra-fast MRI of the human brain with simultaneous multi-slice imaging. *J. Magn. Reson.* **229**, 90–100 (2013).

18. Assaf, Y. Imaging laminar structures in the gray matter with diffusion MRI. *Neuroimage* 1–12 (2018) doi:10.1016/j.neuroimage.2017.12.096.

19. Leuze, C. W. U. *et al.* Layer-specific intracortical connectivity revealed with diffusion MRI. *Cereb. Cortex* **24**, 328–339 (2014).

20. Aggarwal, M., Nauen, D. W., Troncoso, J. C. & Mori, S. Probing region-specific microstructure of human cortical areas using high angular and spatial resolution diffusion MRI. *Neuroimage* **105**, 198–207 (2015).

21. McNab, J. A. *et al.* Surface based analysis of diffusion orientation for identifying architectonic domains in the in vivo human cortex. *Neuroimage* **69**, 87–100 (2013).

22. Setsompop, K. *et al.* High-Resolution In Vivo Diffusion Imaging of the Human Brain With Generalized Slice Dithered Enhanced Resolution : Simultaneous Multislice (gSlider-SMS). **151**, 141–151 (2018).

23. Jeurissen, B., Tournier, J.-D., Dhollander, T., Connelly, A. & Sijbers, J. Multi-tissue constrained spherical deconvolution for improved analysis of multi-shell diffusion MRI data. *Neuroimage* (2014) doi:10.1016/j.neuroimage.2014.07.061.

24. Dell'acqua, F. *et al.* A modified damped Richardson-Lucy algorithm to reduce isotropic background effects in spherical deconvolution. *Neuroimage* **49**, 1446–58 (2010).

25. Pietsch, M. *et al.* A framework for multi-component analysis of diffusion MRI data over the neonatal period. *Neuroimage* **186**, 321–337 (2019).

26. Kamagata, K. *et al.* Connectome analysis with diffusion MRI in idiopathic Parkinson's disease: Evaluation using multi-shell, multi-tissue, constrained spherical deconvolution. *NeuroImage Clin.* **17**, 518–529 (2018).

27. Tournier, J. D., Calamante, F. & Connelly, A. Robust determination of the fibre orientation distribution in diffusion MRI: Non-negativity constrained super-resolved spherical deconvolution. *Neuroimage* **35**, 1459–1472 (2007).

28. Schilling, K. G. *et al.* Histologically derived fiber response functions for diffusion MRI vary across white matter fibers—An ex vivo validation study in the squirrel monkey brain. *NMR Biomed.* **32**, 1–17 (2019).

29. De Luca, A., Guo, F. & Leemans, A. Estimation of multiple fiber orientation distributions (mFODs) from diffusion MRI data using spherical deconvolution. in *International Society for Magnetic Resonance in Medicine* (2019).

30. Guo, F., Leemans, A., Viergever, Max A, Dell'acqua, F. & De Luca, A. Damped Richardson-Lucy Deconvolution for Multi-Shell Diffusion MRI. in *International Society for Magnetic Resonance in Medicine2* (2018).

31. Guo, F., Leemans, A., Viergever, M. A., Dell'Acqua, F. & De Luca, A. Generalized Richardson-Lucy (GRL) for analyzing multi-shell diffusion MRI data. (2019).

32. De Luca, A., Leemans, A., Bertoldo, A., Arrigoni, F. & Froeling, M. A robust deconvolution method to disentangle multiple water pools in diffusion MRI. *NMR Biomed.* 1–17 (2018) doi:10.1002/nbm.3965.

33.   Jensen, J. H., Helpern, J. a, Ramani, A., Lu, H. & Kaczynski, K. Diffusional kurtosis imaging: the quantification of non-gaussian water diffusion by means of magnetic resonance imaging. *Magn. Reson. Med.* **53**, 1432–40 (2005).

34.   De Luca, A., Bertoldo, A. & Froeling, M. Effects of perfusion on DTI and DKI estimates in the skeletal muscle. *Magn. Reson. Med.* **78**, 233–246 (2017).

35.   Zhang, H. G., Schneider, T., Wheeler-Kingshott, C. a & Alexander, D. C. NODDI: practical in vivo neurite orientation dispersion and density imaging of the human brain. *Neuroimage* **61**, 1000–16 (2012).

36.   Kleinnijenhuis, M. *Imaging fibres in the brain*. (2014). doi:10.13140/2.1.4788.7689.

37.   Van Essen, D. C. *et al.* The WU-Minn Human Connectome Project: An overview. *Neuroimage* **80**, 62–79 (2013).

38.   Zhang, Y., Brady, M. & Smith, S. Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE Trans. Med. Imaging* **20**, 45–57 (2001).

39.   Leemans, A., Jeurissen, B., Sijbers, J. & Jones, D. K. ExploreDTI: a graphical toolbox for processing, analyzing, and visualizing diffusion MR data. in *17th annual meeting of the International Society for Magnetic Resonance in Medicine, Honolulu, Hawaii, USA* 3537 (2009).

40.   Kleinnijenhuis, M., Barth, M., Alexander, D. C., van Cappellen van Walsum, A. M. & Norris, D. G. Structure Tensor Informed Fiber Tractography (STIFT) by combining gradient echo MRI and diffusion weighted imaging. *Neuroimage* **59**, 3941–3954 (2012).

41.   Pajevic, S. & Pierpaoli, C. Color schemes to represent the orientation of anisotropic tissues from diffusion tensor data: application to white matter fiber tract mapping in the human brain. *Magn. Reson. Med.* **42**, 526–40 (1999).

42.   Seiger, R., Ganger, S., Kranz, G. S., Hahn, A. & Lanzenberger, R. Cortical Thickness Estimations of FreeSurfer and the CAT12 Toolbox in Patients with Alzheimer's Disease and Healthy Controls. *J. Neuroimaging* **28**, 515–523 (2018).

43.   Veraart, J. *et al.* Denoising of diffusion MRI using random matrix theory. *Neuroimage* **142**, 394–406 (2016).

44.   Koppers, S., Haarburger, C. & Merhof, D. Diffusion MRI Signal Augmentation: From Single Shell to Multi Shell with Deep Learning. in 61–70 (2017). doi:10.1007/978-3-319-54130-3_5.

45.   Budde, M. D. & Annese, J. Quantification of anisotropy and fiber orientation in human brain histological sections. *Front. Integr. Neurosci.* **7**, 1–8 (2013).

46.   Schilling, K. *et al.* Confirmation of a gyral bias in diffusion MRI fiber tractography. *Hum. Brain Mapp.* **39**, 1449–1466 (2018).

47.   Tuch, D. S. Q-ball imaging. *Magn. Reson. Med.* **52**, 1358–72 (2004).

48.   Dell'Acqua, F. *et al.* A model-based deconvolution approach to solve fiber crossing in diffusion-weighted MR imaging. *IEEE Trans. Biomed. Eng.* **54**, 462–72 (2007).

49.   Tax, C. M. W., Jeurissen, B., Vos, S. B., Viergever, M. A. & Leemans, A. Recursive calibration of the

fiber response function for spherical deconvolution of diffusion MRI data. *Neuroimage* **86**, 67–80 (2014).

50. Westin, C. F. *et al.* Q-space trajectory imaging for multidimensional diffusion MRI of the human brain. *Neuroimage* **135**, 345–362 (2016).

51. Sen, P. N. & Basser, P. J. A model for diffusion in white matter in the brain. *Biophys. J.* **89**, 2927–2938 (2005).

52. Daducci, A., Van De Ville, D., Thiran, J. P. & Wiaux, Y. Sparse regularization for fiber ODF reconstruction: From the suboptimality of ℓ2 and ℓ1 priors to ℓ0. *Med. Image Anal.* **18**, 820–833 (2014).

53. Aerts, H., Dhollander, T. & Marinazzo, D. Evaluating the performance of 3-tissue constrained spherical deconvolution pipelines for within-tumor tractography. *bioRxiv* (2019).

54. Parker, G. D. *et al.* A pitfall in the reconstruction of fibre ODFs using spherical deconvolution of diffusion MRI data. *Neuroimage* **65**, 433–448 (2013).

55. Mormina, E. *et al.* Optic radiations evaluation in patients affected by high-grade gliomas: a side-by-side constrained spherical deconvolution and diffusion tensor imaging study. *Neuroradiology* **58**, 1067–1075 (2016).

56. Beare, R. *et al.* Altered structural connectivity in ADHD: a network based analysis. *Brain Imaging Behav.* **11**, 846–858 (2017).

57. Maier-Hein, K. H. *et al.* The challenge of mapping the human connectome based on diffusion tractography. *Nat. Commun.* **8**, (2017).

58. Smith, R. E., Tournier, J.-D., Calamante, F. & Connelly, A. Anatomically-constrained tractography: improved diffusion MRI streamlines tractography through effective use of anatomical information. *Neuroimage* **62**, 1924–38 (2012).

59. Wei, K., Cieslak, M., Greene, C., Grafton, S. T. & Carlson, J. M. Sensitivity analysis of human brain structural network construction. *Netw. Neurosci.* **1**, 446–467 (2017).

60. Ferizi, U. *et al.* A ranking of diffusion MRI compartment models with in vivo human brain data. *Magn. Reson. Med.* **72**, 1785–1792 (2014).