Genome sequence and comparative analysis of reindeer (*Rangifer tarandus*) in northern Eurasia.

Melak Weldenegodguad[1,2*], Kisun Pokharel[1*], Yao Ming[3*], Mervi Honkatukia[1,4], Jaana Peippo[1], Tiina Reilas[1], Knut H. Røed[5] & Juha Kantanen[1**]

[1]Natural Resources Institute Finland, FI-31600 Jokioinen, Finland,

[2]Department of Environmental and Biological Sciences, University of Eastern Finland, FI-70201 Kuopio, Finland,

[3]BGI-Shenzhen, Shenzhen 518083, China,

[4]Nordic Genetics Resource Center – NordGen, P.O. Box 115, NO-1431 Ås, Norway,

[5]Department of Basic Science and Aquatic Medicine, Norwegian University of Life Sciences, PO-8146 Dep, NO-0033 Oslo, Norway;

*These authors contributed equally to this work.

**Correspondence and requests for materials should be addressed to J.K. (email: juha.kantanen@luke.fi)

Keywords: *Cervidae*, *de novo* assembly, resequencing, adaptation, domestication, positive selection

Abstract

Reindeer are semi-domesticated ruminants that have adapted to the challenging northern Eurasian environment characterized by long winters and marked annual fluctuations in daylight. We explored the genetic makeup behind their unique characteristics by *de novo* sequencing the genome of a male reindeer and conducted gene family analyses with nine other mammalian species. We performed a population genomics study of 23 additional reindeer representing both domestic and wild populations and several ecotypes from various geographic locations. We assembled 2.66 Gb (N50 scaffold of 5 Mb) of the estimated 2.92 Gb reindeer genome, comprising 27,332 genes. The results from the demographic history analysis suggested marked changes in the effective population size of reindeer during the Pleistocene period. We detected 160 reindeer-specific and expanded genes, of which zinc finger proteins (n=42) and olfactory receptors (n=13) were the most abundant. Comparative genome analyses revealed several genes that may have promoted the adaptation of reindeer, such as those involved in recombination and speciation (*PRDM9*), vitamin D metabolism (*TRPV5*, *TRPV6*), retinal development (*PRDM1*, *OPN4B*), circadian rhythm (*GRIA1*), immunity (*CXCR1*, *CXCR2*, *CXCR4*, *IFNW1*), tolerance to cold-triggered pain (*SCN11A*) and antler development (*SILT2*). The majority of these characteristic reindeer genes have been reported for the first time here. Moreover, our population genomics analysis suggested at least two independent reindeer domestication events with genetic lineages originating from different refugial regions after the Last Glacial Maximum. Taken together, our study has provided new insights into the domestication, evolution and adaptation of reindeer and has promoted novel genomic research of reindeer.

Introduction

Reindeer (*Rangifer tarandus*) have pivotal economic, societal, cultural and ecological values for indigenous people and pastoralists in northern and subarctic regions of Eurasia. Reindeer are a source of meat, hide and occasionally milk and have been used for transportation. Reindeer were crucial for the colonization of the northernmost parts of Eurasia and have a central symbolic role for the indigenous Sami, Nenets, and Evenki cultures and several other North Eurasian cultures[1,2].

During the evolutionary time scale and in recent demographic history, reindeer were distributed across the northern Eurasian regions, except during the glacial periods[3], and adapted to a challenging northern environment characterized by short daylight and limited vegetation for feeding during the long winter and prolonged daylight during the short summer period[4,5]. The marked fluctuations in daylight may have led to weakened circadian rhythms[6–8]. Similarly, reindeer exhibit retinal structural adaptation to the extreme seasonal light change to increase retinal sensitivity in dim light[9]. Moreover, reindeer have developed unique features, such as fat metabolism processes, a low resting metabolic rate[5] and the annual growth and loss of antlers both in males and females[10].

The genetic diversity, population structure, and prehistorical demographic events of *Rangifer* populations have been investigated, and inferences of domestication have been drawn using mitochondrial DNA (mtDNA) D-loop polymorphisms and autosomal microsatellites as genetic markers[3,11–14]. For example, mtDNA and microsatellite diversity patterns between domestic and

wild Eurasian tundra reindeer (*R. t. tarandus*), domestic and wild boreal forest reindeer (*R. t.*
fennicus), and Arctic Svalbard reindeer (*R. t. platyrhynchus*), wild North American Caribou (*R. t.*
caribou) and several other populations appear to reflect their refugial origins and colonization of
circumpolar regions after the Last Glacial Maximum rather than following the classic division of
Rangifer populations into three main ecological groups adapted to a particular set of
environmental conditions (forest, tundra and arctic environments) and sedentary or migratory
life-history strategies[3,13,15]. Moreover, mtDNA analyses in ancient and modern *Rangifer*
populations indicate at least three different domestication events in the prehistory of reindeer and
suggest that one of the domestication sites previously assumed may not have been located in
northern Fennoscandia[13,15].

For most domesticated animal species (e.g., chicken[16], taurine cattle[17], pig[18], sheep[19] and
horse[20]), several genomic tools, such as annotated *de novo* genome assemblies, resequencing
data sets, and whole-genome single nucleotide polymorphism (SNP) panels, are available for
genomics studies. These whole-genome research approaches have provided critical, new
knowledge of the evolution, domestication, genetic diversity, selection patterns and adaptation of
domestic animal species. For reindeer genetics studies, however, these genomics tools have
become available only recently: Li et al. (2017)[21] reported the *de novo* sequencing of one female
reindeer (*R. tarandus*) originating from Inner Mongolia, PR of China. Subsequently, the reindeer
reference genome was compared with the genomes of 50 additional ruminant species, and the
genomic data from three Chinese domestic reindeer and three Norwegian wild tundra reindeer
were resequenced in the study by Lin et al. (2019)[4]. The genetic basis of specific *R. tarandus*
characteristics, such as adaptation to marked annual regulation in daylight, exceptional vitamin D
metabolism and behavioural traits, was identified. In the present study, we provide an alternative
draft reference genome for reindeer studies: we deep-sequenced and *de novo* assembled the
genome of a male reindeer originating from Sodankylä, northern Finland by using the Illumina
HiSeq platform. In addition, we analysed the whole transcriptome of six tissues of the reference
animal to improve the annotation. Moreover, we resequenced 23 *Rangifer* from Norway and
Russia (wild and domestic), together with wild reindeer from Svalbard and Alaska, representing
different ecotypes, and report here the largest reindeer genome data sequenced thus far. With
these data, we examined the genetic diversity, demographic evolution, adaptation and
domestication of reindeer and compared the genome structure of reindeer with that of several
other mammalian species, e.g., with several domesticated ruminant species having a *de novo*
reference genome sequenced.

Results

Genome assembly and annotation

Genomic DNA extracted from the blood sample of a male reindeer was sequenced using the
Illumina HiSeq 2500 and 4000 platforms. Libraries with insert sizes ranging from 170 bp to 20
kb were sequenced and generated a total of 513 Gb (170X) raw sequence data (Supplementary
Table S1). After filtering low-quality reads, 300.55 Gb (100X) clean data was retained for
assembly (Table 1 and Supplementary Table S1). The genome size of reindeer was estimated to
be 2.92 Gb using k-mer analysis[22] (Supplementary Table S2 and Supplementary Fig. S1). High-
quality reads were assembled using SOAPdenovo V2.04[23], and an assembly spanning 2.66 Gb of

an estimated 2.92 Gb was generated, with N50 scaffolds and contig sizes of 5 Mb and 48.8 kb,
respectively (Table 1, Supplementary Table S3). The final assembly comprised 23,450 scaffolds
(170 kb–28.2 Mb), which represented 2.62 Gb and 107, 910 contigs (200 bp–22.5 kb) that in turn
represented an additional 44.9 Mb of the assembled genome. The assemblies covered ~91% of
the estimated reindeer genome size. The GC content of the assembled genome was
approximately 41.4% (Supplementary Fig. S2 and Supplementary Fig. S3), which is similar to
the GC content of the genomes of other related species[24,25].

We evaluated the quality of the draft assembly by aligning high-quality reads from the short
insert-size libraries against the assembly using Burrows-Wheeler Alignment (BWA)[26] and found
that the mapping rate was 98.3%. In addition, assembly quality was assessed by aligning the *de*
novo assembled transcriptome against the assembly using BLAT[27]. A total of 99% of the
assembled transcripts were mapped to the corresponding assembly (Supplementary Table S4).
Furthermore, to assess the assembly quality, we also used Benchmarking Universal Single-Copy
Orthologs (BUSCO)[28] genes and found that the assembly contained 94.9% (3,895 of 4,104)
complete BUSCOs (Supplementary Table S5).

We predicted a total of 27,332 protein-coding genes with an average of 7.4 exons, 1,305 bp
coding sequences (CDS) and 28,808 bp transcripts per gene in the reindeer genome by
employing a homology-based and an RNA-assisted approach as described by Curwen et al.
(2004)[29] (Table 1, Supplementary Table S7). In total, 26,838 (98.19%) of the total predicted
genes were functionally annotated according to public databases (InterPro, Kyoto Encyclopedia
of Genes and Genomes (KEGG), Swiss-Prot, TrEMBL), while a total of 494 (1.81%) genes
remained unannotated (Supplementary Table S8). To assess the quality of gene prediction, the
gene length distribution, the length of the CDS, exons and introns, and the distribution of exon
number per gene were compared with those of the five mammalian genomes in a homology-
based prediction. No significant differences in exon length and intron length distribution were
observed among the six species (Supplementary Fig. S4). Among the compared species, the gene
models derived for reindeer were similar to those for dog with respect to all of these key
parameters (Supplementary Fig. S4). The annotated gene set was evaluated by BUSCO, and our
gene set mapped 96.8% (3,973 of 4,104) complete BUSCOs (Supplementary Table S6),
indicating the high quality of our draft genome annotation. We also annotated 6,580 non-coding
RNAs (ncRNAs), of which 551 were microRNAs (miRNAs), 329 were rRNAs, 3,994 were
tRNAs, and 1,706 were small nuclear RNAs (snRNAs) (Supplementary Table S9). Compared
with the analyses of cattle, yak, sheep, goat, horse, human and the first reindeer genome
assembly, with less than 900 tRNAs predicted, our analysis revealed an exceptionally high
number of tRNAs (the analysis was repeated for confirmation)[21]. This unique finding may be due
to some unknown bioinformatics bias or could be related to unique characteristics of the
Fennoscandian reindeer.

In total, 38.02% of the assembled reindeer genome comprised repetitive sequences (transposable
elements (TEs) and tandem repeats; Table 1, Supplementary Table S10), most of which were
transposon-derived repeats. All TEs obtained by different methods (Repbase library[30], alignment
with known transposable-element-related genes, and *de novo* RepeatModeller[31] identification)
were combined, and the result indicated that 36.78% of the reindeer genome consisted of TE
sequences (Supplementary Table S11), which is similar to the results obtained for the panda

(36.2%)[24] and dog genomes (36.1%)[32], while lower than the results obtained for the cattle
(46.5%)[17], goat (42.2%)[33] and sheep genomes (42.7%)[19].

Table 1. Assembly and annotation statistics

Sequencing	Insert size	Clean data (Gb)	Sequence coverage (X)
Paired-end libraries	170, 500, 800 bp	181.39	60.46
	2, 5, 10, 20 kb	119.16	39.72
	Total	300.55	100.18
Assembly	**N50 (Kb)**	**Longest (Kb)**	**Size (Gb)**
Contig	48.8	441.2	2.62
Scaffold	5,023.6	28,190.2	2.66
Annotation	**Number**	**Total length (Mb)**	**Percentage of genome**
Genes	27,332	787.3	-
Repeats	-	1,012.6	38.02

Evolution of the reindeer genome

We compared the novel reindeer reference genome to nine different mammalian species
(*Camelus dromedarius, Capra hircus, Ovis aries, Bos taurus, Bos grunniens, Equus caballus,*
Canis familiaris, Ursus maritimus and *Homo sapiens*) to examine the evolution of genes and
gene families. The *R. tarandus* genome revealed 17,709 orthologous gene families that were
shared by at least two species (pairwise comparison to reindeer) and a total of 11,640
orthologous gene families that were shared by all ten species (Fig. 1A, Supplementary Table
S12).

Reindeer-specific gene families. A total of 14,520 orthologous gene families were shared
between the domestic ruminant species, while 547 gene families were specific to reindeer (Fig.
1B, Supplementary Data 1). The reindeer-specific gene families contained 1,090 genes, of which
530 were associated with 945 known InterPro domains (Supplementary Data 2). A number of

olfactory receptors (n=29 genes), zinc finger proteins (n=71 genes) and ribosomal proteins (n=29
genes) were present in the reindeer-specific gene families (Supplementary Data 1). Out of 1,090
reindeer-specific genes, 691 genes lacked Gene Ontology (GO) annotations. Analysis of the
remaining 399 genes revealed 39 significant (false discovery rate (FDR) < 0.05) GO terms, of
which 23 were associated with biological processes (BP) and the rest were categorized in
molecular function (MF) (Supplementary Table S13). The specific gene families enriched in
biological processes revealed several GO terms associated with biosynthetic processes (n=10 GO
terms) and metabolic processes (n=7 GO terms) (Supplementary Table S13). The biological
processes associated with reindeer-specific gene families also included "GO:0097659, nucleic
acid-templated transcription, P=6 x 10^{-6}", "GO:0010468, regulation of gene expression, P=6.2 x
10^{-6}" and "GO:0007186, G protein–coupled receptor signaling pathway, P=0.00028"
(Supplementary Table S13). Similarly, the molecular function category of the GO terms revealed
a number of receptor activities (n=7 GO terms), such as "GO:0099600, transmembrane receptor
activity, P= 3.3 x 10^{-6}", "GO:0004888, transmembrane signaling receptor activity, P= 1.8 x 10^{-5}
", "GO:0004984, olfactory receptor activity, P= 5.9 x 10^{-5}", "GO:0038023, signaling receptor
activity, P= 6.5 x 10^{-5}" and "GO:0001637, G protein–coupled chemoattractant receptor activity,
P=2.9 x 10^{-5}" (Supplementary Table S13). Moreover, three molecular function categories
associated with NADH dehydrogenase activity were significantly enriched in reindeer-specific
gene families (Supplementary Table S13).

We further examined reindeer-specific gene families to search for genes related to adaptation to
subarctic climatic and other environmental factors, such as cold tolerance, marked fluctuations in
daylight and feed shortage during winter. Several genes (*AKAP5, CACNB2, CEBPB, CEBPE,*
COL1A2, CYSLTR1, DNAJB6, EIF2B2, FOXE3, GATA3, GNAO1, MICB, PKD1, PRDM7,
PRDM9, RPS6, TRPV5 and *TRPV6*) (Supplementary Data 1) from our list of reindeer-specific
gene families have been reported to be associated with cold adaptation in Siberian human
populations[34]. We also identified genes related to unique characteristics of the reindeer species[4],
such as vitamin D metabolism (*TRPV5* and *TRPV6*) and antler development (*SILT2*). In addition,
we found that the melanopsin gene (*OPN4B*), which is a photoreceptor, has an important
function in the retina[35–37].

Fig. 1. Gene ontology comparison among genomes of reindeer and nine mammalian species. A) Distribution of gene clusters among the compared mammalian species. **B**) Venn diagram showing unique and shared gene families among five ruminants (cattle, goat, reindeer, sheep and yak). **C**) Gene expansion and contraction in the reindeer genome. The numbers of gene families that have expanded (green, +) and contracted (red, -) after splitting are shown on the corresponding branch. **D**) Phylogenetic tree based on 4-fold degenerate sites of 7,951 single-copy orthologous genes. Estimates of divergence time with 95% CI are shown at each node.

Expanded and contracted gene families. Based on the comparison of orthologous gene families among the 10 species, the reindeer genome revealed 3,312 and 1,840 expanded and contracted gene families, respectively (Fig. 1C). The significantly (P<0.01) expanded (n=368) and contracted (n=15) gene families contained 2,683 and 19 genes, respectively (Supplementary Data 3 and Data 4). A high number of expanded genes were associated with ribosomal proteins (n=658 genes), zinc finger proteins (n=155 genes) and olfactory receptors (n=35 genes) (Supplementary Data 3). We performed functional enrichment analysis for only the 368 expanded (2,688 genes) gene families (Supplementary Table S14). The expanded genes

revealed 163 significant (FDR <0.05) GO terms, of which 122 were categorized into biological
processes and 41 were associated with molecular function (Supplementary Table S14 and Data
5). The expanded gene families enriched in biological processes were mainly associated with
metabolic processes (n=33 GO terms), biosynthetic processes (n=23 GO terms), ion
homeostasis (n=14 GO terms) and transport (n=14 GO terms) (Supplementary Table S14 and
Data 5). Similarly, the expanded gene families enriched in the molecular function category were
dominated by binding activities (n=16 GO terms), such as "GO:0008199, ferric iron binding,
P=4.2 x 10^{-57}", "GO:0005506, iron ion binding, P=2.5 x 10^{-17}", "GO:0031492, nucleosomal
DNA binding, P=9.2 x 10^{-13}" and "GO:0043566, structure-specific DNA binding, P=9.2 x 10^{-13}".
214

We further explored expanded gene families to search for genes related to arctic and subarctic
adaptation and unique reindeer characteristics. Several genes (*ADRA2A, ADRA2B, ADRA2C,*
CIDEA, CRYAB, CYCS, DNAJA1, DRD3, EPHA3, EPHB1, FOXC1, FOXC2, GNG5, HMGN3,
HOXA5, HSF1, HSPE1, HTR1B, HTR2A, PARK2, PPP2R1A, PRDM1, PRDM7, PRDM9,
RAP1B, RHOC, RPS6, SLC25A5, SLC8A1, TRPV5 and *TWIST1)* (Supplementary Data 3) that
were predicted to be in expanded gene families have been reported to be associated with cold
adaptation in Siberian human populations[34]. Moreover, we also identified genes related to
vitamin D metabolism (*TRPV5*) among the list of expanded gene families[4].

Genes under positive selection in reindeer. We conducted an analysis of the signature of
positive selection by assessing dN/dS ratios of 7,951 single-copy orthologous gene sets found in
the genomes of *R. tarandus* and the other nine mammalian species. A total of 418 of the 7,951
(5.26%) single-copy orthologous genes in reindeer revealed a significant signature of positive
selection (P<0.01) (Supplementary Data 6). Analysis of the GO enrichment for the genes under
positive selection in reindeer revealed seven statistically significant (FDR ≤ 0.05) enriched GO
terms particularly related to channel activities (n=6 GO terms) (Supplementary Table S15).

A number of genes (*BDKRB2, CORIN, GCG, GNB1, IL6, KCNMA1, KCNMB2, MED1,*
MLXIPL, NPY, NTS, PRDM10, PRKCQ, RNPEP, SHC1, SORBS3, STRADA, TCF7L2, TRPC1
and *VEGFA*) (Supplementary Data 6) showing positive selection in reindeer have been reported
to be associated with cold adaptation in Siberian human populations[34]. Moreover, we noticed that
a gene regulating circadian rhythm (*GRIA1*)[4] showed signs of positive selection.

Evolutionary divergence analysis. We also estimated the evolutionary divergence of reindeer
by comparing the assembled reindeer genome with four other domesticated ruminant species
(goat, sheep, yak and taurine cattle), dromedary camel, horse, polar bear, dog and human. A
phylogenetic tree was constructed based on 4-fold degenerate codon sites extracted from 7,951
single-copy orthologous genes identified by TreeFam[38]. The estimated divergence time showed
that reindeer shared a common ancestor with other domesticated ruminant species approximately
36 million years ago (95% CI 32-42 million years ago) (Fig. 1D). Our divergence estimate is
longer than the divergence estimate from a previous reindeer study[21]. Our analysis suggests that
the divergence between the domesticated ruminant species and the dromedary camel occurred in
the early phase of *Artiodactyla* evolution[39], 56 million years ago.

Demographic history

We applied the pairwise sequentially Markovian coalescent (PSMC)[40] method to the reindeer
reference genome to examine the changes in the effective population size (N_e) of the ancestral
population of the Fennoscandian reindeer in the course of the Pleistocene period in the world's
history (Fig. 2). The N_e of the ancestor of reindeer gradually decreased between 1 million years
ago and 500 thousand years ago (Kya). The ancestral N_e of reindeer showed peaks at 150 Kya
and 20 Kya, while the population underwent three major bottlenecks at 600 Kya, 40 Kya and 11
Kya.

**Fig. 2. Demographic history of *R. tarandus* reconstructed from the draft reference genome
by using PSMC.** The X axis shows the time in thousands of years (Kyr), and the Y axis shows
the effective population size.

Heterozygosity estimation

Using the assembled draft reindeer genome sequence as a reference, we mapped the reads of
short insert-size libraries to the genome assembly of reindeer and found high coverage, ensuring
a high level of accuracy at the nucleotide level. Approximately 98.34% of the reads were
successfully mapped to the genome, and the sequencing coverage was approximately 60X. We
used the unique map reads to call variants using the Genome Analysis Toolkit (GATK)[41]
best practice pipeline. We identified a total of 5.46 million (M) heterozygous SNPs in the

264 genome. The estimated heterozygosity rate of reindeer was 2.05 x 10e-3, which is 3.48 and 2.3
265 times higher than that of cattle (0.59×10^{-3}) and yak (0.89×10^{-3})[25].

Whole-genome sequence analysis

The raw sequences for the 23 reindeer samples were generated at Beijing Genomics Institute
(BGI) using the Illumina HighSeq 4000 platform. The raw reads were preprocessed; adapters and
low-quality data were removed. After the data were processed, we generated a total of 680 Gb
clean paired-end resequence data (Supplementary Table S16). On average per individual, we
achieved 197 Mb and 29.57 Gb clean reads and bases, respectively (Supplementary Table S16.)
On average per individual, 98.78% of the clean reads were successfully mapped to the reindeer
genome assembly (Supplementary Table S17) and represented 9.46-fold coverage.

A total of 28.6 M high-quality SNPs were detected in the mapped reads across all 23 individuals.
The average number of SNPs detected per individual was 7.24 M (Supplementary Table S18). In
the wild tundra reindeer in Russia (four samples collected at two sites), 8.00 M SNPs were
identified on average per individual; in the wild Norwegian tundra reindeer, 7.09 M (five
animals) SNPs were identified; and in the domestic tundra reindeer, 6.79 M (four animals from
two sites) SNPs were identified. Three arctic reindeer samples from Svalbard Norway showed
low diversity, with an average of 1.37 M (20.42%) heterozygous and 5.35 M (79.58%)
homozygous SNPs per individual, while in the rest of the samples, the respective estimates were
3.81 M (52.14%) and 3.50 M (47.86%). We also detected a total of 3.93 M indels across all 23
samples, and on average, we detected approximately 895,386 indels per individual.

Functional annotation of SNPs. SnpEff[42] was used to annotate all filtered SNPs identified
across all samples, and the proportions of SNPs in the intergenic, intronic, downstream and
upstream, and transcript regions were 58.27%, 14.0%, 12.52% and 14.61%, respectively, while a
small number of variants were annotated in exonic (0.57%) regions (Supplementary Table S19 A
and B). The overall estimated missense to silent ratio for all samples was 0.7963. The average
SNP-variant rate was 1 variant every 90 bases for all samples. The transition to transversion
(Ts/Tv) ratio was calculated for each sample, and the average Ts/Tv ratio was 1.97 for all
reindeer samples, which is slightly lower than that of human (2.1)[43] and lower than that of bovine
(2.2)[44–46].

Population genetic analysis. Genetic relationships between the 23 resequenced animals and the
reference animal were investigated with a principal component analysis (PCA). The PCA plot
revealed two major groups Fennoscandia reindeer versus forest and tundra reindeer/caribou from
Russia and Alaska, as shown in Fig. 3. The reindeer from Svalbard and Novaja Zemlya, both
wild arctic ecotype, grouped separately from the main clusters. We computed genetic diversity
parameters for the two main clusters. The overall genome-wide genetic diversity, as measured by
Watterson's (θ) and pairwise nucleotide diversity (π), was smaller in the Fennoscandian group
(0.002022733 and 0.002096655, respectively) than in the Russian – Alaskan group
(0.002441424 and 0.002265819, respectively). The genetic differentiation between the two main
groups, as measured by F_{st}, was 0.039786, suggesting that these two cluster populations exhibit
low differentiation.

Fig. 3. PCA of 23 resequenced animals and the reference individual. PCA plot. (FI-R-R, Finnish reindeer (i.e. the reference animal); Fe-W-T, Fennoscandia wild tundra reindeer; Fe-D-T, Fennoscandia domestic tundra reindeer; Ru-W-T, Russian wild tundra reindeer; Ru-D-T, Russian domestic tundra reindeer; Ru-D-F, Russian domestic forest reindeer; Ru-W-A, Russian wild arctic reindeer (i.e. Novaia Zemlia); Sv-W-A, Svalbard wild arctic reindeer; Al-D-T, Alaska Domestic tundra reindeer; Al-W-T, Alaska wild tundra caribou)

Assembly and annotation of the reindeer mitochondrial genome

The reindeer mitochondrial genome was assembled using the mitochondrial baiting and iterative mapping (MITObim)[47] approach. A total of 16,962 reads were utilized in the assembly, corresponding to 0.036% of the reads generated from the 170 bp insert-size library. The reads used in the assembly mapped with 84% of bases mapping with quality >Q20. The assembled mitochondrial genome is 16,357 bp in length with an average conversion of 131.84X and a GC content of 36.28%. The reindeer mitogenome consists of 22 tRNAs, 13 protein-coding genes and two rRNAs (Supplementary Table S20).

Discussion

With our present study and the recent publications by Li et al. (2017)[21] and Lin et al. (2019)[4], a new genomic era in the study of reindeer has been established. Here, we have generated and described an alternative draft genome assembly for reindeer and published the largest resequenced *Rangifer* sp. dataset for the investigation of genetic diversity, domestication, demographic aspects and genomic adaptation.

Our reindeer genome assembly is 2.66 Gb in size and represents ~91% of the estimated 2.92 Gb genome size according to the k-mer analysis. The estimated genome size of reindeer in our study

is slightly higher than that in two recent studies, i.e., 2.86[48] Gb and 2.76 Gb[21]. Species belonging
to the *Cervidae* family[48], such as Indian muntjac (2.92 Gb), Chinese muntjac (2.99 Gb), Milu
(3.09 Gb), Black muntjac (3.24 Gb) and White-lipped deer (3.23 Gb), appear to have larger
genome sizes than domesticated mammalian species in general (~2.6–2.7 Gb), such as pig (*Sus*
scrofa[18]), yak (*B. grunniens*[25]), dromedary camel (*C. dromedarius*[49]) and sheep (*O. aries*[19]). In
addition, the number of protein-coding annotated genes (>27,000) in our genome assembly draft
is higher than that typically predicted in the latest assemblies of several other domesticated
mammalian species (~21,000)[18–20,25,33] and in the first genome draft of reindeer (21,555)[21]. We
assume that the differences in the genome size and the gene numbers may be due to the
annotation approach (homology-based approach) used in this study and will require further
validation. However, the BUSCO assessment revealed 97% orthologous genes in the present
assembly, which is better than the BUSCO assessment results in a previous study[21], thus
indicating the high quality of our reindeer gene annotation. In addition, the key gene parameters
(see Supplementary Table S7 and Fig. S4) also revealed no significant differences between the
reindeer genome and the five species that were used in homology predication related to gene and
exon length distribution.

We applied the present draft assembly to estimate the evolutionary divergence times between
reindeer and nine other mammalian species and to reveal recent demographic events in the
history of reindeer that occurred during the last 1 million years. Our phylogeny suggests that
reindeer diverged from four other domestic ruminants ca. 36 million years ago, which is earlier
than previously estimated [21,48]. The demographic analysis (Fig. 2) showed that there have been
marked fluctuations in the N_e of the reindeer species: two population expansions and three
bottlenecks. The decline in the N_e occurred during the mid-Pleistocene period, and the peak
during Last Glacial Maximum period may have been associated with global temporal changes in
climate [50–52]. The pattern of the N_e of the reindeer genome during the Pleistocene period is
consistent with the previously suggested hypotheses of the responses of other domestic
mammalian species, such as pig (*S. scrofa*[18]), horse (*E. caballus*[53]), sheep (*O. aries*[54]) and cattle
(*B. taurus*[55]), to temporal climate changes in the past. However, the heterozygote rate (2.05 X 10$^-$
^{3}) of the reindeer genome estimated in our analysis was 3.48 and 2.3 times higher than that of *B.*
taurus-cattle and *B. grunniens*-yak, respectively[25], suggesting a larger founder population size of
the contemporary semi-domesticated reindeer. In addition, possible gene flow from wild reindeer
populations, a less intensive artificial selection history and the early phase of reindeer
domestication history in general may have contributed to higher diversity estimates for reindeer
than for the two other ruminant species[15].

To explore the genetic distinctiveness of reindeer in the present research context, we focused on
reindeer-specific and expanded orthologous gene families and their genes and gene functions.
The changes in genetic makeup (i.e., gene copy and/or protein domain number changes) are
potentially linked to unique phenotypic and functional changes, suggesting a mechanism for
adaptive divergence of closely related species[24,25,56]. Gene family analysis revealed that 160
genes were shared between 547 reindeer-specific gene families and 368 expanded gene families.
Therefore, the 160 genes that we identified as unique to reindeer compared to four other
ruminants are also rapidly evolving (among nine other mammals) and may thus represent
characteristic features of reindeer. Zinc finger proteins (n=42) dominated the list of uniquely
expanding genes followed by olfactory receptors (n=13). Zinc finger proteins have multi-

functional roles, including lipid binding and the differentiation of adipose tissues[57,58].
Interestingly, among the list of zinc finger proteins was PR domain zinc finger protein 9
(*PRDM9*), which is considered one of the fastest evolving genes in mammalian species. *PRDM9*,
also referred to as the speciation gene, has an important function in enhancing recombination and
epigenetic modification[59,60]. We speculate that the *PRDM9* gene has played a vital role in the
evolution and adaptation of *R. tarandus* to challenging northernmost biogeographic and climatic
conditions. Another member of the zinc finger protein, PR domain zinc finger protein 1
(*PRDM1*), is associated with retinal development[61–63] and is a candidate gene promoting
adaptation to extreme seasonal light changes. Furthermore, the improved sensory systems of
reindeer were also reflected in our study, where we observed several genes associated with
olfactory receptors and G protein–coupled receptors in specific and expanded gene families
(Supplementary Data 1 and Data 3). The mammalian genome contains 4–5% of olfactory
receptor genes, which are sensors to the extracellular environment and essential for the survival
of animals[64]. G protein–coupled receptors are also known to be involved in sensing of the
extracellular environment[64]. During the long winter season, the reindeer sense of smell or
olfaction has an important role in finding feed covered with snow.

Moreover, the examination of genes in the reindeer-specific and expanded orthologous gene
families provided information on genes associated with specific phenotypic characteristics of
reindeer. We identified two genes belonging to the transient receptor potential (TRP) cation
channel subfamily, *TRPV5* and *TRPV6,* among the reindeer-specific gene families. *TRPV5* and
TRPV6 are involved in calcium reabsorption and maintenance of blood calcium levels in higher
organisms[65,66]. Among these, *TRPV5* has been found to be associated with vitamin D
metabolism[4]. Previous studies have reported that reindeer require efficient calcium metabolism
and reabsorption for antler growth during the period of continuous winter darkness and low solar
energy[67,68]. Studies also indicated that efficient vitamin D metabolism is needed to sustain
calcium metabolism and reabsorption[67,68]. The nucleotide sequences of both the *TRPV5* and
TRPV6 genes harbour 15 exons, encode proteins of approximately 730 amino acids and have
75% sequence similarity[65,66]. Both genes have similar functional properties and regulation
mechanisms[65]. We speculate that in addition to *TRPV5, TRPV6* also has a significant role in
vitamin D metabolism in reindeer. In addition, three adaptive immune-related chemokines
(*CXCR1*, *CXCR2*, *CXCR4*)[69–71] and three interferons (*IFNT1*, *IFNT3*, *IFNW1*) that have antiviral
functions[72,73] may have promoted the adaptation of reindeer to the given environment. Finally,
insulin-like growth factor 1 receptor (*IGF1R*) promotes the activity of *IGF1,* which has a pivotal
role in antler formation (*IGF1* regulates *RUNX1* expression via IRS1/2 signalling for antler
growth)[74,75].

We also observed large numbers of ribosomal proteins present in the reindeer-specific and
expanded gene families. A number of reports have suggested that ribosomal proteins are
essential for survival and adaptation to environmental stress[76–78]. For example, one of the
ribosomal proteins we found was the ribosomal large subunit protein 7 (*RPL7*), which has been
previously reported in Russian cattle breeds[79] to be related to adaptation to harsh (cold)
environments. The ribosomal large subunit protein 7 (*RPL7*) gene was found to be upregulated in
the skin of freeze-tolerant wood frogs compared to non-tolerant species, showing that this gene is
cold-responsive and that the gene was upregulated in skeletal muscle and the brain under
freezing stress[78]. Moreover, in the list of reindeer-specific gene families, we found opsin 4B

(*OPN4B)*, which is a photoreceptor that appears to be involved in the regulation of the circadian
clock[35–37]. Several studies[37,80–82] have reported the existence of a melanopsin-associated
photoreceptive system in the mammalian retina that helps to transmit photic information and is
also involved in the regulation of photoentrainment of the circadian clock. A previous study[9]
reported that reindeer possess a tapetum lucidum (reflective surface behind the central retina) to
cope with winter darkness, which may help in changing wavelength reflection and increase the
sensitivity of the animal's vision in dim light. In winter, the reindeer tapetum lucidum changes to
blue, which may help scatter light through photoreceptors, and less light is reflected[9]. We
speculate that this melanopsin-related gene might have important functions for the adaptation to
the light system in circumpolar environments. Furthermore, we found Slit homolog 2 protein
(*SLIT2*) in specific gene families, which has been reported to be associated with antler
development in a recent reindeer study[4].

In our study, signs of positive selection were found in genes in the genome of *R. tarandus* mainly
enriched in GO terms related to channel activity (Supplementary Table S15), such as sodium
channel activity and ion channel activity, which are thought to be relevant in cold sensing
mechanisms[83]. The ability to detect and adapt to cold temperature is crucial for the survival of an
organism[84,85]. The process of sensory transduction and a large array of ion channels, such as Na^+
and K^+ channels, are involved in cold temperature detection[83]. Among the genes categorized into
these GO terms, interestingly, *SCN11A1* has an essential role in pain tolerance caused by cold
stress[86,87], suggesting the gene's role in survival in extremely cold environments. One additional
gene showing signatures of positive selection, glutamate ionotropic receptor AMPA type subunit
1 (GRIA1), appears to have a role in the circadian clock[4].

Finally, we investigated the genetic relationships between the *de novo* reference animal and the
23 resequenced individuals, and two main genetic clusters (Fennoscandia and Russia-Alaska)
were identified in our data (Fig. 3). We suggest that this observation reflects the historical spread
of reindeer populations in northern Eurasia and reveals the domestication history of reindeer. The
Russian/northern American cluster probably reflects the Euro-Beringian lineage evolved from
Pleistocene population in northern Eurasia[3,88] while the Fennoscandia cluster may have
descended from some refugia populations in southern Europe partly isolated from the Euro-
Beringian lineage[89]. In these two main genetic clusters, we found both domestic and wild
reindeer from respective Fennoscandia and Russia, which gives support to previously indicated
independent domestication origin for the breeds in these regions[13,90]. Our findings are consistent
with the previous study[15]. Our data suggest that the semi-domesticated tundra reindeer show
lower genetic diversity in terms of the number of SNPs per individual than the wild tundra
reindeer, implying a genetic bottleneck effect caused by domestication as also seen in temporal
change in mitochondrial DNA[13,91]. Moreover, our data clearly indicated the effects of geographic
isolation and genetic bottleneck in the Svalbard reindeer: the subspecies (or an arctic ecotype)
displayed exceptionally low diversity compared to the other resequenced Rangifer populations.

Here, we have reported a 2.66 Gb genome assembly of a Finnish male reindeer originating from
Sodankylä (67.34°N, 26.83°E), where the annual mean temperature is 0 °C (the mean
temperature in January and February -13 °C), the snow cover is on average 202 days per year,

the daylight is less than 3 h from December until mid-January (during a short period in
December, the sun is down all day) and the sun is up all day from June until mid-July
(https://www.timeanddate.com/). The first draft assembly of the reindeer genome was obtained
from a female animal[21] from Inner Mongolia Autonomous Region, China (50.77°N, 121.47°E),
where the climate is continental, but the annual daylight rhythm (at least 8 h of daylight and no
periods of total darkness or "nightless nights" exist) is different from that typically found in
traditional reindeer herding regions in northern Eurasia. In addition to these climatic differences
in their geographic origins, we assume that the Finnish reindeer and Inner Mongolian reindeer do
not share their genetic origins in the same refugial, ancestral populations. The Finnish reference
reindeer was found to belong to the cluster of northern Fennoscandia reindeer distinct from the
"eastern" and North American genetic cluster. It is recommended that phylogenetically different
animal populations have their own reference assemblies[55]. With our GO family-based
comparisons with nine other mammalian genomes, we identified hundreds of genes and gene
families that are unique, under positive selection and rapidly expanding in reindeer. Exploring
these genes and gene families revealed several reindeer-specific characteristics that have helped
these animals survive in the arctic and subarctic conditions. We identified genes that are
important for vitamin D metabolism (*TRPV5*, *TRPV6*), antler formation (*SLIT2*), and circadian
rhythm (*OPN4B*). While many of the findings from our study complement reports from other
species, the functions of *TRPV6* and *OPN4B* have been reported here for the first time.
Population genomics analyses based on 23 individuals representing domestic and wild reindeer
suggested two domestication events in *R. tarandus* subspecies, but additional larger scale studies
are required for validating this finding. Collectively, our study provides new insight into the
genomic and evolutionary characteristics of reindeer, and the resequencing data (the most
comprehensive in reindeer thus far) will serve as an invaluable resource for future studies
involving reindeer and other cervids.

Materials and methods

Sampling and genome sequencing

All protocols and sample collections were performed in accordance with the legislations
approved by the Animal Experiment Board in Finland (ESAVI/7034/04.10.07.2015).

A one-year-old male reindeer *(R. tarandus)* from Sodankylä, Finland, was selected, and blood
samples were collected for sequencing. Genomic DNA was extracted from the blood using a
standard phenol/chloroform method and sequenced at high coverage on the Illumina HiSeq 2500
and 4000 platforms using a shotgun-sequencing approach. We constructed seven paired-end
DNA libraries with insert sizes ranging from 170 bp to 20 kb. Summary statistics of the
generated reads are shown in Table 1 and Supplementary Table S1. The short reads with low-
quality bases were removed before assembly by filtering as follows:

1. Reads from short insert-size libraries having 'N' over 2% of its length, and the reads from
large insert-size libraries having 'N' over 5% of its length.
2. Reads from short insert-size libraries having more than 40% bases with Q20 (quality) less
than or equal to 7, and the reads from large insert-size libraries having more than 30%
bases with Q20 less than or equal to 7.

3. Reads with adapter sequences.
4. Paired-end reads of read 1 and read 2 that were completely identical.

Estimation of genome size using k-mer analysis

For the short reads with an insert size of 500 bp, a total of 64.7 Gb (approximately 21.6X) was
used to estimate the genome size using the k-mer method[22], as described in previous studies[24].
We used the following formula to estimate the genome size (G): G=K_num/K_depth, where
K_num is the total number of k-mers, and K_depth is the frequency occurring more frequently
than other frequencies. In this study, the size of k-mer for reindeer was set to 17, K_num was
55,451,706,382 and K_depth was 19. Hence, the genome size was estimated to be approximately
2.92 Gb (Supplementary Table S2).

Genome assembly

After the data were filtered, clean data were retained for assembly (Table 1, Supplementary
Table S1). The high-quality clean reads were assembled first into contigs and subsequently into
scaffolds using SOAPdenovo V2.04 software[23].

The quality of the genome assembly was evaluated by aligning high-quality reads from short
insert-size libraries to the *de novo* genome assembly using BWA[26] (version. 0.7.15) with no
more than five mismatches and then by determining the percentage of total aligned reads. Reads
from the transcriptome were assembled by trinityrnaseq-2.0.6[92]; all sequences were mapped to
the assembly by BLAT[27] with the default parameters. We also employed BUSCO (version
3.0.2)[28] to assess the completeness of the genome assembly and annotated gene set using the
conserved 4,104 mammalian genes.

RNA sequencing

To improve the reference genome annotation, we extracted RNA using the RNeasy Plus
Universal Kit (Qiagen, Valencia, CA, USA) from six tissues (scapular adipose, metacarpal
adipose, tailhead adipose, perirenal adipose, liver and musculus gluteobiceps) from the same
male reindeer subjected to *de novo* sequencing. The tissue samples were collected at slaughter
and stored in RNA*later*® Solution (Ambion/Qiagen, Valencia, CA, USA) according to the
manufacturer's instructions. mRNA libraries were sequenced using Illumina HiSeq 2500 with a
paired-end (2x150 bp) strategy. The raw data were preprocessed, and adapters and low-quality
reads were filtered out using the cutadapt tool[93].

Genome annotation

Repeat annotation

Tandem repeats were detected in the generated genome assembly using Tandem Repeats Finder
(TRF) software[94]. TEs were predicted in the genome using the combination of both known and
de novo strategies. For the known-based strategy, we identified known TEs in the genome using
RepeatMasker[95] and RepeatProteinMask[96] by homology search against the Repbase[30] library

with default parameters. For the *de novo* way, we constructed a *de novo* repeat library using
RepeatModeler[31], with the default parameters. RepeatMasker was used again with the *de novo*
libraries to identify new TEs in the genome. All repeats obtained by the different methods were
combined according to the coordinate in the genome and overlapping TEs belonging to the same
repeat class to form a non-redundant list of reindeer repeats.

Protein-coding gene prediction

We employed both a homology-based and an RNA-assisted approach to predict protein-coding
genes as described by Curwen et al. (2004)[29]. First, we downloaded protein sets from five
sequenced animal genomes (*H. sapiens*, *M. musculus, B. taurus, C. dromedarius and C.*
familiaris) from Ensembl (release 89)[97] and mapped them onto the reindeer genome using
tblastn[98]. Second, high-score segment pairs (HSPs) were concatenated between the same pair of
proteins by Solar (in-house software, version 0.9.6). Third, homologous genome sequences were
aligned against the matching proteins using GeneWise[99] to define accurate gene models. Then,
we filtered redundancy based on the score from GeneWise. Finally, we combined the annotated
results from the homologue-based method to form the final genome annotation as described by
Curwen et al. (2004)[29]. Functions of genes were assigned based on the best match derived from
the alignments to proteins annotated in four protein databases: InterPro[100], KEGG[101], Swiss-
Prot[102] and TrEMBL[103].

ncRNA annotation

We used INFERNAL[104] and tRNAscan-SE[105] to predict ncRNA. Four types of ncRNAs were
annotated in our analysis: tRNA, rRNA, miRNA and snRNA. tRNA genes were predicted by
tRNAscan-SE with eukaryotic parameters. rRNA fragments were identified by aligning the
rRNA template sequences from human genomes using BLASTN with an E-value cutoff of $1e^{-5}$.
miRNA and snRNA genes were inferred by the INFERNAL software against the Rfam database
(v11.0)[106].

Gene family construction

To define gene family evolution in the reindeer genome, we used the TreeFam methodology[38] as
follows: BLAST was used to compare all protein sequences from 10 species, *R. tarandus, C.*
dromedarius, C. hircus, O. aries, B. taurus, B. grunniens, E. caballus, C. familiaris, U.
maritimus and *H. sapiens*, with the E-value threshold set to 1e-7. After that, HSPs of each
protein pair were concatenated by Solar software. H-scores were computed based on Bit-scores,
and these were taken to evaluate the similarity among genes. Finally, gene families were
obtained by clustering homologous gene sequences using Hcluster_sg (version 0.5.0). If these
genes had functional motifs, they were annotated by GO[107].

Phylogenetic tree construction and divergence time estimation

The phylogenetic tree was constructed based on the single-copy orthologous genes from the ten
species identified by gene family analysis. CDS from each single-copy gene cluster were aligned
using MUSCLE[108], and their protein sequences were concatenated to form one super gene for

every species. Four-fold degenerate sites (4d sites) of aligned CDS were extracted for subsequent
analysis. The phylogenetic relationships were inferred using PhyML software[109,110], with the
GTR substitution model and gamma distribution rates model as the chosen parameters. We
constructed a phylogenetic framework including six *Artiodactyla* and four other vertebrates, with
H. sapiens as the outgroup. Divergence times were estimated using PAML mcmctree (PAML
version 4.5)[111–113] by implementing the approximate likelihood calculation method.

Expansion and contraction of gene families

We analysed the expansion and contraction of gene families using the CAFE program[114], which
employs a random birth and death model across a user-specified phylogeny. The global
parameter λ, which describes both the gene birth (λ) and death ($\mu = -\lambda$) rate across all branches
in the tree for all gene families, was estimated through maximum likelihood. A conditional P-
value was calculated for each gene family, and families with conditional P-values less than the
threshold (0.01) were considered to have notable gains or losses. Finally, the branches
responsible for low overall P-values were defined as significant families.

Detection of positively selected genes (PSGs)

To estimate positive selection, dN/dS ratios were calculated for all single-copy orthologues of *R.*
tarandus and nine other vertebrates (*C. dromedarius, C. hircus, O. aries, B. taurus, B. grunniens,*
E. caballus, C. familiaris, U. maritimus and *H. sapiens*). Orthologous genes were first aligned by
PRANK[115], then Gblocks 0.91b was used to remove ambiguously aligned blocks within PRANK
alignments and 'codeml' in the PAML package[113] was employed with the free-ratio model to
estimate Ka, Ks, and Ka/Ks ratios of different branches. The difference in mean Ka/Ks ratios for
single-copy genes between *R. tarandus* and each of the other species were compared with paired
Wilcoxon rank-sum tests. The genes that showed values of Ka/Ks higher than 1 along the branch
leading to *R. tarandus* were reanalysed using the codon-based branch-site tests implemented in
PAML. The branch-site model, which allowed ω to vary both at sites in the protein and across
branches, was used to detect episodic positive selection.

SNP calling and heterozygosity estimation

We used high-quality reads from the short insert-size libraries (170, 500 and 800 bp) to call
SNPs. The quality reads were mapped against the assembly using BWA[26] (version. 0.7.15)
software with the default parameter. After mapping, we employed Picard tools
(http://broadinstitute.github.io/picard/) (version.2.5.0) to preprocess alignment reads and remove
PCR duplicates, and then the uniquely mapped reads were used for SNP calling. Further, we
used GATK (version. 3.6)[41] to identify poorly mapped regions nearby indels, realigned and
performed SNP calling according to the GATK best practice pipeline[116]. We estimated the
heterozygosity rate of the reindeer genome; first, we counted the identified heterozygous SNPs
and then divided the total heterozygous SNPs by the effective genome size.

Population history estimation

We inferred a marked population bottleneck in the demographic history of *R. tarandus* using the pairwise sequentially Markovian coalescent (PSMC) model[40] with generation times and mutation rates.

Assembly and annotation of the reindeer mitochondrial genome

Complete mitochondrial genomes are useful materials for phylogenetic studies. The complete mitochondrial genome for reindeer was assembled using MITObim v 1.9 software[47]. Among the reads generated from the seven libraries on the reference individual, for the mitochondrial genome assembly, we used only reads from the 170 bp insert-size library (10% of 66.2 Gb, Table S1) for MITObim input. The short DNA sequence (506 bp), cytochrome oxidase subunit 1 (COI) (GenBank accession number COI: KX085230.1)[12] from *R. tarandus* was used to seed the initial baiting of mitochondrial reads. Following the assembly, the *de novo* assembly was annotated using the MITOs web server[117] with the default parameters.

Whole-genome resequencing and variant calling of 23 domestic and wild reindeer

To investigate genome variation and perform population analyses, we resequenced 23 domesticated and semi-domesticated reindeer individuals from Russia, Norway, Alaska and USA. Resequenced samples originated from the following populations and subspecies: domesticated forest reindeer from Russia (n=2), domesticated tundra reindeer from Norway (n=4), wild island tundra-mountain reindeer from Russia (n=2), wild tundra reindeer from Russia (n=4), wild tundra reindeer from Norway (n=5), wild arctic reindeer from Svalbard Norway (n=3), Alaskan domestic reindeer from Alaska-USA (n=2) and Alaskan wild caribou from Alaska-USA (n=1). Genomic DNA was extracted from 23 blood samples (obtained from the previous study by Flagstad and Røed 2003[3]), and sequence libraries with an average fragment size of 500 bp were constructed for each individual. Paired-end reads were generated using Illumina HiSeq 4000 at BGI. After sequencing, the raw reads were filtered to remove adapter sequences, contamination and low-quality reads. Following the data treatment, we achieved an average of 197 Mb and 29.57 Gb clean reads and bases, respectively (Supplementary Table S16.)

The clean reads were mapped against the present draft reindeer assembly genome using BWA with the default parameters. Using Picard tools, we preprocessed and filtered alignments for SNP calling, including the removal of low-quality alignments, the sorting of alignments and the removal of PCR duplicates. Further, using GATK, we identified poorly mapped regions nearby indels from the alignments and performed a realignment and base quality recalibration on the bases that were disrupted by indel sites. Finally, using the GATK best practice pipeline, we performed SNP discovery and genotyping across the 23 samples.

Functional annotation of SNPs

SnpEff V4.3T[42] was used to annotate the SNPs and categorize them into coding (synonymous and non-synonymous), upstream/downstream and intronic/intergenic classes. For this analysis,

we constructed the SnpEff databases using the draft reindeer reference genome and the
corresponding GTF genome annotation file.

Population genetics statistics

First, to examine the genetic relationships across the 23 reindeer samples, we conducted PCA
with smartpca in EIGENSOFT3.0 software[118] using the detected SNPs in each individual.
Second, using the PCA plot result, we divided the 23 samples according to their grouping in the
PCA plot and computed the average pairwise nucleotide diversity within a population π and the
proportion of polymorphic sites Watterson's θ for the main clusters using the
Bio::PopGen::Statistics package in BioPerl (v1.6.924)[119]. The same program was used to
compute the population differentiation estimate, with pairwise F_{st} between the two population
groups (i.e., the PCA cluster result).

References

1. Helskog, K. & Indrelid, S. Humans and reindeer. *Quat. Int.* **238,** 1–3 (2011).

2. Bjørklund, I. Domestication, Reindeer Husbandry and the Development of Sámi
Pastoralism. *Acta Boreal.* **30,** 174–189 (2013).

3. Flagstad, O. & Roed, K. H. Refugial origins of reindeer (Rangifer tarandus L.) inferred
from mitochondrial DNA sequences. *Evolution* **57,** 658–670 (2003).

4. Lin, Z. *et al.* Biological adaptations in the Arctic cervid, the reindeer (*Rangifer tarandus*).
Science (80-.). **364,** eaav6312 (2019).

5. Blix, A. S. Adaptations to polar life in mammals and birds. *J. Exp. Biol.* **219,** 1093 LP-
1105 (2016).

6. Van Oort, B. E. H. *et al.* Circadian organization in reindeer. *Nature* **438,** 1095–1096
(2005).

7. Lu, W., Meng, Q.-J., Tyler, N. J. C., Stokkan, K.-A. & Loudon, A. S. I. A Circadian Clock
Is Not Required in an Arctic Mammal. *Curr. Biol.* **20,** 533–537 (2010).

8. Stokkan, K.-A., Van Oort, B. E. H., Tyler, N. J. C. & Loudon, A. S. I. Adaptations for life
in the Arctic: evidence that melatonin rhythms in reindeer are not driven by a circadian
oscillator but remain acutely sensitive to environmental photoperiod. *J. Pineal Res.* **43,**
289–293 (2007).

9. Karl-Arne, S. *et al.* Shifting mirrors: adaptive changes in retinal reflections to winter
darkness in Arctic reindeer. *Proc. R. Soc. B Biol. Sci.* **280,** 20132451 (2013).

10. H. Miller, J., Patrick, D. & Volker, B. Antlers on the Arctic Refuge: capturing multi-
generational patterns of calving ground use from bones on the landscape. *Proc. R. Soc. B*

Biol. Sci. **280,** 20130275 (2013).

11. Anderson, D. G., Kvie, K. S., Davydov, V. N. & Røed, K. H. Maintaining genetic
integrity of coexisting wild and domestic populations: Genetic differentiation between
wild and domestic Rangifer with long traditions of intentional interbreeding. *Ecol. Evol.* **7,**
6790–6802 (2017).

12. Kvie, K. S., Heggenes, J. & Røed H., K. Merging and comparing three mitochondrial
markers for phylogenetic studies of Eurasian reindeer (Rangifer tarandus). *Ecol. Evol.* **6,**
4347–4358 (2016).

13. Røed, K. H., Bjørklund, I. & Olsen, B. J. From wild to domestic reindeer – Genetic
evidence of a non-native origin of reindeer pastoralism in northern Fennoscandia. *J.*
Archaeol. Sci. Reports **19,** 279–286 (2018).

14. Weckworth, B. v, Musiani, M., McDevitt, A. D., Hebblewhite, M. & Mariani, S.
Reconstruction of caribou evolutionary history in Western North America and its
implications for conservation. *Mol. Ecol.* **21,** 3610–3624 (2012).

15. Røed, K. H. *et al.* Genetic analyses reveal independent domestication origins of Eurasian
reindeer. *Proceedings. Biol. Sci.* **275,** 1849–55 (2008).

16. Hillier, L. W. *et al.* Sequence and comparative analysis of the chicken genome provide
unique perspectives on vertebrate evolution. *Nature* **432,** 695–716 (2004).

17. Bovine Genome Sequencing and Analysis Consortium, C. G. *et al.* The genome sequence
of taurine cattle: a window to ruminant biology and evolution. *Science* **324,** 522–8 (2009).

18. Groenen, M. A. M. *et al.* Analyses of pig genomes provide insight into porcine
demography and evolution. *Nature* **491,** 393–398 (2012).

19. Jiang, Y. *et al.* The sheep genome illuminates biology of the rumen and lipid metabolism.
Science **344,** 1168–1173 (2014).

20. Wade, C. M. *et al.* Genome sequence, comparative analysis, and population genetics of
the domestic horse. *Science* **326,** 865–7 (2009).

21. Li, Z. *et al.* Draft genome of the reindeer (Rangifer tarandus). *Gigascience* **6,** 1–5 (2017).

22. Liu, B. *et al.* Estimation of genomic characteristics by analyzing k-mer frequency in de
novo genome projects. *ArXiv e-prints* **1308,** arXiv:1308.2012 (2013).

23. Luo, R. *et al.* SOAPdenovo2: an empirically improved memory-efficient short-read de
novo assembler. *Gigascience* **1,** 18 (2012).

24. Li, R. *et al.* The sequence and de novo assembly of the giant panda genome. *Nature* **463,**

311–7 (2010).

25.   Qiu, Q. *et al.* The yak genome and adaptation to life at high altitude. *Nat. Genet.* **44,** 946–
949 (2012).

26.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25,** 1754–60 (2009).

27.   Kent, W. J. BLAT--the BLAST-like alignment tool. *Genome Res.* **12,** 656–64 (2002).

28.   Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V & Zdobnov, E. M.
BUSCO: assessing genome assembly and annotation completeness with single-copy
orthologs.

29.   Curwen, V. *et al.* The Ensembl Automatic Gene Annotation System. *Genome Res.* **14,**
942–950 (2004).

30.   Jurka, J. *et al.* Repbase Update, a database of eukaryotic repetitive elements. *Cytogenet.*
Genome Res. **110,** 462–7 (2005).

31.   Smit, A. F. A. & Hubley, R. RepeatModeler Open-1.0. 2008-2010.

32.   Lindblad-Toh, K. *et al.* Genome sequence, comparative analysis and haplotype structure
of the domestic dog. *Nature* **438,** 803–819 (2005).

33.   Dong, Y. *et al.* Sequencing and automated whole-genome optical mapping of the genome
of a domestic goat (Capra hircus). *Nat. Biotechnol.* **31,** 135–141 (2012).

34.   Cardona, A. *et al.* Genome-Wide Analysis of Cold Adaptation in Indigenous Siberian
Populations. *PLoS One* **9,** e98076 (2014).

35.   Drivenes, Ø. *et al.* Isolation and characterization of two teleost melanopsin genes and their
differential expression within the inner retina and brain. *J. Comp. Neurol.* **456,** 84–93
(2003).

36.   Berson, D. M., Dunn, F. A. & Takao, M. Phototransduction by Retinal Ganglion Cells
That Set the Circadian Clock. *Science (80-.).* **295,** 1070 LP-1073 (2002).

37.   Hannibal, J., Hindersson, P., Knudsen, S. M., Georg, B. & Fahrenkrug, J. The
Photopigment Melanopsin Is Exclusively Present in Pituitary Adenylate Cyclase-
Activating Polypeptide-Containing Retinal Ganglion Cells of the Retinohypothalamic
Tract. *J. Neurosci.* **22,** RC191 LP-RC191 (2002).

38.   Li, H. *et al.* TreeFam: a curated database of phylogenetic trees of animal gene families.
Nucleic Acids Res. **34,** D572–D580 (2005).

39.   Consortium, T. B. C. G. S. and A. *et al.* Genome sequences of wild and domestic bactrian
743   camels. *Nat. Commun.* **3,** 1202 (2012).

744   40.   Li, H. & Durbin, R. Inference of human population history from individual whole-genome
745   sequences. *Nature* **475,** 493–496 (2011).

746   41.   McKenna, A. *et al.* The Genome Analysis Toolkit: A MapReduce framework for
747   analyzing next-generation DNA sequencing data. *Genome Res.* **20,** 1297–1303 (2010).

748   42.   Cingolani, P. *et al.* A program for annotating and predicting the effects of single
749   nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster
750   strain w(1118); iso-2; iso-3. *Fly (Austin).* **6,** 80–92 (2012).

751   43.   Lachance, J. *et al.* Evolutionary history and adaptation from high-coverage whole-genome
752   sequences of diverse African hunter-gatherers. *Cell* **150,** 457–469 (2012).

753   44.   Choi, J.-W. *et al.* Whole-Genome Analyses of Korean Native and Holstein Cattle Breeds
754   by Massively Parallel Sequencing. *PLoS One* **9,** e101127 (2014).

755   45.   Choi, J.-W. *et al.* Whole-Genome Resequencing Analysis of Hanwoo and Yanbian Cattle
756   to Identify Genome-Wide SNPs and Signatures of Selection. *Mol. Cells* **38,** 466–473
757   (2015).

758   46.   Stothard, P. *et al.* Whole genome resequencing of black Angus and Holstein cattle for
759   SNP and CNV discovery. *BMC Genomics* **12,** 559 (2011).

760   47.   Hahn, C., Bachmann, L. & Chevreux, B. Reconstructing mitochondrial genomes directly
761   from genomic next-generation sequencing reads-- a baiting and iterative mapping
762   approach. *Nucleic Acids Res.* **41,** e129–e129 (2013).

763   48.   Chen, L. *et al.* Large-scale ruminant genome sequencing provides insights into their
764   evolution and distinct traits. *Science (80-.).* **364,** eaav6202 (2019).

765   49.   Wu, H. *et al.* Erratum: Camelid genomes reveal evolution and adaptation to desert
766   environments. *Nat. Commun.* **6,** 6107 (2015).

767   50.   Barnosky, A. D., Koch, P. L., Feranec, R. S., Wing, S. L. & Shabel, A. B. Assessing the
768   Causes of Late Pleistocene Extinctions on the Continents. *Science (80-.).* **306,** 70 (2004).

769   51.   Hughes, P. D., Woodward, J. C. & Gibbard, P. L. Middle Pleistocene cold stage climates
770   in the Mediterranean: New evidence from the glacial record. *Earth and Planetary Science
771   Letters* **253,** 50–56 (2007).

772   52.   Yokoyama, Y., Lambeck, K., De Deckker, P., Johnston, P. & Fifield, L. K. Timing of the
773   Last Glacial Maximum from observed sea-level minima. *Nature* **406,** 713 (2000).

774   53.   Librado, P. *et al.* The Evolutionary Origin and Genetic Makeup of Domestic Horses.
775   *Genetics* **204,** 423 (2016).

776   54.   Yang, J. *et al.* Whole-Genome Sequencing of Native Sheep Provides Insights into Rapid
777   Adaptations to Extreme Environments. *Mol. Biol. Evol.* **33,** 2576–2592 (2016).

778   55.   Weldenegodguad, M. *et al.* Whole-Genome Sequencing of Three Native Cattle Breeds
779   Originating From the Northernmost Cattle Farming Regions. *Frontiers in Genetics* **9,** 728
780   (2019).

781   56.   Sudmant, P. H. *et al.* Diversity of human copy number variation and multicopy genes.
782   *Science* **330,** 641–6 (2010).

783   57.   Laity, J. H., Lee, B. M. & Wright, P. E. Zinc finger proteins: new insights into structural
784   and functional diversity. *Curr. Opin. Struct. Biol.* **11,** 39–46 (2001).

785   58.   Wei, S. *et al.* Emerging roles of zinc finger proteins in regulating adipogenesis. *Cell. Mol.*
786   *Life Sci.* **70,** 4569–4584 (2013).

787   59.   Grey, C., Baudat, F. & de Massy, B. PRDM9, a driver of the genetic map. *PLOS Genet.*
788   **14,** e1007479 (2018).

789   60.   Hohenauer, T. & Moore, A. W. The Prdm family: expanding roles in stem cells and
790   development. *Development* **139,** 2267 LP-2282 (2012).

791   61.   Brzezinski, J. A., Lamba, D. A. & Reh, T. A. Blimp1 controls photoreceptor versus
792   bipolar cell fate choice during retinal development. *Development* **137,** 619 LP-629 (2010).

793   62.   Brzezinski, J. A., Uoon Park, K. & Reh, T. A. Blimp1 (Prdm1) prevents re-specification
794   of photoreceptors into retinal bipolar cells by restricting competence. *Dev. Biol.* **384,** 194–
795   204 (2013).

796   63.   Katoh, K. *et al.* Blimp1 Suppresses Chx10 Expression in
797   Differentiating Retinal Photoreceptor Precursors to Ensure Proper Photoreceptor
798   Development. *J. Neurosci.* **30,** 6515 LP-6526 (2010).

799   64.   Niimura, Y. Olfactory receptor multigene family in vertebrates: from the viewpoint of
800   evolutionary genomics. *Curr. Genomics* **13,** 103–14 (2012).

801   65.   Peng, J.-B., Suzuki, Y., Gyimesi, G. & Hediger, M. A. TRPV5 and TRPV6 Calcium-
802   Selective Channels. (2018). doi:10.1201/9781315152592-13

803   66.   Van Abel, M., Hoenderop, J. G. J. & Bindels, R. J. M. The epithelial calcium channels
804   TRPV5 and TRPV6: regulation and implications for disease. *Naunyn. Schmiedebergs.*
805   *Arch. Pharmacol.* **371,** 295–306 (2005).

806   67.   Van der Eems, K. L., Brown, R. D. & Gundberg, C. M. Circulating levels of 1,25
807   dihydroxyvitamin D, alkaline phosphatase, hydroxyproline, and osteocalcin associated
808   with antler growth in white-tailed deer. *Acta Endocrinol. (Copenh)*. **118,** 407–414 (1988).

809   68.   Sempere, A. J., Grimberg, R., Silve, C., Tau, C. & Garabedian, M. Evidence for
810   Extrarenal Production of 1,25-Dihydroxyvitamin During Physiological Bone Growth: In
811   Vivo and in Vitro Production by Deer Antler Cells. *Endocrinology* **125,** 2312–2319
812   (1989).

813   69.   Susek, K. H., Karvouni, M., Alici, E. & Lundqvist, A. The Role of CXC Chemokine
814   Receptors 1–4 on Immune Cells in the Tumor Microenvironment   . *Frontiers in
815   Immunology*   **9,** 2159 (2018).

816   70.   Zarbock, A. & Stadtmann, A. CXCR2: From Bench to Bedside   . *Frontiers in
817   Immunology*   **3,** 263 (2012).

818   71.   Costa, M. J. *et al.* Optimal design, anti-tumour efficacy and tolerability of anti-CXCR4
819   antibody drug conjugates. *Sci. Rep.* **9,** 2443 (2019).

820   72.   Samuel, C. E. Antiviral actions of interferons. *Clin. Microbiol. Rev.* **14,** 778–809 (2001).

821   73.   Goodbourn, S., Didcock, L. & Randall, R. E. Interferons: cell signalling, immune
822   modulation, antiviral response and virus countermeasures. *J. Gen. Virol.* **81,** 2341–2364
823   (2000).

824   74.   Yang, Z.-Q. *et al.* IGF1 regulates RUNX1 expression via IRS1/2: Implications for antler
825   chondrocyte differentiation. *Cell Cycle* **16,** 522–532 (2017).

826   75.   Hu, W. *et al.* MicroRNA let-7a and let-7f as novel regulatory factors of the sika deer
827   (Cervus nippon) IGF-1R gene. *Growth Factors* **32,** 27–33 (2014).

828   76.   Jones, P. G. & Inouye, M. The cold-shock response — a hot topic. *Mol. Microbiol.* **11,**
829   811–818 (1994).

830   77.   Thieringer, H. A., Jones, P. G. & Inouye, M. Cold shock and adaptation. *BioEssays* **20,**
831   49–57 (1998).

832   78.   Wu, S., De Croos, J. N. A. & Storey, K. B. Cold acclimation-induced up-regulation of the
833   ribosomal protein L7 gene in the freeze tolerant wood frog, Rana sylvatica. *Gene* **424,** 48–
834   55 (2008).

835   79.   Yurchenko, A. A. *et al.* Scans for signatures of selection in Russian cattle breed genomes
836   reveal new candidate genes for environmental adaptation and acclimation. *Sci. Rep.* **8,**
837   12984 (2018).

838   80.   Provencio, I., Jiang, G., De Grip, W. J., Hayes, W. P. & Rollag, M. D. Melanopsin: An

839   opsin in melanophores, brain, and eye. *Proc. Natl. Acad. Sci.* **95,** 340 LP-345 (1998).

840   81.   Provencio, I. *et al.* A Novel Human Opsin in the Inner Retina. *J. Neurosci.* **20,** 600 LP-
841   605 (2000).

842   82.   Gooley, J. J., Lu, J., Chou, T. C., Scammell, T. E. & Saper, C. B. Melanopsin in cells of
843   origin of the retinohypothalamic tract. *Nat. Neurosci.* **4,** 1165 (2001).

844   83.   Lolignier, S. *et al.* New Insight in Cold Pain: Role of Ion Channels, Modulation, and
845   Clinical Perspectives. *J. Neurosci.* **36,** 11435 LP-11439 (2016).

846   84.   Luiz, A. P. *et al.* Cold sensing by $Na_V1.8$-positive and $Na_V1.8$-negative sensory neurons.
847   *Proc. Natl. Acad. Sci.* **116,** 3811 LP-3816 (2019).

848   85.   Zimmermann, K. *et al.* Transient receptor potential cation channel, subfamily C, member
849   5 (TRPC5) is a cold-transducer in the peripheral nervous system. *Proc. Natl. Acad. Sci.*
850   **108,** 18114 LP-18119 (2011).

851   86.   Lolignier, S. *et al.* The Nav1.9 Channel Is a Key Determinant of Cold Pain Sensation and
852   Cold Allodynia. *Cell Rep.* **11,** 1067–1078 (2015).

853   87.   Leipold, E. *et al.* Cold-aggravated pain in humans caused by a hyperactive NaV1.9
854   channel mutant. *Nat. Commun.* **6,** 10049 (2015).

855   88.   Yannic, G. *et al.* Genetic diversity in caribou linked to past and future climate change.
856   *Nat. Clim. Chang.* **4,** 132 (2013).

857   89.   Røed Knut H. A4 - Bjørnstad, Gro A4 - Flagstad, Øystein A4 - Haanes, Hallvard A4 -
858   Hufthammer, Anne K. A4 - Jordhøy, Per A4 - Rosvold, Jørgen, K. H. A.-R. Ancient
859   DNA reveals prehistoric habitat fragmentation and recent domestic introgression into
860   native wild reindeer. *Conserv. Genet.* **v. 15,** 1137-1149–2014 v.15 no.5 (2014).

861   90.   Røed, K. H. *et al.* Male phenotypic quality influences offspring sex ratio in a polygynous
862   ungulate. *Proc. R. Soc. B Biol. Sci.* **274,** 727–733 (2007).

863   91.   Bjørnstad, G., Flagstad, Ø., Hufthammer, A. K. & Røed, K. H. Ancient DNA reveals a
864   major genetic change during the transition from hunting economy to reindeer husbandry in
865   northern Scandinavia. *J. Archaeol. Sci.* **39,** 102–108 (2012).

866   92.   Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a
867   reference genome. *Nat Biotech* **29,** 644–652 (2011).

868   93.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
869   *EMBnet.journal* **17,** 10–12 (2011).

94.   Benson, G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids*

Res. **27,** 573–580 (1999).

95.   Tempel, S. Using and understanding repeatMasker. *Methods Mol. Biol.* **859,** 29–51
(2012).

96.   Maja, T. & Nansheng, C. Using RepeatMasker to Identify Repetitive Elements in
Genomic Sequences. *Curr. Protoc. Bioinforma.* **25,** 4.10.1-4.10.14 (2009).

97.   Aken, B. L. *et al.* The Ensembl gene annotation system. *Database (Oxford).* **2016,** (2016).

98.   Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment
search tool. *Journal of Molecular Biology* **215,** 403–410 (1990).

99.   Birney, E., Clamp, M. & Durbin, R. GeneWise and Genomewise. *Genome Res.* **14,** 988–
95 (2004).

100.   Mulder, N. J. *et al.* New developments in the InterPro database. *Nucleic Acids Res.* **35,**
D224-8 (2007).

101.   Kanehisa, M. & Goto, S. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic
Acids Res.* **28,** 27–30 (1999).

102.   Bairoch, A. & Apweiler, R. The SWISS-PROT protein sequence database and its
supplement TrEMBL in 2000. *Nucleic Acids Res.* **28,** 45–48 (1999).

103.   The UniProt Consortium. UniProt: the universal protein knowledgebase. *Nucleic Acids
Res.* **45,** D158–D169 (2016).

104.   Nawrocki, E. P., Kolbe, D. L. & Eddy, S. R. Infernal 1.0: inference of RNA alignments.
Bioinformatics **25,** 1335–1337 (2009).

105.   Lowe, T. M. & Eddy, S. R. tRNAscan-SE: a program for improved detection of transfer
RNA genes in genomic sequence. *Nucleic Acids Res.* **25,** 955–964 (1997).

106.   Gardner, P. P. *et al.* Rfam: updates to the RNA families database. *Nucleic Acids Res.* **37,**
D136–D140 (2008).

107.   The Gene, O. C. *et al.* Gene Ontology: tool for the unification of biology. *Nat. Genet.* **25,**
25–29 (2000).

108.   Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res.* **32,** 1792–1797 (2004).

109.   Guindon, S. & Gascuel, O. A Simple, Fast, and Accurate Algorithm to Estimate Large
Phylogenies by Maximum Likelihood. *Syst. Biol.* **52,** 696–704 (2003).

110. Guindon, S. *et al.* New Algorithms and Methods to Estimate Maximum-Likelihood
Phylogenies: Assessing the Performance of PhyML 3.0. *Syst. Biol.* **59,** 307–321 (2010).

111. Rannala, B., Yang, Z. & Anderson, F. Inferring Speciation Times under an Episodic
Molecular Clock. *Syst. Biol.* **56,** 453–466 (2007).

112. Yang, Z. & Rannala, B. Bayesian Estimation of Species Divergence Times Under a
Molecular Clock Using Multiple Fossil Calibrations with Soft Bounds. *Mol. Biol. Evol.*
23, 212–226 (2006).

113. Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24,**
1586–91 (2007).

114. De Bie, T., Cristianini, N., Demuth, J. P. & Hahn, M. W. CAFE: a computational tool for
the study of gene family evolution. *Bioinformatics* **22,** 1269–71 (2006).

115. Löytynoja, A. & Goldman, N. An algorithm for progressive multiple alignment of
sequences with insertions. *Proc. Natl. Acad. Sci. U. S. A.* **102,** 10557–62 (2005).

116. DePristo, M. A. *et al.* A framework for variation discovery and genotyping using next-
generation DNA sequencing data. *Nat. Genet.* **43,** 491–498 (2011).

117. Bernt, M. *et al.* MITOS: Improved de novo metazoan mitochondrial genome annotation.
Molecular Phylogenetics and Evolution **69,** 313–319 (2013).

118. Patterson, N., Price, A. L. & Reich, D. Population Structure and Eigenanalysis. *PLoS*
Genet **2,** e190 (2006).

119. Stajich, J. E. *et al.* The Bioperl Toolkit: Perl Modules for the Life Sciences. *Genome Res.*
12, 1611–1618 (2002).

Abbreviations

Gb   Gigabases

kb   Kilobases

Mb   Megabases

ncRNA   Non-coding RNA

tRNA   Transfer RNA

930	rRNA	Ribosomal RNA
931	miRNA	MicroRNA
932	snRNA	Small nuclear RNA
933	BP	Biological process
934	MF	Molecular function
935	CDS	Coding sequences
936	PSMC	Pairwise Sequentially Markovian Coalescent
937	SNPs	Single nucleotide polymorphisms
938	BWA	Burrows-Wheeler alignment
939	GATK	Genome Analysis Toolkit
940	PCA	Principal component analysis
941	mtDNA	Mitochondrial DNA
942	N_e	Effective population size
943	Kya	Thousand years ago
944	M	Million

Acknowledgements

This study was financially supported by the Academy of Finland in the Arctic Research
Programme ARKTIKO (decision number 286040). M. B acknowledges partial funding from the
Finnish cultural foundation. We are grateful to Juhani Maijala for providing the reindeer that was
used as a reference genome. The authors thank Nuccio Mazzullo, Päivi Soppela and Anna
Stammler-Gossmann from the Arctic Centre, University of Lapland, Rovaniemi, Finland for
collaboration in the sampling of the Finnish reference reindeer. The authors wish to acknowledge
the CSC-IT Center for Science, Finland, for computational resources. The owners of the animals
included in the study are acknowledged for providing samples for this study.

Contributions

J. K conceived and designed the project. J. K, T. R, M. H, J. P and K. R collected the samples. M. W, K. P and M. Y performed bioinformatics analyses. M. W prepared the manuscript draft with substantial contribution from J. K and K.P. All authors read, reviewed and approved the final manuscript.

Competing interests

The authors declare that there are no competing financial interests.