NbCycB2 represses Nbwo activity via a negative feedback loop in the tobacco

trichome developmemt

Minliang Wu[1, 3], Yuchao Cui[1], Li Ge[1], Lipeng Cui[1], Zhichao Xu[1], Hongying Zhang[2],

Zhaojun Wang[2], Dan Zhou[1], Shuang Wu[3], Liang Chen[1,]*, Hong Cui[2,]*

1. Xiamen Key Laboratory for Plant Genetics, School of Life Sciences, Xiamen

University, Xiamen 361102, China

2. Key Laboratory for Cultivation of Tobacco Industry, College of Tobacco Science,

Henan Agricultural University, Zhengzhou, 450002, China

3. FAFU-UCR Joint Center and Fujian Provincial Key Laboratory of Haixia Applied Plant

Systems Biology, College of Horticulture, Fujian Agriculture and Forestry University,

Fuzhou 350002, China

* Co-Corresponding author (email: chenlg@xmu.edu.cn, cuihonger_13@163.com)

Running title: Reciprocal regulation between *Nbwo* and *NbCycB2*

Highlight

NbCycB2 is specifically expressed in trichomes of *Nicotiana benthamiana* and
represses the Nbwo activity via a negative feedback loop in tobacco trichome
developmemt.

Abstract:

The wo protein and its downstream gene, *SlCycB2* have been demonstrated to
regulate the trichome development in tomato. It was shown that only
gain-of-function mutant form of *wo*, *Wo^V* (*wo* woolly motif mutant allele) could
induce the increase of trichome density. However, it is still unclear the relationships
between wo, *Wo^V* and *SlCycB2* in trichome regulation. In this study, we demonstrated
Nbwo (NbWo^V) directly regulated the expressions *NbCycB2* by binding to the
promoter of *NbCycB2* and its genomic sequences. As a feedback regulation, NbCycB2
negatively regulates the trichome formation by repressing Nbwo activity at protein
level. We further found that the mutations of Nbwo woolly motif could prevent
repression of NbWo^V by NbCycB2, which results in the significant increase of active
Nbwo proteins, trichome density and branches. Our results revealed a novel
reciprocal mechanism between *NbCycB2* and *Nbwo* during the trichome formation in
Nicotiana benthamiana.

Key word: Feedback loop; Trichome formation; Nbwo; NbCycB2; Woolly motif;
L1-like box

Introduction

Trichomes are the specialized epidermal protuberances locating on aerial parts of
nearly all terrestrial plants. They can be classified into various types by cell numbers

and shapes -- unicellular/multicellular, and glandular/non-glandular. In *Arabidopsis*, it

has been well-elucidated that the development of trichomes (unicellular and

non-glandular), is regulated by the trimeric MYB-bHLH-WDR protein activators

complex (GL1 (Oppenheimer et al., 1991)-GL3/EGL3 (Payne et al., 2000)-TTG1(Walker

et al., 1999)). This transcriptional complex activates the expression of the

homeodomain protein GLABROUS2 (GL2) to induce the formation of trichomes

(Rerie et al., 1994; Grebe, 2012). In addition, it triggers the expression of the single

repeat R3 MYBs (including TRY (Schnittger et al., 1999), CPC (Wada et al., 1997), ETC1,

ETC2, ETC3 (Kirik et al., 2004; Wester et al., 2009) and TCL2 (Gan et al., 2011)) which

act as negative regulators of GL3 or EGL3 by forming a repressor complex

(GL3/EGL3-TRY/CPC-TTG1) in trichome development (Wang et al., 2008; Wester et al.,

2009). Thus the control of trichome development in *Arabidopsis* requires a

regulatory loop including both activators and repressors (Grebe, 2012; Pattanaik et

al., 2014).

The trichomes are multicellular glandular (GSTs) structure in approximately 30% of

all vascular plants (Glas et al., 2012). Since many phytochemicals and compounds

with economical values can be synthesized and secreted by multicellular GSTs

(Mauricio and Rausher, 1997; Hollósy, 2002; Valkama et al., 2003; Freeman and

Beattie, 2008), multicellular glandular trichomes have considerable economic

pontential (Sallets et al., 2014; Huchelmann et al., 2017). However, it has been

demonstrated that the networks regulating unicellular trichomes did not work in the

development of multicellular trichomes (Serna and Martin, 2006; Yang et al., 2011;

Kang et al., 2016; Yan et al., 2016).

In tomato, a HD-ZIP IV transcriptional factor, wo protein has been demonstrated to

regulate the trichome initiation (Yang et al., 2011). This HD-ZIP IV member contains

four conserved domains including homeodomain domain (HD), the leucine zipper

domain (LZ), the steroidogenic acute regulatory protein-related lipid transfer (START)

and the START-adjacent domain (SAD). However, overexpression of *wo* failed to

induce the change of trichome density, and only ectopic expression of its

gain-of-function mutant alleles, Wo^V, could cause higher density of trichomes in

tomato (*Solanum lycopersicum*) and tobacco (*Nicotiana tabacum*) (Yang et al., 2011;

Yang et al., 2015). The Wo^V allele has two point mutations at the C-terminal domain

(Since this motif was conserved in the most wo alleles, we name it as woolly motif in

this study). The sequence analysis revealed that wo protein is more similar to

PROTODERMAL FACTOR2 (PDF2) and the PDF2 redundant protein -- ARABIDOPSIS

THALIANA MERISTEM L1 (ATML1), both of which are involved in shoot epidermal cell

differentiation (Abe et al., 2001; Ogawa et al., 2015), than GL2 in *Arabidopsis*.

In *Arabidopsis*, the ectopic expression of a constitutive active B-type cyclin induced

mitotic divisions and resulted in the increase of multicellular trichomes (Schnittger et

al., 2005). *SlCycB2*, a hypothetical B-type cyclin, was reported to directly interact with

wo to promote the development of type I trichome (Yang et al., 2011; Yang et al.,

2015). Its homologous protein in *Arabidopsis* (AT5G06270) was also found to interact

with GL2 or co-repressor TOPLESS proteins (Wu and Citovsky, 2017b, a). However, as

reported in a recent study, overexpression of *SlCycB2* resulted in non-trichome

phenotype, while suppression of *SlCycB2* promoted trichomes formation in tomato

(Gao et al., 2017). These inconsistent results raise the important questions: what is

the function of *SlCycB2* in trichome formation and why the mutation of woolly motif

can promote trichome formation?

Similar to tomato, trichomes in *Nicotiana benthamiana* are typically multicellular

structures, and almost all of the trichomes in *N. benthamiana* are glandular (Fig. S1),

making it a better system for studying glandular trichomes than tomato. In addition,

the genome map of *N. benthamiana* has been constructed (Bombarely et al., 2012).

Thus tobacco represents an excellent model plant to study the molecular mechanism

of multicellular trichome formation (Goodin et al., 2008). In this study, we cloned the

homologues of wo and *SlCycB2* in *N. benthamiana* (named *Nbwo* and *NbCycB2*), and

constructed a two-point mutantion *Nbwo* allele, $NbWo^V$. To investigate their

biological functions in trichome development, we constructed overexpression and

suppression transgenic lines of all genes. We demonstrated that Nbwo could

positively regulate the expression of *NbCycB2* through targeting to the cis-element in

NbCycB2 promoter and its genomic DNA sequence. On the other hand, NbCycB2

could be a negative regulator of multicellular trichomes by directly binding and

inhibiting Nbwo activity. The previous identified mutation in woolly motif (NbWo^{V})

blocked the interaction between NbCycB2 and Nbwo, removing the repression of

Nbwo by NbCycB2 and resulting in increased trichome density. Our results revealed

the mechanisms of the interaction between *Nbwo* and *NbCycB2* in regulating the

development of glandular trichomes.

Materials and Methods

Plant materials and growth conditions

Sterilized seeds of *N. benthamiana* were germinated and grown to seedlings on

MS medium, which solidified with 0.8% (w/v) gellan gum under 26 °C, 14 h light/10 h

dark conditions. Two-week-old plants were transferred to other sterilized bottle (for

genetic transformation) or soil in pots to grow to maturity. All wild type and

transgenic plants were grown in greenhouse under 26 °C, 14 h light/10 h dark

condition.

Sequence analysis

The protein sequences of the homologues of *wo* and *SlCycB2* were downloaded

from NCBI database (http://www.ncbi.nlm.nih.gov/) and Sol genomic network

(Fernandez-Pozo et al., 2015) (https://solgenomics.net/). In order to further analyze

the grouping and relatedness, the aligned sequences were used to construct the

phylogenetic trees in MEGA 5 by using the maximum-likelihood (ML) criterion with

100 bootstrap analysis. In addition, the relative conservation for each amino acid

position in the protein sequences of Nbwo and NbCycB2 was evaluated via WebLogo

(https://weblogo.berkeley.edu/) (Crooks et al., 2004), followed by the prediction of

their conserved domains in SMART program (https://smart.embl-heidelberg.de/)

(Letunic and Bork, 2018).

RNA extraction and Real-time PCR

Total of RNA was extracted from various tissue of plants by using the Eastep®Super

Total RNA Extraction Kit (Promega). The cDNA was synthesized from Dnase I treated

total RNA using M-MLV 1st Strand Kit (Invitrogen). Real-time PCR (qRT-PCR) was

determined by using SYBR Premix Ex Taq II (TaKaRa) and performed on ABI Stepone

real-time PCR system (Applied Biosystems). L25 ribosomal protein (L18908) was used

as an endogenous control (Schmidt and Delaney, 2010). Relative expression levels

were determined as described previously (Guo et al., 2016). Primers are listed in

Table S1.

Plasmid construction and *N. benthamiana* transformation

To investigate the biological functions of *wo* and *SlCycB2* in *N. benthamiana*, the

full-length coding sequences of two alleles of *Nbwo* and *NbCycB2* were amplified

with NheI and BamHI, XbaI and BglII cloning sites respectively from the general cDNA

of leaves (All the primers used in current study are provided in Table S1). In addition,

to confirm the function of the tomato *Wo^V* gene in tobacoo trichome formation (Yang

et al., 2015), an allele *NbWo^V* with two point mutations at the locus 2084 (T was

replaced with G, result in Ile-697 changed to Arg) and 2092 (G was replaced with T,

result in Asp-700 changed to Tyr, Fig. S2c) of *Nbwo* was generated in *N. benthamiana*

by using a Mutagenesis Kit (Toyobo).

To construct the overexpression (OE) lines of *Nbwo*, *NbWo^V* and *NbCycB2*, these

fragments were inserted into pCXSN-HA (Nbwo and NbWo^V fused with HA tag) and

pCXSN-FLAG (NbCycB2 fused with Flag tag) vectors respectively under the control of

CaMV 35S promoter (Chen et al., 2009). The suppression expression constructs of

Nbwo and *NbCycB2* were performed by recombining with the RNAi vector

pH7GWIWGII with LR Clonase II enzyme (Invitrogen).

To comprehensively understand the expression sites of *NbCycB2*, approximate

2880bp upstream promoter fragments were amplified by PCR using the primers

shown in Table S1. The promoter fragments were then inserted into the

corresponding site of pH2GW7 vector to create the promoter-driven GFP-GUS

transformation by using Pro-*NbCycB2*: GFP-GUS gene fusion (Cui et al., 2015).

All of these constructs were transferred into *Agrobacterium tumefaciens* strain

GV3101, which were used to generate transgenic lines by using

Agrobacterium-mediated transformation. The gene expression levels in the

transgenic lines were examined by real-time PCR and western blot.

Subcellular localization and tissue distribution

The pCXDG vector (Chen et al., 2009) was used to analyse the subcellular

localization of *Nbwo*, *NbWo^V* and *NbCycB2*, which were fused with GFP driven by the

CaMV 35S promoter (p35s:GFP-*Nbwo*, p35s:GFP-*NbWo^V*, p35s:GFP-*NbCycB2*).

Transformants carrying these constructs were obtained as described above and then

transiently transformed into 4-week-old *N. benthamiana* leaves. After cultivation in

lowlight conditions for 48-72 h, GFP was observed using confocal microscopy (LSM

780, Carl Zeiss, Jena, Germany) with staining in DAPI solution (1 mg/ml) for 15 min

before observation.

The tissue distribution assays were performed as described in the previous study

(Jefferson et al., 1987). GUS staining was repeated at least three independent

transgenic lines.

Yeast hybrid assays

Yeast one-hybrid assays were performed to test the specific function area of

NbCycB2 promoter binding Nbwo. The promoter of *NbCycB2*, separated into 5

fragments (E: -1027 ~ -831 bp, D: -830 ~ -631 bp, C: - 630 ~ -411 bp, B: -410 ~ -201 bp,

A: -200 ~ -1bp, Fig. 2a), were amplified and inserted into the pHIS 2 vector (Clontech)

(*NbCycB2*proE, *NbCycB2*proD, *NbCycB2*proC, *NbCycB2*proB, *NbCycB2*proA). Further

investigation of the targeted sequences in *NbCycB2* promoter were conducted by

point mutations in the two L1-like boxes in the D fragment (*NbCycB2*proD-m1,

mutant one L1-like box, changed 5'-GCAAATATTTACTC-3' to 5'-GCGGGTGACTC-3';

*NbCycB2*proD-m2, mutant two L1-like boxes, changed 5'-GCAAATATTTACTC-3' to

5'-GCGGGTGACTC-3', and 5'-ATTTACTC-3' to 5'-GGGACTCC-3'). To test the specific

region of *Nbwo* genomic sequence binding itself, four genomic fragments of *Nbwo*

(G1, -8 ~ 251 bp, include T3 fragment; G2, 2169 ~ 2522 bp, include T4 fragment; G3,

3485 ~ 3780 bp, include T5 fragment; G4, 4333 ~ 4660 bp, include T6 fragment, Fig.

7a), were amplified and inserted into the pHIS 2 vector (Clontech) (*Nbwo-G1*,

Nbwo-G2, *Nbwo-G3*, *Nbwo-G4*). In addition, the CDSs of *Nbwo* and *NbWo^V* were

inserted into vectors pGADT7 containing the GAL4 activation domain (AD) (Clontech).

The plasmids were co-transformed into the Y187 yeast strain, empty AD vector was

provided as the negative control, and cultivated on SD/-Leu/-Trp (-L-W) medium and

tested on SD/-Leu/-His/-Trp (-L-W-H) with 60 mM 3-amino-1,2,4-triazole (Sangon

Biotech (Shanghai) Co., Ltd) medium.

The yeast two-hybrid was performed to understand the interaction between *Nbwo*

and *NbCycB2*. We truncated *Nbwo* into four segments containing: HD, LZ, START and

SAD domains, and fused them into AD vectors to find the target region of the

interaction in *Nbwo*. In addition, the CDS of *NbWo^V* was also amplified and inserted

into AD vector. Each pair of AD and BD plasmids were co-transformed into the

Y2HGold yeast strain. BD-53 and AD-T constructs were co-transformed in to Y2HGold

as positive control, and BD-Lam and AD-T as negative control. The transformants

were then cultivated on SD/-Leu/-Trp medium (DDO) and tested on SD/-Ade/

-Leu/-His/-Trp with 40 mg/L X-a-Gal (QDO/X) or SD/-Ade/-Leu/-His/-Trp with 40 mg/L

X-a-Gal and 400 µg/L Aureobasidin A medium (QDO/X/A).

Yeast three-hybrid was conducted to analyze the binding competition between

Nbwo LZ domain (Nbwo-LZ) and NbCycB2 to Nbwo. The Nbwo-LZ was fused with BD

(Clontech). The *NbCycB2* was inserted into the downstream of methionine

repressible promoter (PMet25: *NbCycB2*) in pBridge vector. The plasmid of

BD-*Nbwo-LZ* was transferred with AD-*Nbwo* as positive control, and empty pBridge

vector was transferred to with AD-*Nbwo* as negative control. The transformants

were then tested on SD/-Ade/-Leu/-His/-Trp mediums with different concentrations

of methionine (0, 250 µM).

Bimolecular fluorescence complementation Assay (BiFc)

In order to determine the interaction of NbCycB2 and Nbwo (or NbWo^V) in *N.*

benthamiana protoplasts. The CDSs of *Nbwo*, *NbWo^V* and *NbCycB2* were inserted

into the pSAT6-cEYFP-C1-B vector (2x35s: YFP^c-*Nbwo*, 2x35s: YFP^c-*NbWo^V*) and the

pSAT6-n(1-174)EYFP-C1 vector (2x35s: YFP^n-*NbCycB2*) separately (Citovsky et al.,

2006). Each pair of the two plasmids were then transiently transformed into the

protoplasts via PEG–calcium transfection method as described in the previous study

(Yoo et al., 2007).

Moreover, to determine the interactions between Nbwo and NbCycB2 in vivo, the

CDSs of *Nbwo*, *NbWo*[V] and *NbCycB2* were fused separately with the C-terminal

fragment of YFP in p2YC vector; *Nbwo*-LZ and *NbWo*[V] were also fused with the

N-terminal fragment of YFP in p2YN vector respectively (Shen et al., 2011). Different

plasmid combinations were co-infiltrated into leaves of *N. benthamiana* as described

in previous study (Shen et al., 2011).

The YFP fluorescence was observed by using confocal microscopy (LSM 780, Carl

Zeiss, Jena, Germany). Before the observation, the transformed protoplasts were

cultivation in 26 °C for 12 h, and the transformed leaves of *N. benthamiana* were

cultivation in darkness for 48-72 h. Three biological repeats were observed

independently for each samples.

251 Dual-luciferase (Dual-Luc) assay

The regulatory effectors of 2x35s: *HA-Nbwo*, 2x35s: *HA-NbWo*[V] and 2x35s:

Flag-NbCycB2 were generated by using the DNA sequences with NcoI (5'-end) and

BglII (3'-end) cloning sites.

The firefly luciferase reporters were created by inserting the B and D fragments of

the *NbCycB2* promoter, and the *Renilla* luciferase was driven by the 35S promoter in

the pGreen-0800-II report vector (35S: REN-*NbCycB2*proB: LUC, 35S:

REN-*NbCycB2*proD: LUC). The mutants of the *NbCycB2* promoter D fragments were

also constructed using a Mutagenesis Kit (35S: REN-*NbCycB2*proD-m2: LUC). The

regulatory effector and the reporter were used at the ratio of 5:1 or 5:5:1 for the

expression test of two or three plasmids.

262 Immunoblotting, CO-IP and pull-down assay

The *N. benthamiana* leaves (~0.5 g) were homogenized with liquid nitrogen and

then solubilized with 0.4 ml of lysis buffer (25 mM Tris-HCl, 2.5 mM EDTA, pH 8.0,

0.05% v/v NP-40, 5% glycerol, 150 mM NaCl, 1 mM phenylmethylsulfonyl fluoride

and 20 uM MG132) for 30 min at 4 °C. After solubilization, the protein extract was

centrifuged at 13,000g for 10 min at 4 °C to separate the solubilized (supernatant)

and non-solubilized material. Total protein (~80 μg) was then used for the

immunoblotting assay. After SDS–PAGE separation, the proteins were

electrophoretically transferred to a PVDF membrane for immunodetection.

The interaction of NbCycB2 and Nbwo dimers were determined by using co-IP

assay in *N. benthamiana* protoplasts. In the expression vectors, the Nbwo protein

was fused to the HA and Flag tags, respectively, and the NbCycB2 was fused to the

GFP tag. Each pairs of the plasmids were transformed into the protoplasts via

PEG–calcium transfection method as described in the previous study (Yoo et al.,

2007). The total proteins were extracted by using lysis buffer and then incubated for

3 h at 4°C with 20 ul of Anti-HA Affinity Gel (Millipore). The immunoprecipitates were

washed five times with lysis buffer. The isolated proteins were detected by

immunoblotting with anti-Flag or anti-GFP antibodies.

In the pull-down assay, the CDSs of *Nbwo*, *NbWo*[V] and *NbCycB2* were respectively

inserted into the pET22b and PGEX-4T-1 vectors to create the fusion proteins

(His-Nbwo, His-NbWo[V] and GST-NbCycB2), and then transformed into the *Escherichia*

coli BL21 strain. The purified recombinant bait proteins (2 mg His-Nbwo or

His-NbWo[V]) and 2 mg of prey proteins (GST-NbCycB2) were then mixed with 1 ml

binding buffer (50 mM Tris-HCl pH 7.5, 0.6% Triton and X-100, 100 mM NaCl,). After

incubation at 4 °C for 2 h, 50 μl of glutathione agarose was added to the mixtures

followed by the incubation for additional 1 h. The immunoprecipitates were washed

five times with binding buffer. The isolated proteins were detected by

immunoblotting with anti-His or anti-GST antibodies.

Chromatin immunoprecipitation (ChIP PCR) assay

Four weeks old *NbWo*[V]-*OE* plants were used for ChIP assay as described in previous

study (Gendrel et al., 2005). The NbWo[V] proteins were precipitated by using HA

antibody (Santa Cruz). Primers were designed to amplify 3 fragments (length ~ 120 -

210 bp) within the 1.7k bp upstream sequence of the *NbCycB2* transcription start site

(Fig. 2a), and 7 fragments within ~ 8.7k bp of genomic DNA sequence of *Nbwo* (Fig.

7a). After the immuno-precipitation, the purified DNA was analyzed by real-time PCR

with the primers of *NbCycB2* promoter and *Nbwo* genomic DNA sequence fragments

(Table S1). Enrichment was calculated from the ratio of immuno-precipitated

sequences.

Phenotype observation

Images of the transgenic plants stem were obtained using Nikon camera. The

observation of abaxial leaves and root hair were used an Axioplan 2 microscope (Carl

Zeiss AG).

Leaf of transgenic line and wild type seedlings, which used for scanning electron

microscope analysis, were fixed with 2% glutaraldehyde (0.1M phosphate buffer, PH

7.4) at 4 □ for 12h. Then, the samples were dehydrated with a series of alcohol (10%,

20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% and 100%) for 20 min each time.

Finally, the samples were dried in a critical point drying device (Leica EMCPD030),

and coated with gold particles. The samples were observed by using a JSM-6390/LV

scanning electron microscope.

Data deposition

The sequences reported in this article have been deposited in the Sol genomic

network (Fernandez-Pozo et al., 2015) (https://solgenomics.net/) with the accession

numbers   as   follows: *Nbwo*   (Niben101Scf07790g01007.1),   *Nbwo*-allele

(Niben101Scf00176g11005.1),   *NbCycB2*   (Niben101Scf10299g00003.1),

NbCycB2-allele (Niben101Scf10396g00002.1), *NbML1* (Niben101Scf00703g00003.1),

NbML1-allele (Niben101Scf01158g03010.1).

Results

Expression and cellular analysis of *Nbwo* and *NbCycB2*

We obtained 14 Nbwo and 8 NbCycB2 protein-related sequences from online

databases (Fig. S2a, b). The full-length coding sequences of *Nbwo* and *NbCycB2*

obtained in the current study contained 2,199 bp and 333 bp, respectively. One allele

of each of the *Nbwo* and *NbCycB2* genes were identified via BLAST with 95.82% and

95.20% identity, respectively. The amino acid analysis determined the two-point

mutations at 2,084 and 2,092 of $NbWo^V$ could cause two amino acid replacements in

the woolly motif (Ile to Arg, Asp to Tyr, Fig. S1c). Conserved domain analysis revealed

that NbCycB2 contained a WD40-like domain in the N-terminal (NbCycB2-WD40,

including an EAR like motif) and a RING-like domain in the C-terminal (NbCycB2-RING)

(Fig. S2d). However, no conserved domain of B-Type Cyclin protein was found in

NbCycB2 protein sequences.

The visualization of subcellular localization revealed that NbCycB2, Nbwo and

NbWo^V all localized to the nucleus (Fig. S3a). As shown in the self-activation assay,

clones of AD-Nbwo and AD-NbWo^V can grow on QDO/X/A medium compared to the

positive control, which means they have strong self-activating ability, while the

mutations of woolly motif did not affect Nbwo's transactivation ability (Fig. S3b).

As shown in Fig. S4a, the spatial expression pattern assays of *NbCycB2* and *Nbwo*

indicated they were lowly expressed in roots but at a high level in the trichome

containing organs. Further investigation revealed the expression of the *NbCycB2*

promoter-driven GFP-GUS transgenic lines was only detected in the trichomes of

leaves and stems (Fig. S4b-c). However, GUS of the Nbwo promoter-driven GUS

transgenic line was strongly expressed in the basal and venous regions of young

leaves (Fig. S4d).

NbCycB2 negatively regulates trichome initiation

Most of *NbCycB2* overexpression (OE) transgenic lines underwent a dramatic

reduction of trichomes on leaves and stems (Fig. S5a, Fig. 1c, 1f). The root length and

number of branch root significantly increased in *NbCycB2-OE* lines (Fig. S6). Western

blot and qRT-PCR analysis showed the *NbCycB2* transcripts significantly accumulated

in *NbCycB2-OE* lines, while the expression levels of *Nbwo* and endogenous *NbCycB2*

were significantly reduced (Fig. S5b, c).

In contrast, the density of trichomes increased significantly on the leaves and

stems of 16 *NbCycB2* knockdown lines (*NbCycB2-RNAi*) (Fig. S5d, Fig. 1b, 1f). The

qRT-PCR analysis showed the number of trichomes negatively correlated with the

expression level of *NbCycB2* (Fig. S5e), suggesting that *NbCycB2* may play a negative

role in trichome initiation.

NbWo^V positively regulates trichome initiation

To confirm the function of *Nbwo*, we also generated 22 *Nbwo* knockdown

transgenic plants (*Nbwo-RNAi*). Compared with the wild type, trichome densities

were clearly decreased on the leaves and stems of most *Nbwo* knockdown plants

(Fig. S7a, e). The efficiency of RNAi mediated knockdown was confirmed by qRT-PCR

in two independent lines, in which the expression of *Nbwo* and *NbCycB2* was

significantly reduced (Fig. S7b).

As in the previous study, dramatic increases in the density and branching of

trichomes were found on the leaves and stems of *NbWo^V-OE* plants (Fig. 1d, 1f, S7d,

S7h). Our qRT-PCR assay showed the expression levels of the *NbWo^V*, endogenous

Nbwo and *NbCycB2* genes were all significantly upregulated in transgenic lines (Fig.

S7f-g). The dwarfism phenotypes were observed in the T1 plants of *NbWo^V-OE* (Fig.

S8).

Over-expression of *Nbwo* also induces the dwarfism

Twenty transgenic plants with overexpression of *Nbwo* (*Nbwo-OE*) were

generated. Interestingly, the density of trichomes was shown to negatively correlate

with the expression level of *Nbwo* in the T0 of *Nbwo-OE* plants (Fig. 2a, b).

Additionally, the expression levels of *NbCycB2* were significantly reduced in trichome

reduced plants (Fig. 2b). However, the decreased trichome phenotypes were not

observed in T1 plants, but the higher expression level of exogenous *Nbwo* (for

example, *Nbwo-OE #3*) also resulted in dwarfism similar to *NbWo^V-OE* lines (Fig. 2c).

Except for the dwarfism, these two lines, however had a great difference in both

trichomes and root hairs. In the *Nbwo-OE #3*, the development of glandular

trichomes and the development of root hairs were normal. In *NbWo^V-OE #1*, the

density of glandular trichomes as opposed to root hairs increased significantly (Fig.

2d-f).

Nbwo and NbWo^V directly targeted the L1-like box of the *NbCycB2* promoter

The expression of *NbCycB2* was significantly upregulated in the *NbWo^V*-OE lines,

and decreased in the *Nbwo*-RNAi lines (Fig. S7b, f). These results indicated that

NbCycB2 was positively regulated by Nbwo and NbWo^V. To test whether Nbwo

directly bound to the promoter of *NbCycB2*, *NbWo^V-OE* (*NbWo^V* fused with HA tag)

plants were analyzed by ChIP qRT-PCR assay with an HA antibody. Strong enrichment

of NbWo^V was observed in the P2 region of the *NbCycB2* promoter in *NbWo^V-OE*

plants (Fig. 3a, b).

To further determine the specific area of the *NbCycB2* promoter binding to Nbwo,

we performed yeast one-hybrid (Y1H) assays. The five truncated fragments of

NbCycB2 promoter were shown in Fig. 3a. The yeast colonies containing

NbCycB2-proD-pHIS 2 and AD-*Nbwo* constructs were grown on the selection medium

with 3-aminotriazole (60 mM) (Fig. 3c).

To investigate whether Nbwo and NbWo^V directly affect the expression of the D

fragment in vivo, we performed dual-Luc assays. The reporters 35S:

REN-*NbCycB2*proD: LUC, 35S: REN-*NbCycB2*proB: LUC and the effectors were shown

in Fig. 3d. As shown in Figure 3e, each pair of reporter and effector was transiently

co-expressed in *N. benthamiana* protoplasts. Compared to the B fragment, when the

Nbwo or NbWo^V was transiently co-expressed, the D fragment-driven LUC

expression accumulated significantly in the protoplasts, indicating that the D

fragment of the *NbCycB2* promoter should be a specific site for Nbwo and NbWo^V

binding.

Further analysis of the targeting sequence of *NbCycB2proD* revealed this sequence

contained two L1-like boxes (5'-ATTTACTC-3') (Fig. 4a). In the Y1H assay and the in

vivo LUC assay, when two L1-like boxes were mutated (*NbCycB2proD-m2*), the

interaction with the Nbwo protein is abolished (Fig. 4b, c). Based on these results,

we inferred the L1-like boxes may be the binding target of the Nbwo and NbWo^V

proteins.

NbCycB2 represses the activity of Nbwo rather than NbWo^V

Since the *NbCycB2-OE* and *Nbwo-RNAi* transgenic lines shared the non-trichome

phenotype, and the expression of *Nbwo* was not enhanced in the *NbCycB2-RNAi*

lines (Fig. S5e), we suspect that NbCycB2 could affect Nbwo transactivation ability at

protein levels. To confirm this, constructs of overexpression *Nbwo*, *NbWo^V* and

NbCycB2 were transiently co-expressed in the leaves of the *NbCycB2*pro: *GFP-GUS*

transgenic line using the *Agrobacterium*-mediated method (Fig. 4d). The GUS

induced by the transient expression of *Nbwo-OE* was only detected in the area

where there was no *NbCycB2-OE* Expression. However, the transient expression of

NbWo^V caused a strong GUS staining which is independent of *NbCycB2-OE*

expression. These results were further supported by the LUC assay, in which

co-expression with *NbCycB2* could clearly repress the transactivation activity of

Nbwo, but did not affect the NbWo^V protein (Fig. 4e). These results suggest that

NbCycB2 may act as a negative regulator of Nbwo rather than NbWo^V.

NbCycB2 suppresses the transactivation ability of Nbwo via the direct interaction

The interaction between NbCycB2 and Nbwo was reported in a previous study

(Yang et al., 2011). To explore which domain was involved in the physical interaction

between NbCycB2 and Nbwo, four truncated fragments of Nbwo containing HD, LZ,

START and SAD were used (Fig. 5a). Y2H assays suggested that the LZ domain of

Nbwo interacted with NbCycB2 (Fig. 5b). The BiFC assay was further used to verify

the interaction between the Nbwo LZ domain and NbCycB2 in vivo (Fig. 5c).

To further examine if NbCycB2 also interacts with NbWo^V, we used yeast

two-hybrid (Y2H) assays. Our results indicate that NbCycB2 physically interacts with

Nbwo but did not interact with NbWo^V, suggesting that the interaction between

NbCycB2 and Nbwo can be reduced by mutations in the woolly motif (Fig. 6a).

Consistent with the Y2H result, the interaction between NbCycB2 and Nbwo or

NbWo^V was further confirmed by the pull-down assay (Fig. 6b) and the bimolecular

fluorescence complementation (BiFC) assay (Fig. S10a).

Nbwo was restrained by NbCycB2 through forming a homodimer

It is known that the HD-Zip proteins bind to DNA as dimers by the LZ domain (Ariel

et al., 2007). To verify whether Nbwo can be dimerized via the LZ domain, we first

demonstrated in the Y2H that the LZ domain could bind to the Nbwo protein (Fig.

S10b). In addition, BiFC assays were used to confirm the interaction between the LZ

domain and Nbwo (or NbWo^V) proteins in vivo (Fig. 5c).

To test whether the NbCycB2 could bind at the Nbwo homodimer, we conducted

the yeast three-hybrid (Y3H) and co-IP assays. As shown in Fig. 6c-d, NbCycB2 could

interact with the Nbwo dimers.

Nbwo could bind to its own genomic DNA

The endogenous expression level of *Nbwo* reduced in *NbCycb2-OE* lines (Fig. S5b)

and increased in *NbWo^V-OE* plants (Fig. 1d), indicated that Nbwo may be able to
regulate self expression. In order to prove this hypothesis, ChIP assay was carried out
to check whether Nbwo could bind to its genomic DNA sequence in the leaf of
NbWo^V-OE transgenic line. Interestingly, enrichment of NbWo^V was found in the T5
fragment in *NbWo^V-OE* plants (Fig. 7a, b). This result was further demonstrated by
using Y1H assay. Only the clones with AD-Nbwo (or AD-NbWo^V) and Nbwo-G3-pHIS 2
constructs could grow on the resistant medium, suggesting that Nbwo or NbWo^V
could bind to the G3 fragments (include T5 fragment, Fig. 7a) of its own genomic
DNA sequences (Fig. 7c).

Overexpression of *NbCycB2* can reduce the dwarf phenotype of *Nbwo-OE* plants

To determine whether *NbCycB2* can inhibit the transactivation activity of *Nbwo* in
vivo, we crossed *NbCycB2-OE #2* T1 plant to *Nbwo-OE #3* T0 plants. As shown in Fig.
8 a-b, the dwarf and short-root phenotypes of *Nbwo-OE #3* were indeed reduced by
the *NbCycB2-OE #2*. The crossed F1 plants were tested using PCR (Fig. 8 c), and the
expression of *NbCycB2* and *Nbwo* was also verified by qRT-PCR. Compared with T1
Nbwo-OE #3 plants, the balance between *NbCycB2* and *Nbwo* expression was
restored in *NbCycB2-OE* #2 and *Nbwo-OE* #3 crossed F1 plants (Fig. 8 d).

Discussion

***NbCycB2* negatively regulates trichome initiation**

It is well known that ectopic expression of a constitutive active B-type cyclin
promoted the unicellular trichome differentiate to multicellular in *Arabidopsis*
(Schnittger et al., 2002, 2005). The *SlCycB2* gene has been reported to be a
hypothesis B-type cyclin gene participating in trichome formation in tomato (Yang et
al., 2011; Gao et al., 2017). However, the function of *SlCycB2* during trichome
development has not been well-studied.

In this study, we found that overexpression of *NbCycB2* caused a non-trichome
phenotype, whereas inhibition of *NbCycB2* significantly increased the density of
trichomes instead of branching on stems and leaves (Fig.1b, S5). Consistent with the
qRT-PCR results (Fig. S4a), GUS staining assay indicated *NbCycB2* is specifically

expressed in the trichomes of leaves and stems (Fig. S4b-c), and *Nbwo* is expressed

in the basal and venous regions of young leaves (Fig. S4d). These results suggested

NbCycB2 serves as a negative regulator of trichome initiation. None of B-type cyclin

conserve domains were found in the SlCycB2 and NbCycB2 protein sequences (Fig.

S2d). Thus, whether SlCycB2 has the function of B type cyclin protein requires

further study.

NbWo^V and Nbwo directly regulate the expression of *NbCycB2* through binding to

the L1-like boxes in the promoter

In previous study, *SlCycB2* has been reported to be indirectly regulated by *Wo^V*

(Yang et al., 2011; Yang et al., 2015). The expression level of *NbCycB2* was

upregulated in overexpression of *NbWo^V* plants and downregulated in *Nbwo-RNAi*

lines (Fig. S7), indicating that *NbCycB2* may be the downstream gene of *Nbwo*.

Additionally, the D fragments of the *NbCycB2* promoter have been shown to be the

binding target of Nbwo and NbWo^V through ChIP, Y1H and LUC assays (Fig. 3).

Mutation of the two L1-like box sequences in *NbCycB2*proD inhibited the binding of

Nbwo or NbWo^V in vitro and vivo (Fig. 4b). Therefore, we proved that *NbCycB2* was

directly regulated by Nbwo or NbWo^V through the binding of L1-like boxes in the

NbCycB2 promoter (Fig. 4a-c). In addition, we also proved that Nbwo and NbWo^V can

self-regulate the endogenous expressions by binding to its own genomic DNA

sequence by ChIp and YIH assay (Fig. 7).

The increase of trichome density and plant dwarfism was regulated by Nbwo

through different pathways

The *Wo* homozygous plants have been shown to cause the embryo lethality in

seeds (Yang et al., 2011). We also found the overexpression of *Nbwo* or *NbWo^V* genes

causes abnormal embryonic development and results in dwarf phenotype in its

offspring (Fig. 2c, S6). However, the trichome phenotype of *Nbwo-OE* and *NbWo^V-OE*

was completely different (Fig. 2e and 2f), suggesting that Nbwo was involved in both

the regulation of the development of trihcome and embryo. The trichomes' density

was decreased with the expression of Nbwo in T0 of *Nbwo-OE* lines (Fig.2a and 2e),

which suggested that high expression of *Nbwo* in wild type background could repress

trichome development. Thus, we suspected the woolly motif represses *Nbwo* activity.

However, the trichome density was not increased when the SAD domain deleted

Nbwo CDS (including woolly motif) was overexpressed in the wild type plants (Fig.

S9), which suggesting that woolly motif did not have repressing activity. Therefore,

the decreased trichome density in high expression of *wo* gene plants requires further

investigation.

NbCycB2 represses the transactivation activity of Nbwo at protein level

The *NbCycB2* gene was proven to be directly regulated by Nbwo in our study.

However, the similar non-trichome phenotypes of *NbCycB2-OE* and *Nbwo-RNAi*

transgenic lines were found (Fig. S5a, S7a). Additionally, the expression of *Nbwo* was

not increased in *NbCycB2-RNAi* plants (Fig. S5e). These results suggested that

NbCycB2 might repress the transactivation activity of Nbwo at protein levels. The

GUS activity of the *NbCycB2pro*: *GFP-GUS* transgenic line was upregulated by the

expression of *Nbwo* and inhibited by the co-expression of *NbCycB2* (Fig. 4d), and the

same result was found in the LUC assay (Fig. 4e). Additionally, Hybridization with

NbCycB2-OE can attenuate the dwarf phenotype of T1 *Nbwo-OE* (Fig. 8a-b). Further

studies revealed that the expression of endogenous *NbCycB2* and *Nbwo* were

reduced in the *NbCycB2-OE* lines (Fig. S5b). They were shown to be downstream

regulatory genes of Nbwo (Figures 3 and 7). These results altogether supported our

hypotheses that NbCycB2 may act as a negative regulator of Nbwo at protein levels.

In a previous study, SlCycB2 was reported to interact with wo protein (Yang et al.,

2011). Further investigation of the interaction between Nbwo and NbCycB2 revealed

the dimerized LZ domain of Nbwo binds to NbCycB2 (Fig. 5b, c). Through Y2H, BiFC,

Y3H and co-IP assays, we also found that Nbwo could form a homodimer through the

LZ domain, and the NbCycB2 protein could bind to LZ domain of Nbwo dimers (Fig.

5b-c, 6 c-d, S10b). These results indicated that NbCycB2 may bind to Nbwo protein

via its LZ domain to form a complex, which inhibits its transactivation ability.

However, further study is required to determine whether *NbCycB2* functions

similarly to its homologue gene --AT5G06270.1 which also interacts with the

co-repressor TOPLESS (Long et al., 2006; Szemenyei et al., 2008; Pauwels et al., 2010;

Wu and Citovsky, 2017a).

543 The interaction between Nbwo and NbCycB2 was blocked by the mutation in the

544 Nbwo woolly motif

In *Arabidopsis thaliana*, the feedback loop regulation mechanisms of the R3 MYB

(TRY, CPC and so on) is through competitively binding to GL3/EGL3 to form a

non-functional trimeric protein complex (MYB-bHLH-WDR), inhibiting the formation

of trichomes (Wang et al., 2008; Wester et al., 2009). Feedback loop regulation has

been reported as an effective strategy to maintain normal organism development by

many HD-ZIP proteins (Ohgishi et al., 2001; Williams and Fletcher, 2005; Kim et al.,

2008; San-Bento et al., 2014). However, the trichome formation was not repressed

by the high expression level of *NbCycB2* in the $NbWo^V$-OE plants (Fig.1d, S7d, f),

which suggest that the negative effect of *NbCycB2* could be eliminated by the

mutation in $NbWo^V$. The LUC assay and GUS activity assay in the leaves of

NbCycB2pro: GFP-GUS transgenic lines also supported this conclusion (Fig. 4d, e).

Further investigation demonstrated the interaction between NbCycB2 and Nbwo

could be blocked by the mutation of woolly motif in $NbWo^V$ protein in vitro and in

vivo (Fig. 6a-b, S10a), indicating that $NbWo^V$ abolishes interaction with NbCycB2 to

prevent inhibition of NbCycB2. The high expressions of *NbCycB2* and endogenous

Nbwo in $NbWo^V$-OE lines also supported this conclusion (Fig. S7f).

In summary, through this study, we found NbCycB2 is specifically expressed in the

trichomes of *N. benthamiana* and negatively affects trichome formation. Further

study revealed the Nbwo and $NbWo^V$ were demonstrated to directly regulate

NbCycB2 and *Nbwo* expressions by binding to the L1-like box in the *NbCycB2*

promoter and its own genomic DNA sequences. In addition, the NbCycB2 protein

may via binding to the LZ domain of Nbwo dimers, which represses the activity of

Nbwo and reduces the expression of *Nbwo* downstream genes, eventually leading to

the inhibition of trichome initiation. By contrast, the interaction between NbCycB2

and Nbwo could be blocked by the mutation in woolly motif ($NbWo^V$), which

prevents the repression by NbCycB2, and results in the dramatic increase of

trichome density and branching. In previous studies, since *SlCycB2* is highly
expressed in the *Wo* (wo gain-of-function mutant alleles) and *Wo*^{V} lines and
underexpressed in the *Wo-RNAi* lines, SlCycB2 is believed to promote the
development of type I trichomes in tomato (Yang et al., 2011). However, the
function and detailed molecular mechanisms of NbCycB2 in regulating trichome
development have not been studied. Our findings provide further insights into the
regulatory network of *NbCycB2* and *Nbwo* or *NbWo*^{V} in multicellular trichomes
development of *N. benthamiana* (Fig. S11).

Acknowledgements:
We would like to thank Prof. Xi Huang for kindly providing pBridge and
pGreen-0800-II report vectors and thank Dr. Yongjia Zhong, Fujian Agriculture and
Forestry university, for kindly providing p2YN and p2YC vectors, and thank Prof. Tao
Huang, Prof. Yi Tao and Prof. Hongrui Wang for the help in the implement of
experiments. The authors declare no competing financial interests. This study was
supported by the State Tobacco Monopoly Administration of China [grant No.
110201401003 (JY-03)], XMU Training Program of Innovation and Enterpreneurship
for Undergraduates (2016Y0635) and Guizhou Science and Technology Major Project
[Grant No: (2019)3001-2].

Author contributions:
Liang Chen, Hong Cui, Shuang Wu and Minliang Wu conceived and designed the
experiments. Minliang Wu, Yuchao Cui, Li Ge, Lipeng Cui, Zhichao Xu, Hongying
Zhang, Zhaojun Wang and Dan Zhou performed the experiments and analysed the
results. Minliang Wu and Shuang Wu wrote the main manuscript text. Liang Chen
supervised the project.

Supplemental Data
Supplemental Fig. S1: The Scanning electron micrographs (SEMs) of trichomes in the
leaf of *N. benthamiana.*

Supplemental Fig. S2: Sequence analysis of *Nbwo, NbCycB2* and their similar

proteins.

Supplemental Fig. S3: The subcellular localization and auto activation test of

NbCycB2, Nbwo and *NbWo^V*.

Supplemental Fig. S4: The expression pattern of *NbCycB2* and *Nbwo* in *N.*

benthamiana.

Supplemental Fig. S5: Overexpression of *NbCycB2* and RNA interference of *NbCycB2*

in *N. benthamiana.*

Supplemental Fig. S6: The root phenotypes of wild type, *NbCycB2-RNAi* #7 T1,

NbCycB2-OE #2 T1, *NbWo^V-OE* #1 T1, *Nbwo-RNAi #2* T1 seedlings

Supplemental Fig. S7: RNA interference of *Nbwo* and overexpression of *NbWo^V* in *N.*

benthamiana.

Supplemental Fig. S8: The phenotype of *NbWo^V-OE* lines.

Supplemental Fig. S9: The phenotype of overexpressing *Nbwo-SAD-mutant* in *N.*

benthamiana.

Supplemental Fig. S10: The interaction between Nbwo and NbCycb2, Nbwo and

Nbwo LZ domain.

Supplemental Fig. S11: A simplified model for regulation between *Nbwo* and

NbCycb2.

Supplemental Table S1: Primers used in this study.

Reference:

Abe, M., Takahashi, T., and Komeda, Y. (2001). Identification of a cis-regulatory element for

L1 layer-specific gene expression, which is targeted by an L1-specific homeodomain

protein. The Plant Journal **26,** 487-494.

Ariel, F.D., Manavella, P.A., Dezar, C.A., and Chan, R.L. (2007). The true story of the HD-Zip

family. Trends in plant science **12,** 419-426.

Bombarely, A., Rosli, H.G., Vrebalov, J., Moffett, P., Mueller, L.A., and Martin, G.B. (2012). A

draft genome sequence of Nicotiana benthamiana to enhance molecular

plant-microbe biology research. Molecular Plant-Microbe Interactions **25**, 1523-1530.

Chen, S., Songkumarn, P., Liu, J., and Wang, G.L. (2009). A versatile zero background

T-vector system for gene cloning and functional genomics. Plant physiology **150**,

1111-1121.

Citovsky, V., Lee, L.-Y., Vyas, S., Glick, E., Chen, M.-H., Vainstein, A., Gafni, Y., Gelvin, S.B.,

and Tzfira, T. (2006). Subcellular localization of interacting proteins by bimolecular

fluorescence complementation in planta. Journal of molecular biology **362**,

1120-1131.

Crooks, G.E., Hon, G., Chandonia, J.M., and Brenner, S.E. (2004). WebLogo: a sequence

logo generator. Genome research **14**, 1188-1190.

Cui, Y., Rao, S., Chang, B., Wang, X., Zhang, K., Hou, X., Zhu, X., Wu, H., Tian, Z., Zhao, Z.,

Yang, C., and Huang, T. (2015). AtLa1 protein initiates IRES-dependent translation of

WUSCHEL mRNA and regulates the stem cell homeostasis of Arabidopsis in response

to environmental hazards. Plant, Cell & Environment **38**, 2098-2114.

Fernandez-Pozo, N., Menda, N., Edwards, J.D., Saha, S., Tecle, I.Y., Strickler, S.R.,

Bombarely, A., Fisher-York, T., Pujar, A., Foerster, H., Yan, A., and Mueller, L.A.

(2015). The Sol Genomics Network (SGN)—from genotype to phenotype to breeding.

Nucleic acids research **43**, D1036-D1041.

Freeman, B.C., and Beattie, G.A. (2008). An overview of plant defenses against pathogens

and   herbivores.   The   Plant   Health   Instructor

http://dx.doi.org/10.1094/PHI-I-2008-0226-01.

Gan, L., Xia, K., Chen, J.G., and Wang, S. (2011). Functional characterization of

653   TRICHOMELESS2, a new single-repeat R3 MYB transcription factor in the regulation

654   of trichome patterning in Arabidopsis. BMC plant biology **11**, 1-12.

655   **Gao, S., Gao, Y., Xiong, C., Yu, G., Chang, J., Yang, Q., Yang, C., and Ye, Z.** (2017). The

656   tomato B-type cyclin gene, SlCycB2, plays key roles in reproductive organ

657   development, trichome initiation, terpenoids biosynthesis and Prodenia litura defense.

658   Plant Science **262**, 103-114.

659   **Gendrel, A.V., Lippman, Z., Martienssen, R., and Colot, V.** (2005). Profiling histone

660   modification patterns in plants using genomic tiling microarrays. Nature Methods **2**,

661   213-218.

662   **Glas, J.J., Schimmel, B.C., Alba, J.M., Escobar-Bravo, R., Schuurink, R.C., and Kant, M.R.**

663   (2012). Plant glandular trichomes as targets for breeding or engineering of resistance

664   to herbivores. International journal of molecular sciences **13**, 17077-17103.

665   **Goodin, M.M., Zaitlin, D., Naidu, R.A., and Lommel, S.A.** (2008). Nicotiana benthamiana: its

666   history and future as a model for plant–pathogen interactions. Molecular plant-microbe

667   interactions **21**, 1015-1026.

668   **Grebe, M.** (2012). The patterning of epidermal hairs in Arabidopsis--updated. Current opinion

669   in plant biology **15**, 31-37.

670   **Guo, C., Luo, C., Guo, L., Li, M., Guo, X., Zhang, Y., Wang, L., and Chen, L.** (2016).

671   OsSIDP366, a DUF1644 gene, positively regulates responses to drought and salt

672   stresses in rice. Journal of integrative plant biology **58**, 492-502.

Hollósy, F. (2002). Effects of ultraviolet radiation on plant cells. Micron **33**, 179-197.

Huchelmann, A., Boutry, M., and Hachez, C. (2017). Plant Glandular Trichomes: Natural Cell

Factories of High Biotechnological Interest. Plant physiology **175**, 6-22.

Jefferson, R.A., Kavanagh, T.A., and Bevan, M.W. (1987). GUS fusions: beta-glucuronidase

as a sensitive and versatile gene fusion marker in higher plants. The EMBO journal **6**,

3901.

Kang, J.H., Campos, M.L., Zemelis-Durfee, S., Al-Haddad, J.M., Jones, A.D., Telewski, F.W.,

Brandizzi, F., and Howe, G.A. (2016). Molecular cloning of the tomato Hairless gene

implicates actin dynamics in trichome-mediated defense and mechanical properties of

stem tissue. Journal of experimental botany **67**, 5313-5324.

Kim, Y.S., Kim, S.G., Lee, M., Lee, I., Park, H.Y., Seo, P.J., Jung, J.H., Kwon, E.J., Suh, S.W.,

Paek, K.H., and Park, C.M. (2008). HD-ZIP III activity is modulated by competitive

inhibitors via a feedback loop in Arabidopsis shoot apical meristem development. The

Plant cell **20**, 920-933.

Kirik, V., Simon, M., Huelskamp, M., and Schiefelbein, J. (2004). The ENHANCER OF TRY

AND CPC1 gene acts redundantly with TRIPTYCHON and CAPRICE in trichome and

root hair cell patterning in Arabidopsis. Developmental Biology **268**, 506-513.

Letunic, I., and Bork, P. (2018). 20 years of the SMART protein domain annotation resource.

Nucleic acids research **46**, D493-D496.

Long, J.A., Ohno, C., Smith, Z.R., and Meyerowitz, E.M. (2006). TOPLESS Regulates Apical

Embryonic Fate in Arabidopsis **312**, 1520-1523.

Mauricio, R., and Rausher, M.D. (1997). Experimental manipulation of putative selective

agents provides evidence for the role of natural enemies in the evolution of plant

defense. Evolution, 1435-1444.

Ogawa, E., Yamada, Y., Sezaki, N., Kosaka, S., Kondo, H., Kamata, N., Abe, M., Komeda, Y.,

and Takahashi, T. (2015). ATML1 and PDF2 Play a Redundant and Essential Role in

Arabidopsis Embryo Development. Plant & cell physiology **56**, 1183-1192.

Ohgishi, M., Oka, A., Morelli, G., Ruberti, I., and Aoyama, T. (2001). Negative autoregulation

of the Arabidopsis homeobox gene ATHB-2. The Plant Journal **25**, 389-398.

Oppenheimer, D.G., Herman, P.L., Shan, S., Esch, J., and Marks, M.D. (1991). A myb gene

required for leaf trichome differentiation in Arabidopsis is expressed in stipules. Cell **67,**

483-493.

Pattanaik, S., Patra, B., Singh, S.K., and Yuan, L. (2014). An overview of the gene regulatory

network controlling trichome development in the model plant, Arabidopsis. Frontiers in

Plant Science **5**, 259.

Pauwels, L., Barbero, G.F., Geerinck, J., Tilleman, S., Grunewald, W., Perez, A.C., Chico,

J.M., Bossche, R.V., Sewell, J., Gil, E., Garcia-Casado, G., Witters, E., Inze, D., Long,

J.A., De Jaeger, G., Solano, R., and Goossens, A. (2010). NINJA connects the

co-repressor TOPLESS to jasmonate signalling. Nature **464**, 788-791.

Payne, C.T., Zhang, F., and Lloyd, A.M. (2000). GL3 encodes a bHLH protein that regulates

trichome development in arabidopsis through interaction with GL1 and TTG1.

Genetics **156**, 1349-1362.

Rerie, W.G., Feldmann, K.A., and Marks, M.D. (1994). The GLABRA2 gene encodes a homeo

domain protein required for normal trichome development in Arabidopsis. Genes &

development **8**, 1388-1399.

Sallets, A., Beyaert, M., Boutry, M., and Champagne, A. (2014). Comparative proteomics of

short and tall glandular trichomes of Nicotiana tabacum reveals differential metabolic

activities. Journal of proteome research **13**, 3386-3396.

San-Bento, R., Farcot, E., Galletti, R., Creff, A., and Ingram, G. (2014). Epidermal identity is

maintained by cell-cell communication via a universally active feedback loop in

Arabidopsis thaliana. The Plant Journal **77**, 46-58.

Schmidt, G.W., and Delaney, S.K. (2010). Stable internal reference genes for normalization of

real-time RT-PCR in tobacco (Nicotiana tabacum) during development and abiotic

stress. Molecular genetics and genomics : MGG **283**, 233-241.

Schnittger, A., Schöbinger, U., Stierhof, Y.-D., and Hülskamp, M. (2002). Ectopic B-Type

Cyclin Expression Induces Mitotic Cycles in Endoreduplicating Arabidopsis Trichomes.

Current Biology **12**, 415-420.

Schnittger, A., Schöbinger, U., Stierhof, Y.-D., and Hülskamp, M. (2005). Ectopic B-Type

Cyclin Expression Induces Mitotic Cycles in Endoreduplicating Arabidopsis Trichomes.

Current Biology **15**, 980.

Schnittger, A., Folkers, U., Schwab, B., Jürgens, G., and Hülskamp, M. (1999). Generation of

a spacing pattern: the role of triptychon in trichome patterning in Arabidopsis. The

Plant cell **11**, 1105-1116.

Serna, L., and Martin, C. (2006). Trichomes: different regulatory networks lead to convergent

structures. Trends in plant science **11**, 274-280.

Shen, Q., Liu, Z., Song, F., Xie, Q., Hanley-Bowdoin, L., and Zhou, X. (2011). Tomato

SlSnRK1 protein interacts with and phosphorylates βC1, a pathogenesis protein

encoded by a geminivirus β-satellite. Plant physiology **157,** 1394-1406.

Szemenyei, H., Hannon, M., and Long, J.A. (2008). TOPLESS Mediates Auxin-Dependent

Transcriptional Repression During Arabidopsis Embryogenesis **319,**

1384-1386.

Valkama, E., SALMINEN, J.P., Koricheva, J., and Pihlaja, K. (2003). Comparative analysis of

leaf trichome structure and composition of epicuticular flavonoids in Finnish birch

species. Annals of Botany **91,** 643-655.

Wada, T., Tachibana, T., Shimura, Y., and Okada, K. (1997). Epidermal cell differentiation in

Arabidopsis determined by a Myb homolog, CPC. Science **277,** 1113-1116.

Walker, A.R., Davison, P.A., Bolognesi-Winfield, A.C., James, C.M., Srinivasan, N., Blundell,

T.L., Esch, J.J., Marks, M.D., and Gray, J.C. (1999). The TRANSPARENT TESTA

GLABRA1 locus, which regulates trichome differentiation and anthocyanin

biosynthesis in Arabidopsis, encodes a WD40 repeat protein. The Plant cell **11,**

1337-1349.

Wang, S., Hubbard, L., Chang, Y., Guo, J., Schiefelbein, J., and Chen, J.G. (2008).

Comprehensive analysis of single-repeat R3 MYB proteins in epidermal cell patterning

and their transcriptional regulation in Arabidopsis. BMC plant biology **8,** 1-13.

Wester, K., Digiuni, S., Geier, F., Timmer, J., Fleck, C., and Hülskamp, M. (2009). Functional

diversity of R3 single-repeat genes in trichome development. Development **136,**

1487-1496.

Williams, L., and Fletcher, J.C. (2005). Stem cell regulation in the Arabidopsis shoot apical

meristem. Current opinion in plant biology **8**, 582-586.

Wu, R., and Citovsky, V. (2017a). Adaptor proteins GIR1 and GIR2. II. Interaction with the

co-repressor TOPLESS and promotion of histone deacetylation of target chromatin.

Biochemical and biophysical research communications **488**, 609-613.

Wu, R., and Citovsky, V. (2017b). Adaptor proteins GIR1 and GIR2. I. Interaction with the

repressor GLABRA2 and regulation of root hair development. Biochemical and

biophysical research communications **488**, 547-553.

Yan, T., Chen, M., Shen, Q., Li, L., Fu, X., Pan, Q., Tang, Y., Shi, P., Lv, Z., Jiang, W., Ma,

Y.N., Hao, X., Sun, X., and Tang, K. (2016). HOMEODOMAIN PROTEIN 1 is required

for jasmonate-mediated glandular trichome initiation in Artemisia annua. New

Phytologist **213**, 1145-1155.

Yang, C., Gao, Y., Gao, S., Yu, G., Xiong, C., Chang, J., Li, H., and Ye, Z. (2015).

Transcriptome profile analysis of cell proliferation molecular processes during

multicellular trichome formation induced by tomato Wo (v) gene in tobacco. BMC

genomics **16**, 868.

Yang, C., Li, H., Zhang, J., Luo, Z., Gong, P., Zhang, C., Li, J., Wang, T., Zhang, Y., and Lu,

Y.E. (2011). A regulatory gene induces trichome formation and embryo lethality in

tomato. Proceedings of the National Academy of Sciences of the United States of

America **108**, 11836-11841.

Yoo, S.D., Cho, Y.H., and Sheen, J. (2007). Arabidopsis mesophyll protoplasts: a versatile cell

system for transient gene expression analysis. Nature protocols **2**, 1565-1572.

Figure legends

Fig. 1: The trichome phenotypes of *NbCycB2*, *Nbwo* and *NbWo*^{V} transgenic seedlings

(a), (b), (c), (d), (e), are the trichomes SEMs of wild type, *NbCycB2-RNAi* #7 T1,

NbCycB2-OE #2 T1, *NbWo*^{V}*-OE* #1 T1, *Nbwo-RNAi* #2 T1 10d-old-seedlings

respectively. The white bar is 500 μm. (f) The trichome density of the wild type,

NbCycB2-RNAi #7 T1, *NbCycB2-OE* #2 T1, *NbWo*^{V}*-OE* #1 T1, *Nbwo-RNAi* #2 T1

10d-old-seedlings leaves were shown. "*" indicates a difference at P<0.05 by

Student's t test compared to WT. "**" represent significant difference against WT at

P < 0.01. Error bars represent SD (n = 3).

Fig. 2: Over expression of *Nbwo* also induce the dwarf phenotype in *N. benthamiana*

(a) Over expression of *Nbwo* was drivered by P35S promoter. The trichomes density

reduced in the stems and leaves of transgenic lines. (b) The relative expression level

of *Nbwo* and*NbCycB2* in the transgenic lines measured by qRT-PCR. Error bars

represent SD (n = 3). (c) The root lengths were measured in the Wild type,

NbCycB2-OE #1 T1, *Nbwo-OE* #3 T1, *Nbwo-RNAi* #2 T1 two weeks-old-seedlings. The

white bar is 1 cm. (d) The root hairs of *Nbwo-OE* #3 T1 and *NbWo*^{V}*-OE* #1 T1 two

weeks-old-seedlings were detected by microscope. The red bar is 1 mm. (e), (f) are

the SEMs of the *Nbwo-OE* #3 T1 and *NbWo*^{V}*-OE* #1 T1 10d-old-seedlings respectively.

The white bar is 500 μm.

Fig. 3: Nbwo and NbWo^V protein can bind *NbCycB2* promoter in vitro and vivo

(a) The fragments of *NbCycB2* promoter were used to CiHP (P1, P2 and P3) and yeast

one-hybrid assays (A, B, C, D and E). The numbers indicate positions of the *NbCycB2*

promoter truncations. (b) Graphs show the ratio of bound promoter fragments

(P1-P3) versus total input detected by qRT-PCR after immuno-precipitation in

HA-NbWo^V by HA antibodies. Data shown are mean ± SE (n = 3). (c) One-hybrid (Y1H)

assays were used to determine the interaction of *NbCycB2* promoter fragments (A, B,

C, D, E) bait constructs and AD-*Nbwo* or empty pGADT7 constructs in Y187 yeast

strains. (d) The schematic diagram of the effectors and reporters constructs were

used in the LUC assay. (e) The relative reporter activities were measured in *N.*

benthamiana protoplasts after transiently transformed the effector and reporter

constructs. The relative LUC activities normalized to the REN activity are shown

(LUC/REN). The difference between combinations were detected by Student's t test.

"**" present that the LUC activity is significantly different (P< 0.01). Error bars

represent SD (n = 3).

Fig. 4: NbCycB2 can suppress the function of Nbwo, but has no effect on NbWo^V

(a) The DNA sequences of wild type *NbCycB2* promoter (ProD) and mutant fragments

were shown. L1-like boxes were indicated by using gray arrow. The mutated bases in

the m1 and m2 sequences are shown by a dividing line. (b) One-hybrid (Y1H) assays

were used to detected the interaction between mutant D fragments of *NbCycB2*

promoter bait constructs and AD-*Nbwo* or AD-*NbWo^V* in Y187 yeast strains, the

empty pGADT7 construct as the control. (c) Relative reporter activity was measured

in *N. benthamiana* protoplasts after transiently co-transformed the effector and

reporter constructs. The relative LUC activities normalized to the REN activity are

shown (LUC/REN). "**" present that the LUC activity has significant difference

between the D and m2 fragments reporter when their co-transformed with *Nbwo* or

NbWo^V effector respectively (P< 0.01, Student's t test). Error bars represent SD (n =

3). (d) The GUS staining of the *proNbCycB2: GFP-GUS* transgenic line leaves, when

co-expressing *P35S: Nbwo* and *P35S: NbCycB2*, *P35S: NbWo^V* and *P35S: NbCycB2*. The

area of red circle was injected with *P35S: Nbwo* GV3101 strain; the area of Green

circle was injected with *P35S: NbWo^V* GV3101 strain; the area of blue circles were

injected with *P35S: NbCycB2* GV3101 strain. (e) The LUC activity of co-transforming

the effector and reporter constructs were measured. "**" present that the LUC

activity is significantly different between *Nbwo* and *NbWo^V* effector when they

co-transformed with *NbCycB2* effector respectively (P< 0.01, Student's t test). Error

bars represent SD (n = 3).

Fig. 5: NbCycB2 protein interact with LZ domain of the Nbwo protein in vitro and vivo

(a) Schematic diagrams of Nbwo protein domain constructs. The numbers indicate

positions of the first and the last amino acid of the Nbwo truncations. (b) Interaction

between NbCycB2 and domains of Nbwo protein were determined by YH2 system. (c)

The interaction between Nbwo and NbCycB2 or itself were demonstrated by BiFC

assay. Each indicated pair of constructs were co-infiltrated into leaves of *N.*

benthamiana (Bar, 50 µm).

Fig. 6: Mutation of woolly motif reduce the interaction between Nbwo and NbCycB2

(a) Interaction between NbCycB2 and Nbwo or NbWo^V proteins were determined by

YH2 system. Blue clones grown on the QDO/X/A medium indicates positive

protein-protein interactions. (b) The pull down assay between Nbwo or NbWo^V with

NbCycB2 proteins. Only recombinant HIS-Nbwo protein can co-precipitate with

GST-NbCycB2 protein. (c) The competitive binding between NbCycB2 and LZ domain,

LZ domain and Nbwo were determined by Yeast three-hybrid assays. (d) The co-IP

assay between NbCycB2 and Nbwo dimers. The total protein immunoprecipitation

was performed by using anti-HA beads.

Fig. 7: Nbwo and NbWo^V can bind at itself genomic DNA sequences

(a) The fragments of *Nbwo* genomic sequences were used to CHiP (T1, T2, T3, T4, T5,

T6 and T7) and yeast one-hybrid assays (G1, G2, G3 and G4). The bar present 1 kb

DNA sequences. (b) Graphs show the ratio of bound genomic fragments (T1-T7)

versus total input detected by real-time PCR after immuno-precipitation in

NbWo^V-OE lines by HA antibodies. Data shown are mean ± SE (n = 3). (c) One-hybrid

(Y1H) assays were used to determine the interaction of *Nbwo* genomic sequence

fragments (G1, G2, G3, G4) bait constructs and AD-Nbwo, AD-NbWo^V or empty

pGADT7 constructs in Y187 yeast strains. The clones grown on the SD/-Leu/-His/-Trp

(-L-W-H) with 60 mM 3-AT medium indicates the interaction between DNA fragment

and Nbwo or NbWo^V proteins.

Fig. 8: Hybridization between *Nbwo-OE* and *NbCycB2-OE* plants

(a), (b) The phenotype of *NbCycB2-OE* #2 and *Nbwo-OE* #3 hybridization F1 two

weeks old (a) and mature (b) plants were shown. "I"present wild type *N.*

benthamiana, "II"present *NbCycB2-OE/WT* hybrid T1 plants, "III"present *NbCycB2-OE*

#2 and *Nbwo-OE #3* hybridization F1 plants, "IV"present *Nbwo-OE #3/WT* hybrid T1

plants. (c) F1 plants were tested by using PCR. Wild type as a negative control,

Nbwo-OE #3 and *NbCycB2-OE* #2 as positive controls. No DNA band was detected in

the wild type *N. benthamiana*. A ~330 bp DNA band was detected in *NbCycB2-OE* #2

lines. A ~2199 bp DNA band was detected in *Nbwo-OE #3* lines. In contrary, two

bands were detected in *NbCycB2-OE #2* and *Nbwo-OE #3* hybridization F1 plants. (d)

The relative expression levels of *Nbwo* and *NbCycb2* were measured by qRT-PCR in

F1 plants. Data are given as means SD (n = 3).

(a) WT

(b) *NbCycB2-RNAi #7*

(c) *NbCycB2-OE #2*

(d) *NbWo^V-OE #1*

(e) *Nbwo-RNAi #2*

(f) The number of trichome (/0.5 mm²)

(a)

WT | **Nbwo-OE #1** | **Nbwo-OE #2** | **Nbwo-OE #3**

Adaxial leaf

Stem

(b)

Relative expression

□ Nbwo
■ NbCycB2

WT | Nbwo-OE #1 | Nbwo-OE #2 | Nbwo-OE #3

(c)

WT | **NbCycB2-OE #1** | **Nbwo-OE #3** | **Nbwo-RNAi #2**

(d)

Nbwo-OE #3 | *NbWo^V-OE #1*

(e)

Nbwo-OE #3

(f)

NbWo^V-OE #1

(a)

ProD −815 ... L1-like box ... L1-like box ... −695

ProD-m1

ProD-m2

(b)

	PREYS	−L−W		−L−W−H+3-AT		
		1 OD	0.1 OD	1 OD	0.1 OD	
Control						Pro-D
AD-Nbwo						
AD-NbWo^V						
AD-Nbwo						m1
AD-Nbwo						m2
AD-NbWo^V						

BAITS

(c)

LUC/REN

ProD	+	−	+	−
ProD-m2	−	+	−	+
Nbwo	+	+	−	−
NbWo^V	−	−	+	+

** **

(d)

P35S:Nbwo *P35S:NbWo^V*

P35S:NbCycB2 *P35S:NbCycB2*

(e)

LUC/REN

ProD	+	+
NbCycB2	+	+
Nbwo	+	−
NbWo^V	−	+

**

(a)

I: *WT*
II: *NbcycB2-OE #2 / WT*
III: *Nbwo-OE #3 / NbcycB2-OE #2*
IV: *Nbwo-OE #3 /WT*

(b)

(c)

(d)