Neuromuscular fatigue and recovery after strenuous exercise

depends on skeletal muscle size and stem cell characteristics

Baumert, P.[1,2]*, Temple, S.[1], Stanley, J.M.[1], Cocks M.[1], Strauss, J.A.[1], Shepherd

S.O.[1], Drust, B.[1], Lake, M.J.[1], Stewart, C.E.[1], Erskine, R.M.[1,3]

[1]Research Institute for Sport & Exercise Sciences, Liverpool John Moores University,
Liverpool, United Kingdom.
[2]Exercise Biology Group, Faculty of Sport and Health Sciences, Technical University of Munich,
Munich, Germany

[3]Institute of Sport, Exercise & Health, University College London, London, UK

Running title: Neuromuscular Response following muscle damage *in vivo* and *in vitro*

Address for reprint requests and all other correspondence:

Philipp Baumert, Exercise Biology Group, Faculty of Sport and Health Sciences, Technical University of Munich, Munich, Germany; Email: philipp.baumert@tum.de

Key words: Exercise-induced muscle damage (EIMD); biceps femoris architecture; extracellular matrix (ECM); myoblast; fibroblast

This is an original research article

ABSTRACT

Hamstring muscle injury is highly prevalent in sports involving repeated maximal sprinting. Although neuromuscular fatigue is thought to be a risk factor, the physiological and cellular mechanisms underlying the fatigue response to repeated maximal sprints are unclear. Here, we show that repeated maximal sprints induce neuromuscular fatigue accompanied with a prolonged strength loss in hamstring muscles. The immediate hamstring strength loss was linked to both central and peripheral fatigue, while prolonged strength loss was associated with indicators of muscle damage. The kinematic changes immediately after sprinting likely protected fatigued hamstrings from excess elongation stress and larger hamstring muscle physiological cross-sectional area appeared to protect against fatigue/damage. Specific skeletal muscle stem cell composition (i.e lower *myoblast:fibroblast* ratio) was linked to improved muscle recovery within 48 h after sprinting. Contrastingly, a high *myoblast:fibroblast* ratio appears crucial for the latter stage of muscle regeneration. We have therefore identified novel mechanisms that likely regulate the fatigue/damage response and recovery following repeated maximal sprinting in humans.

INTRODUCTION

Hamstring strain is the most frequently occurring injury in sport (Crema *et al.*, 2017), particularly in those sports that involve high-speed running (Ekstrand *et al.*, 2011; Opar *et al.*, 2012). Although the aetiology is unclear, numerous risk factors have been proposed, such as short fascicle length (Timmins *et al.*, 2015), poor flexibility (Jonhagen *et al.*, 1994), poor hamstring strength (Opar *et al.*, 2014), and inadequate warm-up (Woods *et al.*, 2007). Further, it is unknown whether hamstring strain is the result of a single event that exceeds the physiological range of hamstring muscle extensibility and contractility, or as a result of an accumulation of eccentric contractions during repeated maximal sprints, causing neuromuscular fatigue (Opar *et al.*, 2012). Neuromuscular fatigue is responsible for acute, as well as prolonged, impairment of muscle function, classified as central fatigue (i.e. originating in the central nervous system), or as peripheral fatigue (i.e. distal to the neuromuscular junction) (Byrne *et al.*, 2004). Although it was recently reported that both central and peripheral fatigue contribute to impaired hamstring muscle function immediately after repeated maximal sprint-related interventions (Marshall *et al.*, 2014; Timmins *et al.*, 2014), the contribution of neuromuscular fatigue to hamstring muscle impairment and recovery, following repeated maximal sprints over time, is insufficiently studied (Verma *et al.*, 2015). An understanding of hamstring neuromuscular fatigue following repeated maximal sprints may be crucial for understanding hamstring strain aetiology (van der Horst *et al.*, 2015).

Peripheral fatigue may be caused by ultrastructural muscle damage, which is indicated by Z-line disturbance (Newham *et al.*, 1983) as well as disruption of the extracellular matrix (Stauber *et al.*, 1990). The extracellular matrix provides structural scaffolding for muscle remodelling and plays an integral role in force transmission (Wang *et al.*, 2009; Gillies & Lieber, 2011). This is referred to as exercise-induced muscle damage and it is exhibited by prolonged strength loss and delayed-onset muscle soreness, as well as the release of muscle-specific proteins [e.g. creatine kinase (CK)] into the circulation over the following days (Howatson & Van Someren, 2008). After substantial muscle damage, myogenic satellite cells (skeletal muscle

stem cells), play a key role in skeletal muscle regeneration and remodelling (Tidball, 2011).

Activated satellite cells (myoblasts) proliferate and migrate from their niche along the basal

lamina to the injury site before terminally differentiating and fusing with damaged myofibrils to

repair injury. There is increasing evidence that fibroblasts, the main cell type of muscle

connective tissue, also play a critical role in supporting muscle regeneration (Murphy *et al.*,

2011; Fry *et al.*, 2017; Mackey *et al.*, 2017). Following damage, infiltrating inflammatory cells

activate muscle fibroblasts, which proliferate and migrate to the area of the myotrauma and

produce extracellular matrix components in an orchestrated and regulated fashion to support

healthy muscle remodelling (Joe *et al.*, 2010; Murphy *et al.*, 2011). However, little is known

about the role of fibroblasts during the initial response and recovery following *physiological*

exercise-induced muscle damage, e.g. following repeated sprinting.

Acute damage to the muscle-tendon complex may facilitate hamstring strain, which is thought

to occurs in the late swing phase of sprinting, when the hamstring muscles contract

eccentrically, i.e. trying to shorten while being lengthened in an attempt to decelerate the shaft

before initial foot-ground contact (Thelen *et al.*, 2005; Chumanov *et al.*, 2012). Therefore, a

short biceps femoris long head (BF_{LH}) fascicle length has been suggested to increase

hamstring strain risk (Timmins *et al.*, 2016a), as the BF_{LH} is thought to be relatively more

eccentrically stretched during the late swing phase of sprinting compared to the other

hamstring muscles (Thelen *et al.*, 2005). However, no study has investigated the relationship

between BF_{LH} architecture (including muscle fascicle length and cross-sectional area), and the

prolonged hamstring muscle response to exercise-induced neuromuscular fatigue. Finally,

lower limb neuromuscular fatigue might cause a number of biomechanical alterations in

running kinematics (Paschalis *et al.*, 2007). However, it is not known whether repeated

maximal sprints influence kinematic patterns, and whether this can lead to prolonged changes

in lower-limb kinematics, which may play a role in the development of muscle strain following

insufficient recovery (Opar *et al.*, 2012).

Here we demonstrated that both central and peripheral fatigue, assessed via surface

electromyography (sEMG) and electrical stimulation, contributed to the immediate loss of

muscle function in both the quadriceps and hamstring muscle groups, but that peripheral

factors mainly contributed to the sustained loss of hamstring muscle function. Moreover, we

established that a lower myoblast:fibroblast ratio in isolated primary human muscle stem cells

correlated with improved recovery from both repeated maximal sprints and an *in vitro* artificial

wounding assay within the first 48 h. However, at seven days post *in vitro* damage, a higher

myoblast:fibroblast ratio appeared important for optimal muscle regeneration. We also report

that BF_{LH} architecture (i.e. PCSA) was associated with hamstring fatigue, and that

neuromuscular fatigue led to reduced hip flexion and knee extension during the late swing

phase of steady-state running. Thus, with this interdisciplinary study, we identify novel cellular

and neuromuscular mechanisms underpinning central and peripheral fatigue following

repeated sprinting, which ultimately led to kinematic changes during the running stride phase

associated with hamstring strain injury*.*

RESULTS

Effect of the Repeated Maximal Sprint Intervention on Neuromuscular Fatigue

To examine the effect of repeated maximal sprints on neuromuscular fatigue, we measured

different fatigue parameters before (PRE), immediately after (POST), and 48 h after (POST48)

the repeated maximal sprint intervention. The average 30 m sprinting speed was 6.48 ± 0.33

m s^{-1}. There was a main effect of time for heart rate, 30 m sprinting time, rating of perceived

exertion and lactate concentration, with all parameters increasing from PRE to POST (all

$P<0.001$). Blood lactate concentration increased from PRE (1.63 ± 0.45 mmol/L) to POST (9.82

± 3.62 mmol/L; $P<0.001$). The sprinting performance (measured via the performance

decrement score (Glaister *et al.*, 2008)) decreased by 3.98 ± 2.99% during the run, and rating

of perceived exertion increased by 96.5 ± 35.2% from PRE-to-POST, indicating fatigue had

occurred.

We then performed *in vivo* functional analysis to assess if repeated maximal sprints resulted in an increase in central and/or peripheral fatigue. We, therefore, measured BF_{LH} muscle activation via normalised surface electromyography (sEMG) during hamstring maximum voluntary contraction (MVC). Assessment revealed that a change had occurred in sEMG ($F_{F2,24}$=4.35, P=0.022), with post-hoc pairwise comparisons revealing a decrease from PRE-to-POST (-24.3%; P=0.019). However, this change was no longer evident at POST48 (P=0.157, Table 1), suggesting that central fatigue occurred immediately after repeated maximal sprints. No other changes in muscle (co)activation were observed at any time point (P>0.05).

Table 1 *Effect of the intermitted sprint intervention on measures of muscle activation. Data are presented as mean ± SD. One-way ANOVA, F- and P-values are reported.*

Assessment [unit]	n	PRE	POST	POST48	F-Test	P Value
Hamstring Muscle Voluntary Activation [ITT, %]	20	98.5 ± 2.64	94.1 ± 7.83	96.9 ± 5.96	F(1.4,38) = 2.75	0.099
Normalised BF_{LH} knee flexion sEMG$_{max}$ [%]	14	3.32 ± 1.33	2.27 ± 0.72*	2.85 ± 1.16	F(2,24) = 4.35	**0.022**
Vastus lateralis knee extension sEMG$_{max}$ [mV]	15	0.50 ± 0.29	0.47 ± 0.34	0.51 ± 0.34	F(2,24) = 0.17	0.772
Quadriceps CoA during 30° hamstring MVC [%]	12	5.74 ± 7.23	4.08 ± 6.86	4.33 ± 8.76	F(2,22) = 0.50	0.613
Hamstring CoA during 80° quadriceps MVC [%]	10	4.79 ± 3.37	6.62 ± 3.79	7.51 ± 4.16	F(2,18) = 1.64	0.223

ITT – interpolated twitch technique; BF_{LH} – Biceps femoris long head; CoA – Co-Activation; sEMG – surface Electromyography. * Different to PRE (P<0.05).

We also assessed the torque-frequency relationship *in vivo* via electrical stimulation to indicate

peripheral (muscle) fatigue. There was an interaction between time x stimulation frequency

(n=19; $F_{4.9,216}$=6.62, P<0.001; Figure 1A). Post-hoc paired t-tests revealed differences PRE-to-

POST for 10-50 Hz (P<0.05), but lower frequencies between 10-20 Hz reverted to baseline

values POST48 (P>0.05), while the frequencies of 30 and 50 Hz were still decreased POST48

compared to their baseline values (P<0.05), providing evidence that peripheral fatigue

occurred immediately after the repeated sprints and remained for 48 h

Figure 1 (A) *Torque-frequency relationship, all frequencies were normalised to 100 Hz. * significant*
differences between before (PRE) and immediadtely after (POST) the repeated maximal sprint
intervention, P<0.05; # significant differences between PRE and POST, and between PRE and POST48,
*P<0.05. Data are presented as mean ± SEM. **(B)** Comparison of relative maximal voluntary contraction*
(MVC) loss between hamstring and quadriceps muscle group before (PRE), immediately after (POST)
*and 48h after (POST48) the repeated maximal sprint intervention. * signifcant differences compared to*
PRE, P< 0.001; # significant differences between quadriceps and hamstring MVC, P< 0.05. Data are
expressed as mean ± SEM.

Effect of the Repeated Maximal Sprint Intervention on MVC Strength, Muscle

Soreness and Serum Markers of Exercise-Induced Muscle Damage

To investigate the effect of intermittent sprints on biomarkers of exercise-induced muscle

damage, we assessed hamstring (knee flexion) and quadriceps (knee extension) MVC, muscle

soreness, serum creatine kinase (CK) activity and interleukin-6 (IL-6) concentrations PRE,

POST and POST48. Isometric hamstring and quadriceps MVC, muscle soreness (all P<0.001)

and serum CK activity ($F_{1.3,34}$=5.98, P=0.017), as well as IL-6 concentration ($F_{1.3,34}$=5.96,

P=0.018), showed a main effect of time, which are indicators of muscle damage (Table 2) that was similar to other studies (Howatson & Milak, 2009; Verma *et al.*, 2015; Chen *et al.*, 2017). Post-hoc pairwise comparisons revealed that, compared to PRE, both serum CK activity (+93.0%) and serum IL-6 concentration (+307%) were elevated at POST (both P=0.027), and CK activity further increased at POST48 (+256%; P=0.012), while serum IL-6 concentration reverted to baseline values (P>0.05).

Table 2 *Effect of the Repeated Maximal Sprint Intervention on Muscle Damage-Biomarkers. Values are mean ± SD. One-way ANOVA, F- and P-values are reported.*

Assessment [unit]	PRE	POST	POST48	F-Test	P Value
Quadriceps MVC [N·m]	270.5 ± 51.6*	222.4 ± 52.5*	243.0 ± 71.3*	$F_{(2,38)}$ = 16.55	**<0.001**
Hamstring MVC [N·m]	142.5 ± 25.0*	124.8 ± 29.9*	112.4 ± 30.1*	$F_{(2,38)}$ = 25.12	**<0.001**
Squat Muscle soreness [cm]	0.20 ± 0.41*	1.95 ± 1.61*	2.87 ± 1.71*	$F_{(2,38)}$ = 28.62	**<0.001**
Lunge Muscle soreness [cm]	0.30 ± 0.57	2.30 ± 2.08†	3.48 ± 2.07†	$F_{(2,38)}$ = 17.02	**<0.001**
Range of Motion [°]	120.3 ± 6.76	115.7 ± 6.77†	116.0 ± 6.27†	$F_{(2,38)}$ = 9.33	**<0.001**
CK activity [mU/mL]	27.9 ± 23.3	53.8 ± 45.3†	99.3 ± 104.5†	$F_{(1.3,34)}$ = 5.98	**0.017**
IL-6 concentration [pg/mL]	1.89 ± 3.10	7.68 ± 9.95#	1.59 ± 3.46	$F_{(1.3,34)}$ = 5.96	**0.018**

MVC – Maximal voluntary contraction; CK – Creatine Kinase; IL-6 – Interleukin-6; * Significant differences between all time points; † differences PRE–to-POST and PRE-to-POST48 (P<0.05); # differences PRE-to-POST and POST-to-POST48 (P<0.05).

Further, there was an interaction between time and muscle groups concerning relative MVC torque loss (percentage change from PRE MVC) ($F_{1.4,38}$=7.92, P=0.004). Relative MVC decreased similarly in both quadriceps and hamstring muscle groups PRE-to-POST (Figure

1B). However, at POST48, hamstring MVC continued to decrease from POST (-9.26%;

P=0.010), while quadriceps MVC began to return to PRE values (+8.47%; P=0.016) and was

higher than hamstring MVC at POST48 (P=0.038).

Effect of the Repeated Maximal Sprint Intervention on Lower-Limb Kinematics

To assess the consequential effect of neuromuscular fatigue on lower-limb kinematics, we

captured treadmill running (4.17 m s^{-1}) kinematics with an eight-camera motion capture system

over time. Three-dimensional motion analysis demonstrated, that despite significance not

being achieved, there was a tendency towards a longer running cycle time POST (+1.01%)

and POST48 (+1.86%), compared to PRE (P=0.080; Table 3). Further, treadmill running

demonstrated decreased peak knee extension (P=0.047) during the late swing phase at POST

(-10.9%) compared to PRE, but this reverted to baseline POST48. The percentage change in

peak knee extension correlated with the percentage change in relative hamstring MVC torque

both measured POST-to-POST48 (r^2=0.258, $F_{1,2}$=5.673, P=0.031).

Table 3 *Effect of the Repeated Maximal Sprint Intervention on Kinematics of Treadmill Running at 4.17 m s⁻¹ Values are mean ± SD. One-way ANOVA, F- and P-values are reported.*

Kinematics [unit]	PRE	POST	POST48	F-Test	P Value
Peak knee flexion (swing phase) [°]	-103 ± 13.9	-110 ± 12.4	-108 ± 12.6	$F_{(2,20)} = 2.84$	0.082
Peak knee extension (swing phase) [°]	-3.66 ± 5.32	-7.08 ± 5.07*	-4.29 ± 7.06	$F_{(2,20)} = 3.57$	**0.047**
Contact hip flexion (toe strike) [°]	25.2 ± 5.42	29.3 ± 7.62	17.6 ± 15.6	$F_{(1.2,20)} = 4.05$	0.062
Contact knee flexion (toe strike) [°]	-15.0 ± 6.25	-17.6 ± 6.79	-13.3 ± 9.64	$F_{(2,22)} = 2.79$	0.083
Duration Running cycle [s]	0.67 ± 0.03	0.68 ± 0.03	0.69 ± 0.02	$F_{(2,20)} = 2.88$	0.080
Stance phase duration [s]	0.18 ± 0.02	0.19 ± 0.02	0.19 ± 0.03	$F_{(2,20)} = 1.01$	0.356
Swing Phase [s]	0.50 ± 0.04	0.50 ± 0.05	0.50 ± 0.03	$F_{(1.2,20)} = 0.03$	0.899

Knee fully extended=0°; negative value indicates a flexed knee; * different to PRE (P<0.05).

Architecture of the Biceps Femoris Long Head Muscle

To assess whether architectural parameters of the BF_{LH} muscle (Figure 2A and B) were

associated with markers of peripheral fatigue, we performed ultrasound measurements of the

BF_{LH} muscle (Table supplement 5). Muscle fascicle length and pennation angle of the BF_{LH},

which have previously been linked to hamstring muscle strain risk (Timmins *et al.*, 2016b), did

not correlate with any outcome variable of neuromuscular fatigue. However, BF_{LH} PCSA (mean

±SD: 23.4 ± 4.62 cm²) correlated inversely with relative hamstring MVC loss PRE-to-POST

(R^2=0.421, $F_{1,17}$=12.37, P=0.003, Figure 2C).

Figure 2 (A) *Longitudinal image of biceps femoris long head, assessment of the biceps femoris long head is highlighted (total muscle length and fascicle length together with pennation angle at 50% of total muscle length).* **(B)** *Cross-sectional image at 60% muscle length (=100% proximal myotendinous junction), biceps femoris long head is highlighted.* **(C)** *Correlation between biceps femoris long head muscle physiological cross-sectional area and % hamstring maximum isometric voluntary contraction (MIVC) decrease from before (PRE) to immediately after (POST) the repeated maximal sprint intervention (P=0.003).*

Artificial Wound Healing Assay to Investigate Repair and Regeneration Regarding Myoblast:Fibroblast Ratio

Preliminary data from our laboratory demonstrated that skeletal muscle stem cell composition (i.e. myoblast:fibroblast ratio), derived from two volunteers with ratios representing the extreme conditions of the myoblast:fibroblast percentage, played a role in the success of artificial wound healing *in vitro* (Owens, 2015). Human primary skeletal muscle stem cells with a high myoblast:fibroblast ratio resulted in reduced cell migration into an artificial wound (assessed by cell number within the wound) when compared with stem cells with a low myoblast:fibroblast ratio. No significant differences were observed in the relative proportion of migrating cells (i.e. between myoblasts and fibroblasts) over 48 hours. Further, recent investigations have revealed that the skeletal muscle stem cell ratio does not change during *in vitro* cell culturing (Foulstone *et al.*, 2004; Grohmann *et al.*, 2005). We, therefore, further assessed the effect of the myoblast:fibroblast ratio on skeletal muscle recovery following *in vitro* artificial wounding to

extend the preliminary *in vitro* results from our laboratory. To be able to demonstrate a

coefficient of determination of ≥ 0.50 between myoblast:fibroblast ratio and our dependent

variables, a priori power calculations for a Pearson's r correlation was run using G*Power

(version 3.1.9.1), and it revealed that 11 persons provided alpha = 5%, and power = 80%.

Thus, we used primary human skeletal muscle stem cells derived from 12 participants, six who

participated in both the repeated maximal sprint intervention and also volunteered to provide

a muscle biopsy at least three weeks before the repeated maximal sprint intervention, and

another six (two male and four females), who did not participate in the repeated maximal sprint

intervention (to increase the power of the *in vitro* study). The muscle stem cells were isolated,

cultured and then characterized by immunofluorescence staining. The mean ± SD

myoblast:fibroblast ratio of the twelve participants was 1.26 ± 1.00 (range: 0.276 to 2.93). We

did not detect any differences in muscle stem cell characteristics between cells obtained from

females and males (data not shown). We, therefore, combined the data from all muscle cells

and correlated the muscle characteristics with individual myoblast:fibroblast ratios. We

observed no correlations regarding myoblast:fibroblast ratio and the total number of myoblasts

and fibroblasts migrating into the artificial wound within all three segments combined at 24 h

(R^2=0.20, $F_{1,10}$=2.56, P=0.141) or 48 h (R^2=0.02, $F_{1,10}$=0.19, P=0.671) after the scratch assay.

Figure 3 (A) *Representative images for cell migration of muscle cells with a high myoblast:fibroblast ratio (2.4; left) and with a low percentage of myoblasts (0.3; right) into the artificial wound. The wound area is about 900 μm in width and split into 3 x 300 μm segments (one inner and two outer segments). Magnification is x 10.5, and scale bar is 100 μm.* **(B)** *Inverse correlations between the myoblast:fibroblast ratio and the migration dynamics of 12 different primary muscle stem cells 24 h (P=0.011) and a trend 48 h (P=0.064) after the artificial wound healing assay.*

However, there was an inverse correlation between myoblast:fibroblast ratio and migration dynamics for the 12 participants (Figure 3 B). Muscle stem cells with a low myoblast:fibroblast ratio demonstrated more cells in the inner segment than to the outer segment compared to muscle stem cells with high myoblast:fibroblast ratio at 24 h (R^2=0.49, $F_{1,10}$=9.53, P=0.011) and with a non-significant trend at 48 h (R^2=0.30, $F_{1,10}$=4.33, P=0.064) after the artificial wound healing assay. Further, there were positive correlations between the myoblast:fibroblast ratio and all parameters at seven (myotube formation) and 10 days (CK activity) after the artificial wound healing assay (all P<0.05; Table supplement 6). Biopsies with a higher myoblast:fibroblast ratio showed more myotubes per field, which had a higher diameter and area compared to biopsies with a lower myoblast:fibroblast ratio (Figure supplement 5).

Comparison of the Muscle Response between the Repeated Maximal Sprint Protocol and the Muscle Stem Cell Study

In order to determine whether skeletal muscle stem cell ratio (i.e. myoblast:fibroblast ratio) played a role in muscle strength recovery *in vivo*, further studies were performed. Previous investigations have shown that skeletal muscles of different origin, but with similar physiological functions and fibre type composition demonstrate similar transcriptome expression patterns of up to 99% (Evangelidis *et al.*, 2016; Terry *et al.*, 2018). Further, all limb muscles arise developmentally from the ventrolateral dermomyotome of the segmented paraxial mesoderm (Randolph & Pavlath, 2015) and these muscles with a similar fibre type composition show low intra-subject (Collins *et al.*, 2005; Jakobsen *et al.*, 2018) but high inter-subject variability (Collins *et al.*, 2005; Owens *et al.*, 2015) regarding the total amount of stem cells within these muscle fibres. Thus, muscle biopsies were obtained from the vastus lateralis of the six participants, who both performed the repeated maximal sprint intervention and volunteered to provide a muscle biopsy at least three weeks before the repeated maximal sprint intervention. As fibre-type composition is similar between the quadriceps and hamstrings (Evangelidis *et al.*, 2016), the stem cell composition of the vastus lateralis was considered representative of both the quadriceps and hamstring muscle groups. The mean ± SD myoblast:fibroblast ratio of the six participants was 1.46 ± 1.06 (range: 0.299 to 2.93). There was an inverse correlation between myoblast:fibroblast ratio and the percentage change in relative hamstring MVC torque measured PRE-to-POST48 *in vivo* (R=-0.945, $F_{1,4}$=33.73, P=0.004; Figure 4A). Thus, participants with a high myoblast:fibroblast ratio showed a delayed hamstring strength recovery 48 h after repeated maximal sprints compared to those with a low myoblast:fibroblast ratio. Further, there was an inverse correlation between myoblast:fibroblast ratio and relative hamstring MVC torque measured POST-to-POST48 (R=-0.943, $F_{1,4}$=17.08, P=0.014; Figure 4B). No correlations were found between the myoblast:fibroblast ratio and changes in quadriceps MVC torque or with any other muscle damage and fatigue biomarker following repeated maximal sprints. These inverse correlations could be confounded by the

265 model assumptions and the relative low participant number. However, post hoc power

266 calculations demonstrate that the bivariate correlation had adequate power (0.97).

268 **Figure 4** *Inverse correlation between the myoblast:fibroblast ratio, assessed in the current in vitro study*
269 *and the change of hamstring MVC torque measured **(A)** before and 48 h after (P=0.004), and **(B)***
270 *measured immediately after and 48 h after (P=0.014) (B) the repeated maximal sprint intervention.*

DISCUSSION

In this study, we have used an interdisciplinary approach to investigate the potential

biomechanical, physiological and cellular factors underpinning neuromuscular fatigue following

repeated maximal sprints. We have shown that immediate strength loss was associated with

reduced hamstring sEMG activity (indicating impaired hamstring motor unit recruitment) and

markers of peripheral fatigue, but the magnitude and sustained changes in MVC torque over

time (especially of the hamstrings) was largely associated with indicators of peripheral fatigue.

Muscle damage biomarkers indicated that the peripheral fatigue might have been caused

predominantly by ultrastructural damage within the hamstring muscle tissue. Further, both

central and peripheral fatigue caused by repeated maximal sprints appeared to affect the

neuromuscular control of running patterns and BF_{LH} PCSA correlated inversely with the change

in hamstring MVC torque immediately after the repeated maximal sprint intervention. Our

results suggest that a high myoblast:fibroblast ratio might lead to a delayed wound closure *in*

vitro and to a delayed MVC torque recovery following repeated maximal sprints *in vivo* within

the first 48 h, indicating that stem cells of the non-contractile muscle tissue might positively

affect the response to muscle damaging exercises. However, a higher myoblast:fibroblast ratio

led to better myotube formation at seven days, and higher CK activity at ten days, after the

scratch assay, suggesting that increased myoblast numbers might be important for the latter

stage of muscle regeneration.

Fatigue and Muscle Damage Following Repeated Maximal Sprints

We showed a decreased activity of normalised BF_{LH} sEMG activity immediately after the

repeated maximal sprint intervention, but no significant changes in neuromuscular activation

using the interpolated twitch technique. The discrepancy between these two methods might

be explained by the fact that voluntary activation measured via the interpolated twitch

technique investigates all of the hamstring muscles, whilst the normalised EMG analysis was

confined solely to the BF_{LH}, which is in line with a previous study (Marshall *et al.*, 2014). The

BF_{LH} might fatigue to a greater degree immediately after repeated maximal sprint related

interventions compared to the other hamstring muscles. However, we also provide evidence

for peripheral fatigue occurring immediately after the repeated maximal sprints and a delayed

recovery at higher frequencies (30-50 Hz) after observing a right shift in the torque-frequency

relationship. This may be due to ultrastructural damage predominantly in fast-twitch (which fire

at rates from 30-50 Hz) compared to slow-twitch muscle fibres (discharge rates 10-25 Hz)

(Friden *et al.*, 1983; Jones *et al.*, 1986), leading to impaired force generation rather than simply

fatiguing the muscle fibres.

Both the quadriceps and hamstring muscle groups showed similar strength loss immediately

after the repeated maximal sprints, but the hamstring muscle group showed further strength

loss 48 h later compared to the quadriceps. Other studies did not show this additional strength

loss for the hamstring muscle group POST48. Differences in the training status of the

participants (Verma *et al.*, 2015; Chen *et al.*, 2017) and in the methodological approaches

(Balsom *et al.*, 1992) might partly explain the different outcomes. The peak BF_{LH} EMG activity

occurs at a more extended knee angle during hamstring isokinetic muscle contraction

compared to the peak EMG occurring at a more flexed knee angle for the other hamstring

muscles, such as the semitendinosus (Onishi *et al.*, 2002). Therefore, we suggest that the BF_{LH}

is the key hamstring muscle responsible for decelerating the shaft at the end of the late swing

phase. After repeated bouts of high-speed running, the semitendinosus might fatigue

prematurely (Schuermans *et al.*, 2014) and the BF_{LH} would need to substitute the impaired

function of the preceding semitendinosus to decelerate the shaft.

During high-speed running, eccentric contractions occur in the hamstrings during the late

swing phase (i.e during deceleration of the knee extension) (Chumanov *et al.*, 2007) and in the

quadriceps during the early/mid swing phase (= deceleration of hip extension) (Jönhagen *et*

al., 1996; Mendiguchia *et al.*, 2013). In general, the likelihood of sustaining a quadriceps strain

injury is significantly lower compared to a hamstring strain injury (Ekstrand *et al.*, 2011). If

repeated eccentric contractions are one of the main causes for strain injuries, as indicated by

several investigations (Higashihara *et al.*, 2010; Malone *et al.*, 2017), then the difference in

strength loss between the quadriceps and hamstrings observed in our study might be

explained by the different levels of eccentric force generated by the muscles. Muscle damage

in the quadriceps muscle presumably occurs during the deceleration phase of sprinting and

during the backswing phase, when the quadriceps muscle works eccentrically to decelerate

the leg with a flexed knee during the early swing phase of high-speed running (Howatson &

Milak, 2009; Mendiguchia *et al.*, 2013). Hamstring strain injury, however, occurs with an almost

extended leg during the late swing phase (Chumanov *et al.*, 2007). This extended lever arm

might cause higher eccentric force in the hamstring muscles compared to the shorter lever arm

with a flexed knee on the quadriceps muscle. Continuously repeated eccentric contractions

with the longer lever arm will potentially induce more muscle damage within the hamstrings in

total during e.g. a soccer match, compared to the quadriceps muscle group, which might

explain the significant different strength loss POST48 repeated sprints in the current study,

and potentially the different injury rates between these two muscle groups.

Kinematic Analysis

Our *in vivo* intervention caused changes in the running kinematics with reduced knee extension

in the late swing phase immediately after the repeated maximal sprints. Reduced hamstring

muscle strength due to neuromuscular fatigue might trigger a protective mechanism directly

after repeated maximal sprints. Afferent signals from the fatigued and damaged hamstrings

might activate the Golgi tendon organ (Byrne *et al.*, 2004), thus limiting hamstring muscle fibre

strain in an attempt to minimise further muscle damage. These kinematic changes were not

evident 48 h after the repeated maximal sprints. However, there was a non-significant tendency

for prolonged stride duration during running (P=0.08, data not shown) 48 h later and the

percentage change of knee extension in the late swing phase of running correlated with

changes in hamstring strength both measured from POST to POST48. This indicates that

participants with delayed hamstring strength recovery were still not able to fully control running.

As hamstring MVC continued to deteriorate 48 h after intermittent sprints but quadriceps MVC

started to improve, it could be that lower-limb kinematics in the sagittal plane are controlled by

the hamstrings more than the quadriceps. That outcome could also have implications for the

underlying mechanisms of knee injuries, such as an anterior cruciate ligament injury (Alentorn-

Geli *et al.*, 2009), which would need further investigation. Summarised, ultrastructural damage

in the hamstring muscles might lead to decelerated movement patterns over time, which could

increase the risk for hamstring strain injury during sprinting (Opar *et al.*, 2012).

The Role of the Extracellular Matrix on the Muscle Response Following Repeated

maximal sprinting

Recent investigations have suggested that hamstring maximum eccentric strength and BF_{LH}

fascicle length are predictors of hamstring strain injury (Timmins *et al.*, 2016a). Further, a

fatigued muscle is likely to accentuate the risk of muscle strain (Ekstrand *et al.*, 2011). However,

we could not find any correlation between BF_{LH} fascicle length and any biomarker of fatigue

but BF_{LH} PCSA correlated inversely with hamstring strength loss from PRE to POST. During

the late swing phase of sprinting, the hamstring muscles contract eccentrically to decelerate

the shaft and to enhance the subsequent concentring shortening contraction for maximal

sprinting by using stored elastic energy from the muscle-tendon unit. In comparison to other

conventional muscle-damaging interventions (Franchi & Maffiuletti, 2019), this dynamic

(stretch-shortening) movement might lead to an additional damage of the hamstring muscle

connective tissue structure. Therefore, a larger BF_{LH} PCSA might protect against immediate

hamstring MVC loss due to a greater ability to transmit the ground reaction forces laterally

(from fibre to fibre) (Turrina *et al.*, 2013), which might disperse the force more efficiently to the

tendon, while the muscle fibres themselves undergo less strain. Further, a greater BF_{LH} PCSA

reflects more fibres aligned in parallel, which would be accompanied by more muscle

connective tissue of the extracellular matrix, thus potentially protecting the muscle fibres from

excessive damage during eccentric contractions.

The stem cells of the extracellular matrix also demonstrated an important role for muscle

strength recovery in the subgroup of participants, as there was a strong inverse correlation

between myoblast:fibroblast ratio and hamstring MVC torque recovery POST48. Skeletal

muscles with a higher availability of fibroblasts around the area of myotrauma might have a

better capacity to reorganise the complex extracellular matrix, thus restoring (lateral) force

transmission, which results in a faster recovery of muscle strength after muscle damage. This

was in line with the myoblast:fibroblast ratio effect on cellular aspects of muscle regeneration

and remodelling assessed in primary muscle stem cells *in vitro*. Muscle stem cells with a low

myoblast:fibroblast ratio revealed a faster wound closure (i.e. more cells migrated to the inner

part of the artificial injury compared to the outer part), in particular 24 h after performing the

scratch assay. However, this effect was less significant 48 h after the scratch protocol, and, at

day seven, muscle primary cells with a higher myoblast:fibroblast ratio showed an improved

myotube formation, which is in line with the investigation of Fry *et al.* (2014). Together, these

results suggest that the abundance and activity of fibroblasts and myoblasts may play different

roles, depending on the time points during muscle repair. A larger abundance of fibroblasts

has a positive effect at the beginning of muscle repair, but a larger number of myoblasts is

more important for the latter stage of muscle regeneration or hypertrophy, when myoblasts

differentiate and fuse to become myotubes. We, therefore, assume that repeated maximal

sprints with insufficient recovery of previously fatigued and damage muscles (where the fatigue

and damage response is modulated by the muscle size and stem cell composition, respectively)

might augment the risk of muscle strains, as appropriate damage to the muscle connective

tissue is thought to differentiate between exercise-induced muscle damage and muscle strains

(Valle *et al.*, 2017; Balius *et al.*, 2018).

Limitations

The current study observed a relationship between human primary muscle cell type (*in vitro*)

and physiological biomarkers of skeletal muscle damage/recovery following strenuous

exercise. Further research is necessary to confirm these results with a larger sample size

regarding the *in vitro* study. However, given the scarcity of data that have examined human

muscle stem cell characteristics in association with muscle damage/recovery *in vivo*, and that

most of our results were sufficiently powered, we believe that our study represents an important

advancement in our understanding of how skeletal muscle recovers following strenuous

exercise. There was no relationship between the myoblast:fibroblast ratio and any

physiological variables regarding the quadriceps femoris, from which the muscle biopsies were

obtained. It has previously been shown that skeletal muscles of different origin, but with similar

physiological functions and fibre type composition, demonstrate similar transcriptome

expression patterns of up to 99% (Evangelidis *et al.*, 2016; Terry *et al.*, 2018). For

immnunohistochemistry analysis, the ICC (3, k) of 0.83 (95% CIs: 0.59-0.95) indicates a good

reliability for the characterisation and the quantification of myoblasts and fibroblasts. Therefore,

it is likely, that the correlation between myoblast:fibroblast ratio and the muscle damage-

response of the hamstrings but not the quadriceps muscles is explained by more severe

ultrastructural damage in the hamstrings than quadriceps. Furthermore, peripheral fatigue can

be caused by metabolic perturbations, such as the depletion of intramuscular glycogen

(Howatson & Van Someren, 2008). Therefore, because we did not control diet throughout the

study, it is possible that inter-individual differences in baseline muscle glycogen may have

influenced the ability to maintain maximal intensity throughout the sprints. However,

participants were instructed to eat and drink similar foods two hours before each laboratory

visit, and to avoid strenuous exercise for at least 48 h prior to the testing. Further, participants

were given sufficient recovery between sprint repetitions and there was a low decrement in

sprint performance, indicating that glycogen depletion was probably only a minor factor.

CONCLUSION

Repeated maximal sprints induces a greater and more prolonged strength loss in the

hamstrings compared to the quadriceps muscles. The immediate loss of hamstring function

appears to be due to both central (particularly reduced neuromuscular activation of the biceps

femoris long head) and peripheral fatigue, while prolonged hamstring strength loss is

predominantly linked to peripheral fatigue. Thigh neuromuscular fatigue following repeated

maximal sprints alters hip and knee kinematics during running immediately after the repeated

maximal sprints, but this (likely) protective effect is less evident 48 hours after the intervention,

which may lead to an increased hamstring muscle injury risk. Furthermore, biceps femoris long

head PCSA was inversely related to hamstring strength loss immediately after repeated

maximal sprinting, suggesting that the structure of the muscle, including non-contractile tissue

(e.g. the extracellular matrix) protects against neuromuscular fatigue. Skeletal muscles with an

increased number of fibroblasts might have a better capacity to reorganise the complex

extracellular matrix, which results in a faster wound closure after substantial muscle damage.

However, a larger number of myoblasts seems to be more important for the latter stage of

muscle regeneration. These novel findings improve our understanding of the physiological,

biomechanical and cellular causes and effects of exercise-induced neuromuscular fatigue, but

these results need to be confirmed in studies with a larger sample size. The practical

implications are that a 48 h recovery period following repeated maximal sprinting is insufficient,

and might increase hamstring strain injury risk. Furthermore, increasing hamstring PCSA via

resistance training is likely to reduce peripheral fatigue following repeated maximal sprinting,

thereby reducing hamstring strain injury risk.

MATERIALS AND METHODS

Participants

In vivo repeated maximal sprint intervention: Twenty recreationally active and healthy young men (*mean ± SD*; age 20.3 ± 2.87 years; height 1.79 ± 0.05 m; body mass 75.0 ± 7.89 kg) participated in the repeated maximal sprint intervention.

In vitro muscle stem cell component: Eight healthy young male (age 21.25 ± 4.27 years; height 1.77 ± 0.05 m; body mass 73.78 ± 5.68 kg) and four healthy young female (age 25.5 ± 1.29 years; height 1.67 ± 0.08 m; body mass 61.40 ± 2.57 kg) participants provided a biopsy of the vastus lateralis muscle for the *in vitro* muscle stem cell component of this study. Six of the eight males also participated in the repeated maximal sprint intervention at least three weeks after providing a muscle biopsy. Only males were recruited for the repeated maximal sprint intervention, as there is some evidence of sex differences in neuromuscular fatigue (Wüst *et al.*, 2008). However, both men and women were recruited for the muscle stem cell component due to there being no reported sex differences in stem cell properties and none within our own pilot studies (data not shown). Prior to starting the study, written informed consent was obtained from each participant and pre-biopsy screening was performed by a physician for those participants who volunteered a muscle biopsy. The study was approved by the Research Ethics Committee of Liverpool John Moores University and complied with the Declaration of Helsinki. Volunteers were physically active but were ineligible to participate if they had performed strength training of the lower limbs within 6 months prior to participation in the study, which was determined during pre-participation screening. Further exclusion criteria were: (i) any lower limb injury in the past 12 months; (ii) age under 18 or above 35 years; and (iii) more than three structured exercise sessions per week.

Experimental Design of the Repeated Maximal Sprint Intervention *in vivo*

Participants were required to visit the temperature-controlled (22-24°C) laboratory on three occasions: (i) familiarisation, (ii) testing day including the PRE and POST assessments; and

(iii) POST48 assessments after the repeated maximal sprint intervention. One week prior to

the testing day, participants were familiarised with the assessments as well as with repeated

maximal sprints (by performing 2-3 submaximal sprints) and BF_{LH} architecture of the hamstring

muscle group was assessed via ultrasound. On the test day, participants performed the

repeated maximal sprint intervention of 15 x 30 m sprints to induce neuromuscular

fatigue/damage in both the quadriceps femoris and hamstring muscle groups. All tests were

performed at the same time of the day for each participant. Further, participants were instructed

to maintain their normal routine (including eating habits), to refrain from drinking alcohol and

to avoid any strenuous exercise 48 h prior to testing and throughout the study, and to refrain

from consuming caffeine on testing days. Nothing was consumed throughout the testing

sessions except water, which was available ad libitum.

The test battery was always performed in the same order with the right leg of each participant

and comprised (i) venous blood sampling [for analysing serum interleukin-6 (IL-6)

concentration and creatine kinase (CK) activity]; (ii) hamstrings and quadriceps muscle

soreness via visual analogue scale; (iii) isometric maximum voluntary contraction (MVC)

torque of the quadriceps, as well as both voluntary and involuntary muscle activation and

torque-frequency relationship via electrical stimulation] MVC torque of the hamstring together

with normalised BF_{LH} sEMG (see below); and (iv) treadmill running (4.17 m s^{-1}) kinematics of

the right leg (via an eight-camera motion capture system) PRE and POST48 following the

repeated maximal sprint intervention. At POST, kinematic assessments were performed first

followed by the aforementioned order of the assessment for practical reasons.

Maximal Repeated Sprint Protocol

496 Many athletes in team sports are required to perform repeated maximal sprints, which are

497 characterised by short-duration (<10 s) and relatively longer recovery times (>60 s) between

498 maximal sprint bouts, and have a different physiological demand compared to repeated-sprint

499 exercises with shorter recovery times (<60 s) (Bradley *et al.*, 2010; Girard *et al.*, 2011).

500 Therefore, the IS intervention consisted of 15 repetitions of 30 m maximal sprints with a

deceleration zone of 12 m. The 30 m distance was chosen as the upper average of both the total sprinting distance (346 ± 115 m) of wide-midfielders and the mean recovery time (70.2 ± 25.1 s) between sprint bouts in soccer (Bradley *et al.*, 2009; Bradley *et al.*, 2010), which allows the athlete to maintain the performance of the sprint bouts. Similar protocols have been used elsewhere (Timmins *et al.*, 2014; Verma *et al.*, 2015; Chen *et al.*, 2017). Prior to the repeated maximal sprint intervention, a five-minute warm-up was performed, comprising jogging, dynamic stretching and three self-paced 20 m runs at 60%, 80%, 100% of perceived top speed. During the repeated maximal sprint intervention, the participants were instructed to sprint maximally (verbal encouragement) and to stop within the deceleration zone. Further, they were instructed to move slowly back to the start line and to sit on a chair for the remaining time until the next sprint. The recovery comprised 90 s between repetitions and after every 5th repetition, the participants were allowed to rest for 3 min. Sprinting time during trials was measured and controlled with timing gates (Brower Timing Systems, Draper, UT, USA), which were placed on the start and finish line. Participants started 30 cm before the start line to avoid interfering with timing gates with the arms upon initial acceleration (Howatson & Milak, 2009). Further, heart rate (Polar Oy, Kempele, Finland) and rating of perceived exertion (Borg, 1982) were recorded before and after each repetition. Participants were instructed to wear the same footwear for each testing day. As the fastest sprint time does not necessarily need to occur at the first of the 15 sprint bouts and there was an upsurge in speed of the final sprint, fatigue was assessed with the performance decrement score using the following formula (Glaister *et al.*, 2008):

$$Fatigue = \big(100 \; X \; (total \; sprint \; time \; \div \; ideal \; sprint \; time)\big) - 100$$

Where total sprint time = sum of time from all 15 sprints; and ideal sprint time = total number of sprints (15) x fastest repetition sprint time. The calculation of this decrement score was also used to quantify changes in heart rate and rating of perceived exertion during the repeated maximal sprint intervention.

Maximal Voluntary Contraction (MVC)

Three test sessions were conducted with an isokinetic dynamometer (Humac Norm, CSMI Solutions, Massachusetts, USA). As *isokinetic* maximum voluntary contractions (MVC) torque tests are only weak predictors of hamstring strain injury (van Dyk *et al.*, 2016), we decided to focus on *isometric* MVC quadriceps and hamstring torque at optimal knee strength angles (optimal torque-joint angle relationship, see below) to avoid further fatiguing the participants. The torque signal was interfaced with an acquisition system (AcqKnowledge, Biopac Systems, Santa Barbara, USA) for analogue-to-digital conversion and sampled at a frequency of 2 kHz. The participant was seated in an upright position and securely fastened with inextensible straps at the chest and waist while the arms were held crossed above the chest. The tibiofemoral epicondyle was aligned with the lever arm rotation axis, and the lever arm shin pad was strapped to the leg, 2 cm above the centre of the lateral malleolus. A Velcro strap secured the distal thigh just above the knee. The hip joint angle was set to 85° (180° = supine position) in order to analyse the knee flexor muscle group at a sprint specific angle associated with the late swing phase of sprinting (Guex *et al.*, 2012). Participants were instructed to maximally extend and flex their leg to measure knee range of motion. Quadriceps MVC was measured at 80° knee flexion (0° = full knee extension), as this is the optimal joint angle for peak quadriceps MVC in healthy young men (Erskine *et al.*, 2009). Hamstring MVC was measured at 30° knee flexion based on this being the optimal joint angle for peak hamstring MVC during our pilot work. Published studies during the time of data collection used a similar angle of hamstring MVC torque (Nedelec *et al.*, 2014; Kirk & Rice, 2016). This was also in line with sprinting kinematics, demonstrating that maximal hamstring muscle lengths during sprinting occur during the late swing phase when the knee is flexed between 30° and 45° (Thelen *et al.*, 2005). Prior to isometric MVC assessments, participants underwent a standardised warm up consisting of 10 submaximal isokinetic leg extensions ($60°·s^{-1}$). Participants then performed three isometric knee extension (quadriceps) and flexion (hamstring) at both joint angles (each MVC lasting 2-3 s), with 60 s rest between MVC of a given muscle group. The highest MVC of

the three attempts for each muscle group at each angle was used for subsequent analyses. Throughout the tests, participants received verbal encouragement and biofeedback (MVC outputs) were projected onto a screen in front of the participant.

Hamstring Muscle Voluntary Activation

To measure hamstring muscle voluntary activation capacity via the interpolated twitch technique, stimulation electrodes (12.5 mm x 7.5 mm self-adhesive electrodes (DJO Global, California, USA) were used. The general procedure has been described elsewhere (Erskine *et al.*, 2009; Erskine *et al.*, 2010; Marshall *et al.*, 2014). Briefly, the anode was placed proximal to the popliteal fossa, and the cathode was placed beneath the gluteal fold and slightly medial to avoid activation of the vastus lateralis. Protocols were completed with electrical stimulation pads carefully taped down during the sprinting protocol and were additionally marked on the skin with a permanent marker, to ensure a precise relocation for the POST and POST48 tests. Stimulation was delivered by a high-voltage stimulator (DS7AH; Digitimer Ltd., Welwyn Garden City, United Kingdom), and consisted of a doublet using two 240-V rectangular pulses (200 μs pulse width) with an inter-pulse duration of 10 ms (100 Hz stimulation). During each experimental session, relaxed hamstring muscles were stimulated while participants were fixed in the isokinetic dynamometer with the same setting for knee flexion MVC (85º hip angle, 30 º knee flexion). The amplitude started with 50 mA to familiarise the participants to the stimulation and was gradually increased in 20 mA increments until a plateau in doublet torque was achieved. We decided to use the individual maximal stimulation (100%) intensity despite the fact that other publications used supramaximal stimulation (110-130%) (Marshall *et al.*, 2014) as we experienced lower MVC knee flexion torque output beyond 100%. That individual amplitude (162.0 ± 17.4 mA; range: 130–200 mA) was applied during all maximal contractions in the experimental session.

The maximal doublet stimulation was used two minutes later to elicit resting maximal doublet torque in the resting state (control doublet), followed 2.5 s later by a second (superimposed) doublet during an isometric knee flexion MVC. The superimposed doublet torque was always

calculated manually from careful selection and inspection of the respective time periods compared to a normal increase in voluntary torque. Voluntary activation was calculated according to the following equation:

$$VA\ (\%) = [1- (superimposed\ doublet\ torque/\ control\ doublet\ torque)]$$

Surface Electromyography and Antagonist Muscle Co-activation

Surface electromyographic (sEMG) activity was recorded from the vastus lateralis and BF_{LH} to determine the extent of antagonist muscle co-activation during MVCs of the respective muscle group. Previous reports have shown that the vastus lateralis (Reeves *et al.*, 2004) and BF_{LH} (Kellis & Baltzopoulos, 1999) are representative muscles for the quadriceps femoris and hamstring muscle group, respectively. This procedure has been reported in detail elsewhere (Reeves *et al.*, 2004). Briefly, once the muscles were identified via palpation, and the skin surface was shaved and cleaned with 70% ethanol, two bipolar Ag-AgCl surface electrodes with an inter-electrode distance of 2 cm (Noraxon duel sEMG electrode, Noraxon, Scottsdale, USA) were placed along the sagittal axis over the muscle belly at 33% of the respective muscle length from the distal end [according to SENIAM guidelines (Hermens *et al.*, 2000)] and one reference electrode (Ambu Blue, Ambu, Copenhagen, Denmark) was positioned over the medial tibial condyle. The exact location of the electrodes were marked on the participant's skin with a permanent marker to ensure precise electrode repositioning for the following assessments.

Surface EMG signals were sampled at 2000 Hz (Biopac Systems, Santa Barbara, USA) and then band-pass filtered between 10–500 Hz (AcqKnowledge, Biopac Systems, Santa Barbara, USA). Surface EMG activity of both the agonist and antagonist muscles were analysed by calculating the root mean square of the sEMG signal of a 500-ms epoch around peak MVC. To compare BF_{LH} sEMG activity at all three time points, BF_{LH} sEMG of the hamstring MVC at 30° was normalised to the evoked maximum compound muscle action potential (M-wave) of the BF_{LH} (see below). Antagonist muscle co-activation (i.e. quadriceps activation during

hamstring MVC at 30° knee flexion, or hamstring activation during quadriceps MVC at 80°

knee flexion) was calculated with the following formula (where EMG_{max} is the maximum sEMG

of the antagonist muscle when acting as an agonist at the same knee joint angle):

$$Antagonist\ muscle\ co-activation = \frac{EMGantagonist}{EMGmax} \times 100$$

Torque signals, electrical stimuli, and sEMG activity were displayed on a computer screen,

interfaced with an acquisition system (AcqKnowledge, Biopac Systems, Santa Barbara, USA)

used for analogue-to-digital conversion. Due to technical issues, co-activation data were

available for hamstring n = 12; and quadriceps n=10.

Hamstring Muscle Maximal Compound Muscle Action Potential

The hamstring muscle group was stimulated with single square wave twitch pulses (200 µs

duration) using the same electrical stimulator and stimulating electrodes, as described above.

While the participant sat resting on the isokinetic dynamometer with the knee angle set at 30°

knee flexion, compound muscle action potentials (*M*-waves) were evoked with 10 to 20 mA

incremental amplitudes until a maximal *M*-wave (M_{max}) was achieved. The average amplitude

necessary to evoke a maximal *M*-wave was 166.8 ± 19.8 mA; range: 130–210 mA). The

maximal *M*-wave was defined as the mean peak-to-peak sEMG response from the three

highest observed *M*-waves. Due to inter-individual differences in subcutaneous fat and the

(re)location of small sEMG electrodes over a relatively large muscle belly, sEMG amplitude is

notoriously variable (Araujo *et al.*, 2000). To reduce this inter- and intra-individual variability,

we normalised absolute BF_{LH} sEMG to the individual's BF_{LH} maximal *M*-wave, determined at

each testing session (Lanza *et al.*, 2018). Due to technical issues, data of BF_{LH} sEMG

normalised to maximal *M*-wave was only available for n = 13.

Torque-frequency Relationship

The torque-frequency relationship was determined by stimulating the hamstring muscle group

with single square wave twitch pulses (200 µs duration) at 1, 10, 15, 20, 30, 50 and 100 Hz for

1 s each in a random order and with 15 s rest between each stimulation, using the same

electrical stimulator and stimulating electrodes (and location), as described above. The

stimulus intensity for 100-Hz stimulation was the amplitude necessary to elicit ~20% knee

flexion MVC torque at PRE, and the same amplitude was used for the same test at POST and

48POST. The absolute peak torque at each frequency was normalised to the peak torque at

100 Hz for each time point (PRE, POST and POST48).

Delayed Onset Muscle Soreness

Using a visual analogue scale that consisted of a 100 mm line (scale 0-10 cm; 0 cm=no

soreness; 10 cm= unbearably painful), in conjunction with both a three-repetition bilateral squat

(predominantly to determine quadriceps femoris muscle soreness) (Scott & Huskisson, 1979)

and lunges (predominantly to determine hamstring muscle soreness), participants rated their

perceived lower limb muscle soreness along the muscle length immediately after each

movement. Muscle soreness was also measured by recording the force required to elicit

tenderness at nine fixed sites on the skin over the quadriceps (distal, central and proximal

locations of the three superficial quadriceps heads, vastus lateralis, vastus medialis and rectus

femoris) and six sites on the hamstrings (distal, central and proximal locations of both BF_{LH}

and the medial hamstrings), which were previously marked with a permanent marker to ensure

the same measuring position PRE, POST and POST48. At each site, a gradually increasing

force was applied by the investigator with an algometer (FPK/FPN Mechanical Algometer,

Wagner Instruments, Greenwich, USA) with a maximum of 10 kg/cm^2. Lying in the prone

position with the hip and knee fully extended and muscles relaxed, the participant was asked

to indicate when the sensation of pressure changed to discomfort, and the force at that point

was recorded (Newham *et al.*, 1987).

Ultrasound

Architectural parameters of the BF_{LH} were assessed using B-mode ultrasound imaging.

Participants were in the prone position with the hip and knee fully extended and muscles

relaxed. The BF_{LH} was investigated, as this muscle is the most commonly injured hamstring

muscle in team sports, particularly during sprinting (Ekstrand *et al.*, 2011). Longitudinal and

cross-sectional panoramic ultrasound images of the right BF_{LH} were obtained (Philips EPIQ 7

Ultrasound System, Bothel, USA). The linear transducer (5-18 MHz; aperture 38.9 mm) was

carefully placed on the skin with transmission gel and BF_{LH} was scanned (i) longitudinally from

its distal (=0% muscle length) to proximal (=100% muscle length) myotendinous junction along

a line drawn with a permanent marker to mark the central pathway between the medial and

lateral aspects of the muscle (incorporating the intra-muscular aponeurosis (Evangelidis *et al.*,

2014) (Figure 2A); and (ii) cross-sectionally at 20, 40, 60 and 80% along the total muscle length,

measured on the skin using a tape measure (Seca, Hamburg, Germany) (Figure 2B).

All images were analysed offline (ImageJ, version 1.51s, National Institutes of Health,

Bethesda, USA). Two images for each of the four cross-sectional points were recorded and

the image of best quality was used to calculate BF_{LH} muscle volume. The volume of the

muscular portion between every two consecutive scans was calculated with the following

equation:

673
$$Volume = \frac{1}{3} * d * \left(a + \sqrt{(ab) + b} \right)$$

Where *a* and *b* are the anatomical cross-sectional areas of the muscle of two consecutive

cross-sectional scans and *d* is the interval distance between the cross-sectional area

measurements. The volume of the entire muscle was calculated by summing up all of the inter-

scan muscular volumes (Erskine *et al.*, 2016). Two full-length sagittal images were then

recorded to allow for the measurement of resting BF_{LH} muscle fascicle length and pennation

angle, which were both assessed in 3 fascicles at 50% of the total length of BF_{LH}. This point

was measured offline (ImageJ). A comparison between offline (sagittal ultrasound) and tape

measurements of the total BF_{LH} length revealed a very high correlation (R^2=0.958, P<0.001).

Fascicle length was measured by tracing the fascicular path from the upper aponeurosis to the

intra-muscular aponeurosis of the BF_{LH}. Muscle fascicle pennation angle was determined as

the angle between the muscle fascicular paths and their insertion into the intra-muscular

aponeurosis. The mean of the three measurements for each parameter were used to

determine both fascicle length and pennation angle of the BF_{LH} muscle. PCSA was calculated

by dividing BF_{LH} volume by its fascicle length. During the time of data collection, a similar

methodological approach was published elsewhere (Seymore *et al.*, 2017). One longitudinal

image of one participant in the present study was not analysed due to low image quality.

Ultrasound scans and image analysis was performed by the same investigator.

Kinematic and Kinetic Data

Three-dimensional kinematic and kinetic data were synchronously collected at 500 Hz using

an eight-camera motion analysis system (Qqus 300+; Qualisys, Gothenburg, Sweden),

together with a ground-embedded force plate (90 x 60 cm, 9287B; Kistler Holding, Winterthur,

Switzerland) at 1,500 Hz. The data were filtered with a digital dual low-pass Butterworth filter

at 20 Hz for motion and 60 Hz for force, as previously described (Verheul *et al.*, 2017).

Retroreflective markers (12 mm diameter) were placed on anatomical landmarks on the right

leg and pelvis, as previously described (McClay & Manal, 1999). One standing static and two

functional motion calibration trials were recorded of the participant PRE, POST and POST48.

For the static trial, participants stood with their feet approximately shoulder width apart and

knees fully extended. This static trial determined local coordinate systems, the location of joint

centres, and the foot, shank, thigh, and pelvis segment lengths of each participant. The

functional trials defined functional hip joint centres (Schwartz & Rozumalski, 2005) and knee

joint axes (Robinson & Vanrenterghem, 2012). Kinematic data were tracked using Qualisys

Track Manager Software (Qualisys). Data processing and analysis were undertaken in

Visual3D (C-Motion, Germantown, MD). To examine any changes between the time points,

joint angles were normalised relative to the static trial of the accompanying time point for

minimising the influence of potential slightly different marker positions between the trials.

Lower extremity 3D joint angles and angular velocities were calculated using an X-Y-Z Cardan

angle rotation sequence. Investigated variables included peak knee and hip angles, as well as

range of motion and time during stance and swing phase of the treadmill run, for all three

planes, were calculated as described in previous studies (Verheul *et al.*, 2017).

Motorised Treadmill Run

Participants ran on a motorised treadmill (HP Cosmos Pulsar; Nussdorf, Germany) for 30 s at

4.17 m s^{-1} (0° incline), as high-speed running was of interest. The selected speed was based

on pilot testing, which demonstrated that 15 km/h (4.17 m/s) was the fastest speed on a

motorized treadmill where the participants still felt comfortable. Motion analysis data were

recorded for the last 10 s of the run and data were analysed for at least 6 consecutive strides.

Peak knee and hip angle data, for all three planes, were calculated (i) between the initial

contact and terminal stance of foot; and (ii) between initial and terminal swing phase. The

touchdown of the foot during the treadmill run was determined from the kinematic data as

occurring at the local minima and the toe-off during running as the local maxima of the vertical

velocity of the head of the fifth metatarsal marker on the foot (Maiwald *et al.*, 2009).

Blood Samples

A 10 mL blood sample was drawn from an antecubital vein in the forearm and collected into a

serum vacutainer (BD Vacutainer systems, Plymouth, UK). The blood samples were obtained

at each time point and left at 22-24°C for 30 min to allow clotting, and then kept on ice when

necessary. Serum samples were centrifuged at 1,300 g for 15 min at 4°C. All samples were

then aliquoted into 1.5 mL microcentrifuge tubes [Axygen (Corning), New York, USA] and

stored at -80°C until subsequent analysis (see below).

Serum Interleukin-6 (IL-6) Concentration

Serum samples were assayed for IL-6 concentration using commercially available human IL-

6 enzyme linked immunosorbent assay kits (Quantikine®, R&D systems, Minneapolis, MN,

USA) according to the manufacturer's instructions. The intensity of the colour produced after

20 min was measured with a Thermo Multiskan Spectrum microplate reader (Thermo Fisher

Scientific. Waltham, MA. USA) at 450 nm and values were calculated with Excel 365 (Microsoft, v. 365, USA) by generating a four-parameter logistic curve fit. The minimum detectable dose of human IL-6 was 0.70 pg/mL.

Serum Creatine Kinase Activity

Creatine kinase (CK) activity was assayed using a commercially available CK assay (Catachem Inc., Connecticut, NE, USA), as described in detail elsewhere (Owens *et al.*, 2015). Briefly, 10 μL blood serum were loaded onto a 96-well UV plate. The CK reaction reagent and diluent (Catachem) were prepared as per the manufacturer's instructions and added to the samples and the change in absorbance monitored continuously over 20 min in a Thermo Multiskan Spectrum plate reader (Thermo Fisher Scientific. Waltham, MA. USA) at a wavelength of 340 nm.

Capillary Blood Lactate Concentration

Capillary blood samples were taken from the finger-tip via a Safety-Lancet Extra 18G needle (Sarstedt; Nümbrecht, Germany) at rest before and immediately after the repeated maximal sprint intervention. Blood samples were analysed within 60 seconds of collection using a portable blood lactate analyser (Arkray Lactate Pro; Kyoto, Japan).

Reagents, Chemicals, and Solvents for Muscle Cell Culture *in vitro*

Growth media used for the expansion of human muscle-derived cell populations consisted of Hams F-10 nutrient mix (Lonza, Basel, Switzerland) with added L-glutamine (2.5 mM), 10% heat-inactivated fetal bovine serum (hiFBS; Gibco, Thermo Fisher Scientific, Altincham, UK), 1% penicillin-streptomycin (Life Technologies, Warrington, UK), and 1% L-Glutamine (Gibco). Differentiation media consisted of α-MEM (Lonza), 1% hiFBS, 1% penicillin-streptomycin, and 1% L-glutamine. Phosphate-buffered saline (PBS; Sigma-Aldrich) was used to wash cell monolayers. Desmin polyclonal rabbit anti-human antibody (Cat# ab15200, RRID: AB_301744) was used (1:200) from Abcam (Abcam, Cambridge, UK), and secondary antibody (TRITC

761 polyclonal goat anti-rabbit; Cat# A16101, RRID: AB_2534775) was used (1:200) from Fisher

762 Scientific.

Muscle Biopsy Procedure

764 Participants were instructed to avoid any strenuous exercise 48 h prior to the biopsy procedure.

765 Biopsies from the vastus lateralis muscle were obtained under local anaesthesia from each

766 participant, using the Weil-Blakesley conchotome technique as described previously

767 (Baczynska *et al.*, 2016). The conchotome was inserted through the incision into the muscle

768 belly to obtain the 134 ± 82.7 mg muscle biopsy.

Extraction of Human Muscle-Derived Cells

The muscle biopsies analysed in this study were isolated (Blau & Webster, 1981; Crown *et al.*,

2000) and cultured (Owens *et al.*, 2015), as reported previously. Briefly, biopsy samples were

transferred with precooled growth media from the muscle biopsy suite to the sterile tissue

culture hood (Kojair Biowizard Silverline class II hood; Kojair, Vippula, Finland) within 40 min

and muscle biopsy samples were washed three times with ice-cold PBS (0.01 M phosphate

buffer, 0.0027 M KCl, and 0.137 M NaCl, pH 7.4, in dH2O). Visible fibrous and fat tissue was

removed using sterile scissor and forceps. Samples were cut in small pieces (1 mm^3) and

digested in 5 ml of trypsin-EDTA for 15 min on a magnetic stirring platform at 37°C to dissociate

muscle cells. The trypsinisation process was repeated twice. Supernatant derived following

each treatment was collected and pooled with hiFBS at a concentration of 10% of the total

volume to inhibit further protease activity. Cell supernatant was centrifuged at 450 g for 5 min.

Supernatant was discarded and the cell pellet was resuspended in growth media and plated

on a T25 cm^2 culture flask (Corning, Life Sciences, New York, USA) for cell population

expansion. Culture flasks were previously coated with a 2 mg/l porcine gelatin solution (90–

110 g, Bloom; Sigma-Aldrich, Dorset, UK) to support cell adhesion.

Expansion of Extracted Cells

The medium was refreshed on the fourth day after the extraction procedure and subsequently

every 48 h following two brief washes with PBS. Cells were incubated in a HERAcell 150i CO_2

Incubator (Thermo Scientific, Cheshire, UK). T25 cm^2 culture flasks reached 80% confluence

after aproximately 10 days and were passaged via trypsinisation. Cells were counted using

Trypan Blue exclusion and re-plated on gelatinised T75 cm^2 culture flasks (Nunc, Roskilde,

Denmark). The cells were expanded until passage 3 and then frozen in GM with 10% dimethyl

sulfoxide (DMSO) in liquid N_2 as a cryopreservant. All experiments were performed on cells

between passages 3 and 6 to avoid potential issues of senescence (Foulstone *et al.*, 2004).

Characterization of Human Muscle-Derived Cells

The mixed population of human skeletal muscle-derived myoblasts and fibroblasts were

characterised by immunofluorescent staining for the detection of desmin expressed by

myoblasts (desmin positive) and non-myoblasts (desmin negative) to determine the

percentage of myoblasts and fibroblasts. Previous investigations have determined that the

non-myoblast (desmin negative) fraction is highly enriched in fibroblasts, with up to 99% of this

fraction being fibroblasts (Stewart *et al.*, 2003; Agley *et al.*, 2013), as our group also observed

(Owens *et al.*, 2015). Therefore, non-myoblasts (desmin negative) were referred to here as

fibroblasts.

Grohmann *et al.* (2005) showed that passaging does not change the percentage of myoblast

and fibroblasts and all populations were included for analysis. Monolayers were incubated with

25% [vol/vol methanol in Tris-buffered saline (10 mmTris-HCl, pH 7.8, 150 mm NaCl)], 50%

and 100% for 5-min to fix the cells and stored at 4°C wet in Tris-buffered saline until further

analysis. Fixed monolayers were permeabilised and blocked for 2 h with 5% goat serum and

0.2% Triton X-100 in Tris-buffered saline, prior to staining. Cells were incubated overnight at

4°C with anti-Desmin antibody (1:200). After overnight incubation, the primary antibody was

removed, and the cells were washed three times with Tris-buffered saline. Secondary TRITC

polyclonal goat anti-rabbit antibody (1:200) was then applied and incubated for 2 h at 4°C.

Fluorescent images were captured using live imaging microscopy (Leica DMB 6000;

Magnification x 10.5) and analysed via ImageJ cell counter plug-in. A total of four randomly

selected areas per well were analysed per individual.

Wound-Healing Assay, Migration and Differentiation Analysis

For the wound healing assay, 100,000 cells/ml were seeded in gelatinised six-well plates (Nunc,

Roskilde, Denmark). Cells were expanded as described above until cell monolayers reached

a confluent state, Growth media was removed, monolayers were washed with PBS and cells

were damaged by a vertical scrape with a 1-ml pipette tip (width of the wound area, *mean ±*

S.E.M.: 896.4 ± 21.24 *μ*m), as previously reported by our group (Owens *et al.*, 2015; Brown *et*

al., 2017). PBS was aspirated, damaged cell monolayers were washed twice with PBS to

remove cell debris and 2 ml differentiation media was added. Monolayers were imaged with a

live imaging microscopy (Leica) for the analysis of cell migration immediately, 24h, 48h and

seven days after the wound healing assay. Additional 500 μl differentiation media was added

in each well at day 4. TIF images were exported from Leica Application Suite and loaded as

TIF image stacks in ImageJ with a cell counter plug-in. Cells in the outer and inner segments

were then counted (Figure 3A).

Damaged monolayers were imaged at two sites per well in the wound site immediately post-

damage (0 h). These image coordinates were tracked and stored to allow subsequent

monitoring of the same sites on the wound to reduce this experimental bias. Myotube formation

was captured on day 7. Captured images were exported as TIF image files, and analysed in

ImageJ. Muscle cell fusion/differentiation was assessed as the number of myotubes per field

of view and myotube hypertrophy was determined by measuring myotube length, myotube

diameter (the average of three diameters along the length of the myotube), myotube area

(determined by manually drawing a line around the sarcolemma of each myotube) and total

myotube number via ImageJ cell counter plug-in. By normalising the pixel scale to the micron

scale of each image, a value expressed as µm^2 was obtained for myotube area. A total of two

images per well were analysed.

Creatine Kinase (CK) Activity

At 0 and 10 days following the mechanical scrape insult, CK activity was analysed as a marker

of muscle cell differentiation/fusion into myotubes. Cell monolayers were first lysed with 300

µl/well of 50 M Tris-mes and 1% Triton-X 100, pH 7.8 (TMT). Ten microliters of TMT cell lysate

was loaded in duplicate wells on a 96-well UV plate and used for quantification of CK activity.

The CK reaction reagent and diluent (Catachem, Bridgeport, CT, USA) were prepared as per

the manufacturer's instructions as previously described (please see subsection Serum

Creatine Kinase).

Statistical Analysis

One-way repeated-measures analysis of variances (ANOVA)s were performed to determine

whether there was a significant main effect for time (within subject factor) for the following

dependent variables: MVC torque, voluntary muscle activation, antagonist muscle co-

activation, muscle soreness (for squat lunge via measured via visual analogue scale as well

as algometer), rating of perceived exertion, CK activity, IL-6 concentration, and for kinematics

data (hip and knee angle parameters). MVC torque data were analysed for interactions and

main effects for muscle group and time using two-way mixed design ANOVAs, comparing

differences between muscle groups across 3-time points; PRE, POST, and POST48. For

within test comparisons, either, independent t-tests, or one-way ANOVAs were used where

appropriate. For the torque-frequency relationship, normalised torque at each frequency was

analysed using a two-way repeated measures ANOVA, with stimulation frequency (1-100 Hz)

and time (PRE, POST and POST48) as the within-groups independent variables. Post-hoc

one-way repeated measures ANOVAs were used to determine if the normalised torque at each

frequency differed between time points. Bivariate correlations were used to analyse the relation

between architectural parameters of the BF$_{LH}$ (volume, fascicle length, fascicle pennation angle

and PCSA) and fatigue biomarkers (relative MVC loss normalised to PRE MVC), serum CK

activity, serum IL-6 concentration, muscle soreness, knee joint range of motion or changes in

range of motion during treadmill running.

Bivariate correlations were used to analyse the relation between myoblast:fibroblast ratio and

quadriceps and hamstring MVC, migration dynamics (total cell migration, cell proportion of

inner to outer segment) and myotube formation (total myotube number, myotube length,

average diameter, myotube area and CK activity) of the muscle stem cells. Standard guidelines

concerning violation of the sphericity assumption to adjust the degree of freedom of the F-test

by the Huynh-Felt epsilon if epsilon is greater than 0.75 and to use the more stringent

Greenhouse-Geisser adjustment if epsilon is less than 0.75 were followed. Results were

expressed as mean ± SD, unless otherwise stated, with statistical significance set at $P<0.05$.

All MVC data were analysed with AcqKnowledge software 4.4 (Biopac-Systems Inc., Goleta,

USA) and SPSS 23 Software (IBM Inc., Armonk, NY: IBM Corp) was used for statistical

analysis. Occasional missing data are reflected in the reported degrees of freedom.

GRANTS

878 This study was supported by a Liverpool John Moores University PhD studentship (P.B.) and

879 the Wellcome Trust Biomedical Vacation Scholarships (R.M.E., M.S.).

DISCLOSURES

881 No conflicts of interest, financial or otherwise, are declared by the authors.

AUTHOR CONTRIBUTION

883 R.M.E, P.B., B.D., C.E.S., and M.L. conceived and designed the research; P.B., S.T., M.S.,
884 M.C., J.A.S., and S.O.S performed the experiments; P.B., S.T. and R.M.E analyzed the data;
885 R.M.E., P.B, M.L. and C.E.S. interpreted the results of the experiments; P.B. prepared the
886 figures; P.B. drafted the manuscript; R.M.E., P.B., S.T., M.S., M.L., B.D., C.E.S., M.C., J.A.S.,
887 and S.O.S edited and approved the final version of the manuscript.

REFERENCES

Agley CC, Rowlerson AM, Velloso CP, Lazarus NR & Harridge SD. (2013). Human skeletal muscle fibroblasts, but not myogenic cells, readily undergo adipogenic differentiation. *Journal of cell science*, jcs. 132563.

Alentorn-Geli E, Myer GD, Silvers HJ, Samitier G, Romero D, Lázaro-Haro C & Cugat R. (2009). Prevention of non-contact anterior cruciate ligament injuries in soccer players. Part 1: Mechanisms of injury and underlying risk factors. *Knee Surgery, Sports Traumatology, Arthroscopy* **17,** 705-729.

Araujo RC, Duarte M & Amadio AC. (2000). On the inter-and intra-subject variability of the electromyographic signal in isometric contractions. *Electromyogr Clin Neurophysiol* **40,** 225-230.

Baczynska AM, Shaw S, Roberts HC, Cooper C, Sayer AA & Patel HP. (2016). Human Vastus Lateralis Skeletal Muscle Biopsy Using the Weil-Blakesley Conchotome. *Journal of Visualized Experiments*, e53075-e53075.

Balius R, Alomar X, Pedret C, Blasi M, Rodas G, Pruna R, Peña-Amaro J & Fernández-Jaén T. (2018). Role of the Extracellular Matrix in Muscle Injuries: Histoarchitectural Considerations for Muscle Injuries. *Orthopaedic Journal of Sports Medicine* **6,** 2325967118795863.

Balsom P, Seger J, Sjödin B & Ekblom B. (1992). Maximal-intensity intermittent exercise: effect of recovery duration. *Int J Sports Med* **13,** 528-528.

Blau HM & Webster C. (1981). Isolation and characterization of human muscle cells. *Proc Natl Acad Sci* **78,** 5623-5627.

Borg GA. (1982). Psychophysical bases of perceived exertion. *Medicine and science in sports and exercise* **14,** 377-381.

Bradley PS, Di Mascio M, Peart D, Olsen P & Sheldon B. (2010). High-intensity activity profiles of elite soccer players at different performance levels. *J Strength Cond* **24,** 2343-2351.

Bradley PS, Sheldon W, Wooster B, Olsen P, Boanas P & Krustrup P. (2009). High-intensity running in English FA Premier League soccer matches. *Journal of sports sciences* **27,** 159-168.

Brown AD, Close GL, Sharples AP & Stewart CE. (2017). Murine myoblast migration: influence of replicative ageing and nutrition. *Biogerontology* **18,** 947-964.

Byrne C, Twist C & Eston R. (2004). Neuromuscular function after exercise-induced muscle damage. *Sports Med* **34,** 49-69.

Chen C-H, Ye X, Wang Y-T, Chen Y-S & Tseng W-C. (2017). Differential Effects of Different Warm-up Protocols on Repeated Sprints-Induced Muscle Damage. *J Strength Cond*.

Chumanov ES, Heiderscheit BC & Thelen DG. (2007). The effect of speed and influence of individual muscles on hamstring mechanics during the swing phase of sprinting. *Journal of biomechanics* **40,** 3555-3562.

939 Chumanov ES, Schache AG, Heiderscheit BC & Thelen DG. (2012). Hamstrings are most susceptible to
940 injury during the late swing phase of sprinting. *Br J Sports Med* **46,** 90.

941
942 Collins CA, Olsen I, Zammit PS, Heslop L, Petrie A, Partridge TA & Morgan JE. (2005). Stem cell
943 function, self-renewal, and behavioral heterogeneity of cells from the adult muscle satellite
944 cell niche. *Cell* **122,** 289-301.

945
946 Crema MD, Jarraya M, Engebretsen L, Roemer FW, Hayashi D, Domingues R, Skaf AY & Guermazi A.
947 (2017). Imaging-detected acute muscle injuries in athletes participating in the Rio de Janeiro
948 2016 Summer Olympic Games. *Br J Sports Med***,** bjsports-2017-098247.

949
950 Crown A, He X, Holly J, Lightman S & Stewart C. (2000). Characterisation of the IGF system in a
951 primary adult human skeletal muscle cell model, and comparison of the effects of insulin and
952 IGF-I on protein metabolism. *J Endocrinol* **167,** 403-415.

953
954 Ekstrand J, Hägglund M & Waldén M. (2011). Epidemiology of muscle injuries in professional football
955 (soccer). *The American journal of sports medicine* **39,** 1226-1232.

956
957 Erskine R, Tomlinson D, Morse C, Winwood K, Hampson P, Lord J & Onambélé G. (2016). The
958 individual and combined effects of obesity-and ageing-induced systemic inflammation on
959 human skeletal muscle properties. *Int J Obes (Lond)*.

960
961 Erskine RM, Jones DA, Maganaris CN & Degens H. (2009). In vivo specific tension of the human
962 quadriceps femoris muscle. *Eur J Appl Physiol* **106,** 827-838.

963
964 Erskine RM, Jones DA, Williams AG, Stewart CE & Degens H. (2010). Resistance training increases in
965 vivo quadriceps femoris muscle specific tension in young men. *Acta physiologica* **199,** 83-89.

966
967 Evangelidis P, Massey GJ, Ferguson RA, Wheeler PC, Pain MT & Folland JP. (2016). The functional
968 significance of hamstrings composition: is it really a 'fast'muscle group? *Scandinavian journal
969 of medicine & science in sports* **27,** 1181-1189.

970
971 Evangelidis PE, Massey GJ, Pain M & Folland JP. (2014). Biceps Femoris Aponeurosis Size: A Potential
972 Risk Factor for Strain Injury? *Medicine and science in sports and exercise*.

973
974 Foulstone EJ, Huser C, Crown AL, Holly JM & Stewart CE. (2004). Differential signalling mechanisms
975 predisposing primary human skeletal muscle cells to altered proliferation and differentiation:
976 roles of IGF-I and TNFα. *Experimental cell research* **294,** 223-235.

977
978 Franchi MV & Maffiuletti NA. (2019). Distinct modalities of eccentric exercise: different recipes, not
979 the same dish. *J Appl Physiol* **0,** null.

980
981 Friden J, Sjöström M & Ekblom B. (1983). Myofibrillar damage following intense eccentric exercise in
982 man. *Int J Sports Med* **4,** 170-176.

983
984 Fry CS, Kirby TJ, Kosmac K, McCarthy JJ & Peterson CA. (2017). Myogenic progenitor cells control
985 extracellular matrix production by fibroblasts during skeletal muscle hypertrophy. *Cell stem
986 cell* **20,** 56-69.

987
Fry CS, Lee JD, Jackson JR, Kirby TJ, Stasko SA, Liu H, Dupont-Versteegden EE, McCarthy JJ & Peterson
CA. (2014). Regulation of the muscle fiber microenvironment by activated satellite cells
during hypertrophy. *FASEB J* **28,** 1654-1665.

Gillies AR & Lieber RL. (2011). Structure and function of the skeletal muscle extracellular matrix.
Muscle Nerve **44,** 318-331.

Girard O, Mendez-Villanueva A & Bishop D. (2011). Repeated-Sprint Ability—Part I. *Sports Med* **41,**
673-694.

Glaister M, Howatson G, Pattison JR & McInnes G. (2008). The reliability and validity of fatigue
measures during multiple-sprint work: an issue revisited. *J Strength Cond* **22,** 1597-1601.

1000

1001 Grohmann M, Foulstone E, Welsh G, Holly J, Shield J, Crowne E & Stewart C. (2005). Isolation and
1002   validation of human prepubertal skeletal muscle cells: maturation and metabolic effects of
1003   IGF-I, IGFBP-3 and TNFα. *J Physiol* **568,** 229-242.

1004

1005 Guex K, Gojanovic B & Millet GP. (2012). Influence of hip-flexion angle on hamstrings isokinetic
1006   activity in sprinters. *Journal of athletic training* **47,** 390-395.

1007

1008 Hermens HJ, Freriks B, Disselhorst-Klug C & Rau G. (2000). Development of recommendations for
1009   SEMG sensors and sensor placement procedures. *Journal of electromyography and*
1010   *Kinesiology* **10,** 361-374.

1011

1012 Higashihara A, Ono T, Kubota J, Okuwaki T & Fukubayashi T. (2010). Functional differences in the
1013   activity of the hamstring muscles with increasing running speed. *Journal of sports sciences*
1014   **28,** 1085-1092.

1015

1016 Howatson G & Milak A. (2009). Exercise-induced muscle damage following a bout of sport specific
1017   repeated sprints. *J Strength Cond* **23,** 2419-2424.

1018

1019 Howatson G & Van Someren KA. (2008). The prevention and treatment of exercise-induced muscle
1020   damage. *Sports Med* **38,** 483-503.

1021

1022 Jakobsen JR, Jakobsen NR, Mackey AL, Koch M, Kjaer M & Krogsgaard MR. (2018). Remodeling of
1023   muscle fibers approaching the human myotendinous junction. *Scandinavian journal of*
1024   *medicine & science in sports*.

1025

1026 Joe AW, Yi L, Natarajan A, Le Grand F, So L, Wang J, Rudnicki MA & Rossi FM. (2010). Muscle injury
1027   activates resident fibro/adipogenic progenitors that facilitate myogenesis. *Nat Cell Biol* **12,**
1028   153.

1029

1030 Jones D, Newham D, Round J & Tolfree S. (1986). Experimental human muscle damage:
1031   morphological changes in relation to other indices of damage. *J Physiol* **375,** 435-448.

1032

1033 Jönhagen S, Ericson MO, Nemeth G & Eriksson E. (1996). Amplitude and timing of electromyographic
1034   activity during sprinting. *Scandinavian journal of medicine & science in sports* **6,** 15-21.

1035

1036 Jonhagen S, Nemeth G & Eriksson E. (1994). Hamstring injuries in sprinters the role of concentric and
1037   eccentric hamstring muscle strength and flexibility. *The American Journal of Sports Medicine*
1038   **22,** 262-266.

1039

1040 Kellis E & Baltzopoulos V. (1999). In vivo determination of the patella tendon and hamstrings
1041   moment arms in adult males using videofluoroscopy during submaximal knee extension and
1042   flexion. *Clinical Biomechanics* **14,** 118-124.

1043

1044 Kirk EA & Rice CL. (2016). Contractile function and motor unit firing rates of the human hamstrings. *J*
1045   *Neurophysiol***,** jn. 00620.02016.

1046

1047 Lanza MB, Balshaw TG, Massey GJ & Folland JP. (2018). Does normalization of voluntary EMG
1048   amplitude to MMAX account for the influence of electrode location and adiposity? *Scand J*
1049   *Med Sci Sports*.

1050

1051 Mackey AL, Magnan M, Chazaud B & Kjaer M. (2017). Human skeletal muscle fibroblasts stimulate in
1052   vitro myogenesis and in vivo muscle regeneration. *J Physiol* **595,** 5115-5127.

1053

1054 Maiwald C, Sterzing T, Mayer T & Milani T. (2009). Detecting foot-to-ground contact from kinematic
1055   data in running. *Footwear Science* **1,** 111-118.

1056

1057 Malone S, Roe M, Doran DA, Gabbett TJ & Collins K. (2017). High chronic training loads and exposure
1058   to bouts of maximal velocity running reduce injury risk in elite Gaelic football. *Journal of*
1059   *Science and Medicine in Sport* **20,** 250-254.

1060

1061 Marshall PW, Lovell R, Jeppesen GK, Andersen K & Siegler JC. (2014). Hamstring muscle fatigue and
1062   central motor output during a simulated soccer match. *PLoS One* **10,** e102753.

1063

1064 McClay I & Manal K. (1999). Three-dimensional kinetic analysis of running: significance of secondary
1065   planes of motion. *Medicine and Science in Sports and Exercise* **31,** 1629-1637.

1066

1067 Mendiguchia J, Alentorn-Geli E, Idoate F & Myer GD. (2013). Rectus femoris muscle injuries in
1068   football: a clinically relevant review of mechanisms of injury, risk factors and preventive
1069   strategies. *Br J Sports Med* **47,** 359-366.

1070

1071 Murphy MM, Lawson JA, Mathew SJ, Hutcheson DA & Kardon G. (2011). Satellite cells, connective
1072   tissue fibroblasts and their interactions are crucial for muscle regeneration. *Development*
1073   **138,** 3625-3637.

1074

1075 Nedelec M, McCall A, Carling C, Legall F, Berthoin S & Dupont G. (2014). The influence of soccer
1076   playing actions on the recovery kinetics after a soccer match. *J Strength Cond* **28,** 1517-1523.

1077

1078 Newham D, Jones D & Clarkson P. (1987). Repeated high-force eccentric exercise: effects on muscle
1079   pain and damage. *J Appl Physiol* **63,** 1381-1386.

1080

1081 Newham D, McPhail G, Mills K & Edwards R. (1983). Ultrastructural changes after concentric and
1082   eccentric contractions of human muscle. *J Neurol Sci* **61,** 109-122.

1083

1084 Onishi H, Yagi R, Oyama M, Akasaka K, Ihashi K & Handa Y. (2002). EMG-angle relationship of the
1085   hamstring muscles during maximum knee flexion. *Journal of Electromyography and*
1086   *Kinesiology* **12,** 399-406.

1087

1088 Opar DA, Williams M, Timmins R, Hickey J, Duhig S & Shield A. (2014). Eccentric hamstring strength
1089   and hamstring injury risk in Australian footballers. *Medicine & Science in Sports & Exercise* **46**.

1090

1091 Opar DA, Williams MD & Shield AJ. (2012). Hamstring strain injuries. *Sports Med* **42,** 209-226.

1092

1093 Owens DJ. (2015). The Role of Vitamin D in Skeletal Muscle Function and Regeneration. Liverpool
1094   John Moores University.

1095

1096 Owens DJ, Sharples AP, Polydorou I, Alwan N, Donovan T, Tang J, Fraser WD, Cooper RG, Morton JP &
1097   Stewart C. (2015). A systems-based investigation into vitamin D and skeletal muscle repair,
1098   regeneration, and hypertrophy. *Am J Physiol-Endoc M* **309,** E1019-E1031.
1099

1100 Paschalis V, Giakas G, Baltzopoulos V, Jamurtas AZ, Theoharis V, Kotzamanidis C & Koutedakis Y.
1101   (2007). The effects of muscle damage following eccentric exercise on gait biomechanics. *Gait*
1102   *& posture* **25,** 236-242.
1103

1104 Randolph ME & Pavlath GK. (2015). A muscle stem cell for every muscle: variability of satellite cell
1105   biology among different muscle groups. *Front Aging Neurosci* **7,** 190.
1106

1107 Reeves ND, Narici MV & Maganaris CN. (2004). In vivo human muscle structure and function:
1108   adaptations to resistance training in old age. *Experimental physiology* **89,** 675-689.
1109

1110 Robinson MA & Vanrenterghem J. (2012). An evaluation of anatomical and functional knee axis
1111   definition in the context of side-cutting. *Journal of biomechanics* **45,** 1941-1946.
1112

1113 Schuermans J, Van Tiggelen D, Danneels L & Witvrouw E. (2014). Biceps femoris and
1114   semitendinosus—teammates or competitors? New insights into hamstring injury
1115   mechanisms in male football players: a muscle functional MRI study. *Br J Sports Med* **48,**
1116   1599-1606.
1117

1118 Schwartz MH & Rozumalski A. (2005). A new method for estimating joint parameters from motion
1119   data. *Journal of biomechanics* **38,** 107-116.
1120

1121 Scott J & Huskisson E. (1979). Vertical or horizontal visual analogue scales. *Ann Rheum Dis* **38,** 560.
1122

1123 Seymore KD, Domire ZJ, DeVita P, Rider PM & Kulas AS. (2017). The effect of Nordic hamstring
1124   strength training on muscle architecture, stiffness, and strength. *Eur J Appl Physiol* **117,** 943-
1125   953.
1126

1127 Stauber W, Clarkson P, Fritz V & Evans W. (1990). Extracellular matrix disruption and pain after
1128   eccentric muscle action. *J Appl Physiol* **69,** 868-874.
1129

1130 Stewart JD, Masi TL, Cumming AE, Molnar GM, Wentworth BM, Sampath K, McPherson JM & Yaeger
1131   PC. (2003). Characterization of proliferating human skeletal muscle-derived cells in vitro:
1132   Differential modulation of myoblast markers by TGF-β2. *J Cell Physio* **196,** 70-78.
1133

1134 Terry EE, Zhang X, Hoffmann C, Hughes LD, Lewis SA, Li J, Wallace MJ, Riley LA, Douglas CM &
1135   Gutierrez-Monreal MA. (2018). Transcriptional profiling reveals extraordinary diversity
1136   among skeletal muscle tissues. *eLife* **7**.
1137

1138 Thelen DG, Chumanov ES, Hoerth DM, Best TM, Swanson SC, Li L, Young M & Heiderscheit BC. (2005).
1139   Hamstring muscle kinematics during treadmill sprinting. *Medicine and science in sports and*
1140   *exercise* **37,** 108-114.
1141

1142 Tidball JG. (2011). Mechanisms of muscle injury, repair, and regeneration. *Comprehensive Physiology*
1143   **1,** 2029-2062.
1144

1145 Timmins RG, Bourne MN, Shield AJ, Williams MD, Lorenzen C & Opar DA. (2015). Short biceps femoris
1146   fascicles and eccentric knee flexor weakness increase the risk of hamstring injury in elite
1147   football (soccer): a prospective cohort study. *Br J Sports Med*, bjsports-2015-095362.
1148
1149 Timmins RG, Bourne MN, Shield AJ, Williams MD, Lorenzen C & Opar DA. (2016a). Short biceps
1150   femoris fascicles and eccentric knee flexor weakness increase the risk of hamstring injury in
1151   elite football (soccer): a prospective cohort study. *Br J Sports Med* **50,** 1524-1535.
1152
1153 Timmins RG, Opar DA, WIlliams MD, Schache AG, Dear NM & Shield AJ. (2014). Reduced biceps
1154   femoris myoelectrical activity influences eccentric knee flexor weakness after repeat sprint
1155   running. *Scandinavian journal of medicine & science in sports* **24,** e299-e305.
1156
1157 Timmins RG, Shield AJ, Williams MD, Lorenzen C & Opar DA. (2016b). Architectural adaptations of
1158   muscle to training and injury: a narrative review outlining the contributions by fascicle
1159   length, pennation angle and muscle thickness. *Br J Sports Med*, bjsports-2015-094881.
1160
1161 Turrina A, Martínez-González MA & Stecco C. (2013). The muscular force transmission system: role of
1162   the intramuscular connective tissue. *J Bodywork Movement Ther* **17,** 95-102.
1163
1164 Valle X, Alentorn-Geli E, Tol JL, Hamilton B, Garrett WE, Pruna R, Til L, Gutierrez JA, Alomar X & Balius
1165   R. (2017). Muscle injuries in sports: a new evidence-informed and expert consensus-based
1166   classification with clinical application. *Sports Med* **47,** 1241-1253.
1167
1168 van der Horst N, Smits D-W, Petersen J, Goedhart EA & Backx FJ. (2015). The Preventive Effect of the
1169   Nordic Hamstring Exercise on Hamstring Injuries in Amateur Soccer Players A Randomized
1170   Controlled Trial. *The American journal of sports medicine*, 0363546515574057.
1171
1172 van Dyk N, Bahr R, Whiteley R, Tol JL, Kumar BD, Hamilton B, Farooq A & Witvrouw E. (2016).
1173   Hamstring and Quadriceps Isokinetic Strength Deficits Are Weak Risk Factors for Hamstring
1174   Strain Injuries A 4-Year Cohort Study. *The American journal of sports medicine*,
1175   0363546516632526.
1176
1177 Verheul J, Clansey AC & Lake MJ. (2017). Adjustments with running speed reveal neuromuscular
1178   adaptations during landing associated with high mileage running training. *Journal of applied
1179   physiology* **122,** 653-665.
1180
1181 Verma S, Moiz J, Shareef M & Husain M. (2015). Physical performance and markers of muscle
1182   damage following sports specific sprints in male collegiate soccer players: repeated bout
1183   effect. *The Journal of sports medicine and physical fitness*.
1184
1185 Wang N, Tytell JD & Ingber DE. (2009). Mechanotransduction at a distance: mechanically coupling the
1186   extracellular matrix with the nucleus. *Nature reviews Molecular cell biology* **10,** 75-82.
1187
1188 Woods K, Bishop P & Jones E. (2007). Warm-up and stretching in the prevention of muscular injury.
1189   *Sports Med* **37,** 1089-1099.
1190
1191 Wüst RC, Morse CI, De Haan A, Jones DA & Degens H. (2008). Sex differences in contractile properties
1192   and fatigue resistance of human skeletal muscle. *Experimental physiology* **93,** 843-850.
1193

1194 # SUPPLEMENTARY INFORMATION

1195

1196 ***Figure supplement 5*** *Representative images for muscle cell differentiation with a high*
1197 *myoblast:fibroblast ratio (2.4; left) and with a low myoblast:fibroblast ratio (0.3; right) at day seven.*
1198 *Magnification is x 10.5, and scale bar is 100 µm.*

1199

1200

1201 **Table supplement 4** Architectural parameters of biceps femoris long head (mean ± SD).

Muscle length [cm]	Fascicle length [cm]	PCSA [cm^2]	Volume [cm^3]	Fascicle Pennation Angle [°]
27.86 ± 2.13	7.94 ± 1.38	23.4 ± 4.62	182.2 ± 29.5	12.7 ± 2.77

Fascicle length at 50% of BF_{LH} muscle length; PCSA – Physiological cross-sectional area.

1202

1203 ***Table supplement 6*** *Linear regression analysis between the myoblast:fibroblast ratio and fusion*
1204 *parameters at day 7 and CK activity at day 10 in vitro.*

	R^2	F-Test	P Value
Total myotubes [n]	0.35	$F(1,10) = 5.27$	0.045*
Myotube Length [µm]	0.34	$F(1,10) = 5.13$	0.047*
Myotube Diameter [µm]	0.39	$F(1,10) = 5.56$	0.031*
Myotube Area [µm^2]	0.40	$F(1,10) = 6.79$	0.026*
CK activity [mU/mL]	0.41	$F(1,10) = 6.23$	0.034*

* significant (P<0.05);

1205