Neuromuscular fatigue and recovery after strenuous exercise depends on skeletal muscle size and stem cell characteristics

Baumert, P.[1,2]*, Temple, S.[1], Stanley, J.M.[1], Cocks M. [1], Strauss, J.A. [1], Shepherd

S.O.[1], Drust, B.[1], Lake, M.J.[1], Stewart, C.E.[1], Erskine, R.M.[1,3]

[1]Research Institute for Sport & Exercise Sciences, Liverpool John Moores University,
Liverpool, United Kingdom.

[2]Exercise Biology Group, Faculty of Sport and Health Sciences, Technical University of Munich,
Munich, Germany

[3]Institute of Sport, Exercise & Health, University College London, London, UK

Running title: Neuromuscular Response following muscle damage *in vivo* and *in vitro*

Address for reprint requests and all other correspondence:

Philipp Baumert, Exercise Biology Group, Faculty of Sport and Health Sciences, Technical University of Munich, Munich, Germany; Email: philipp.baumert@tum.de

Key words: Biceps femoris architecture; exercise-induced muscle damage (EIMD); extracellular matrix (ECM); fibroblast; muscle strain injury; myoblast

This is an original research article

ABSTRACT

Hamstring muscle injury is highly prevalent in sports involving repeated maximal sprinting. Although neuromuscular fatigue is thought to be a risk factor, the mechanisms underlying the fatigue response to repeated maximal sprints are unclear. Here, we show that repeated maximal sprints induce neuromuscular fatigue accompanied with a prolonged strength loss in hamstring muscles. The immediate hamstring strength loss was linked to both central and peripheral fatigue, while prolonged strength loss was associated with indicators of muscle damage. The kinematic changes immediately after sprinting likely protected fatigued hamstrings from excess elongation stress, while larger hamstring muscle physiological cross-sectional area and lower *myoblast:fibroblast* ratio appeared to protect against fatigue/damage and improve muscle recovery within the first 48 h after sprinting. We have therefore identified novel mechanisms that likely regulate the fatigue/damage response and initial recovery following repeated maximal sprinting in humans.

INTRODUCTION

Hamstring strain is the most frequently occurring injury in sport [1], particularly in those sports that involve high-speed running [2]. Although the aetiology is unclear, numerous risk factors have been proposed, such as short fascicle length, poor flexibility, poor hamstring strength, and inadequate warm-up [3]. Further, it is unknown whether hamstring strain is the result of a single event that exceeds the physiological range of hamstring muscle extensibility and contractility, or as a result of an accumulation of eccentric contractions during repeated maximal sprints, causing neuromuscular fatigue [3]. Neuromuscular fatigue is responsible for acute, as well as prolonged, impairment of muscle function, classified as central fatigue (i.e. originating in the central nervous system), or peripheral fatigue (i.e. distal to the neuromuscular junction) [4]. Although it was recently reported that both central and peripheral fatigue contribute to impaired hamstring muscle function immediately after repeated maximal sprint-related interventions [5,6], the contribution of neuromuscular fatigue to hamstring muscle impairment and recovery, following repeated maximal sprints over time, is insufficiently studied [7]. An understanding of hamstring neuromuscular fatigue following repeated maximal sprints may be crucial for understanding hamstring strain aetiology.

Peripheral fatigue may be caused by ultrastructural muscle damage, which is indicated by Z-line disturbance [8] as well as disruption of the extracellular matrix [9]. The extracellular matrix provides structural scaffolding for muscle remodelling and plays an integral role in force transmission [10]. This is referred to as exercise-induced muscle damage and it is exhibited by prolonged strength loss and delayed-onset muscle soreness, as well as the release of muscle-specific proteins [e.g. creatine kinase (CK)] into the circulation over the following days [11]. After substantial muscle damage, myogenic satellite cells (skeletal muscle stem cells), play a key role in skeletal muscle regeneration and remodelling [12]. Activated satellite cells (myoblasts) proliferate and migrate from their niche along the basal lamina to the injury site before terminally differentiating and fusing with damaged myofibrils to repair injury. There is increasing evidence that fibroblasts, the main cell type of muscle connective tissue, also play

a critical role in supporting muscle regeneration [13,14]. Following damage, infiltrating

inflammatory cells activate muscle fibroblasts, which proliferate and migrate to the area of the

myotrauma and produce extracellular matrix components in an orchestrated and regulated

fashion to support healthy muscle remodelling [14,15]. However, little is known about the role of

fibroblasts during the initial response and recovery following *physiological* exercise-induced

muscle damage, e.g. following repeated sprinting.

Acute damage to the muscle-tendon complex may facilitate hamstring strain, which is thought

to occur in the late swing phase of sprinting, when the hamstring muscles contract eccentrically,

i.e. trying to shorten while being lengthened in an attempt to decelerate the shaft before initial

foot-ground contact [16]. Therefore, a short biceps femoris long head (BF_{LH}) fascicle length has

been suggested to increase hamstring strain risk [17], as the BF_{LH} is thought to be relatively more

eccentrically stretched during the late swing phase of sprinting compared to the other

hamstring muscles [16]. However, no study has investigated the relationship between BF_{LH}

architecture (including muscle fascicle length and cross-sectional area), and the prolonged

hamstring muscle response to exercise-induced neuromuscular fatigue. Finally, lower limb

neuromuscular fatigue might cause a number of biomechanical alterations in running

kinematics [18]. However, it is not known whether repeated maximal sprints influence kinematic

patterns, and whether this can lead to prolonged changes in lower-limb kinematics, which may

play a role in the development of muscle strain following insufficient recovery [3].

Here we demonstrated that both central and peripheral fatigue, assessed via surface

electromyography (sEMG) and electrical stimulation, contributed to the immediate loss of

muscle function in both the quadriceps and hamstring muscle groups, but that peripheral

factors mainly contributed to the sustained loss of hamstring muscle function. Moreover, we

established that a lower myoblast:fibroblast ratio in isolated primary human muscle stem cells

correlated with improved recovery from both repeated maximal sprints and an *in vitro* artificial

wounding assay within the first 48 h. We also report that BF_{LH} architecture (i.e. physiological

cross-sectional area, PCSA) was associated with hamstring fatigue, and that neuromuscular

fatigue led to reduced hip flexion and knee extension during the late swing phase of steady-

state running. Thus, with this interdisciplinary study, we identify novel cellular and

neuromuscular mechanisms underpinning central and peripheral fatigue following repeated

sprinting, which ultimately led to kinematic changes during the running stride phase associated

with hamstring strain injury.

RESULTS

Effect of the Repeated Maximal Sprint Intervention on Neuromuscular Fatigue

To examine the effect of repeated maximal sprints on neuromuscular fatigue, we measured

different fatigue parameters before (PRE), immediately after (POST), and 48 h after (POST48)

the repeated maximal sprint intervention. The average 30 m sprinting speed was 6.48 ± 0.33

m s^{-1}. There was a main effect of time for heart rate, 30 m sprinting time, rating of perceived

exertion and lactate concentration, with all parameters increasing from PRE to POST (all

$P<0.001$). Blood lactate concentration increased from PRE (1.63 ± 0.45 mmol/L) to POST (9.82

± 3.62 mmol/L; $P<0.001$). The sprinting performance (measured via the performance

decrement score [19]) decreased by $3.98 \pm 2.99\%$ during the run, and rating of perceived exertion

increased by $96.5 \pm 35.2\%$ from PRE-to-POST, indicating fatigue had occurred.

We then performed *in vivo* functional analysis to assess if repeated maximal sprints resulted

in an increase in central and/or peripheral fatigue. We, therefore, measured BF_{LH} muscle

activation via normalised surface electromyography (sEMG) during hamstring maximum

voluntary contraction (MVC). We observed a change in sEMG ($F_{F2,24}=4.35$, $P=0.022$), with

post-hoc pairwise comparisons revealing a decrease from PRE-to-POST (-24.3%; $P=0.019$).

However, this change was no longer evident at POST48 ($P=0.157$, Table 1), suggesting that

central fatigue occurred immediately after repeated maximal sprints. No other changes in

muscle (co)activation were observed at any time point ($P>0.05$).

[Please insert Table 1 near here]

We also assessed the torque-frequency relationship *in vivo* via electrical stimulation to indicate

peripheral (muscle) fatigue. There was an interaction between time x stimulation frequency

(n=19; $F_{4.9,216}$=6.62, P<0.001; Figure 1A). Post-hoc paired t-tests revealed differences PRE-to-

POST for 10-50 Hz (P<0.05), but lower frequencies between 10-20 Hz reverted to baseline

values POST48 (P>0.05), while the frequencies of 30 and 50 Hz were still decreased POST48

compared to their baseline values (P<0.05), providing evidence that peripheral fatigue

occurred immediately after the repeated sprints and remained for 48 h

[Please inside Figure 1 near here]

Effect of the Repeated Maximal Sprint Intervention on MVC Strength, Muscle

Soreness and Serum Markers of Exercise-Induced Muscle Damage

To investigate the effect of repeated maximal sprints on biomarkers of exercise-induced

muscle damage, we assessed hamstring (knee flexion) and quadriceps (knee extension) MVC,

muscle soreness, serum creatine kinase (CK) activity and interleukin-6 (IL-6) concentrations

PRE, POST and POST48. Isometric hamstring and quadriceps MVC, muscle soreness (all

P<0.001) and serum CK activity ($F_{1.3,34}$=5.98, P=0.017), as well as IL-6 concentration

($F_{1.3,34}$=5.96, P=0.018), showed a main effect of time, which are indicators of muscle damage

[Please insert Table 2 near here]

) that was similar to other studies [7,20,21]. Post-hoc pairwise comparisons revealed that,

compared to PRE, both serum CK activity (+93.0%) and serum IL-6 concentration (+307%)

were elevated at POST (both P=0.027), and CK activity further increased at POST48 (+256%;

P=0.012), while serum IL-6 concentration reverted to baseline values (P>0.05).

[Please insert Table 2 near here]

Further, there was an interaction between time and muscle groups concerning relative MVC

torque loss (percentage change from PRE MVC) ($F_{1.4,38}$=7.92, P=0.004). Relative MVC

decreased similarly in both quadriceps and hamstring muscle groups PRE-to-POST (Figure

1B). However, at POST48, hamstring MVC continued to decrease from POST (-9.26%;

P=0.010), while quadriceps MVC began to return to PRE values (+8.47%; P=0.016) and was

higher than hamstring MVC at POST48 (P=0.038).

Effect of the Repeated Maximal Sprint Intervention on Lower-Limb Kinematics

To assess the consequential effect of neuromuscular fatigue on lower-limb kinematics, we

captured treadmill running (4.17 m s^{-1}) kinematics with an eight-camera motion capture system

over time. Three-dimensional motion analysis demonstrated, that despite significance not

being achieved, there was a tendency towards a longer running cycle time POST (+1.01%)

and POST48 (+1.86%), compared to PRE (P=0.080; Table 3). Further, treadmill running

demonstrated decreased peak knee extension (P=0.047) during the late swing phase at POST

(-10.9%) compared to PRE, but this reverted to baseline POST48. The percentage change in

peak knee extension correlated with the percentage change in relative hamstring MVC torque

both measured POST-to-POST48 (R^2=0.26, $F_{1,2}$=5.673, P=0.031).

[Please insert Table 3 near here]

Architecture of the Biceps Femoris Long Head Muscle

To assess whether architectural parameters of the BF_{LH} muscle (Figure 2) were associated

with markers of peripheral fatigue, we performed ultrasound measurements of the BF_{LH} muscle

(Table supplement 4). Muscle fascicle length and pennation angle of the BF_{LH}, which have

previously been linked to hamstring muscle strain risk [22], did not correlate with any outcome

variable of neuromuscular fatigue. However, BF_{LH} PCSA (mean ±SD: 23.4 ± 4.62 cm^2)

correlated inversely with relative hamstring MVC loss PRE-to-POST (R^2=0.42, $F_{1,17}$=12.37,

P=0.003, Figure 2C).

[Please insert Figure 2 near here]

Artificial Wound Healing Assay to Investigate Repair and Regeneration Regarding Myoblast:Fibroblast Ratio

Preliminary data from our laboratory demonstrated that skeletal muscle stem cell composition

(i.e. myoblast:fibroblast ratio), derived from two volunteers with ratios representing the extreme

conditions of the myoblast:fibroblast percentage, played a role in the success of artificial wound

healing *in vitro* [23]. Human primary skeletal muscle stem cells with a high myoblast:fibroblast

ratio resulted in reduced cell migration into an artificial wound (assessed by cell number within

the wound) when compared with stem cells with a low myoblast:fibroblast ratio. No significant

differences were observed in the relative proportion of migrating cells (i.e. between myoblasts

and fibroblasts) over 48 hours. Further, recent investigations have revealed that the skeletal

muscle stem cell ratio does not change during *in vitro* cell culturing [24,25]. We, therefore, further

assessed the effect of the myoblast:fibroblast ratio on skeletal muscle recovery following *in*

vitro artificial wounding to extend the preliminary *in vitro* results from our laboratory. To be able

to demonstrate a coefficient of determination of ≥ 0.50 between myoblast:fibroblast ratio and

our dependent variables, a priori power calculations for a Pearson's r correlation was run using

G*Power (version 3.1.9.1), and it revealed that 11 persons provided alpha = 5%, and power =

80%. Thus, we used primary human skeletal muscle stem cells derived from 12 participants,

six who participated in both the repeated maximal sprint intervention and also volunteered to

provide a muscle biopsy at least three weeks before the repeated maximal sprint intervention,

and another six (two male and four females), who did not participate in the repeated maximal

sprint intervention (to increase the power of the *in vitro* study). The muscle stem cells were

isolated, cultured and then characterized by immunofluorescence staining. The mean ± SD

myoblast:fibroblast ratio of the twelve participants was 1.26 ± 1.00 (range: 0.276 to 2.93). We

did not detect any differences in muscle stem cell characteristics between cells obtained from

females and males (data not shown). We, therefore, combined the data from all muscle cells

and correlated the muscle characteristics with individual myoblast:fibroblast ratios. We

observed no correlations regarding myoblast:fibroblast ratio and the total number of skeletal

muscle stem cells (combined number of myoblasts, fibroblasts and other stem cells) migrating

into the artificial wound within all three segments combined at 24 h (R^2=0.20, $F_{1,10}$=2.56,

P=0.141) or 48 h (R^2=0.02, $F_{1,10}$=0.19, P=0.671) after the scratch assay.

[Please insert Figure 3 near here]

However, there was an inverse correlation between myoblast:fibroblast ratio and migration

dynamics for the 12 participants (Figure 3B). Muscle stem cells with a low myoblast:fibroblast

ratio demonstrated more cells in the inner segment than to the outer segment compared to

muscle stem cells with high myoblast:fibroblast ratio at 24 h (R^2=0.49, $F_{1,10}$=9.53, P=0.011)

and with a non-significant trend at 48 h (R^2=0.30, $F_{1,10}$=4.33, P=0.064) after the artificial wound

healing assay reinforcing the preliminary *in vitro* results from our laboratory

Comparison of the Muscle Response between the Repeated Maximal Sprint

Protocol and the Muscle Stem Cell Study

In order to determine whether skeletal muscle stem cell ratio (i.e. myoblast:fibroblast ratio)

played a role in muscle strength recovery *in vivo*, further studies were performed. Previous

investigations have shown that skeletal muscles of different origin, but with similar

physiological functions and fibre type composition demonstrate similar transcriptome

expression patterns of up to 99% [26,27]. Further, all limb muscles arise developmentally from the

ventrolateral dermomyotome of the segmented paraxial mesoderm [28] and these muscles with

a similar fibre type composition show low intra-subject [29,30] but high inter-subject variability [30,31]

regarding the total amount of stem cells within these muscle fibres. Thus, muscle biopsies

were obtained from the vastus lateralis of six participants, who also performed the repeated

maximal sprint intervention (they volunteered to provide a muscle biopsy at least three weeks

before completing the repeated maximal sprint intervention). As muscle fibre-type composition

is similar between the quadriceps and hamstrings [26], the stem cell composition of the vastus

lateralis was considered representative of both the quadriceps and hamstring muscle groups.

The mean ± SD myoblast:fibroblast ratio of the six participants was 1.46 ± 1.06 (range: 0.299

to 2.93). There was an inverse correlation between myoblast:fibroblast ratio and the

percentage change in relative hamstring MVC torque measured PRE-to-POST48 *in vivo* (R=-

0.945, $F_{1,4}$=33.73, P=0.004; Figure 4A). Thus, participants with a high myoblast:fibroblast ratio

showed a delayed hamstring strength recovery 48 h after repeated maximal sprints compared

to those with a low myoblast:fibroblast ratio. Further, there was an inverse correlation between

myoblast:fibroblast ratio and relative hamstring MVC torque measured POST-to-POST48 (R=-

0.943, $F_{1,4}$=17.08, P=0.014; Figure 4B). No correlations were found between the

myoblast:fibroblast ratio and changes in quadriceps MVC torque or with any other muscle

damage and fatigue biomarker following repeated maximal sprints. These inverse correlations

could be confounded by the model assumptions and the relative low participant number.

However, post hoc power calculations demonstrate that the bivariate correlation had adequate

power (0.97).

[Please insert Figure 4 near here]

DISCUSSION

In this study, we have used an interdisciplinary approach to investigate the potential

biomechanical, physiological and cellular factors underpinning neuromuscular fatigue following

repeated maximal sprints. We have shown that immediate strength loss was associated with

reduced hamstring sEMG activity (indicating impaired hamstring motor unit recruitment) and

markers of peripheral fatigue, but the magnitude and sustained changes in MVC torque over

time (especially in the hamstrings) was largely associated with indicators of peripheral fatigue.

Muscle damage biomarkers indicated that the hamstring peripheral fatigue might have been

caused predominantly by ultrastructural damage within the muscle tissue. Further, both central

and peripheral fatigue caused by repeated maximal sprints appeared to affect the

neuromuscular control of running patterns, while a larger BF_{LH} PCSA was related to attenuated

hamstring strength loss immediately after the repeated maximal sprint intervention. Finally, our

results suggest that a high myoblast:fibroblast ratio leads to a delayed wound closure *in vitro*

and to a delayed MVC torque recovery following repeated maximal sprints *in vivo* within the

first 48 h, indicating that stem cells of the non-contractile muscle tissue might positively affect

the response to muscle damaging exercise.

Fatigue and Muscle Damage Following Repeated Maximal Sprints

We showed a decreased activity of normalised BF_{LH} sEMG activity immediately after the

repeated maximal sprint intervention, but no significant changes in neuromuscular activation

using the interpolated twitch technique. The discrepancy between these two methods might

be explained by the fact that voluntary activation measured via the interpolated twitch

technique investigates all of the hamstring muscles, whilst the normalised EMG analysis was

confined solely to the BF_{LH}, which is in line with a previous study [5]. The BF_{LH} might fatigue to a

greater degree immediately after repeated maximal sprint related interventions compared to

the other hamstring muscles. However, we also provide evidence for peripheral fatigue

occurring immediately after the repeated maximal sprints and a delayed recovery at higher

frequencies (30-50 Hz) after observing a right shift in the torque-frequency relationship. This

may be due to ultrastructural damage predominantly in fast-twitch (which fire at rates from 30-

50 Hz) compared to slow-twitch muscle fibres (discharge rates 10-25 Hz) [32,33], leading to

impaired force generation rather than simply fatiguing the muscle fibres.

Both the quadriceps and hamstring muscle groups showed similar strength loss immediately

after the repeated maximal sprints, but the hamstring muscle group showed further strength

loss 48 h later compared to the quadriceps. Other studies did not show this additional strength

loss for the hamstring muscle group POST48. Differences in the training status of the

participants [7,20] and in the methodological approaches [34] might partly explain the different

outcomes. The peak BF_{LH} EMG activity occurs at a more extended knee angle during

hamstring isokinetic muscle contraction compared to the peak EMG occurring at a more flexed

knee angle for the other hamstring muscles, such as the semitendinosus [35]. Therefore, we

suggest that the BF_{LH} is the key hamstring muscle responsible for decelerating the shaft at the

end of the late swing phase. After repeated bouts of high-speed running, the semitendinosus

might fatigue prematurely [36] and the BF_{LH} would need to substitute the impaired function of the

preceding semitendinosus to decelerate the shaft.

During high-speed running, eccentric contractions occur in the hamstrings during the late

swing phase (i.e during deceleration of the knee extension) [37] and in the quadriceps during the

early/mid swing phase (= deceleration of hip extension) [38]. In general, the likelihood of

sustaining a quadriceps strain injury is significantly lower compared to a hamstring strain injury

[2]. If repeated eccentric contractions are one of the main causes for strain injuries, as indicated

by several investigations [39,40], then the difference in strength loss between the quadriceps and

hamstrings observed in our study might be explained by the different levels of eccentric force

generated by the muscles. Muscle damage in the quadriceps muscle presumably occurs

during the deceleration phase of sprinting and during the backswing phase, when the

quadriceps muscle works eccentrically to decelerate the leg with a flexed knee during the early

swing phase of high-speed running [21,38]. Hamstring strain injury, however, occurs with an

almost extended leg during the late swing phase [37]. This extended lever arm might cause

higher eccentric force in the hamstring muscles compared to the shorter lever arm with a flexed

knee on the quadriceps muscle. Continuously repeated eccentric contractions with the longer

lever arm will potentially induce more muscle damage within the hamstrings in total during e.g.

a soccer match, compared to the quadriceps muscle group, which might explain the significant

different strength loss POST48 repeated sprints in the current study, and potentially the

different injury rates between these two muscle groups.

Kinematic Analysis

Our *in vivo* intervention caused changes in the running kinematics with reduced knee extension in the late swing phase immediately after the repeated maximal sprints. Reduced hamstring muscle strength due to neuromuscular fatigue might trigger a protective mechanism directly after repeated maximal sprints. Afferent signals from the fatigued and damaged hamstrings might activate the Golgi tendon organ [4], thus limiting hamstring muscle fibre strain in an attempt to minimise further muscle damage. These kinematic changes were not evident 48 h after the repeated maximal sprints. However, there was a non-significant tendency for prolonged stride duration during running (P=0.08, data not shown) 48 h later and the percentage change of knee extension in the late swing phase of running correlated with changes in hamstring strength both measured from POST to POST48. This indicates that participants with delayed hamstring strength recovery were still not able to fully control running. As hamstring MVC continued to deteriorate 48 h after repeated maximal sprints but quadriceps MVC started to improve, it could be that lower-limb kinematics in the sagittal plane are controlled by the hamstrings more than the quadriceps. That outcome could also have implications for the underlying mechanisms of knee injuries, such as an anterior cruciate ligament injury [41], which would need further investigation. Summarised, ultrastructural damage in the hamstring muscles might lead to decelerated movement patterns over time, which could increase the risk for hamstring strain injury during sprinting [3].

The Role of the Extracellular Matrix on the Muscle Response Following Repeated maximal sprinting

Recent investigations have suggested that hamstring maximum eccentric strength and BF_{LH} fascicle length are predictors of hamstring strain injury [17]. Further, a fatigued muscle is likely to accentuate the risk of muscle strain [2]. However, we could not find any correlation between BF_{LH} fascicle length and any biomarker of fatigue but BF_{LH} PCSA correlated inversely with hamstring strength loss from PRE to POST. During the late swing phase of sprinting, the

hamstring muscles contract eccentrically to decelerate the shaft and to enhance the

subsequent concentring shortening contraction for maximal sprinting by using stored elastic

energy from the muscle-tendon unit. In comparison to other conventional muscle-damaging

interventions [42], this dynamic (stretch-shortening) movement might lead to an additional

damage of the hamstring muscle connective tissue structure. Therefore, a larger BF_{LH} PCSA

might protect against immediate hamstring MVC loss due to a greater ability to transmit the

ground reaction forces laterally (from fibre to fibre) [43], which might disperse the force more

efficiently to the tendon, while the muscle fibres themselves undergo less strain. Further, a

greater BF_{LH} PCSA reflects more fibres aligned in parallel, which would be accompanied by

more muscle connective tissue of the extracellular matrix, thus potentially protecting the

muscle fibres from excessive damage during eccentric contractions.

The stem cells of the extracellular matrix also demonstrated an important role for muscle

strength recovery in the subgroup of participants, as there was a strong inverse correlation

between myoblast:fibroblast ratio and hamstring MVC torque recovery POST48. Skeletal

muscles with a higher availability of fibroblasts around the area of myotrauma might have a

better capacity to reorganise the complex extracellular matrix, thus restoring (lateral) force

transmission, which results in a faster recovery of muscle strength after muscle damage. This

was in line with the myoblast:fibroblast ratio effect on cellular aspects of muscle regeneration

and remodelling assessed in primary muscle stem cells *in vitro*. Muscle stem cells with a low

myoblast:fibroblast ratio revealed a faster wound closure (i.e. more cells migrated to the inner

part of the artificial injury compared to the outer part), in particular 24 h after performing the

scratch assay. These results suggest that a larger abundance of fibroblasts has a positive

effect at the beginning of muscle repair. We, therefore, assume that repeated maximal sprints

with insufficient recovery of previously fatigued and damaged muscles (where the fatigue and

damage response is modulated by the muscle size and stem cell composition, respectively)

might augment the risk of muscle strains, as appropriate damage to the muscle connective

tissue is thought to differentiate between exercise-induced muscle damage and muscle strains

[44,45].

The practical implications of our study are that a 48 h recovery period following repeated

maximal sprinting is insufficient, and might increase hamstring strain injury risk. Furthermore,

increasing hamstring PCSA via resistance training is likely to reduce peripheral fatigue

following repeated maximal sprinting, thereby reducing hamstring strain injury risk.

Limitations

The current study observed a relationship between human primary muscle cell type (*in vitro*)

and physiological biomarkers of skeletal muscle damage/recovery following strenuous

exercise. Further research is necessary to confirm these results with a larger sample size

regarding the *in vitro* study. However, given the scarcity of data that have examined human

muscle stem cell characteristics in association with muscle damage/recovery *in vivo*, and that

most of our results were sufficiently powered, we believe that our study represents an important

advancement in our understanding of how skeletal muscle recovers following strenuous

exercise. There was no relationship between the myoblast:fibroblast ratio and any

physiological variables regarding the quadriceps femoris, from which the muscle biopsies were

obtained. It has previously been shown that skeletal muscles of different origin, but with similar

physiological functions and fibre type composition, demonstrate similar transcriptome

expression patterns of up to 99% [26,27]. For immnunohistochemistry analysis, the ICC (3, k) of

0.83 (95% CIs: 0.59-0.95) indicates a good reliability for the characterisation and the

quantification of myoblasts and fibroblasts. Therefore, it is likely, that the correlation between

myoblast:fibroblast ratio and the muscle damage-response of the hamstrings but not the

quadriceps muscles is explained by more severe ultrastructural damage in the hamstrings than

quadriceps. Furthermore, peripheral fatigue can be caused by metabolic perturbations, such

as the depletion of intramuscular glycogen [11]. Therefore, because we did not control diet

throughout the study, it is possible that inter-individual differences in baseline muscle glycogen

may have influenced the ability to maintain maximal intensity throughout the sprints. However,

participants were instructed to eat and drink similar foods two hours before each laboratory

visit, and to avoid strenuous exercise for at least 48 h prior to the testing. Further, participants

were given sufficient recovery between sprint repetitions and there was a low decrement in

sprint performance, indicating that glycogen depletion was probably only a minor factor.

CONCLUSION

Repeated maximal sprints induce a greater and more prolonged strength loss in the hamstrings

compared to the quadriceps muscles. The immediate loss of hamstring function appears to be

due to both central (particularly reduced neuromuscular activation of the biceps femoris long

head) and peripheral fatigue, while prolonged hamstring strength loss is predominantly linked

to peripheral fatigue. Thigh neuromuscular fatigue following repeated maximal sprints alters

hip and knee kinematics during running immediately after the repeated maximal sprints, which

may lead to an increased hamstring muscle injury risk. Furthermore, biceps femoris long head

PCSA was inversely related to hamstring strength loss immediately after repeated maximal

sprinting. This suggests a greater PCSA may help transmit more ground reaction force laterally

between muscle fibres, e.g. via the extracellular matrix, thus placing less stress during

eccentric contractions on the individual muscle fibres and protecting against muscle

damage/fatigue. Furthermore, our results suggest that skeletal muscles with an increased

number of fibroblasts might have a better capacity to reorganise the complex extracellular

matrix, which results in a faster wound closure after substantial muscle damage.

MATERIALS AND METHODS

Participants

In vivo repeated maximal sprint intervention: Twenty recreationally active and healthy young men (*mean ± SD*; age 20.3 ± 2.87 years; height 1.79 ± 0.05 m; body mass 75.0 ± 7.89 kg) participated in the repeated maximal sprint intervention.

In vitro muscle stem cell component: Eight healthy young male (age 21.25 ± 4.27 years; height 1.77 ± 0.05 m; body mass 73.78 ± 5.68 kg) and four healthy young female (age 25.5 ± 1.29 years; height 1.67 ± 0.08 m; body mass 61.40 ± 2.57 kg) participants provided a biopsy of the vastus lateralis muscle for the *in vitro* muscle stem cell component of this study. Six of the eight males also participated in the repeated maximal sprint intervention at least three weeks after providing a muscle biopsy. Only males were recruited for the repeated maximal sprint intervention, as there is some evidence of sex differences in neuromuscular fatigue [46]. However, both men and women were recruited for the muscle stem cell component due to there being no reported sex differences in stem cell properties and none within our own pilot studies (data not shown). Prior to starting the study, written informed consent was obtained from each participant and pre-biopsy screening was performed by a physician for those participants who volunteered a muscle biopsy. The study was approved by the Research Ethics Committee of Liverpool John Moores University and complied with the Declaration of Helsinki. Volunteers were physically active but were ineligible to participate if they had performed strength training of the lower limbs within 6 months prior to participation in the study, which was determined during pre-participation screening. Further exclusion criteria were: (i) any lower limb injury in the past 12 months; (ii) age under 18 or above 35 years; and (iii) more than three structured exercise sessions per week.

Experimental Design of the Repeated Maximal Sprint Intervention *in vivo*

Participants were required to visit the temperature-controlled (22-24°C) laboratory on three occasions: (i) familiarisation, (ii) testing day including the PRE and POST assessments; and (iii) POST48 assessments after the repeated maximal sprint intervention. One week prior to

the testing day, participants were familiarised with the assessments as well as with repeated

maximal sprints (by performing 2-3 submaximal sprints) and BF_{LH} architecture of the hamstring

muscle group was assessed via ultrasound. On the test day, participants performed the

repeated maximal sprint intervention of 15 x 30 m sprints to induce neuromuscular

fatigue/damage in both the quadriceps femoris and hamstring muscle groups. All tests were

performed at the same time of the day for each participant. Further, participants were instructed

to maintain their normal routine (including eating habits), to refrain from drinking alcohol and

to avoid any strenuous exercise 48 h prior to testing and throughout the study, and to refrain

from consuming caffeine on testing days. Nothing was consumed throughout the testing

sessions except water, which was available ad libitum.

The test battery was always performed in the same order with the right leg of each participant

and comprised (i) venous blood sampling [for analysing serum interleukin-6 (IL-6)

concentration and creatine kinase (CK) activity]; (ii) hamstrings and quadriceps muscle

soreness via visual analogue scale; (iii) isometric maximum voluntary contraction (MVC)

torque of the quadriceps, as well as both voluntary and involuntary muscle activation and

torque-frequency relationship via electrical stimulation] MVC torque of the hamstring together

with normalised BF_{LH} sEMG (see below); and (iv) treadmill running (4.17 m s^{-1}) kinematics of

the right leg (via an eight-camera motion capture system) PRE and POST48 following the

repeated maximal sprint intervention. At POST, kinematic assessments were performed first

followed by the aforementioned order of the assessment for practical reasons.

Maximal Repeated Sprint Protocol

Many athletes in team sports are required to perform repeated maximal sprints, which are

characterised by short-duration (<10 s) and relatively longer recovery times (>60 s) between

maximal sprint bouts, and have a different physiological demand compared to repeated-sprint

exercises with shorter recovery times (<60 s) [47,48]. Therefore, the repeated maximal sprint

intervention consisted of 15 repetitions of 30 m maximal sprints with a deceleration zone of 12

m. The 30 m distance was chosen as the upper average of both the total sprinting distance

(346 ± 115 m) of wide-midfielders and the mean recovery time (70.2 ± 25.1 s) between sprint

bouts in soccer [47,49], which allows the athlete to maintain the performance of the sprint bouts.

Similar protocols have been used elsewhere [6,7,20]. Prior to the repeated maximal sprint

intervention, a five-minute warm-up was performed, comprising jogging, dynamic stretching

and three self-paced 20 m runs at 60%, 80%, 100% of perceived top speed. During the

repeated maximal sprint intervention, the participants were instructed to sprint maximally

(verbal encouragement) and to stop within the deceleration zone. Further, they were instructed

to move slowly back to the start line and to sit on a chair for the remaining time until the next

sprint. The recovery comprised 90 s between repetitions and after every 5th repetition, the

participants were allowed to rest for 3 min. Sprinting time during trials was measured and

controlled with timing gates (Brower Timing Systems, Draper, UT, USA), which were placed

on the start and finish line. Participants started 30 cm before the start line to avoid interfering

with timing gates with the arms upon initial acceleration [21]. Further, heart rate (Polar Oy,

Kempele, Finland) and rating of perceived exertion were recorded before and after each

repetition. Participants were instructed to wear the same footwear for each testing day. As the

fastest sprint time does not necessarily need to occur at the first of the 15 sprint bouts and

there was an upsurge in speed of the final sprint, fatigue was assessed with the performance

decrement score using the following formula [19]:

457
$$Fatigue = \big(100\, X\, (total\ sprint\ time \div ideal\ sprint\ time)\big) - 100$$

Where total sprint time = sum of time from all 15 sprints; and ideal sprint time = total number

of sprints (15) x fastest repetition sprint time. The calculation of this decrement score was also

used to quantify changes in heart rate and rating of perceived exertion during the repeated

maximal sprint intervention.

Maximal Voluntary Contraction (MVC)

Three test sessions were conducted with an isokinetic dynamometer (Humac Norm, CSMI

Solutions, Massachusetts, USA). As *isokinetic* maximum voluntary contractions (MVC) torque

tests are only weak predictors of hamstring strain injury [50], we decided to focus on *isometric*

MVC quadriceps and hamstring torque at optimal knee strength angles (optimal torque-joint

angle relationship, see below) to avoid further fatiguing the participants. The torque signal was

interfaced with an acquisition system (AcqKnowledge, Biopac Systems, Santa Barbara, USA)

for analogue-to-digital conversion and sampled at a frequency of 2 kHz. The participant was

seated in an upright position and securely fastened with inextensible straps at the chest and

waist while the arms were held crossed above the chest. The tibiofemoral epicondyle was

aligned with the lever arm rotation axis, and the lever arm shin pad was strapped to the leg, 2

cm above the centre of the lateral malleolus. A Velcro strap secured the distal thigh just above

the knee. The hip joint angle was set to 85° (180° = supine position) in order to analyse the

knee flexor muscle group at a sprint specific angle associated with the late swing phase of

sprinting [51]. Participants were instructed to maximally extend and flex their leg to measure knee

range of motion. Quadriceps MVC was measured at 80° knee flexion (0° = full knee extension),

as this is the optimal joint angle for peak quadriceps MVC in healthy young men [52]. Hamstring

MVC was measured at 30° knee flexion based on this being the optimal joint angle for peak

hamstring MVC during our pilot work. Published studies during the time of data collection used

a similar angle of hamstring MVC torque [53,54]. This was also in line with sprinting kinematics,

demonstrating that maximal hamstring muscle lengths during sprinting occur during the late

swing phase when the knee is flexed between 30° and 45° [16]. Prior to isometric MVC

assessments, participants underwent a standardised warm up consisting of 10 submaximal

isokinetic leg extensions ($60°·s^{-1}$). Participants then performed three isometric knee extension

(quadriceps) and flexion (hamstring) at both joint angles (each MVC lasting 2-3 s), with 60 s

rest between MVC of a given muscle group. The highest MVC of the three attempts for each

muscle group at each angle was used for subsequent analyses. Throughout the tests,

participants received verbal encouragement and biofeedback (MVC outputs) were projected

onto a screen in front of the participant.

Hamstring Muscle Voluntary Activation

To measure hamstring muscle voluntary activation capacity via the interpolated twitch technique, stimulation electrodes (12.5 mm x 7.5 mm self-adhesive electrodes (DJO Global, California, USA) were used. The general procedure has been described elsewhere [5,52,55]. Briefly, the anode was placed proximal to the popliteal fossa, and the cathode was placed beneath the gluteal fold and slightly medial to avoid activation of the vastus lateralis. Protocols were completed with electrical stimulation pads carefully taped down during the sprinting protocol and were additionally marked on the skin with a permanent marker, to ensure a precise relocation for the POST and POST48 tests. Stimulation was delivered by a high-voltage stimulator (DS7AH; Digitimer Ltd., Welwyn Garden City, United Kingdom), and consisted of a doublet using two 240-V rectangular pulses (200 μs pulse width) with an inter-pulse duration of 10 ms (100 Hz stimulation). During each experimental session, relaxed hamstring muscles were stimulated while participants were fixed in the isokinetic dynamometer with the same setting for knee flexion MVC (85º hip angle, 30 º knee flexion). The amplitude started with 50 mA to familiarise the participants to the stimulation and was gradually increased in 20 mA increments until a plateau in doublet torque was achieved. We decided to use the individual maximal stimulation (100%) intensity despite the fact that other publications used supramaximal stimulation (110-130%) [5] as we experienced lower MVC knee flexion torque output beyond 100%. That individual amplitude (162.0 ± 17.4 mA; range: 130–200 mA) was applied during all maximal contractions in the experimental session.

The maximal doublet stimulation was used two minutes later to elicit resting maximal doublet torque in the resting state (control doublet), followed 2.5 s later by a second (superimposed) doublet during an isometric knee flexion MVC. The superimposed doublet torque was always calculated manually from careful selection and inspection of the respective time periods compared to a normal increase in voluntary torque. Voluntary activation was calculated according to the following equation:

VA (%) = [1- (superimposed doublet torque/ control doublet torque)]

Surface Electromyography and Antagonist Muscle Co-activation

Surface electromyographic (sEMG) activity was recorded from the vastus lateralis and BF_{LH} to determine the extent of antagonist muscle co-activation during MVCs of the respective muscle group. Previous reports have shown that the vastus lateralis [56] and BF_{LH}[57] are representative muscles for the quadriceps femoris and hamstring muscle group, respectively. This procedure has been reported in detail elsewhere [56]. Briefly, once the muscles were identified via palpation, and the skin surface was shaved and cleaned with 70% ethanol, two bipolar Ag-AgCl surface electrodes with an inter-electrode distance of 2 cm (Noraxon duel sEMG electrode, Noraxon, Scottsdale, USA) were placed along the sagittal axis over the muscle belly at 33% of the respective muscle length from the distal end [according to SENIAM guidelines [58]] and one reference electrode (Ambu Blue, Ambu, Copenhagen, Denmark) was positioned over the medial tibial condyle. The exact location of the electrodes were marked on the participant's skin with a permanent marker to ensure precise electrode repositioning for the following assessments.

Surface EMG signals were sampled at 2000 Hz (Biopac Systems, Santa Barbara, USA) and then band-pass filtered between 10–500 Hz (AcqKnowledge, Biopac Systems, Santa Barbara, USA). Surface EMG activity of both the agonist and antagonist muscles were analysed by calculating the root mean square of the sEMG signal of a 500-ms epoch around peak MVC. To compare BF_{LH} sEMG activity at all three time points, BF_{LH} sEMG of the hamstring MVC at 30° was normalised to the evoked maximum compound muscle action potential (M-wave) of the BF_{LH} (see below). Antagonist muscle co-activation (i.e. quadriceps activation during hamstring MVC at 30° knee flexion, or hamstring activation during quadriceps MVC at 80° knee flexion) was calculated with the following formula (where EMG_{max} is the maximum sEMG of the antagonist muscle when acting as an agonist at the same knee joint angle):

$$Antagonist\ muscle\ co-activation = \frac{EMG antagonist}{EMG max} \times 100$$

543 Torque signals, electrical stimuli, and sEMG activity were displayed on a computer screen,

544 interfaced with an acquisition system (AcqKnowledge, Biopac Systems, Santa Barbara, USA)

545 used for analogue-to-digital conversion. Due to technical issues, co-activation data were

546 available for hamstring n = 12; and quadriceps n=10.

Hamstring Muscle Maximal Compound Muscle Action Potential

The hamstring muscle group was stimulated with single square wave twitch pulses (200 µs

duration) using the same electrical stimulator and stimulating electrodes, as described above.

While the participant sat resting on the isokinetic dynamometer with the knee angle set at 30°

knee flexion, compound muscle action potentials (M-waves) were evoked with 10 to 20 mA

incremental amplitudes until a maximal M-wave (M_{max}) was achieved. The average amplitude

necessary to evoke a maximal M-wave was 166.8 ± 19.8 mA; range: 130–210 mA). The

maximal M-wave was defined as the mean peak-to-peak sEMG response from the three

highest observed M-waves. Due to inter-individual differences in subcutaneous fat and the

(re)location of small sEMG electrodes over a relatively large muscle belly, sEMG amplitude is

notoriously variable [59]. To reduce this inter- and intra-individual variability, we normalised

absolute BF_{LH} sEMG to the individual's BF_{LH} maximal M-wave, determined at each testing

session [60]. Due to technical issues, data of BF_{LH} sEMG normalised to maximal M-wave was

only available for n = 13.

Torque-frequency Relationship

The torque-frequency relationship was determined by stimulating the hamstring muscle group

with single square wave twitch pulses (200 µs duration) at 1, 10, 15, 20, 30, 50 and 100 Hz for

1 s each in a random order and with 15 s rest between each stimulation, using the same

electrical stimulator and stimulating electrodes (and location), as described above. The

stimulus intensity for 100-Hz stimulation was the amplitude necessary to elicit ~20% knee

flexion MVC torque at PRE, and the same amplitude was used for the same test at POST and

48POST. The absolute peak torque at each frequency was normalised to the peak torque at

100 Hz for each time point (PRE, POST and POST48).

Delayed Onset Muscle Soreness

Using a visual analogue scale that consisted of a 100 mm line (scale 0-10 cm; 0 cm=no

soreness; 10 cm= unbearably painful), in conjunction with both a three-repetition bilateral squat

(predominantly to determine quadriceps femoris muscle soreness) [61] and lunges

(predominantly to determine hamstring muscle soreness), participants rated their perceived

lower limb muscle soreness along the muscle length immediately after each movement.

Muscle soreness was also measured by recording the force required to elicit tenderness at

nine fixed sites on the skin over the quadriceps (distal, central and proximal locations of the

three superficial quadriceps heads, vastus lateralis, vastus medialis and rectus femoris) and

six sites on the hamstrings (distal, central and proximal locations of both BF_{LH} and the medial

hamstrings), which were previously marked with a permanent marker to ensure the same

measuring position PRE, POST and POST48. At each site, a gradually increasing force was

applied by the investigator with an algometer (FPK/FPN Mechanical Algometer, Wagner

Instruments, Greenwich, USA) with a maximum of 10 kg/cm^2. Lying in the prone position with

the hip and knee fully extended and muscles relaxed, the participant was asked to indicate

when the sensation of pressure changed to discomfort, and the force at that point was recorded.

Ultrasound

Architectural parameters of the BF_{LH} were assessed using B-mode ultrasound imaging.

Participants were in the prone position with the hip and knee fully extended and muscles

relaxed. The BF_{LH} was investigated, as this muscle is the most commonly injured hamstring

muscle in team sports, particularly during sprinting [2]. Longitudinal and cross-sectional

panoramic ultrasound images of the right BF_{LH} were obtained (Philips EPIQ 7 Ultrasound

System, Bothel, USA). The linear transducer (5-18 MHz; aperture 38.9 mm) was carefully

placed on the skin with transmission gel and BF_{LH} was scanned (i) longitudinally from its distal

(=0% muscle length) to proximal (=100% muscle length) myotendinous junction along a line

drawn with a permanent marker to mark the central pathway between the medial and lateral

aspects of the muscle (incorporating the intra-muscular aponeurosis (Figure **2**2A); and (ii)

cross-sectionally at 20, 40, 60 and 80% along the total muscle length, measured on the skin

using a tape measure (Seca, Hamburg, Germany) (Figure 2B).

All images were analysed offline (ImageJ, version 1.51s, National Institutes of Health,

Bethesda, USA). Two images for each of the four cross-sectional points were recorded and

the image of best quality was used to calculate BF$_{LH}$ muscle volume. The volume of the

muscular portion between every two consecutive scans was calculated with the following

equation:

604
$$Volume = \frac{1}{3} * d * \left(a + \sqrt{(ab) + b} \right)$$

Where *a* and *b* are the anatomical cross-sectional areas of the muscle of two consecutive

cross-sectional scans and *d* is the interval distance between the cross-sectional area

measurements. The volume of the entire muscle was calculated by summing up all of the inter-

scan muscular volumes [62]. Two full-length sagittal images were then recorded to allow for the

measurement of resting BF$_{LH}$ muscle fascicle length and pennation angle, which were both

assessed in 3 fascicles at 50% of the total length of BF$_{LH}$. This point was measured offline

(ImageJ). A comparison between offline (sagittal ultrasound) and tape measurements of the

total BF$_{LH}$ length revealed a very high correlation (R^2=0.96, P<0.001). Fascicle length was

measured by tracing the fascicular path from the upper aponeurosis to the intra-muscular

aponeurosis of the BF$_{LH}$. Muscle fascicle pennation angle was determined as the angle

between the muscle fascicular paths and their insertion into the intra-muscular aponeurosis.

The mean of the three measurements for each parameter were used to determine both fascicle

length and pennation angle of the BF$_{LH}$ muscle. PCSA was calculated by dividing BF$_{LH}$ volume

by its fascicle length. During the time of data collection, a similar methodological approach was

published elsewhere [63]. One longitudinal image of one participant in the present study was not

analysed due to low image quality. Ultrasound scans and image analysis was performed by

the same investigator.

Kinematic and Kinetic Data

Three-dimensional kinematic and kinetic data were synchronously collected at 500 Hz using

an eight-camera motion analysis system (Qqus 300+; Qualisys, Gothenburg, Sweden). The

data were filtered with a digital dual low-pass Butterworth filter at 20 Hz for motion, as

previously described [64]. Retroreflective markers (12 mm diameter) were placed on anatomical

landmarks on the right leg and pelvis, as previously described [65]. One standing static and two

functional motion calibration trials were recorded of the participant PRE, POST and POST48.

For the static trial, participants stood with their feet approximately shoulder width apart and

knees fully extended. This static trial determined local coordinate systems, the location of joint

centres, and the foot, shank, thigh, and pelvis segment lengths of each participant. The

functional trials defined functional hip joint centres [66] and knee joint axes [67]. Kinematic data

were tracked using Qualisys Track Manager Software (Qualisys). Data processing and

analysis were undertaken in Visual3D (C-Motion, Germantown, MD). To examine any changes

between the time points, joint angles were normalised relative to the static trial of the

accompanying time point for minimising the influence of potential slightly different marker

positions between the trials. Lower extremity 3D joint angles and angular velocities were

calculated using an X-Y-Z Cardan angle rotation sequence. Investigated variables included

peak knee and hip angles, as well as range of motion and time during stance and swing phase

of the treadmill run, for all three planes, were calculated as described in previous studies [64].

Motorised Treadmill Run

Participants ran on a motorised treadmill (HP Cosmos Pulsar; Nussdorf, Germany) for 30 s at

4.17 m s^{-1} ($0°$ incline), as high-speed running was of interest. The selected speed was based

on pilot testing, which demonstrated that 15 km/h (4.17 m/s) was the fastest speed on a

motorized treadmill where the participants still felt comfortable. Motion analysis data were

recorded for the last 10 s of the run and data were analysed for at least 6 consecutive strides.

Peak knee and hip angle data, for all three planes, were calculated (i) between the initial

contact and terminal stance of foot; and (ii) between initial and terminal swing phase. The

touchdown of the foot during the treadmill run was determined from the kinematic data as

occurring at the local minima and the toe-off during running as the local maxima of the vertical

velocity of the head of the fifth metatarsal marker on the foot [68].

Blood Samples

A 10 mL blood sample was drawn from an antecubital vein in the forearm and collected into a

serum vacutainer (BD Vacutainer systems, Plymouth, UK). The blood samples were obtained

at each time point and left at 22-24°C for 30 min to allow clotting, and then kept on ice when

necessary. Serum samples were centrifuged at 1,300 g for 15 min at 4°C. All samples were

then aliquoted into 1.5 mL microcentrifuge tubes [Axygen (Corning), New York, USA] and

stored at -80°C until subsequent analysis (see below).

Serum Interleukin-6 (IL-6) Concentration

Serum samples were assayed for IL-6 concentration using commercially available human IL-

6 enzyme linked immunosorbent assay kits (Quantikine®, R&D systems, Minneapolis, MN,

USA) according to the manufacturer's instructions. The intensity of the colour produced after

20 min was measured with a Thermo Multiskan Spectrum microplate reader (Thermo Fisher

Scientific. Waltham, MA. USA) at 450 nm and values were calculated with Excel 365 (Microsoft,

v. 365, USA) by generating a four-parameter logistic curve fit. The minimum detectable dose

of human IL-6 was 0.70 pg/mL.

Serum Creatine Kinase Activity

Creatine kinase (CK) activity was assayed using a commercially available CK assay

(Catachem Inc., Connecticut, NE, USA), as described in detail elsewhere [31]. Briefly, 10 μL

blood serum were loaded onto a 96-well UV plate. The CK reaction reagent and diluent

(Catachem) were prepared as per the manufacturer's instructions and added to the samples

and the change in absorbance monitored continuously over 20 min in a Thermo Multiskan

Spectrum plate reader (Thermo Fisher Scientific. Waltham, MA. USA) at a wavelength of 340

nm.

Capillary Blood Lactate Concentration

676　Capillary blood samples were taken from the finger-tip via a Safety-Lancet Extra 18G needle

677　(Sarstedt; Nümbrecht, Germany) at rest before and immediately after the repeated maximal

678　sprint intervention. Blood samples were analysed within 60 seconds of collection using a

679　portable blood lactate analyser (Arkray Lactate Pro; Kyoto, Japan).

Reagents, Chemicals, and Solvents for Muscle Cell Culture *in vitro*

Growth media used for the expansion of human muscle-derived cell populations consisted of

Hams F-10 nutrient mix (Lonza, Basel, Switzerland) with added L-glutamine (2.5 mM), 10%

heat-inactivated fetal bovine serum (hiFBS; Gibco, Thermo Fisher Scientific, Altincham, UK),

1% penicillin-streptomycin (Life Technologies, Warrington, UK), and 1% L-Glutamine (Gibco).

Differentiation media consisted of α-MEM (Lonza), 1% hiFBS, 1% penicillin-streptomycin, and

1% L-glutamine. Phosphate-buffered saline (PBS; Sigma-Aldrich) was used to wash cell

monolayers. Desmin polyclonal rabbit anti-human antibody (Cat# ab15200, RRID: AB_301744)

was used (1:200) from Abcam (Abcam, Cambridge, UK), and secondary antibody (TRITC

polyclonal goat anti-rabbit; Cat# A16101, RRID: AB_2534775) was used (1:200) from Fisher

Scientific.

Muscle Biopsy Procedure

692　Participants were instructed to avoid any strenuous exercise 48 h prior to the biopsy procedure.

693　Biopsies from the vastus lateralis muscle were obtained under local anaesthesia from each

694　participant, using the Weil-Blakesley conchotome technique as described previously [69]. The

695 conchotome was inserted through the incision into the muscle belly to obtain the 134 ± 82.7

696 mg muscle biopsy.

Extraction of Human Muscle-Derived Cells

The muscle biopsies analysed in this study were isolated and cultured [31], as reported

previously. Briefly, biopsy samples were transferred with precooled growth media from the

muscle biopsy suite to the sterile tissue culture hood (Kojair Biowizard Silverline class II hood;

Kojair, Vippula, Finland) within 40 min and muscle biopsy samples were washed three times

with ice-cold PBS (0.01 M phosphate buffer, 0.0027 M KCl, and 0.137 M NaCl, pH 7.4, in

dH2O). Visible fibrous and fat tissue was removed using sterile scissor and forceps. Samples

were cut in small pieces (1 mm^3) and digested in 5 ml of trypsin-EDTA for 15 min on a magnetic

stirring platform at 37°C to dissociate muscle cells. The trypsinisation process was repeated

twice. Supernatant derived following each treatment was collected and pooled with hiFBS at a

concentration of 10% of the total volume to inhibit further protease activity. Cell supernatant

was centrifuged at 450 g for 5 min. Supernatant was discarded and the cell pellet was

resuspended in growth media and plated on a T25 cm^2 culture flask (Corning, Life Sciences,

New York, USA) for cell population expansion. Culture flasks were previously coated with a 2

mg/l porcine gelatin solution (90–110 g, Bloom; Sigma-Aldrich, Dorset, UK) to support cell

adhesion.

Expansion of Extracted Cells

The medium was refreshed on the fourth day after the extraction procedure and subsequently

every 48 h following two brief washes with PBS. Cells were incubated in a HERAcell 150i CO_2

Incubator (Thermo Scientific, Cheshire, UK). T25 cm^2 culture flasks reached 80% confluence

after aproximately 10 days and were passaged via trypsinisation. Cells were counted using

Trypan Blue exclusion and re-plated on gelatinised T75 cm^2 culture flasks (Nunc, Roskilde,

Denmark). The cells were expanded until passage 3 and then frozen in GM with 10% dimethyl

sulfoxide (DMSO) in liquid N_2 as a cryopreservant. All experiments were performed on cells

between passages 3 and 6 to avoid potential issues of senescence [25].

Characterization of Human Muscle-Derived Cells

The mixed population of human skeletal muscle-derived myoblasts and fibroblasts were

characterised by immunofluorescent staining at passage 3 (about 7-10 days after the muscle

stem cell isolation procedure before we froze the remaining cells down) for the detection of

desmin expressed by myoblasts (desmin positive) and non-myoblasts (desmin negative) to

determine the percentage of myoblasts and fibroblasts. Grohmann, et al. [24] showed that

passaging does not change the percentage of myoblast and fibroblasts and all populations

were included for analysis. Previous investigations have determined that the non-myoblast

(desmin negative) fraction is highly enriched in fibroblasts, with up to 99% of this fraction being

fibroblasts [70,71], as our group also observed [31]. Therefore, non-myoblasts (desmin negative)

were referred to here as fibroblasts.

Monolayers were incubated with 25% [vol/vol methanol in Tris-buffered saline (10 mmTris-HCl,

pH 7.8, 150 mm NaCl)], 50% and 100% for 5-min to fix the cells and stored at 4°C wet in Tris-

buffered saline until further analysis. Fixed monolayers were permeabilised and blocked for 2

h with 5% goat serum and 0.2% Triton X-100 in Tris-buffered saline, prior to staining. Cells

were incubated overnight at 4°C with anti-Desmin antibody (1:200). After overnight incubation,

the primary antibody was removed, and the cells were washed three times with Tris-buffered

saline. Secondary TRITC polyclonal goat anti-rabbit antibody (1:200) was then applied and

incubated for 2 h at 4°C. Fluorescent images were captured using live imaging microscopy

(Leica DMB 6000; Magnification x 10.5) and analysed via ImageJ cell counter plug-in. A total

of four randomly selected areas per well were analysed per individual.

Wound-Healing Assay, Migration and Differentiation Analysis

For the wound healing assay, 100,000 cells/ml were seeded in gelatinised six-well plates (Nunc,

Roskilde, Denmark). Cells were expanded as described above until cell monolayers reached

a confluent state, Growth media was removed, monolayers were washed with PBS and cells

were damaged by a vertical scrape with a 1-ml pipette tip (width of the wound area, *mean ±*

S.E.M.: 896.4 ± 21.24 *µ*m), as previously reported by our group [31]. PBS was aspirated,

damaged cell monolayers were washed twice with PBS to remove cell debris and 2 ml

differentiation media was added. Monolayers were imaged with a live imaging microscopy

(Leica) for the analysis of cell migration immediately, 24h, and 48h. TIF images were exported

from Leica Application Suite and loaded as TIF image stacks in ImageJ with a cell counter

plug-in. Cells in the outer and inner segments were then counted (Figure 3A).

Damaged monolayers were imaged at two sites per well in the wound site immediately post-

damage (0 h). These image coordinates were tracked and stored to allow subsequent

monitoring of the same sites on the wound to reduce this experimental bias. Captured images

were exported as TIF image files, and analysed in ImageJ.

Statistical Analysis

One-way repeated-measures analysis of variances (ANOVA)s were performed to determine

whether there was a significant main effect for time (within subject factor) for the following

dependent variables: MVC torque, voluntary muscle activation, antagonist muscle co-

activation, muscle soreness (for squat lunge via measured via visual analogue scale as well

as algometer), rating of perceived exertion, CK activity, IL-6 concentration, and for kinematics

data (hip and knee angle parameters). MVC torque data were analysed for interactions and

main effects for muscle group and time using two-way mixed design ANOVAs, comparing

differences between muscle groups across 3-time points; PRE, POST, and POST48. For

within test comparisons, either, independent t-tests, or one-way ANOVAs were used where

appropriate. For the torque-frequency relationship, normalised torque at each frequency was

analysed using a two-way repeated measures ANOVA, with stimulation frequency (1-100 Hz)

and time (PRE, POST and POST48) as the within-groups independent variables. Post-hoc

one-way repeated measures ANOVAs were used to determine if the normalised torque at each

frequency differed between time points. Bivariate correlations were used to analyse the relation

between architectural parameters of the BF_{LH} (volume, fascicle length, fascicle pennation angle

and PCSA) and fatigue biomarkers (relative MVC loss normalised to PRE MVC), serum CK

activity, serum IL-6 concentration, muscle soreness, knee joint range of motion or changes in

range of motion during treadmill running.

Bivariate correlations were used to analyse the relation between myoblast:fibroblast ratio and

quadriceps and hamstring MVC, and migration dynamics (total cell migration, cell proportion

of inner to outer segment) of the muscle stem cells. Standard guidelines concerning violation

of the sphericity assumption to adjust the degree of freedom of the F-test by the Huynh-Felt

epsilon if epsilon is greater than 0.75 and to use the more stringent Greenhouse-Geisser

adjustment if epsilon is less than 0.75 were followed. Results were expressed as mean ± SD,

unless otherwise stated, with statistical significance set at $P<0.05$. All MVC data were analysed

with AcqKnowledge software 4.4 (Biopac-Systems Inc., Goleta, USA) and SPSS 23 Software

(IBM Inc., Armonk, NY: IBM Corp) was used for statistical analysis. Occasional missing data

are reflected in the reported degrees of freedom.

GRANTS

788 This study was supported by a Liverpool John Moores University PhD studentship (P.B.) and

789 the Wellcome Trust Biomedical Vacation Scholarships (R.M.E., M.S.).

DISCLOSURES

791 No conflicts of interest, financial or otherwise, are declared by the authors.

AUTHOR CONTRIBUTION

793 R.M.E, P.B., B.D., C.E.S., and M.L. conceived and designed the research; P.B., S.T., M.S.,
794 M.C., J.A.S., and S.O.S performed the experiments; P.B., S.T. and R.M.E analyzed the data;
795 R.M.E., P.B, M.L. and C.E.S. interpreted the results of the experiments; P.B. prepared the
796 figures; P.B. drafted the manuscript; R.M.E., P.B., S.T., M.S., M.L., B.D., C.E.S., M.C., J.A.S.,
797 and S.O.S edited and approved the final version of the manuscript.

DATA SHARING

The data that support the findings of this study are available from the corresponding author upon reasonable request.

REFERENCES

Crema, M. D. *et al.* Imaging-detected acute muscle injuries in athletes participating in the Rio de Janeiro 2016 Summer Olympic Games. *Br. J. Sports Med.*, bjsports-2017-098247 (2017).

Ekstrand, J., Hägglund, M. & Waldén, M. Epidemiology of muscle injuries in professional football (soccer). *The American journal of sports medicine* **39**, 1226-1232 (2011).

Opar, D. A., Williams, M. D. & Shield, A. J. Hamstring strain injuries. *Sports Med* **42**, 209-226 (2012).

Byrne, C., Twist, C. & Eston, R. Neuromuscular function after exercise-induced muscle damage. *Sports Med* **34**, 49-69 (2004).

Marshall, P. W., Lovell, R., Jeppesen, G. K., Andersen, K. & Siegler, J. C. Hamstring muscle fatigue and central motor output during a simulated soccer match. *PLoS One* **10**, e102753 (2014).

Timmins, R. G. *et al.* Reduced biceps femoris myoelectrical activity influences eccentric knee flexor weakness after repeat sprint running. *Scandinavian journal of medicine & science in sports* **24**, e299-e305 (2014).

Verma, S., Moiz, J., Shareef, M. & Husain, M. Physical performance and markers of muscle damage following sports specific sprints in male collegiate soccer players: repeated bout effect. *The Journal of sports medicine and physical fitness* (2015).

Newham, D., McPhail, G., Mills, K. & Edwards, R. Ultrastructural changes after concentric and eccentric contractions of human muscle. *J. Neurol. Sci.* **61**, 109-122 (1983).

Stauber, W., Clarkson, P., Fritz, V. & Evans, W. Extracellular matrix disruption and pain after eccentric muscle action. *J. Appl. Physiol.* **69**, 868-874 (1990).

Wang, N., Tytell, J. D. & Ingber, D. E. Mechanotransduction at a distance: mechanically coupling the extracellular matrix with the nucleus. *Nature reviews Molecular cell biology* **10**, 75-82 (2009).

Howatson, G. & Van Someren, K. A. The prevention and treatment of exercise-induced muscle damage. *Sports Med* **38**, 483-503 (2008).

Tidball, J. G. Mechanisms of muscle injury, repair, and regeneration. *Comprehensive Physiology* **1**, 2029-2062 (2011).

Mackey, A. L., Magnan, M., Chazaud, B. & Kjaer, M. Human skeletal muscle fibroblasts stimulate in vitro myogenesis and in vivo muscle regeneration. *J. Physiol.* **595**, 5115-5127, doi:10.1113/jp273997 (2017).

Murphy, M. M., Lawson, J. A., Mathew, S. J., Hutcheson, D. A. & Kardon, G. Satellite cells, connective tissue fibroblasts and their interactions are crucial for muscle regeneration. *Development* **138**, 3625-3637 (2011).

Joe, A. W. *et al.* Muscle injury activates resident fibro/adipogenic progenitors that facilitate myogenesis. *Nat. Cell Biol.* **12**, 153 (2010).

Thelen, D. G. *et al.* Hamstring muscle kinematics during treadmill sprinting. *Medicine and science in sports and exercise* **37**, 108-114 (2005).

Timmins, R. G. *et al.* Short biceps femoris fascicles and eccentric knee flexor weakness increase the risk of hamstring injury in elite football (soccer): a prospective cohort study. *Br. J. Sports Med.* **50**, 1524-1535, doi:10.1136/bjsports-2015-095362 (2016).

18 Paschalis, V. *et al.* The effects of muscle damage following eccentric exercise on gait
biomechanics. *Gait & posture* **25**, 236-242 (2007).
19 Glaister, M., Howatson, G., Pattison, J. R. & McInnes, G. The reliability and validity of fatigue
measures during multiple-sprint work: an issue revisited. *J. Strength Cond.* **22**, 1597-1601
(2008).
20 Chen, C.-H., Ye, X., Wang, Y.-T., Chen, Y.-S. & Tseng, W.-C. Differential Effects of Different
Warm-up Protocols on Repeated Sprints-Induced Muscle Damage. *J. Strength Cond.* (2017).
21 Howatson, G. & Milak, A. Exercise-induced muscle damage following a bout of sport specific
repeated sprints. *J. Strength Cond.* **23**, 2419-2424 (2009).
22 Timmins, R. G., Shield, A. J., Williams, M. D., Lorenzen, C. & Opar, D. A. Architectural
adaptations of muscle to training and injury: a narrative review outlining the contributions by
fascicle length, pennation angle and muscle thickness. *Br. J. Sports Med.*, bjsports-2015-
094881 (2016).
23 Owens, D. J. *The Role of Vitamin D in Skeletal Muscle Function and Regeneration*, Liverpool
John Moores University, (2015).
24 Grohmann, M. *et al.* Isolation and validation of human prepubertal skeletal muscle cells:
maturation and metabolic effects of IGF-I, IGFBP-3 and TNFα. *J. Physiol.* **568**, 229-242 (2005).
25 Foulstone, E. J., Huser, C., Crown, A. L., Holly, J. M. & Stewart, C. E. Differential signalling
mechanisms predisposing primary human skeletal muscle cells to altered proliferation and
differentiation: roles of IGF-I and TNFα. *Experimental cell research* **294**, 223-235 (2004).
26 Evangelidis, P. *et al.* The functional significance of hamstrings composition: is it really a
'fast'muscle group? *Scandinavian journal of medicine & science in sports* **27**, 1181-1189
(2016).
27 Terry, E. E. *et al.* Transcriptional profiling reveals extraordinary diversity among skeletal
muscle tissues. *eLife* **7** (2018).
28 Randolph, M. E. & Pavlath, G. K. A muscle stem cell for every muscle: variability of satellite
cell biology among different muscle groups. *Front. Aging Neurosci.* **7**, 190 (2015).
29 Jakobsen, J. R. *et al.* Remodeling of muscle fibers approaching the human myotendinous
junction. *Scandinavian journal of medicine & science in sports* (2018).
30 Collins, C. A. *et al.* Stem cell function, self-renewal, and behavioral heterogeneity of cells
from the adult muscle satellite cell niche. *Cell* **122**, 289-301 (2005).
31 Owens, D. J. *et al.* A systems-based investigation into vitamin D and skeletal muscle repair,
regeneration, and hypertrophy. *Am. J. Physiol.-Endoc. M.* **309**, E1019-E1031 (2015).
32 Jones, D., Newham, D., Round, J. & Tolfree, S. Experimental human muscle damage:
morphological changes in relation to other indices of damage. *J. Physiol.* **375**, 435-448 (1986).
33 Friden, J., Sjöström, M. & Ekblom, B. Myofibrillar damage following intense eccentric exercise
in man. *Int. J. Sports Med.* **4**, 170-176 (1983).
34 Balsom, P., Seger, J., Sjödin, B. & Ekblom, B. Maximal-intensity intermittent exercise: effect of
recovery duration. *Int. J. Sports Med.* **13**, 528-528 (1992).
35 Onishi, H. *et al.* EMG-angle relationship of the hamstring muscles during maximum knee
flexion. *Journal of Electromyography and Kinesiology* **12**, 399-406 (2002).
36 Schuermans, J., Van Tiggelen, D., Danneels, L. & Witvrouw, E. Biceps femoris and
semitendinosus—teammates or competitors? New insights into hamstring injury
mechanisms in male football players: a muscle functional MRI study. *Br. J. Sports Med.* **48**,
1599-1606 (2014).
37 Chumanov, E. S., Heiderscheit, B. C. & Thelen, D. G. The effect of speed and influence of
individual muscles on hamstring mechanics during the swing phase of sprinting. *Journal of
biomechanics* **40**, 3555-3562 (2007).
38 Mendiguchia, J., Alentorn-Geli, E., Idoate, F. & Myer, G. D. Rectus femoris muscle injuries in
football: a clinically relevant review of mechanisms of injury, risk factors and preventive
strategies. *Br. J. Sports Med.* **47**, 359-366 (2013).

39　Malone, S., Roe, M., Doran, D. A., Gabbett, T. J. & Collins, K. High chronic training loads and
exposure to bouts of maximal velocity running reduce injury risk in elite Gaelic football.
Journal of Science and Medicine in Sport **20**, 250-254 (2017).

40　Higashihara, A., Ono, T., Kubota, J., Okuwaki, T. & Fukubayashi, T. Functional differences in
the activity of the hamstring muscles with increasing running speed. *Journal of sports*
sciences **28**, 1085-1092 (2010).

41　Alentorn-Geli, E. *et al.* Prevention of non-contact anterior cruciate ligament injuries in soccer
players. Part 1: Mechanisms of injury and underlying risk factors. *Knee Surgery, Sports*
Traumatology, Arthroscopy **17**, 705-729, doi:10.1007/s00167-009-0813-1 (2009).

42　Franchi, M. V. & Maffiuletti, N. A. Distinct modalities of eccentric exercise: different recipes,
not the same dish. *J. Appl. Physiol.* **0**, null, doi:10.1152/japplphysiol.00093.2019 (2019).

43　Turrina, A., Martínez-González, M. A. & Stecco, C. The muscular force transmission system:
role of the intramuscular connective tissue. *J. Bodywork Movement Ther.* **17**, 95-102 (2013).

44　Balius, R. *et al.* Role of the Extracellular Matrix in Muscle Injuries: Histoarchitectural
Considerations for Muscle Injuries. *Orthopaedic Journal of Sports Medicine* **6**,
2325967118795863, doi:10.1177/2325967118795863 (2018).

45　Valle, X. *et al.* Muscle injuries in sports: a new evidence-informed and expert consensus-
based classification with clinical application. *Sports Med* **47**, 1241-1253 (2017).

46　Wüst, R. C., Morse, C. I., De Haan, A., Jones, D. A. & Degens, H. Sex differences in contractile
properties and fatigue resistance of human skeletal muscle. *Experimental physiology* **93**, 843-
850 (2008).

47　Bradley, P. S., Di Mascio, M., Peart, D., Olsen, P. & Sheldon, B. High-intensity activity profiles
of elite soccer players at different performance levels. *J. Strength Cond.* **24**, 2343-2351
(2010).

48　Girard, O., Mendez-Villanueva, A. & Bishop, D. Repeated-Sprint Ability—Part I. *Sports Med*
41, 673-694 (2011).

49　Bradley, P. S. *et al.* High-intensity running in English FA Premier League soccer matches.
Journal of sports sciences **27**, 159-168 (2009).

50　van Dyk, N. *et al.* Hamstring and Quadriceps Isokinetic Strength Deficits Are Weak Risk
Factors for Hamstring Strain Injuries A 4-Year Cohort Study. *The American journal of sports*
medicine, 0363546516632526 (2016).

51　Guex, K., Gojanovic, B. & Millet, G. P. Influence of hip-flexion angle on hamstrings isokinetic
activity in sprinters. *Journal of athletic training* **47**, 390-395, doi:10.4085/1062-6050-47.4.04
(2012).

52　Erskine, R. M., Jones, D. A., Maganaris, C. N. & Degens, H. In vivo specific tension of the
human quadriceps femoris muscle. *Eur. J. Appl. Physiol.* **106**, 827-838 (2009).

53　Nedelec, M. *et al.* The influence of soccer playing actions on the recovery kinetics after a
soccer match. *J. Strength Cond.* **28**, 1517-1523 (2014).

54　Kirk, E. A. & Rice, C. L. Contractile function and motor unit firing rates of the human
hamstrings. *J. Neurophysiol.*, jn. 00620.02016 (2016).

55　Erskine, R. M., Jones, D. A., Williams, A. G., Stewart, C. E. & Degens, H. Resistance training
increases in vivo quadriceps femoris muscle specific tension in young men. *Acta physiologica*
199, 83-89 (2010).

56　Reeves, N. D., Narici, M. V. & Maganaris, C. N. In vivo human muscle structure and function:
adaptations to resistance training in old age. *Experimental physiology* **89**, 675-689 (2004).

57　Kellis, E. & Baltzopoulos, V. In vivo determination of the patella tendon and hamstrings
moment arms in adult males using videofluoroscopy during submaximal knee extension and
flexion. *Clinical Biomechanics* **14**, 118-124 (1999).

58　Hermens, H. J., Freriks, B., Disselhorst-Klug, C. & Rau, G. Development of recommendations
for SEMG sensors and sensor placement procedures. *Journal of electromyography and*
Kinesiology **10**, 361-374 (2000).

59 Araujo, R. C., Duarte, M. & Amadio, A. C. On the inter-and intra-subject variability of the
electromyographic signal in isometric contractions. *Electromyogr. Clin. Neurophysiol.* **40**, 225-
230 (2000).
60 Lanza, M. B., Balshaw, T. G., Massey, G. J. & Folland, J. P. Does normalization of voluntary
EMG amplitude to MMAX account for the influence of electrode location and adiposity?
Scand J Med Sci Sports, doi:10.1111/sms.13270 (2018).
61 Scott, J. & Huskisson, E. Vertical or horizontal visual analogue scales. *Ann. Rheum. Dis.* **38**,
560 (1979).
62 Erskine, R. *et al.* The individual and combined effects of obesity-and ageing-induced systemic
inflammation on human skeletal muscle properties. *Int. J. Obes. (Lond.)* (2016).
63 Seymore, K. D., Domire, Z. J., DeVita, P., Rider, P. M. & Kulas, A. S. The effect of Nordic
hamstring strength training on muscle architecture, stiffness, and strength. *Eur. J. Appl.*
Physiol. **117**, 943-953 (2017).
64 Verheul, J., Clansey, A. C. & Lake, M. J. Adjustments with running speed reveal
neuromuscular adaptations during landing associated with high mileage running training.
Journal of applied physiology **122**, 653-665, doi:10.1152/japplphysiol.00801.2016 (2017).
65 McClay, I. & Manal, K. Three-dimensional kinetic analysis of running: significance of
secondary planes of motion. *Medicine and Science in Sports and Exercise* **31**, 1629-1637
(1999).
66 Schwartz, M. H. & Rozumalski, A. A new method for estimating joint parameters from motion
data. *Journal of biomechanics* **38**, 107-116 (2005).
67 Robinson, M. A. & Vanrenterghem, J. An evaluation of anatomical and functional knee axis
definition in the context of side-cutting. *Journal of biomechanics* **45**, 1941-1946 (2012).
68 Maiwald, C., Sterzing, T., Mayer, T. & Milani, T. Detecting foot-to-ground contact from
kinematic data in running. *Footwear Science* **1**, 111-118 (2009).
69 Baczynska, A. M. *et al.* Human Vastus Lateralis Skeletal Muscle Biopsy Using the Weil-
Blakesley Conchotome. *Journal of Visualized Experiments*, e53075-e53075 (2016).
70 Agley, C. C., Rowlerson, A. M., Velloso, C. P., Lazarus, N. R. & Harridge, S. D. Human skeletal
muscle fibroblasts, but not myogenic cells, readily undergo adipogenic differentiation.
Journal of cell science, jcs. 132563 (2013).
71 Stewart, J. D. *et al.* Characterization of proliferating human skeletal muscle-derived cells in
vitro: Differential modulation of myoblast markers by TGF-β2. *J. Cell. Physio.* **196**, 70-78
(2003).

TABLES

981 *Table 1 Effect of the repeated maximal sprint intervention on measures of muscle activation. Data are*
982 *presented as mean ± SD. One-way ANOVA, F- and P-values are reported.*

Assessment [unit]	n	PRE	POST	POST48	F-Test	P Value
Hamstring Muscle Voluntary Activation [ITT, %]	20	98.5 ± 2.64	94.1 ± 7.83	96.9 ± 5.96	F(1.4,38) = 2.75	0.099
Normalised BF_{LH} knee flexion $sEMG_{max}$ [%]	14	3.32 ± 1.33	2.27 ± 0.72*	2.85 ± 1.16	F(2,24) = 4.35	**0.022**

Vastus lateralis knee extension sEMG$_{max}$ [mV]	15	0.50 ± 0.29	0.47 ± 0.34	0.51 ± 0.34	F(2,24) = 0.17	0.772
Quadriceps CoA during 30° hamstring MVC [%]	12	5.74 ± 7.23	4.08 ± 6.86	4.33 ± 8.76	F(2,22) = 0.50	0.613
Hamstring CoA during 80° quadriceps MVC [%]	10	4.79 ± 3.37	6.62 ± 3.79	7.51 ± 4.16	F(2,18) = 1.64	0.223

ITT – interpolated twitch technique; BF$_{LH}$ – Biceps femoris long head; CoA – Co-Activation; sEMG – surface Electromyography. * Different to PRE (P<0.05).

984 **Table 2** Effect of the Repeated Maximal Sprint Intervention on Muscle Damage-Biomarkers. Values
985 are mean ± SD. One-way ANOVA, F- and P-values are reported.

Assessment [unit]	PRE	POST	POST48	F-Test	P Value
Quadriceps MVC [N·m]	270.5 ± 51.6*	222.4 ± 52.5*	243.0 ± 71.3*	F(2,38) = 16.55	**<0.001**
Hamstring MVC [N·m]	142.5 ± 25.0*	124.8 ± 29.9*	112.4 ± 30.1*	F(2,38) = 25.12	**<0.001**
Squat Muscle soreness [cm]	0.20 ± 0.41*	1.95 ± 1.61*	2.87 ± 1.71*	F(2,38) = 28.62	**<0.001**
Lunge Muscle soreness [cm]	0.30 ± 0.57	2.30 ± 2.08†	3.48 ± 2.07†	F(2,38) = 17.02	**<0.001**
Range of Motion [°]	120.3 ± 6.76	115.7 ± 6.77†	116.0 ± 6.27†	F(2,38) = 9.33	**<0.001**
CK activity [mU/mL]	27.9 ± 23.3	53.8 ± 45.3†	99.3 ± 104.5†	F(1.3,34) = 5.98	**0.017**
IL-6 concentration [pg/mL]	1.89 ± 3.10	7.68 ± 9.95#	1.59 ± 3.46	F(1.3,34) = 5.96	**0.018**

MVC – Maximal voluntary contraction; CK – Creatine Kinase; IL-6 – Interleukin-6; * Significant differences between all time points; † differences PRE–to-POST and PRE-to-POST48 (P<0.05); # differences PRE-to-POST and POST-to-POST48 (P<0.05).

986 **Table 3** *Effect of the Repeated Maximal Sprint Intervention on Kinematics of Treadmill Running*
987 *at 4.17 m s⁻¹ Values are mean ± SD. One-way ANOVA, F- and P-values are reported.*

Kinematics [unit]	PRE	POST	POST48	F-Test	P Value
Peak knee flexion (swing phase) [°]	-103 ± 13.9	-110 ± 12.4	-108 ± 12.6	$F_{(2,20)} = 2.84$	0.082
Peak knee extension (swing phase) [°]	-3.66 ± 5.32	-7.08 ± 5.07*	-4.29 ± 7.06	$F_{(2,20)} = 3.57$	**0.047**
Contact hip flexion (toe strike) [°]	25.2 ± 5.42	29.3 ± 7.62	17.6 ± 15.6	$F_{(1.2,20)} = 4.05$	0.062
Contact knee flexion (toe strike) [°]	-15.0 ± 6.25	-17.6 ± 6.79	-13.3 ± 9.64	$F_{(2,22)} = 2.79$	0.083
Duration Running cycle [s]	0.67 ± 0.03	0.68 ± 0.03	0.69 ± 0.02	$F_{(2,20)} = 2.88$	0.080
Stance phase duration [s]	0.18 ± 0.02	0.19 ± 0.02	0.19 ± 0.03	$F_{(2,20)} = 1.01$	0.356
Swing Phase [s]	0.50 ± 0.04	0.50 ± 0.05	0.50 ± 0.03	$F_{(1.2,20)} = 0.03$	0.899

Knee fully extended=0°; negative value indicates a flexed knee; * different to PRE (P<0.05).

LEGENDS TO FIGURES

Figure 1 (A) *Torque-frequency relationship, all frequencies were normalised to 100 Hz. * significant*
differences between before (PRE) and immediadtely after (POST) the repeated maximal sprint
intervention, P<0.05; # significant differences between PRE and POST, and between PRE and POST48,
P<0.05. Data are presented as mean ± SEM. (B) Comparison of relative maximal voluntary contraction
(MVC) loss between hamstring and quadriceps muscle group before (PRE), immediately after (POST)
*and 48h after (POST48) the repeated maximal sprint intervention. * signifcant differences compared to*
PRE, P< 0.001; # significant differences between quadriceps and hamstring MVC, P< 0.05. Data are
expressed as mean ± SEM.

Figure 2 (A) *Longitudinal image of biceps femoris long head, assessment of the biceps femoris long*
head is highlighted (total muscle length and fascicle length together with pennation angle at 50% of total
muscle length). (B) Cross-sectional image at 60% muscle length (=100% proximal myotendinous
junction), biceps femoris long head is highlighted. (C) Correlation between biceps femoris long head
muscle physiological cross-sectional area and % hamstring maximum isometric voluntary contraction
(MIVC) decrease from before (PRE) to immediately after (POST) the repeated maximal sprint
intervention (P=0.003).

Figure 3 (A) *Representative images for cell migration of muscle cells with a high myoblast:fibroblast*
ratio (2.4; left) and with a low percentage of myoblasts (0.3; right) into the artificial wound. The wound
area is about 900 µm in width and split into 3 x 300 µm segments (one inner and two outer segments).

Magnification is x 10.5, and scale bar is 100 μm. ***(B)*** *Inverse correlations between the myoblast:fibroblast*
ratio and the migration dynamics of 12 different primary muscle stem cells 24 h (P=0.011) and a trend
48 h (P=0.064) after the artificial wound healing assay.

Figure 4 *Inverse correlation between the myoblast:fibroblast ratio, assessed in the current in vitro study*
and the change of hamstring MVC torque measured ***(A)*** *before and 48 h after (P=0.004), and* ***(B)***
measured immediately after and 48 h after (P=0.014) (B) the repeated maximal sprint intervention.

1014

FIGURES

1015

1016

1017 ***Figure 1***

1018

1019

1020 ***Figure 2***

1021

1022 *Figure 3*

1023

1024 *Figure 4*

1025

SUPPLEMENTARY INFORMATION

1026

1027 **Table supplement 4** Architectural parameters of biceps femoris long head (mean ± SD).

Muscle length [cm]	Fascicle length [cm]	PCSA [cm²]	Volume [cm³]	Fascicle Pennation Angle [°]
27.86 ± 2.13	7.94 ± 1.38	23.4 ± 4.62	182.2 ± 29.5	12.7 ± 2.77

Fascicle length at 50% of BF$_{LH}$ muscle length; PCSA – Physiological cross-sectional area.