Gene networks and expression quantitative trait loci associated with platinum-based

chemotherapy response in high-grade serous ovarian cancer

Jihoon Choi[1], Anastasiya Tarnouskaya[2], Sean Nesdoly[2], Danai G. Topouza[1], Madhuri Koti[1],

Qing Ling Duan[1,2]

[1]Department of Biomedical and Molecular Sciences, Queen's University, Kingston, Ontario,

Canada; [2]School of Computing, Queen's University, Kingston, Ontario, Canada;

Correspondence: QL Duan (qingling.duan@queensu.ca)

Abstract

Introduction: A major impediment in the treatment of ovarian cancer is the relapse of

platinum-resistant tumors, which occurs in approximately 25% of patients. A better

understanding of the genetic mechanisms underlying response to platinum-based

chemotherapy will improve treatment efficacy.

Objectives: We used a systems biology approach to identify novel gene networks and

regulatory sequence variants associated with platinum-based chemotherapy response.

Methods: From the Cancer Genome Atlas (TCGA), we classified high-grade serous ovarian

carcinoma (HGSOC) patients who remained cancer-free 12 months following completion of

platinum-based chemotherapy as "chemo-sensitive" (N=160) and those who had cancer

recurrence within six months of chemotherapy as "chemo-resistant" (N=110). Both univariate

and multivariate analysis of gene expression microarrays identified differentially expressed

genes and co-expression networks associated with chemotherapy response. Moreover, we

integrated genomics data to determine expression quantitative trait loci (eQTL).

Results: One differentially expressed gene encoding Valosin-containing protein (VCP) and

five co-expression network modules were associated with chemotherapy response in HGSOC

patients. These genes contribute to protein processing in the endoplasmic reticulum, which

has been previously implicated in variable chemotherapy response. In addition, 192 eQTLs

were associated with co-expressed gene modules as well as genes regulating cholesterol

levels, also previously described to affect chemotherapy response.

Conclusion: This study implicates known and novel gene networks underlying response to

platinum-based chemotherapy among HGSOC patients. A better understanding of the genetic

determinants of chemotherapy response will facilitate genetic testing for predicting drug

response, which will increase treatment efficacy and identify patients for alternative therapies.

Introduction

Ovarian cancer is the most lethal gynecological malignancy and the 8[th] leading cause of cancer death in women around the world.[1] According to the Global Cancer Observatory report in 2012, ovarian cancer accounts for 3.6% of all cancer cases and 4.3% of all cancer related deaths worldwide.[2] High-grade serous ovarian carcinoma (HGSOC) is the most malignant form of ovarian cancer that accounts for up to 70% of all cases.[3] Routine diagnosis is often difficult due to the lack of mass screening methods and heterogenous manifestations of the cancer symptoms, which result in approximately 75% of patients diagnosed with advanced stages.[4] The average 5-year survival rates are 39% for Stage 3 and 17% for Stage 4 cancers.[1]

The current standard of care for ovarian cancer is aggressive cytoreductive surgery followed by platinum-based chemotherapy.[4] However, this standard of care is not effective for all patients, with approximately 25% experiencing relapse within six months following platinum-based therapy, likely due to the development of antineoplastic resistance.[5] The median survival time for recurrent ovarian cancers range from 12-24 months.[6,7] Treatment options for patients with recurrent ovarian cancer include non-platinum-based chemotherapy regimens, immunotherapy, and molecular targeted therapy.[7,8]

Ovarian cancer has a multifactorial etiology that includes genetic and non-genetic risk factors. An estimated 23% of cases are hereditary, but the majority are sporadic with multiple reported risk factors such as history of gravidity, infertility, and late age menopause.[9,10] A better understanding of the etiology of ovarian cancer, as well as the genetic mechanisms underlying variable response to platinum-based chemotherapy is needed for improved diagnosis and treatment. For example, previous studies reported that *BRCA1* and

BRCA2 genes, associated with increased risk of ovarian cancer, harbor mutations associated

with platinum drug sensitivity and survival time.[11] Similarly, tumor suppressor genes such as

RB1, *NF1*, *RAD51B*, *PTEN* have been associated with acquired chemotherapy resistance.[12]

Earlier studies have also highlighted the importance of the immune system in the treatment of

ovarian cancer. For example, loss of chemokines and disruptions to the IFN-γ pathway have

been associated with poor treatment outcomes in HGSOC paients[13] whereas the NFκB

signaling pathway and elevated expression of *STAT1* were associated with increased response

to platinum therapy.[14,15,16] However, these known genetic variations do not account for all of

the variability in chemotherapy response among HGSOC patients and there is currently no

screening method to accurately predict prognosis prior to start of chemotherapy. Thus, further

studies are necessary to determine additional modulators of chemotherapy response, which

can be used as biomarkers for genetic testing.

In this study, we classified HGSOC patients as sensitive to platinum-based

chemotherapy if they did not experience a recurrence one year after completing

chemotherapy. Our definition differs from previous studies which classified patients as

sensitive if they did not relapse after a platinum-free interval of six months or greater (The

Cancer Genome Atlas Research Network, 2011). In addition, we classified patients as

resistant if they experienced tumor progression during chemotherapy or had a recurrence

within the six months following the end of therapy. Our classification of sensitive and

resistant patients are in accordance with the standard Gynecologic Oncology Group (GOG)

criteria.[5] Consequently, we excluded patients who had a recurrence between six to 12 months

after platinum therapy. Our strategy for dichotomizing chemotherapy response using these

"extremes" of response instead of a single threshold aims to enrich for genetic differences

between resistant and sensitive patients.

80   In addition to our novel classification of chemotherapy response in ovarian cancer

patients, we applied a multivariate, systems biology analysis approach to identify and

characterize the gene networks associated with this variable therapeutic outcome. Earlier

studies largely focused on univariate analysis of gene expression data known as differential

gene expression (DGE) analysis[17,18]. This analysis assumes that each gene functions in

isolation within the genome and is limited in statistical power due to multiple testing

corrections to reduce likelihood of false positives. The multivariate approach used in this

study identifies novel connections among genes, which represent potential interactions of

genes within biological pathways or networks. Specifically, we applied Weighted Gene Co-

expression Network Analysis[19] (WGCNA), which uses an unsupervised machine-learning

algorithm to identify clusters of highly correlated or co-expressed genes. Moreover, we

correlated sequence variations with co-expressed gene networks known as expression

Quantitative trait loci (eQTLs). A better understanding of the biological mechanisms

regulating chemotherapy response will enable more effective treatments either by improving

the accuracy of genetic testing or through the identification of novel therapies for cancer

patients.

Methods

Patient Classification:

Using a cohort of 608 high-grade grade serous ovarian carcinoma (HGSOC) patients from The Cancer Genome Atlas Project (TCGA), we retrieved clinical data from the Genomic Data Commons Portal via the TCGAbiolinks R/Bioconductor package[20]. Among the HGSOC cases, 587 had available clinical data; these were filtered and subsequently classified into two distinct groups based on their response to platinum-based chemotherapy. Specifically, we focused on the time interval between a patients' last primary platinum treatment and the onset of a recurrent tumor, or progression of an existing one for those patients whose initial tumor did not undergo complete remission[17]. We excluded patients with insufficient clinical information, were administered non-platinum drugs, and those who had died for any reason within a year of initial diagnosis, as they cannot be defined conclusively as sensitive or resistant to their adjuvant chemotherapy. Of those remaining, a total of 270 patients were classified as either resistant or sensitive to platinum-based chemotherapy. Patients who developed a new tumor in less than six months following their last platinum treatment were defined as resistant (N=110). In contrast, those who did not have a recurrent tumor event more than a year after their last primary platinum treatment were defined as sensitive (N=160). Those who had a recurrent tumour event between six months to one year following therapy were excluded from the study. This strategy was used to enrich for the genetic differences between the two drug response groups.

Transcriptomics Data Processing and Analysis:

Expression Microarrays: Expression data from the Affymetrix ht_hg_u133a microarray chip was used for univariate and multivariate (co-expression) analysis. Of the 270 subjects

classified as sensitive or resistant to chemotherapy, 238 had expression array data (N=138

sensitive, 100 resistant). Array measures were adjusted with robust multi-array average

(RMA)[21], followed by quantile normalization, and log-transformation using the *affy* package

from Bioconductor[22]. Two potential outliers and two duplicated samples were removed from

the study using the *arrayQualityMetrics* package for quality control (see **Supplemental Data**

1 for normalization, transformation and quality control assessments), resulting in 135

sensitive and 99 resistant subjects remaining. Probe sets were sorted by median absolute

deviation (MAD), and the top 50% of the probes with highest variation (n=11,107) were

included for analysis. This non-specific filtering step was used to remove non-variable probes,

therefore, reducing multiple testing and likelihood of false positives.

Differential Gene Expression Analysis: The *Limma*[23] package in Bioconductor[24] was used to

identify differentially expressed genes between chemo-sensitive and resistant groups, using

linear models. We applied the false discovery rate (FDR) method for multiple testing

correction to reduce the likelihood of false-positive results. Age at diagnosis and stage of

tumor were included as covariates in this analysis (**Supplemental Data 2**).

Weighed Gene Co-expression Network Analysis (WGCNA): We performed hierarchical

clustering of genes with the R package *WGCNA*[19], grouping genes based on their similarity in

expression. This was achieved by first creating a similarity matrix using Pearson correlations

of expression among all genes. The resulting matrix was raised to a soft-thresholding power

of 9, as suggested by the soft-thresholding power estimation plot (**Supplemental Figure 1**).

This process amplifies the connection strength between genes and reduces noise in the

adjacency matrix. To account for topological similarity of genes, the number of shared

neighbors among genes was used to give weight to gene-gene connectivity. To avoid

excessive splitting of genes into smaller modules, minimum module size was set to 30, split

sensitivity (deep split) to 4, and modules with similar expression profiles were merged at a

height of 0.5 (**Supplemental Figure 2**). Using principal component analysis, we calculated

module eigengene for each co-expression cluster to summarize module expression into a

single measure. Each module eigengene was tested for association with platinum

chemotherapy response using generalized linear models, while adjusting for age at diagnosis

and stage of cancer as covariates. Finally, we used *Cytoscape*[25], an open source

bioinformatics platform, to visualize gene co-expression networks.

Gene function and pathway annotations: We used the Database for Annotation, Visualization

and Integrated Discovery (DAVID) [26] to identify biological pathways and functions that were

enriched in each significant gene co-expression module. We also screened significant genes

in the GeneMANIA[27] database to identify for functional connections reported in published

literature. Next, we searched the UCSC TFBS conservation sites track with DAVID to

identify enriched motifs of transcription factors that may co-regulate genes within the

associated clusters. Finally, we used the drug–gene interaction database (DGIdb)[28], a public

database with curation of data describing relationships between genes, chemicals, drugs, and

pathological phenotype, to identify genes with prior reported associations with

chemotherapeutic agents.

Genomics Data Processing and Analysis:

Genomics Data: Single nucleotide polymorphisms (SNPs) data from the germline tissues

(blood derived normal or solid tissue normal, based on availability) were obtained from the

TCGA legacy database. The Affymetrix Genome-Wide Human SNP Array 6.0 was used to

capture genetic variations, which detected 906,600 SNPs. Of the 270 subjects classified as

resistant or sensitive to platinum-based chemotherapy, 266 (157 sensitive and 109 resistant)

had genotype data available.

Imputations: The imputation of autosomal chromosomes was performed using the Michigan

imputation server pipeline[29]. We used the 1000 Genome Project phase 3 sequencing data

(version 5)[30] reference panel for the imputation of missing genotypes. We then used Eagle

v.2.3[31] for phasing of the genotypes to their respective chromosomes. For the imputation of

variants on the X chromosome, SHAPEIT[32] was used for phasing in combination with the

1000 genomes project phase 3 (version 5) reference panel (**Supplemental Data 3**).

Quality control:

Subject level: Two pairs of individuals had a relatedness coefficient (pi-hat) > 0.9, which are

likely duplicated samples. One subject from each pair was randomly removed from the

dataset. Next, inbreeding coefficients (F) were computed for each subject using PLINK[33]. A

total of 18 subjects with high homozygosity (F>0.05) or heterozygosity (F<-0.05) rates were

excluded. Moreover, sex was computed based on heterozygosity rates (F) of the X

chromosome. Four subjects whose predicted gender was undefined (F>0.2) were removed

from the study.

SNP level: SNPs with minor allele frequencies (MAF) less than 1% or with genotyping call

rate less than 90% were removed. This step removed 38,430,595 SNPs with MAF < 0.01,

resulting in 9,528,963 SNPs to be used for further analysis.

Genome-wide Association Study: After imputations and quality control filtering, 240 subjects

(N= 142 sensitive, 98 resistant) and a total of 9,528,963 SNPs (MAF > 0.1) remained for

analysis (**Supplemental Data 4**). Plink (v.1.90) was applied with association analysis of

platinum-based chemotherapy response using a logistic regression method. Age at diagnosis

and tumor stages were included as covariates.

 Expression Quantitative Trait Loci (eQTL) Analysis:

191   Common SNPs (MAF > 0.01) were tested for association with gene expression of individual

192   genes and module eigengenes using the *matrixeQTL* R package[34]. Correlated loci indicated

193   potential regulatory function of the SNPs on the expression of the nearby corresponding gene,

194   known as cis-expression Quantitative Trait Loci (cis-eQTL). Cis-eQTL were defined as

195   correlated SNPs within 1 Mb from the gene transcriptional start site (TSS).

197   Results

Differential gene expression (DGE) analysis returned one significant probe mapping to

Valosin Containing Protein (*VCP*) with FDR adjusted p-value < 0.05 (**Figure 1**) as well as

606 probes mapping to 521 unique genes with nominal significance (unadjusted p-value <

0.05; **Supplemental Table 1**). Functional annotation analysis showed that the 521 genes were

enriched for many oncogenic processes, such as cellular response to DNA damage stimulus

(GO:0006974), DNA repair (GO:0006281), positive regulation of cell growth (GO:0030307),

and positive regulation of apoptotic process (GO:0043065). Pathway analysis identified many

immune related pathways, such as mitogen-activated protein kinase (MAPK) pathway and B-

Cell antigen receptor (BCR) signaling pathway. A detailed list of functional annotations and

identified pathways of differentially expressed genes can be found in **Supplemental Data 5**.

The hierarchical clustering of genes using *WGCNA* resulted in 86 unique modules of

co-expressed genes (**Supplemental Table 2**). Each module was assessed for association with

chemotherapy response (**Supplemental Figure 3**). Five gene clusters (*honeydew1, lightcyan1,*

lightpink3, orangered4, and skyblue3) were identified to be correlated with sensitivity to

chemotherapy (p < 0.05); one gene cluster remained significant after Bonferroni correction

(*honeydew1*, p-value=7e-05) (**Figure 2**). The significant modules were annotated using

DAVID, which identified gene enrichment for biological pathways including apoptosis,

negative regulation of the Wnt signaling pathway, transcription, immune responses, and DNA

double-strand break processing involved in repair via single-strand annealing. GeneMANIA

analysis showed that genes in these modules were previously reported in 49 publications,

some of which documented associations with oncogenic pathways and chemotherapeutic

outcomes. We searched in these genes in the gene-drug interaction database (DGIdb) and

found that 35 were associated with chemotherapeutic agents such as carboplatin and

paclitaxel. Finally, we identified over-represented conserved transcription factor binding sites

located in genes from each module. A detailed list of functional annotations, transcription

factors and pathways related to gene modules can be found in **Supplemental Data 6.**

224   Our genome-wide association study (GWAS) of SNPs did not identify significantly

correlated variants for platinum chemotherapy response. The Manhattan plot (**Figure 3**),

shows that all SNPs fell below the genome-wide significance threshold ($p < 5 \times 10^{-8}$), as

indicated by the red horizontal line. As this may due to insufficient statistical power, we

performed a targeted association analysis on two well-known genes associated with

chemotherapeutic outcomes in ovarian cancer: *BRCA1* and *BRCA2*. Of the 237 SNPs in

BRCA1 and 256 in *BRCA2*, we identified 34 SNPs in *BRCA2* and 1 SNP in *BRCA1* that were

significantly associated with chemotherapy response (**Supplemental Table 3**).

232   Next, SNPs were tested for correlation with expression of the 606 differentially

233 expressed probes and 5 co-expression modules. This identified 5,242 cis-eQTL associated

234 with the expression of differentially expressed probes, and 192 cis-eQTL associated with co-

235 expression networks (**Supplemental Data 7**).

Discussion

In this manuscript, we identified novel genes and gene networks correlated with variable

response to platinum-based chemotherapy in HGSOC patients. Using a univariate analysis

approach, we identified a differentially expressed locus encoding the valosin-containing

protein (*VCP*) associated with sensitivity to platinum-based chemotherapy. In addition, we

applied a multivariate, co-expression analysis method for identifying groups of

interconnected genes that could contribute to common biological pathways. This method

identified 5 clusters of co-expressed genes nominally correlated with chemo-response, one of

which remained significant after correction of multiple testing. Genes in these modules have

been annotated to be associated with biological pathways such as apoptosis, transcription,

immune response, negative regulation of the Wnt signaling pathway and DNA double-strand

break processing involved in repair via single-strand annealing.

The most significantly associated probe identified in the DGE analysis was for a

gene encoding Valosin-containing protein (*VCP*, p = 3.91E-06). *VCP* plays a critical role in

disintegrating large polypeptide cellular structures for further degradation by proteolytic

enzymes. It functions to regulate important pathways of DNA repair, replication and cell

cycle progression by removing faulty polypeptide structures from chromatin material,

ribosomes, endoplasmic reticulum and mitochondria. This gene has been previously

identified as a potential biomarker for predicting the success of platinum-based chemotherapy

in lung cancer patients.[35] Drugs that inhibit *VCP*, such as the novel inhibitor CB-5083[36], are

currently being evaluated for their prospective anticancer properties.

Many of the nominally correlated genes from our DGE analysis are enriched for

pathways known to be critical for oncogenesis and chemotherapeutic drug resistance. For

example, probe 213532_at mapping to the A Disintegrin and Metalloproteinase-17 (*ADAM17*)

gene was upregulated in the chemo-resistant group (p = 0.017). This is consistent with prior

studies, which reported that overexpression of *ADAM17* results in reduced cisplatin-induced

apoptosis in hepatocellular carcinoma cells and may contribute to cisplatin resistance via the

EGFR pathway[37]. Moreover, we identified another up-regulated probe 205239_at (p = 0.003)

for the gene encoding Amphiregulin (AREG), a protein found in the EGFR signaling pathway,

which has been reported to promote ovarian cancer stemness and drug resistance to anti-

cancer therapy.[38] Taken together, our findings support a potential role of the AREG-EGFR

signaling pathway in the development of platinum-resistance in ovarian carcinoma.

It is also important to note that one of the most enriched pathways from our DGE

analysis was the protein processing in endoplasmic reticulum (hsa04141) pathway. Using

DAVID, we have identified 18 differentially expressed probes mapping to 15 unique genes

within this pathway. Prior studies have highlighted that endoplasmic reticulum (ER) stress

may cause cisplatin resistance in ovarian carcinoma by inducing autophagy in cancer cells,

allowing them to escape apoptosis.[39,40] Findings from this pathway analysis suggest that

future studies of the contribution of ER stress in cisplatin sensitivity is needed to improve our

understanding of platinum-resistance in ovarian cancer.

In our co-expression network analysis, the gene module "*honeydew1*" showed the

most significant correlation with chemotherapy response (p = 6.53e-05). This module

includes two probes that map to *VCP*, a gene that was associated with platinum-based

chemotherapy response in our DGE analysis. Additional genes in this module were associated

with positive regulation of mitochondrial membrane potential, protein ubiquitination, mitosis,

alternative splicing, and apoptotic processes. Pathway analysis showed that this module is

involved in protein processing in the endoplasmic reticulum. A prior study has found that

VCP is critical for extraction and degradation of unfolded proteins in endoplasmic reticulum

is crucial for cancer cell survival and lower expression of *VCP* is associated with poor

response to platinum☐based chemotherapy.[41] In alignment with this finding, genes co-

expressed in the *honeydew1* module were co-downregulated in chemo-resistant patients.

The *honeydew1* module is composed of 76 probes mapping to 54 unique genes, and

of these, 44 genes are found to be in chromosome 9, demonstrating the importance of

chromosome 9 in the regulation of chemo-resistance in ovarian cancer. These findings

support previous studies, where genetic imbalance and alterations in chromosome 9 have

been associated with progression of ovarian cancer and increased cisplatin resistance.[42]

Analysis of overrepresented transcription factor binding sites demonstrated that genes in this

module may be co-regulated by a common transcription factor known as organic cation

transporter 1 (OCT1). We found that over 96% of genes in this module (49/54 genes) contain

a nucleotide motif bound by OCT1. Prior studies have reported that silencing OCT1 impaired

cisplatin-induced apoptosis in esophageal cancer cells, and that cisplatin-resistant cells were

already expressing significantly reduced levels of OCT1.[43] Taken together, these findings

characterize a network of co-expressed genes that is associated with platinum-resistance in

ovarian cancer. Genes within this module may be co-regulated by the OCT1 transcription

factor, which may be used as a novel potential target for ovarian cancer therapies

The other four co-expression modules associated with platinum-based response also

include genes known to be involved in oncogenic process and drug response outcomes. For

example, the *orangered4* module is composed of genes that are critical for regulation of

immune response, which have been shown to play a role in chemotherapy response in

HGSOC.[1415] Genes in this module are associated with functional annotation terms including

immunoglobulin receptor binding, antigen binding, B cell receptor signaling pathway, and

phagocytosis (DAVID). In addition, 10 of the 58 genes in this module are enriched for a

common transcription factor binding site: acute myeloid leukemia 1 protein (AML1). This

transcription factor is involved in haematopoiesis process and immune functions such as

thymic T-cell development. Studies have reported that the AML1 transcription factor is

overexpressed in ovarian cancer patients, and may contribute to cancer cell proliferation,

migration and invasion.[44] In addition, we found that the *lightpink3* module is strongly

associated with transcription regulation process, which plays a pivotal role in cancer

progression. Finally, genes in modules *lightcyan1* and *skyblue3* are target regions of

chemotherapeutic agents such as tyrosine kinase inhibitors (**Supplemental data 6**).

317   The analysis of *BRCA1* and *BRCA2* SNPs demonstrated that 28 out of 35 variants

318   associated with chemotherapy response were also cis-acting eQTLs, correlated with the

319   expression of *BRCA2* as well as neighboring genes *N4BP2L1*, *N4BP2L2*, *FRY*, and *STARD13*

320   (nominal p-value <0.05). Both *BRCA2* and *STARD13* are well known tumor-suppressors, and

321   upregulation of *N4BP2L1* and *N4BP2L2* is associated with positive prognosis in ovarian

322   cancer cases.[45] This finding shows the potential regulatory effect of the variants in *BRCA2*.

323   In addition, our results show that 47% of variants identified in *BRCA2* are associated

with LDL/HDL cholesterol levels (**Supplemental Table 3**). Prior studies of lung and ovarian

cancers consistently reported that cholesterol levels may affect the efficacy of platinum-based

chemotherapeutic agents.[46,47] Our findings indicate a new link between genetic variants in

BRCA2 and platinum-based chemotherapy response through cholesterol level regulation.

Conclusion

In this study, we identified genes and gene networks correlated with platinum-based

chemotherapy response in high-grade serous ovarian cancer patients, which implicate known

and novel biological mechanisms. Specifically, we identified that reduced expression of *VCP*

is associated with platinum-resistance. This gene is critical for removing unfolded proteins

from the endoplasmic reticulum and previously correlated with cancer cell survival and

platinum-based chemotherapy response. In addition, we report potentially regulatory variants

in the *BRCA2* gene correlated with chemotherapy response and the expression of genes that

determine cholesterol levels. Moreover, we identified a novel group of potentially co-

regulated genes on chromosome 9 that are correlated with platinum resistance using a

machine-learning algorithm. In addition, genes from this module including *VCP* are also

involved in protein processing in the endoplasmic reticulum. This manuscript supports earlier

studies which implicated *VCP* and *BRCA2* genes in chemotherapy response. We also

identified regulatory variants in *BRCA2* and additional genes co-regulated with *VCP* on

chromosome 9 that also contribute to protein removal in the ER. Findings from our study

could facilitate genetic testing through the identification of gene signatures that may predict

chemotherapy response as well as lead to novel drug targets, given a better understanding of

the biological mechanisms underlying chemotherapy response.

Figure Legends

Figure 1. Gene co-expression modules correlated with platinum-based chemotherapy

response. Network plot showing the five significant gene co-expression modules from

WGCNA: honeydew1 - centre, lightcyan1 - left, lightpink3 - top, orangered4 - bottom, and

skyblue3 - right. Nodes represent probes and edges are connections among the probes. Co-

expressed probes (i.e. belonging to a single module) are indicated in the same colour.

Figure 2. Differential expression analysis of platinum-based chemotherapy response in

HGSOC patients. Volcano plot showing univariate analysis results. One probe, 208648_at,

which maps to the Valosin-Containing Protein (*VCP*) gene is significantly correlated with

chemotherapy outcome as indicated by the red line (FDR-corrected p value < 0.05). A total of

606 probes mapping to 521 unique genes with nominal correlate as indicated by the green

line (p = 0.05).

Figure 3. GWAS of chemotherapy response. Manhattan plot shows no SNPs are

significantly associated (i.e. surpasses the Red line indicating a p-value threshold of 5x10^-8).

Acknowledgements

Computations in this manuscript were performed on resources and with support provided by

the Centre for Advanced Computing (CAC) at Queen's University in Kingston, Ontario. The

CAC is funded by: the Canada Foundation for Innovation, the Government of Ontario, and

Queen's University. JC is funded by Queen's University, Faculty of Health Sciences Dean's

Doctoral Award. QLD receives funding from the Canadian Institutes of Health Research and

Queen's University.

Conflict of interest

The authors declare no conflicts of interest.

Author contribution

J.C. performed the data analyses and drafted the manuscript. D.G.T., S.N. and A.T. assisted in

the data analyses. Q.L.D. designed the research project, supervised data analyses and assisted

in the writing of the manuscript. M.K. assisted in the study design and editing of the

manuscript.

References

1. Reid, F. World Ovarian Cancer Coalition 2018. *World Ovarian Cancer Coalit.* (2018).

2. International Agency for Research on Cancer Website. Globocan 2012 - Home.
Webpage (2012). doi:NO:11

3. Brett M., R. *et al.* Epidemiology of ovarian cancer: a review. *Cancer Biol. Med.* (2017).
doi:10.20892/j.issn.2095-3941.2016.0084

4. Cannistra, S. A. Cancer of the Ovary. *N. Engl. J. Med.* **351**, 2519–2529 (2004).

5. Miller, D. S. *et al.* Phase II Evaluation of Pemetrexed in the Treatment of Recurrent or
Persistent Platinum-Resistant Ovarian or Primary Peritoneal Carcinoma: A Study of
the Gynecologic Oncology Group. *J. Clin. Oncol.* **27**, 2686–2691 (2009).

6. Armstrong, D. K. Relapsed ovarian cancer: challenges and management strategies for
a chronic disease. *Oncologist* **7 Suppl 5**, 20–28 (2002).

7. Foley, O. W., Rauh-Hain, J. A. & del Carmen, M. G. Recurrent epithelial ovarian
cancer: an update on treatment. *Oncology (Williston Park).* **27**, 288–94, 298 (2013).

8. Ozols, R. F. Recurrent ovarian cancer: Evidence-based treatment. *Journal of Clinical*
Oncology **20**, 1161–1163 (2002).

9. Walsh, T. *et al.* Mutations in 12 genes for inherited ovarian, fallopian tube, and
peritoneal carcinoma identified by massively parallel sequencing. *Proc. Natl. Acad. Sci.*
(2011). doi:10.1073/pnas.1115052108

10. Booth, M., Beral, V. & Smith, P. Risk factors for ovarian cancer: A case-control study.
Br. J. Cancer (1989). doi:10.1038/bjc.1989.320

11. Vencken, P. M. L. H. *et al.* Chemosensitivity and outcome of BRCA1- and BRCA2-
associated ovarian cancer patients after first-line chemotherapy compared with
sporadic ovarian cancer patients. *Ann. Oncol.* (2011). doi:10.1093/annonc/mdq628

12. Patch, A. M. *et al.* Whole-genome characterization of chemoresistant ovarian cancer.
Nature (2015). doi:10.1038/nature14410

13. Hao, D. *et al.* Immunogenomic analyses of advanced serous ovarian cancer reveal
immune score is a strong prognostic factor and an indicator of chemosensitivity. *Clin.*
Cancer Res. (2018). doi:10.1158/1078-0432.CCR-17-3862

14. Koti, M. *et al.* A distinct pre-existing inflammatory tumour microenvironment is
associated with chemotherapy resistance in high-grade serous epithelial ovarian cancer.
Br. J. Cancer (2015). doi:10.1038/bjc.2015.81

15. Au, K. K. *et al.* STAT1-associated intratumoural TH1 immunity predicts
chemotherapy resistance in high-grade serous ovarian cancer. *J. Pathol. Clin. Res.*
(2016). doi:10.1002/cjp2.55

16. Koti, M. *et al.* Identification of the IGF1/PI3K/NF κB/ERK gene signalling networks
associated with chemotherapy resistance and treatment response in high-grade serous
epithelial ovarian cancer. *BMC Cancer* **13**, 549 (2013).

17. Cancer, T. & Atlas, G. Integrated genomic analyses of ovarian carcinoma. *Nature* **474**,
609–15 (2011).

18. Li, J., Wood, W. H., Becker, K. G., Weeraratna, A. T. & Morin, P. J. Gene expression
response to cisplatin treatment in drug-sensitive and drug-resistant ovarian cancer cells.
Oncogene **26**, 2860–2872 (2007).

19. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network
analysis. *BMC Bioinformatics* **9**, 559 (2008).

20. Colaprico, A. *et al.* TCGAbiolinks: An R/Bioconductor package for integrative
analysis of TCGA data. *Nucleic Acids Res.* (2016). doi:10.1093/nar/gkv1507

21. Irizarry, R. A. *et al.* Exploration, normalization, and summaries of high density
oligonucleotide array probe level data. *Biostatistics* **4**, 249–264 (2003).

22. Gautier, L., Cope, L., Bolstad, B. M. & Irizarry, R. A. Affy - Analysis of Affymetrix
GeneChip data at the probe level. *Bioinformatics* **20**, 307–315 (2004).

23. Smyth, G. limma: Linear Models for Microarray Data. in *Bioinformatics and*
Computational Biology Solutions Using R and Bioconductor 397–420 (2005).
doi:citeulike-article-id:5722720

24. Gentleman, R. *et al.* Bioconductor: open software development for computational
biology and bioinformatics. *Genome Biol.* **5**, R80 (2004).

25. Shannon, P. *et al.* Cytoscape: A software Environment for integrated models of
biomolecular interaction networks. *Genome Res.* **13**, 2498–2504 (2003).

26. Huang, D. W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis
of large gene lists using DAVID bioinformatics resources. *Nat. Protoc.* **4**, 44–57
(2009).

27. Warde-Farley, D. *et al.* The GeneMANIA prediction server: Biological network
integration for gene prioritization and predicting gene function. *Nucleic Acids Res.* **38**,
(2010).

28. Cotto, K. C. *et al.* DGIdb 3.0: a redesign and expansion of the drug–gene interaction
database. *Nucleic Acids Res.* (2017). doi:10.1093/nar/gkx1143

29. Das, S. *et al.* Next-generation genotype imputation service and methods. *Nat. Genet.*
(2016). doi:10.1038/ng.3656

30.   The 1000 Genomes Project Consortium. A global reference for human genetic
448   variation. *Nature* **526**, 68–74 (2015).

449   31.   Loh, P. R. *et al.* Reference-based phasing using the Haplotype Reference Consortium
450   panel. *Nat. Genet.* (2016). doi:10.1038/ng.3679

451   32.   Delaneau, O. *et al.* Integrating sequence and array data to create an improved 1000
452   Genomes Project haplotype reference panel. *Nat. Commun.* (2014).
453   doi:10.1038/ncomms4934

454   33.   Purcell, S. *et al.* PLINK: A tool set for whole-genome association and population-
455   based linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

456   34.   Shabalin, A. A. Matrix eQTL: Ultra fast eQTL analysis via large matrix operations.
457   *Bioinformatics* **28**, 1353–1358 (2012).

458   35.   Peng, J. *et al.* VCP gene variation predicts outcome of advanced non-small-cell lung
459   cancer platinum-based chemotherapy. *Tumor Biol.* **34**, 953–961 (2013).

460   36.   Lan, B., Chai, S., Wang, P. & Wang, K. VCP/p97/Cdc48, A Linking of Protein
461   Homeostasis and Cancer Therapy. *Curr. Mol. Med.* (2018).
462   doi:10.2174/1566524018666180308111238

463   37.   Wang, X. J., Feng, C. W. & Li, M. ADAM17 mediates hypoxia-induced drug
464   resistance in hepatocellular carcinoma cells through activation of EGFR/PI3K/Akt
465   pathway. *Mol. Cell. Biochem.* (2013). doi:10.1007/s11010-013-1657-z

466   38.   Tung, S. L. *et al.* miRNA-34c-5p inhibits amphiregulin-induced ovarian cancer
467   stemness and drug resistance via downregulation of the AREG-EGFR-ERK pathway.
468   *Oncogenesis* (2017). doi:10.1038/oncsis.2017.25

469   39.   Tian, J., Liu, R. & Qu, Q. Role of endoplasmic reticulum stress on cisplatin resistance
470   in ovarian carcinoma. *Oncol. Lett.* (2017). doi:10.3892/ol.2017.5580

471   40.   Xu, Y. *et al.* Tolerance to endoplasmic reticulum stress mediates cisplatin resistance in
472   human ovarian cancer cells by maintaining endoplasmic reticulum and mitochondrial
473   homeostasis. *Oncol. Rep.* (2015). doi:10.3892/or.2015.4283

474   41.   Bastola, P., Neums, L., Schoenen, F. J. & Chien, J. VCP inhibitors induce endoplasmic
475   reticulum stress, cause cell cycle arrest, trigger caspase-mediated cell death and
476   synergistically kill ovarian cancer cells in combination with Salubrinal. *Mol. Oncol.*
477   (2016). doi:10.1016/j.molonc.2016.09.005

478   42.   Devlin, J., Elder, P. A., Gabra, H., Steel, C. M. & Knowles, M. A. High frequency of
479   chromosome 9 deletion in ovarian cancer: Evidence for three tumour-suppressor loci.
480   *Br. J. Cancer* (1996). doi:10.1038/bjc.1996.75

481   43.   Lin, R. *et al.* Long-term cisplatin exposure promotes methylation of the OCT1 gene in
482   human esophageal cancer cells. *Dig. Dis. Sci.* (2013). doi:10.1007/s10620-012-2424-9

483   44.   Keita, M. *et al.* The RUNX1 transcription factor is expressed in serous epithelial
484   ovarian carcinoma and contributes to cell proliferation, migration and invasion. *Cell
485   Cycle* (2013). doi:10.4161/cc.23963

45. Koussounadis, A., Langdon, S. P., Harrison, D. J. & Smith, V. A. Chemotherapy-
induced dynamic gene expression changes in vivo are prognostic in ovarian cancer. *Br.*
J. Cancer **110**, 2975–2984 (2014).

46. Wu, Y. *et al.* Cholesterol reduces the sensitivity to platinum-based chemotherapy via
upregulating ABCG2 in lung adenocarcinoma. *Biochem. Biophys. Res. Commun.*
(2015). doi:10.1016/j.bbrc.2015.01.035

47. Kim, S., Lee, M., Dhanasekaran, D. N. & Song, Y. S. Activation of LXR$\square$/β by
cholesterol in malignant ascites promotes chemoresistance in ovarian cancer. *BMC*
Cancer **18**, 1232 (2018).

Manhattan Plot

Chromosome