Gene networks and expression quantitative trait loci associated with platinum-based

chemotherapy response in high-grade serous ovarian cancer

Jihoon Choi[1], Anastasiya Tarnouskaya[2], Sean Nesdoly[2], Danai G. Topouza[1], Madhuri Koti[1],

Qing Ling Duan[1,2]

[1]Department of Biomedical and Molecular Sciences, Queen's University, Kingston, Ontario,

Canada; [2]School of Computing, Queen's University, Kingston, Ontario, Canada;

Correspondence: QL Duan (qingling.duan@queensu.ca)

Abstract

Background: A major impediment in the treatment of ovarian cancer is the relapse of

platinum-resistant tumors, which occurs in approximately 25% of patients. A better

understanding of the biological mechanisms underlying platinum-based response will

improve treatment efficacy through genetic testing and novel therapies.

Methods: Using data from high-grade serous ovarian carcinoma (HGSOC) patients in the

Cancer Genome Atlas (TCGA), we classified those who remained progression-free for 12

months following platinum-based chemotherapy as "chemo-sensitive" (N=160) and those

who had recurrence within six months as "chemo-resistant" (N=110). Univariate and

multivariate analysis of expression microarrays identified differentially expressed genes and

co-expression gene networks associated with chemotherapy response. Moreover, we

integrated genomics data to determine expression quantitative trait loci (eQTL).

Results: Differential expression of the Valosin-containing protein (VCP) gene and five co-

expression gene networks were associated with chemotherapy response in HGSOC. VCP and

the gene networks contribute to protein processing in the endoplasmic reticulum, which has

been implicated in chemotherapy response. These findings were successfully replicated using

independent replication cohort. Furthermore, 192 QTLs were associated with these gene

networks and *BRCA2* expression.

Conclusion: This study implicates both known and novel genes as well as biological

networks underlying response to platinum-based chemotherapy among HGSOC patients.

Introduction

Ovarian cancer is the most lethal gynecological malignancy and the 8^{th} leading cause of cancer death in women around the world.[1] According to the Global Cancer Observatory report in 2012, ovarian cancer accounts for 3.6% of all cancer cases and 4.3% of all cancer related deaths worldwide.[2] High-grade serous ovarian carcinoma (HGSOC) is the most malignant form of ovarian cancer that accounts for up to 70% of all cases.[3] Routine diagnosis is often difficult due to the lack of mass screening methods and heterogenous manifestations of the cancer symptoms, which result in approximately 75% of patients diagnosed with advanced stages.[4] The average 5-year survival rates are 39% for Stage 3 and 17% for Stage 4 cancers.[1]

The current standard of care for ovarian cancer is aggressive cytoreductive surgery followed by platinum-based chemotherapy.[4] However, this standard of care is not effective for all patients, with approximately 25% experiencing relapse within six months following platinum-based therapy, likely due to the development of antineoplastic resistance.[5] The median survival time for recurrent ovarian cancers range from 12-24 months.[6,7] Treatment options for patients with recurrent ovarian cancer include non-platinum-based chemotherapy regimens, immunotherapy, and molecular targeted therapy.[7,8]

Ovarian cancer has a multifactorial etiology that includes genetic and non-genetic risk factors. An estimated 23% of cases are hereditary, but the majority are sporadic with multiple reported risk factors such as history of gravidity, infertility, and late age menopause.[9,10] A better understanding of the etiology of ovarian cancer, as well as the genetic mechanisms underlying variable response to platinum-based chemotherapy is needed for improved diagnosis and treatment. For example, previous studies reported that *BRCA1* and

BRCA2 genes, associated with increased risk of ovarian cancer, harbor mutations associated

with platinum drug sensitivity and survival time.[11] Similarly, tumor suppressor genes such as

RB1, *NF1*, *RAD51B*, *PTEN* have been associated with acquired chemotherapy resistance.[12]

Earlier studies have also highlighted the importance of the immune system in the treatment of

ovarian cancer. For example, loss of chemokines and disruptions to the IFN-γ pathway have

been associated with poor treatment outcomes in HGSOC paients[13] whereas the NFκB

signaling pathway and elevated expression of *STAT1* were associated with increased response

to platinum therapy.[14,15,16] However, these known genetic variations do not account for all of

the variability in chemotherapy response among HGSOC patients and there is currently no

screening method to accurately predict prognosis prior to start of chemotherapy. Thus, further

studies are necessary to determine additional modulators of chemotherapy response, which

can be used as biomarkers for genetic testing.

The majority of earlier studies of chemotherapy response in ovarian cancer patients

used univariate analysis of gene expression data known as differential gene expression (DGE)

analysis.[17,18] This analysis assumes that each gene functions in isolation within the genome

and fails to capture the effects of complex gene-gene interactions. This study applies a

multivariate approach to identify groups of interconnected genes, which may contribute to

common biological pathways. These genes may each have modest effects that are not

detected by conventional univariate analysis. Specifically, we applied Weighted Gene Co-

expression Network Analysis[19] (WGCNA), which uses an unsupervised machine-learning

algorithm to identify clusters of highly correlated or co-expressed genes. Moreover, we

correlated sequence variations with co-expressed gene networks to identify expression

Quantitative Trait Loci (eQTLs), which are potentially regulatory variants associated with co-

expression gene networks. A better understanding of the biological mechanisms regulating

chemotherapy response will enable more effective treatments by improving the accuracy of

genetic testing and through the identification of novel therapies for cancer patients.

Methods

Patient Classification:

From the Cancer Genome Atlas (TCGA) Genomic Data Commons (GDC) portal[20], we

retrieved 587 high-grade serous ovarian carcinoma (HGSOC) patients with available clinical

data using the TCGAbiolinks R/Bioconductor package.[21] The interval between patient's last

primary platinum treatment and the onset of a recurrent tumor or progression of an existing

tumor was used as a metric for determining chemotherapy sensitivity. Patients who developed

a new tumor in less than six months following their last platinum treatment were defined as

resistant (N=110). In contrast, those who did not have a recurrent tumor event for over a year

after their last primary platinum treatment were defined as sensitive (N=160). Individuals

who had a recurrent tumor event between six months to one year following chemotherapy

were excluded from the study. This strategy for dichotomizing resistant from sensitive

patients was used to enrich for the genetic differences.

Transcriptomics Data Processing and Analysis:

Expression Microarrays: Of the 270 HGSOC subjects classified as sensitive or resistant to

chemotherapy, 238 (138 sensitive, 100 resistant) had microarray expression data available

(Affymetrix ht_hg_u133a chip) in the GDC portal. The robust multi-array average (RMA)

method[22] in the *affy* package from Bioconductor[23] was used for background correction, log-

transformation, and quantile normalization of the probe intensities. Two potential outliers and

two duplicated samples were removed from the study during the quality control step using the

arrayQualityMetrics[24] package (see **Supplemental Data 1** for steps of pre-processing),

resulting in 135 sensitive and 99 resistant HGSOC subjects in the expression set. Next,

probes were filtered using the median absolute deviation (MAD) whereby the top 50% with

highest variation (n=11,107) were selected for analysis. Thus, this non-specific filtering step

removed probes with no or low variability, which are not likely to be differentially expressed

between sensitive and resistant patients. Moreover, this step reduces the number of multiple

testing corrections and therefore, the likelihood of false positives.

Differential Gene Expression Analysis: The *Limma*[25] package in Bioconductor[26] was used to

identify differentially expressed genes between chemo-sensitive and resistant groups using

linear models. The false discovery rate (FDR) method was employed as a measure for

multiple testing correction to control for type I error. Age at diagnosis and stage of tumor

were included as covariates in this analysis (**Supplemental Data 2**).

Weighed Gene Co-expression Network Analysis (WGCNA): We performed hierarchical

clustering of genes using the R package *WGCNA*[19], which groups genes based on their

similarity in expression. This was achieved by first creating a similarity matrix using Pearson

correlations of expression among all genes. The resulting matrix was raised to a power of 9,

as suggested by the soft-thresholding power estimation plot (**Supplemental Figure 1**). This

assumes that raising the correlation matrix to a power will enrich differences between weak

and strong signals, allowing for better quantification of gene-gene interactions. In addition,

because information between two genes are often too weak and noisy to interpret, similarity

matrix is transformed to Topological Overlap Matrix (TOM), where strength of association

between a pair of genes is reinforced by the common neighbors shared by them. To avoid

excessive splitting of genes into smaller modules, minimum module size was set to 30, split

sensitivity (deep split) to 4, and modules with similar expression profiles were merged at a

height of 0.5 (**Supplemental Figure 2**). Using principal component analysis, we calculated

the module eigengene for each co-expression cluster to summarize module expression into a

single measure. Each module eigengene was tested for association with platinum

chemotherapy response using generalized linear models, while adjusting for age at diagnosis

and stage of cancer as covariates. Finally, we used *Cytoscape*[27], an open source

bioinformatics platform, to visualize gene co-expression networks.

Gene function and pathway annotations: The Database for Annotation, Visualization and

Integrated Discovery (DAVID) [28] was applied to identify biological pathways and functions

that were enriched in each significant gene co-expression module. We also screened

significant genes in the GeneMANIA[29] database to identify for functional connections

reported in published literature. Next, we searched the UCSC TFBS conservation sites track

with DAVID to identify enriched motifs of transcription factors that may co-regulate genes

within each cluster. Finally, we used Drug–Gene Interaction database (DGIdb)[30], a public

database with curation of data describing relationships between genes, chemicals, drugs, and

pathological phenotype, to identify genes with prior reported associations with

chemotherapeutic agents.

Validation of differentially expressed gene: The Kaplan–Meier plotter was used to cross-

validate the effect of differentially expressed genes in an independent ovarian cancer cohort

(Geo accession identifier: GSE9891).[31] This replication cohort included gene expression

profiling (Affymetrix Human Genome U133 Plus 2.0 Array) of 285 ovarian tumor samples.

Patients were filtered to include those with cancer histology of serous carcinoma and who

received chemotherapy containing a platin compound to allow close comparison with the

TCGA ovarian cancer cohort. This step omitted a total of 60 subjects from analysis, which

included 21 with endometrioid carcinoma cases and 43 who did not receive platinum therapy

(4 overlapping subjects). Thus, 225 patients remained for replication analysis. Patient

survival was evaluated using Cox proportional hazards model and progression-free survival

(PFS) was the primary outcome used in the replication analysis.[32]

 Validation of co-expression network:

 For validation of co-expression networks, SurvExpress database was used, which allow users

 to validate the combined effect of multiple gene expression measures with a target trait.[33]

 Same cohort and filtering steps were used for validation (GSE9891, N = 225). Survival curve

 was evaluated using Cox proportional hazards model and progression-free survival (PFS) was

 the primary outcome used in the replication analysis.

 Genomics Data Processing and Analysis:

 Genomics Data: Single nucleotide polymorphisms (SNPs) data from the germline tissues

 (DNA extracted from blood or solid non-tumor ovarian tissue) were obtained from the TCGA

 legacy database. The Affymetrix Genome-Wide Human SNP Array 6.0 was used to capture

 genetic variations, which detected 906,600 SNPs. Of the 270 subjects from TCGA classified

 as resistant or sensitive to platinum-based chemotherapy, 266 (157 sensitive and 109 resistant)

 had genotype data available.

 Imputations: The imputation of autosomal chromosomes was performed using the Michigan

 imputation server pipeline[34]. We used the 1000 Genome Project phase 3 sequencing data

 (version 5)[35] reference panel for the imputation of missing genotypes. We then used Eagle

 v.2.3[36] for phasing of the genotypes to their respective chromosomes. For the imputation of

 variants on the X chromosome, SHAPEIT[37] was used for phasing in combination with the

 1000 genomes project phase 3 (version 5) reference panel (**Supplemental Data 3**).

 Quality control:

 Subject level: Two pairs of individuals had a relatedness coefficient (pi-hat) > 0.9, which are

 likely duplicated samples. One subject from each pair was randomly removed from the

dataset. Next, inbreeding coefficients (F) were computed for each subject using PLINK[38]. A

total of 18 subjects with high homozygosity (F>0.05) or heterozygosity (F<-0.05) rates were

excluded. Moreover, genetic sex was estimated based on heterozygosity rates (F) of the X

chromosome, and four subjects who had undefined (F>0.2) genetic sex were removed from

the study.

SNP level: SNPs with minor allele frequencies (MAF) less than 1% or with genotyping call

rate less than 90% were removed. This step removed 38,430,595 SNPs with MAF < 0.01,

resulting in 9,528,963 SNPs to be used for further analysis.

Genome-wide Association Study: After imputations and quality control, 240 subjects (N= 142

sensitive, 98 resistant) and a total of 9,528,963 SNPs (MAF > 0.1) remained available for

analysis (**Supplemental Data 4**). Plink (v.1.90) was used to compute genome wide and

BRCA1/2 targeted association analysis using a logistic regression model. Age at diagnosis

and tumor stages were included as covariates.

Expression Quantitative Trait Loci (eQTL) Analysis:

Common SNPs (MAF > 0.01) were tested for association with differentially expressed genes

and gene networks using the *matrixeQTL* R package[39]. Loci correlated with nearby gene

expression indicates potential regulatory function of the SNP on the corresponding gene,

known as cis-expression Quantitative Trait Loci (cis-eQTL). Cis-eQTLs are defined as

correlated SNPs found within 1 Mb from the gene transcriptional start site (TSS).

Results

Differential gene expression (DGE) analysis identified that low expression of a probe

(208648_at) mapping to the Valosin Containing Protein (*VCP*) gene was significantly

associated with resistance to chemotherapy (FDR adjusted p-value < 0.05; **Figure 1**). In

addition, 606 probes mapping to 521 unique genes were nominally correlated with

chemotherapy response (unadjusted p-value < 0.05; **Supplemental Table 1**). The Kaplan–

Meier survival plotter demonstrated that low expression of VCP is associated with poor

progression-free survival (p = 0.015) and shorter median survival time (**Figure 2**) in an

independent ovarian serous cancer cohort following treatment with platinum antineoplastic

agents. Functional annotation analysis showed that the 521 genes were enriched for numerous

oncogenic processes, such as cellular response to DNA damage stimulus (GO:0006974),

DNA repair (GO:0006281), positive regulation of cell growth (GO:0030307), and positive

regulation of apoptotic process (GO:0043065). Pathway analysis identified many immune

related pathways including the mitogen-activated protein kinase (MAPK) and B-Cell antigen

receptor (BCR) signaling pathways. A detailed list of functional annotations and pathways are

shown in **Supplemental Data 5**.

The hierarchical clustering of genes using *WGCNA* resulted in 86 unique modules of

co-expressed genes (**Supplemental Table 2**). Each module was assessed for association with

chemotherapy response (results shown in **Supplemental Figure 3**). Five gene clusters

(*honeydew1, lightcyan1, lightpink3, orangered4, and skyblue3*) were co-downregulated in

platinum-resistant patients (p < 0.05); one gene cluster remained significant after Bonferroni

correction (*honeydew1*, p-value=7e-05) (**Figure 3**). This signal was validated using

SurvExpress in an independent cohort (Tothill et al.), which demonstrates that

downregulation of the genes in five modules result in significantly reduced survival of

patients (**Supplemental Figure 4**). The five significant modules were annotated using

DAVID, which identified gene enrichment for biological pathways including apoptosis,

negative regulation of the Wnt signaling pathway, transcription, immune responses, and DNA

double-strand break processing involved in repair via single-strand annealing. GeneMANIA

analysis showed that genes in these modules were previously reported in 49 publications,

some of which documented associations with oncogenic pathways and chemotherapeutic

outcomes. Our search of these genes in the gene-drug interaction database (DGIdb) found

that 35 were associated with chemotherapeutic agents such as carboplatin and paclitaxel.

Furthermore, we identified over-represented conserved transcription factor binding sites

located in genes from each module. A detailed list of functional annotations, transcription

factors and pathways related to gene modules can be found in **Supplemental Data 6.**

229   Our GWAS of SNPs did not identify any variants correlated for platinum

chemotherapy response after multiple testing correction. The Manhattan plot (**Supplemental**

Figure 5), demonstrates that none of the SNPs meet the genome-wide significance threshold

($p<5x10-8$), as indicated in the by the red horizontal line. This is likely due to insufficient

statistical power resulting from low number of subjects in TCGA-OV cohort. Next, we

performed a targeted association analysis of two well-known genes associated with ovarian

cancer and chemotherapeutic outcomes: *BRCA1* and *BRCA2*. Of the 238 SNPs in *BRCA1* and

256 in *BRCA2*, we identified 34 SNPs in *BRCA2* and 1 SNP in *BRCA1* that were significantly

associated with chemotherapy response (**Supplemental Table 3**).

238   Next, SNPs were tested for correlation with expression of the 606 differentially

expressed probes and 5 co-expression modules. This identified 5,242 cis-eQTL associated

with the expression of differentially expressed probes, and 192 cis-eQTL associated with co-

expression networks (**Supplemental Data 7**).

Discussion

In this manuscript, we identified novel genes and gene networks correlated with variable

response to platinum-based chemotherapy in HGSOC patients. Using a univariate analysis

approach, we identified a differentially expressed locus encoding the valosin-containing

protein (*VCP*) associated with sensitivity to platinum-based chemotherapy. In addition, we

applied a multivariate, co-expression analysis method for identifying groups of

interconnected genes that could contribute to common biological pathways. This method

identified 5 clusters of co-expressed genes nominally correlated with chemo-response, one of

which remained significant after correction of multiple testing. Genes in these modules have

been annotated to be associated with biological pathways such as apoptosis, transcription,

immune response, negative regulation of the Wnt signaling pathway and DNA double-strand

break processing involved in repair via single-strand annealing.

The most significantly associated probe identified in the DGE analysis was for a

gene encoding Valosin-containing protein (*VCP*, p = 3.91E-06). We have confirmed that this

signal is replicated in an independent ovarian cancer cohort with statistical significance

(p=0.015). *VCP* plays a critical role in disintegrating large polypeptide cellular structures for

further degradation by proteolytic enzymes. It functions to regulate important pathways of

DNA repair, replication and cell cycle progression by removing faulty polypeptide structures

from chromatin material, ribosomes, endoplasmic reticulum and mitochondria. This gene has

been previously identified as a potential biomarker for predicting the success of platinum-

based chemotherapy in lung cancer patients.[40]

Many of the nominally correlated genes from our DGE analysis are enriched for

pathways known to be critical for oncogenesis and chemotherapeutic drug resistance. For

example, probe 213532_at mapping to the A Disintegrin and Metalloproteinase-17 (*ADAM17*)

gene was upregulated in the chemo-resistant group (p = 0.017). This is consistent with prior

studies, which reported that overexpression of *ADAM17* results in reduced cisplatin-induced

apoptosis in hepatocellular carcinoma cells and may contribute to cisplatin resistance via the

EGFR pathway[41]. Moreover, we identified another up-regulated probe 205239_at (p = 0.003)

for the gene encoding Amphiregulin (AREG), a protein found in the EGFR signaling pathway,

which has been reported to promote ovarian cancer stemness and drug resistance to anti-

cancer therapy.[42] Taken together, our findings support a potential role of the AREG-EGFR

signaling pathway in the development of platinum-resistance in ovarian carcinoma.

It is also important to note that one of the most enriched pathways from our DGE

analysis was the protein processing in endoplasmic reticulum (hsa04141) pathway. Using

DAVID, we have identified 18 differentially expressed probes mapping to 15 unique genes

within this pathway. Prior studies have highlighted that endoplasmic reticulum (ER) stress

may cause cisplatin resistance in ovarian carcinoma by inducing autophagy in cancer cells,

allowing them to escape apoptosis.[43,44] Findings from this pathway analysis suggest that

future studies of the contribution of ER stress in cisplatin sensitivity is needed to improve our

understanding of platinum-resistance in ovarian cancer.

In our co-expression network analysis, the gene module "*honeydew1*" showed the

most significant correlation with chemotherapy response (p = 6.53e-05). This association

signal was validated with statistical significance (p = 5.88e-07) in an independent replication

cohort. This module includes two probes that map to *VCP*, a gene that was associated with

platinum-based chemotherapy response in our DGE analysis. Additional genes in this module

were associated with positive regulation of mitochondrial membrane potential, protein

ubiquitination, mitosis, alternative splicing, and apoptotic processes. Pathway analysis

showed that this module is involved in protein processing in the endoplasmic reticulum. A

prior study has found that *VCP* plays a crucial role in cancer cell survival through extraction

and degradation of unfolded proteins in endoplasmic reticulum, and noted that lower

expression of *VCP* was associated with poor response to platinum□based chemotherapy.[45] In

alignment with this finding, genes co-expressed in the *honeydew1* module were co-

downregulated in chemo-resistant patients.

The *honeydew1* module is composed of 76 probes mapping to 54 unique genes, and

of these, 44 genes are located in chromosome 9, demonstrating the importance of

chromosome 9 in the regulation of chemo-resistance in ovarian cancer. These findings

support previous studies, where genetic imbalance and alterations in chromosome 9 have

been associated with progression of ovarian cancer and increased cisplatin resistance.[46]

Analysis of overrepresented transcription factor binding sites demonstrated that genes in this

module may be co-regulated by a common transcription factor known as organic cation

transporter 1 (OCT1). We found that over 96% of genes in this module (49/54 genes) contain

a nucleotide motif bound by OCT1. Prior studies have reported that silencing OCT1 impaired

cisplatin-induced apoptosis in esophageal cancer cells, and that cisplatin-resistant cells were

already expressing significantly reduced levels of OCT1.[47] Taken together, these findings

characterize a network of co-expressed genes that is associated with platinum-resistance in

ovarian cancer. Genes within this module may be co-regulated by the OCT1 transcription

factor, which may be used as a novel potential target for ovarian cancer therapies

The other four co-expression modules, which we were also able to replicate in an

independent cohort, include genes known to be involved in oncogenic process and drug

response outcomes. For example, the *orangered4* module is composed of genes that are

critical for regulation of immune response, which have been shown to play a role in

chemotherapy response in HGSOC.[14,15] Genes in this module are associated with functional

annotation terms including immunoglobulin receptor binding, antigen binding, B cell receptor

signaling pathway, and phagocytosis. In addition, 10 of the 58 genes in this module are

enriched for a common transcription factor binding site: acute myeloid leukemia 1 protein

(AML1). This transcription factor is involved in haematopoiesis process and immune

functions such as thymic T-cell development. Studies have reported that the AML1

transcription factor is overexpressed in ovarian cancer patients, and may contribute to cancer

cell proliferation, migration and invasion.[48] In addition, we found that the *lightpink3* module

is strongly associated with transcription regulation process, which plays a pivotal role in

cancer progression. Finally, genes in modules *lightcyan1* and *skyblue3* are target regions of

chemotherapeutic agents such as tyrosine kinase inhibitors (**Supplemental data 6**).

Targeted analysis of *BRCA1* and *BRCA2* SNPs demonstrated that 28 out of 35

variants associated with chemotherapy response were also cis-acting eQTLs, correlated with

the expression of *BRCA2* as well as neighboring genes *N4BP2L1*, *N4BP2L2*, *FRY*, and

STARD13 (nominal p-value <0.05). Both *BRCA2* and *STARD13* are well known tumor-

suppressors, and upregulation of *N4BP2L1* and *N4BP2L2* have been associated with positive

prognosis in ovarian cancer cases.[49] This finding shows the potential regulatory effect of the

variants in *BRCA2*.

In addition, our annotation results show that 47% of platin-response associated

variants in *BRCA2* are linked with LDL/HDL cholesterol levels (**Supplemental Table 3**).

Prior studies of lung and ovarian cancers consistently reported that cholesterol levels may

affect the efficacy of platinum-based chemotherapeutic agents.[50,51] Our findings indicate a

new link between genetic variants in *BRCA2* and platinum-based chemotherapy response

through cholesterol level regulation.

Conclusion

In this study, we identified genes and gene networks correlated with platinum-based

chemotherapy response in high-grade serous ovarian cancer patients, which implicate known

and novel biological mechanisms. Specifically, we identified that reduced expression of *VCP*

is associated with platinum-resistance. This gene is critical for removing unfolded proteins

from the endoplasmic reticulum and have been correlated with cancer cell survival and

platinum-based chemotherapy response in prior studies. In addition, we report potentially

regulatory variants in the *BRCA2* gene correlated with chemotherapy response and the

expression of genes that determine cholesterol levels. Moreover, we identified a novel group

of potentially co-regulated genes on chromosome 9 that are correlated with platinum

resistance using a machine-learning algorithm. In addition, genes from this module are also

involved in protein processing in the endoplasmic reticulum. This manuscript supports earlier

studies which implicated *VCP* and *BRCA2* genes in chemotherapy response. We also

identified regulatory variants in *BRCA2* and additional genes co-regulated with *VCP* on

chromosome 9 that also contribute to protein removal in the ER. Findings from our study

could facilitate genetic testing through the identification of gene signatures that may predict

chemotherapy response as well as lead to novel drug targets, given a better understanding of

the biological mechanisms underlying chemotherapy response.

Figure Legends

Figure 1. Differential expression analysis of platinum-based chemotherapy response in

HGSOC patients. Volcano plot showing univariate association analysis results. One probe,

208648_at, which maps to the Valosin-Containing Protein (*VCP*) gene is significantly

differentially expressed and correlated with chemotherapy outcome after multiple testing

correction. Red horizontal line demonstrates FDR-corrected p value (< 0.05) threshold. A

total of 606 probes mapping to 521 unique genes are also differentially expressed with

nominal significance, as indicated by the green line ($p = 0.05$).

Figure 2. Kaplan- Meier plot of two ovarian cancer cohorts for cross-validation of

differentially expressed gene VCP. Kaplan-Meier (KM) plot shows the Progression Free

Survival (PFS) of the replication ovarian cancer cohort post platinum-based chemotherapy

treatment. Red line indicates the PFS of patients with high VCP expression, while black line

indicates the PFS of patients with low VCP expression. Patients with high expression of VCP

is associated with better PFS with statistical significance.

Figure 3. Gene co-expression modules correlated with platinum-based chemotherapy

response. Network plot displaying the five significant gene co-expression modules from

WGCNA: honeydew1 - centre, lightcyan1 - left, lightpink3 - top, orangered4 - bottom, and

skyblue3 - right. Nodes represent probes and edges are connections among the probes. Co-

expressed probes (i.e. belonging to a single module) are indicated by the same color.

Acknowledgements

Computations in this manuscript were performed on resources and with support provided by

the Centre for Advanced Computing (CAC) at Queen's University in Kingston, Ontario. The

CAC is funded by: the Canada Foundation for Innovation, the Government of Ontario, and

Queen's University. JC is funded by Queen's University, Faculty of Health Sciences Dean's

Doctoral Award. QLD receives funding from the Canadian Institutes of Health Research and

Queen's University.

Conflict of interest

The authors declare no conflicts of interest.

Author contribution

J.C. performed the data analyses and drafted the manuscript. D.G.T., S.N. and A.T. assisted in

the data analyses. Q.L.D. designed the research project, supervised data analyses and assisted

in the writing of the manuscript. M.K. assisted in the study design and editing of the

manuscript.

Data availability

Transcriptomics, genomics, and clinical data used for the analysis of TCGA-OV cohort can

be accessed/downloaded from the Genomic Data Commons (GDC) Data Portal

(https://portal.gdc.cancer.gov/). Gene expression and clinical data of the replication cohort

can be accessed/downloaded from Gene Expression Omnibus (GEO) database

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE9899).

References

1.   Reid, F. World Ovarian Cancer Coalition 2018. *World Ovarian Cancer Coalit.* (2018).

2.   International Agency for Research on Cancer Website. Globocan 2012 - Home.
Webpage (2012). doi:NO:11

3.   Brett M., R. *et al.* Epidemiology of ovarian cancer: a review. *Cancer Biol. Med.* (2017).
doi:10.20892/j.issn.2095-3941.2016.0084

4.   Cannistra, S. A. Cancer of the Ovary. *N. Engl. J. Med.* **351**, 2519–2529 (2004).

5.   Miller, D. S. *et al.* Phase II Evaluation of Pemetrexed in the Treatment of Recurrent or
Persistent Platinum-Resistant Ovarian or Primary Peritoneal Carcinoma: A Study of
the Gynecologic Oncology Group. *J. Clin. Oncol.* **27**, 2686–2691 (2009).

6.   Armstrong, D. K. Relapsed ovarian cancer: challenges and management strategies for
a chronic disease. *Oncologist* **7 Suppl 5**, 20–28 (2002).

7.   Foley, O. W., Rauh-Hain, J. A. & del Carmen, M. G. Recurrent epithelial ovarian
cancer: an update on treatment. *Oncology (Williston Park).* **27**, 288–94, 298 (2013).

8.   Ozols, R. F. Recurrent ovarian cancer: Evidence-based treatment. *Journal of Clinical*
Oncology **20**, 1161–1163 (2002).

9.   Walsh, T. *et al.* Mutations in 12 genes for inherited ovarian, fallopian tube, and
peritoneal carcinoma identified by massively parallel sequencing. *Proc. Natl. Acad. Sci.*
(2011). doi:10.1073/pnas.1115052108

10.   Booth, M., Beral, V. & Smith, P. Risk factors for ovarian cancer: A case-control study.
Br. J. Cancer (1989). doi:10.1038/bjc.1989.320

11.   Vencken, P. M. L. H. *et al.* Chemosensitivity and outcome of BRCA1- and BRCA2-
associated ovarian cancer patients after first-line chemotherapy compared with
sporadic ovarian cancer patients. *Ann. Oncol.* (2011). doi:10.1093/annonc/mdq628

12.   Patch, A. M. *et al.* Whole-genome characterization of chemoresistant ovarian cancer.
Nature (2015). doi:10.1038/nature14410

13.   Hao, D. *et al.* Immunogenomic analyses of advanced serous ovarian cancer reveal
immune score is a strong prognostic factor and an indicator of chemosensitivity. *Clin.*
Cancer Res. (2018). doi:10.1158/1078-0432.CCR-17-3862

14.   Koti, M. *et al.* A distinct pre-existing inflammatory tumour microenvironment is
associated with chemotherapy resistance in high-grade serous epithelial ovarian cancer.
Br. J. Cancer (2015). doi:10.1038/bjc.2015.81

15.   Au, K. K. *et al.* STAT1-associated intratumoural TH1 immunity predicts
chemotherapy resistance in high-grade serous ovarian cancer. *J. Pathol. Clin. Res.*
(2016). doi:10.1002/cjp2.55

16.   Koti, M. *et al.* Identification of the IGF1/PI3K/NF κB/ERK gene signalling networks
associated with chemotherapy resistance and treatment response in high-grade serous
epithelial ovarian cancer. *BMC Cancer* **13**, 549 (2013).

17.   Cancer, T. & Atlas, G. Integrated genomic analyses of ovarian carcinoma. *Nature* **474**,
609–15 (2011).

18.   Li, J., Wood, W. H., Becker, K. G., Weeraratna, A. T. & Morin, P. J. Gene expression
response to cisplatin treatment in drug-sensitive and drug-resistant ovarian cancer cells.
Oncogene **26**, 2860–2872 (2007).

19.   Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network
analysis. *BMC Bioinformatics* **9**, 559 (2008).

20.   Grossman, R. L. *et al.* Toward a shared vision for cancer genomic data. *New England*

Journal of Medicine (2016). doi:10.1056/NEJMp1607591

21.   Colaprico, A. *et al.* TCGAbiolinks: An R/Bioconductor package for integrative
analysis of TCGA data. *Nucleic Acids Res.* (2016). doi:10.1093/nar/gkv1507

22.   Irizarry, R. A. *et al.* Exploration, normalization, and summaries of high density
oligonucleotide array probe level data. *Biostatistics* **4**, 249–264 (2003).

23.   Gautier, L., Cope, L., Bolstad, B. M. & Irizarry, R. A. Affy - Analysis of Affymetrix
GeneChip data at the probe level. *Bioinformatics* **20**, 307–315 (2004).

24.   Kauffmann, A., Gentleman, R. & Huber, W. arrayQualityMetrics - A bioconductor
package for quality assessment of microarray data. *Bioinformatics* **25**, 415–416 (2009).

25.   Smyth, G. limma: Linear Models for Microarray Data. in *Bioinformatics and*
Computational Biology Solutions Using R and Bioconductor 397–420 (2005).
doi:citeulike-article-id:5722720

26.   Gentleman, R. *et al.* Bioconductor: open software development for computational
biology and bioinformatics. *Genome Biol.* **5**, R80 (2004).

27.   Shannon, P. *et al.* Cytoscape: A software Environment for integrated models of
biomolecular interaction networks. *Genome Res.* **13**, 2498–2504 (2003).

28.   Huang, D. W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis
of large gene lists using DAVID bioinformatics resources. *Nat. Protoc.* **4**, 44–57
(2009).

29.   Warde-Farley, D. *et al.* The GeneMANIA prediction server: Biological network
integration for gene prioritization and predicting gene function. *Nucleic Acids Res.* **38**,
(2010).

30.   Cotto, K. C. *et al.* DGIdb 3.0: a redesign and expansion of the drug–gene interaction
database. *Nucleic Acids Res.* (2017). doi:10.1093/nar/gkx1143

31.   Tothill, R. W. *et al.* Novel molecular subtypes of serous and endometrioid ovarian
cancer linked to clinical outcome. *Clin. Cancer Res.* (2008). doi:10.1158/1078-
0432.CCR-08-0196

32.   Gyorffy, B., Lánczky, A. & Szállási, Z. Implementing an online tool for genomewide
validation of survival-associated biomarkers in ovarian-cancer using microarray data
from 1287 patients. *Endocr. Relat. Cancer* (2012). doi:10.1530/ERC-11-0329

33.   Aguirre-Gamboa, R. *et al.* SurvExpress: An Online Biomarker Validation Tool and
Database for Cancer Gene Expression Data Using Survival Analysis. *PLoS One* (2013).
doi:10.1371/journal.pone.0074250

34.   Das, S. *et al.* Next-generation genotype imputation service and methods. *Nat. Genet.*
(2016). doi:10.1038/ng.3656

35.   The 1000 Genomes Project Consortium. A global reference for human genetic
variation. *Nature* **526**, 68–74 (2015).

36.   Loh, P. R. *et al.* Reference-based phasing using the Haplotype Reference Consortium

panel. *Nat. Genet.* (2016). doi:10.1038/ng.3679

37.   Delaneau, O. *et al.* Integrating sequence and array data to create an improved 1000
Genomes Project haplotype reference panel. *Nat. Commun.* (2014).
doi:10.1038/ncomms4934

38.   Purcell, S. *et al.* PLINK: A tool set for whole-genome association and population-
based linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

39.   Shabalin, A. A. Matrix eQTL: Ultra fast eQTL analysis via large matrix operations.
Bioinformatics **28**, 1353–1358 (2012).

40.   Peng, J. *et al.* VCP gene variation predicts outcome of advanced non-small-cell lung
cancer platinum-based chemotherapy. *Tumor Biol.* **34**, 953–961 (2013).

41.   Wang, X. J., Feng, C. W. & Li, M. ADAM17 mediates hypoxia-induced drug
resistance in hepatocellular carcinoma cells through activation of EGFR/PI3K/Akt
pathway. *Mol. Cell. Biochem.* (2013). doi:10.1007/s11010-013-1657-z

42.   Tung, S. L. *et al.* miRNA-34c-5p inhibits amphiregulin-induced ovarian cancer
stemness and drug resistance via downregulation of the AREG-EGFR-ERK pathway.
Oncogenesis (2017). doi:10.1038/oncsis.2017.25

43.   Tian, J., Liu, R. & Qu, Q. Role of endoplasmic reticulum stress on cisplatin resistance
in ovarian carcinoma. *Oncol. Lett.* (2017). doi:10.3892/ol.2017.5580

44.   Xu, Y. *et al.* Tolerance to endoplasmic reticulum stress mediates cisplatin resistance in
human ovarian cancer cells by maintaining endoplasmic reticulum and mitochondrial
homeostasis. *Oncol. Rep.* (2015). doi:10.3892/or.2015.4283

45.   Bastola, P., Neums, L., Schoenen, F. J. & Chien, J. VCP inhibitors induce endoplasmic
reticulum stress, cause cell cycle arrest, trigger caspase-mediated cell death and
synergistically kill ovarian cancer cells in combination with Salubrinal. *Mol. Oncol.*
(2016). doi:10.1016/j.molonc.2016.09.005

46.   Devlin, J., Elder, P. A., Gabra, H., Steel, C. M. & Knowles, M. A. High frequency of
chromosome 9 deletion in ovarian cancer: Evidence for three tumour-suppressor loci.
Br. J. Cancer (1996). doi:10.1038/bjc.1996.75

47.   Lin, R. *et al.* Long-term cisplatin exposure promotes methylation of the OCT1 gene in
human esophageal cancer cells. *Dig. Dis. Sci.* (2013). doi:10.1007/s10620-012-2424-9

48.   Keita, M. *et al.* The RUNX1 transcription factor is expressed in serous epithelial
ovarian carcinoma and contributes to cell proliferation, migration and invasion. *Cell*
Cycle (2013). doi:10.4161/cc.23963

49.   Koussounadis, A., Langdon, S. P., Harrison, D. J. & Smith, V. A. Chemotherapy-
induced dynamic gene expression changes in vivo are prognostic in ovarian cancer. *Br.*
J. Cancer **110**, 2975–2984 (2014).

50.   Wu, Y. *et al.* Cholesterol reduces the sensitivity to platinum-based chemotherapy via
upregulating ABCG2 in lung adenocarcinoma. *Biochem. Biophys. Res. Commun.*
(2015). doi:10.1016/j.bbrc.2015.01.035

51.   Kim, S., Lee, M., Dhanasekaran, D. N. & Song, Y. S. Activation of LXR$\square$/β by
cholesterol in malignant ascites promotes chemoresistance in ovarian cancer. *BMC*
Cancer **18**, 1232 (2018).

208648_at
(VCP)

HR = 0.69 (0.51 - 0.93)
logrank P = 0.015

Probability

Expression
— low
— high

0   50   100   150

Time (months)