SNP-based quantitative deconvolution of biological mixtures: application to the detection

of cows with subclinical mastitis by whole genome sequencing of tank milk.

Wouter Coppieters[1], Latifa Karim[1], Michel Georges[2].

[1]Genomics Platform, GIGA Institute, University of Liège. [2]Unit of Animal Genomics, GIGA

Institute & Faculty of Veterinary Medicine, University of Liège.

Correspondence: michel.georges@uliege.be

Biological products of importance in food (f.i. milk) and medical (f.i. donor blood derived

products) sciences often correspond to mixtures of samples contributed by multiple

individuals. Identifying which individuals contributed to the mixture and in what

proportions may be of interest in several circumstances. We herein present a method that

allows to do this by shallow whole genome sequencing of the DNA in mixed samples from

hundreds of donors. We demonstrate the efficacy of the approach for the detection of cows

with subclinical mastitis by analysis of farms' tank mixtures containing milk from as many

as 500 cows.

Introduction

Mastitis is the most important health issue in dairy cattle costing European farmers > 1 billion

€ per year in treatment and milk loss[1]. Mastitis is routinely managed by periodically counting

immune cells in milk samples to preemptively identify cows developing subclinical udder

inflammation. As profit margins decrease, farmers tend to forgo milk testing thereby

compromising health management. Cost-effective alternatives for rapid detection of cows

with subclinical mastitis are needed[2]. We previously proposed that somatic cell counts (SCC)

in the milk of individual cows could be estimated if B allele frequencies were measured for

sufficient numbers of SNPs in the tank milk, provided that all cows contributing milk to the

tank be genotyped for the corresponding variants[3]. Thus, the proposed method would allow

to identify a minority of cows with subclinical mastitis by analyzing a single sample containing

a mixture of milk from all the cows on the farm, hence dramatically reducing costs. As

genomic selection (GS) is becoming routine (including for dams)[4], herds that are fully

genotyped with low density SNP arrays (~15K) are becoming standard, and the proposed

method feasible. We herein demonstrate that by combining low density SNP genotyping or

shallow sequencing of the cows and tank milk's DNA with in silico genotype imputation,

individual SCC can be accurately determined and cows with subclinical mastitis effectively

identified even in the largest farms ($\geq$ 500). The proposed method has the potential to

dramatically improve the monitoring of udder health in dairy farms, and to allow the tracing

of the origin of bulk animal food products other than milk.

Results

Principle of the proposed method. Milk of healthy cows typically contains $\leq$ 100,000 somatic

cells per ml. Upon infection leucocytes migrate in the udder and SCC increase rapidly: SCC $\geq$

200,000 / ml are indicative of subclinical mastitis, while SCC into the millions are common for

cows with overt mastitis[5]. Assume that cows and tank (i.e. the reservoir in which the milk of

the cows is collected) milk are genotyped for a collection of SNPs. If all cows contribute

identical amounts of DNA to the milk, the expected "B" allele frequency in the tank milk

corresponds to the frequency of the "B" allele in the farm's cow population. The actual DNA

amount contributed by each cow depends on the volume of milk produced and its SCC.

Unequal DNA contributions will cause slight departures from the expected B allele

frequencies in the tank milk. Integrating these shifts over a large number of SNPs in

conjunction with the known genotypes of individual cows (using f.i. a linear model) allows for

the estimation of the relative DNA contribution of each cow. Accounting for individual milk

volumes and for the SCC in the tank milk allows for the estimation of SCC for individual cows

(Fig. 1 and Methods).

Evaluating the proposed method by simulation. We first evaluated the proposed method by

simulation (cfr. Methods). Genotyping the cows and the tank milk using 10K SNP arrays (i.e.

low-density (LD) arrays as generally used for GS) allowed for the accurate estimation of

individual SCC for farms with up to 100 cows ($r \geq 0.9$, where r is the correlation between

real and estimated SCC) (scheme A). However, farms with > 100 cows are increasingly

common.   Medium- (MD, f.i. 50K) and high-density (HD, f.i. 700K) SNP arrays would be

needed for the approach to be effective in farms with $\geq$ 250 or $\geq$ 500 cows, respectively. Yet

– being too expensive - this is presently not a viable proposition (Fig. 2A).   We therefore

envisaged a second scheme (B) in which the cows would still be genotyped with LD SNP arrays

(as done in practice) yet imputed[6] to whole genome (8 million SNPs in the simulations) using

a sequenced reference population[7], while the DNA of the tank milk would by genotyped by

shallow whole-genome sequencing (SWGS). We found that under this scenario sequencing

the tank milk at a depth of 0.25 was sufficient for farms with 100 cows, 0.5 for farms with 250

cows, and 2 for farms with 500 cows (Fig. 2B). Accuracies were not significantly affected by

the density of the SNP arrays, i.e. the method performed as well with LD as with MD arrays

(data not shown). Anticipating further advances in sequencing technology, we also envisaged

a scheme (C) in which both cows and tank milk would be genotyped by SWGS. We found that

a 1-fold sequencing depth of the tank milk would be sufficient when combined with a 0.25-

fold depth for 100 cows, while a 5-fold sequencing depth of the tank milk would be needed

in combination with 0.25-fold depth for 250 cows and 1-fold depth for 500 cows (Fig. 2C). In

scheme C, allelic dosage in the cows is directly measured from the number of alternative and

reference alleles in the sequence reads. We further explored the effectiveness of augmenting

the cow genotype information from SWGS by imputation (scheme D). This proved to be

effective, reducing the required sequence depth to 0.25-fold for tank milk and 0.25-fold for

100 cows, to 1-fold for tank milk and 0.25-fold for 250 cows, and to 5-fold for tank milk and

0.25-fold for 500 cows (Fig. 2D).

Real-world application of the proposed method. To test the feasibility of our method in the

real world, we first collected cow (blood) and tank (milk) samples from a farm milking 133

Holstein-Friesian cows. When only using genotypes from the Illumina LD arrays (17K SNPs)

for both cows and tank milk (scheme A), correlations between predicted and measured SCC

were 0.91 (or 0.79 when ignoring one cow with SCC > 3 million). We then imputed the cows

to whole genome (13M SNPs) using a reference population of ~750 whole genome

sequenced Holstein-Friesian animals, and sequenced the tank milk at ~3.5-fold depth. The

corresponding correlations (scheme B) were 0.97 (0.95) when using all sequence information,

or 0.96 (0.92) when down-sampling sequence information as low as 0.1-fold depth (Fig. 3A).

We next performed a similar experiment on a farm milking 520 Holstein-Friesian cows.   The

correlation between predicted and measured SCC was 0.78 (or 0.42 when ignoring 23 cows

with SCC > 3 million) when only using information from the LD array for both cows and tank

milk (scheme A).   When imputing the cows to whole genome (13M SNPs) and sequencing

the milk at ~3.5-fold depth (scheme B), the correlation increased to 0.89 (0.83). Down-

sampling the sequence information to 0.1-fold depth reduced the correlation to 0.79 (0.57)

(Fig. 3B).

As shown in both farms, correlation estimates are affected by SCC spread: small numbers of

cows with very high SCC tend to inflate r. We therefore computed accuracies, computed as

the proportion of correctly classified cows for different SCC thresholds, which is how farmers

would likely use the information. It can be seen that for a threshold value of for example

500,000 SCC, accuracies > 0.85 were obtained when sequencing (scheme B) the tank milk at

respectively 0.1x (133 cows) and 3.5x depth (520 cows). Thus - as predicted by the simulations

- scheme A provided adequate precision for the farm with 133 cows, but not for the farm

with 520 cows. However, in this large farm, combining SWGS of the tank milk with whole

genome imputation of the cows (i.e. scheme B) was indeed effective (Fig. 3).

As costs per bp continue to decline, sequencing is likely to replace array-based genotyping in

the future. To test the feasibility of schemes C and D (i.e. genotype the cows by SWGS rather

than with SNP arrays, without (C) and with (D) imputation), we collected samples from a farm

with 120 Holstein-Friesian cows. All cows were genotyped with the Illumina LD array (17K) as

well as sequenced at average 1.08 -fold depth (range: 0.26-1.73). The milk was sequenced at

~3.5-fold depth. The correlation between predicted and measured SCC was 0.97 (or 0.96

when ignoring one cow with SCC > 3 million) under scheme A. Under scheme C, correlations

were 0.82 (0.83) when sequencing the milk at 3.5x and 0.75 (0.76) when down-sampling the

milk to 0.1x. We then imputed the sequenced cows to HD (770K SNPs) using a population of

800 reference animals genotyped with the HD array (scheme D). The correlation increased

to 0.93 (0.94) when sequencing the milk at 3.5x and to 0.83 (0.77) when down-sampling the

milk to 0.1x (Fig. 3C). Accuracies at SCC threshold of 500,000 were 0.96 (scheme A), 0.95

(3.5x) and 0.80 (0.1x) (scheme B), 0.82 (3.5x) and 0.81 (0.1x) (scheme C), and 0.95 (3.5x) and

0.88 (0.1x) (scheme D) (Fig. 3C). In summary, (i) combining cow genotyping using SNP arrays

with genome-wide imputation with SWGS of tank milk allows for cost-effective identification

of cows with subclinical mastitis even in farms with as many as 500 cows per milk tank, and

(ii) as sequencing costs continue to decline, arrays-based targeted SNP genotyping of the

cows could be replaced by genotyping by SWGS and yield comparable results.

Monitoring SCC dynamics with the proposed method. Farmers typically measure individual

SCC once a month or less. Yet, SCC may rapidly change. The SCC measured on the milk testing

date may not be a reliable indicator of the cow's udder health during the intervening period.

To examine the SCC dynamics over time, we collected 20 tank milk samples over a 100-day

period (day -84 to +17 from day of milk testing) for the farm with 120 cows. Milk samples

were genotyped using the Illumina LD array, and individual SCC estimated using scheme A.

Fig. 4A shows the SCC predicted every 5 days on average for the 120 cows, sorted by SCC

measured on day 0 (=milk testing day).   Of note, the correlation between the SCC measured

on day 0 and the average of the SCC estimates for the 21 collection dates was low ($r =$

0.52)(Fig. 4B) and decreased rapidly with the number of days from milk testing day (Fig. 4C).

Discussion

We herein demonstrate that by combining array-based SNP genotyping and whole-genome

imputation for the cows with SWGS of the tank milk, it is possible to accurately estimate SCC

for individual cows and hence effectively identify animals with subclinical mastitis even for

tanks collecting milk for >500 cows, and this by performing a single analysis for the entire

herd. Reagent costs to sequence a mammalian genome at 1-fold depth are now <20€ thus

making this a cost-effective proposition.   As a matter of fact, the method is being deployed

in the field in several countries.

Implementing the method requires all cows on the farm to be genotyped. This will

increasingly correspond to reality as genotyping costs continue to decrease and genomic

selection is more and more used for the selection of cows.   In 2016 more than 1.2 million

dairy cows had been reportedly genotyped in the US alone[8] and present worldwide numbers

are likely ≥ 3 million.   In addition, a reference population of a few hundred animals of the

breed of interest that are either HD genotyped (700K) or better whole-genome sequenced

are required for accurate imputation. Such reference populations are already available for

the most important dairy cattle breeds[7,9], and could be easily generated for the remaining

ones.

We show that SCC are dynamic and rapidly change over time. SCC measured on day 0 are

poor indicators of SCC in previous and future weeks: cows with high SCC on the day of milk

testing may have low SSC a few days later (or earlier) and vice versa. The proposed method

would allow tighter monitoring of SCC hence improving udder health management.   More

frequent monitoring of SCC for large number of cows may reveal interindividual differences

with regards to SCC dynamics that may be correlated with mastitis resistance, heritable and

hence amenable to selection including by GS.

Sequencing of the DNA in the tank milk allows simultaneous characterization of the tank's

microbiome.   As a matter of fact, ~1% of reads in this study mapped to bacterial genomes

(data not shown). This information may be very useful both from a farm health management

point of view as well as from a downstream dairy processing point of view. Whole genome

sequence data of bulk milk also informs about the herd frequency of functional variants such

casein variants affecting consumer health or processing properties[10], or variants causing

inherited defects or embryonic lethality in cows[4].   In many countries, it is not allowed to add

milk from cows being treated with antibiotics to the tank. As suggested before, the proposed

approach can be adapted to verify whether a specific cow did contribute milk to the tank or

not (f.i. by testing the significance of the corresponding cow effect in the linear model)[3].   The

described method may have applications in tracing the origins of bulk animal food products

other than milk, as well as in monitoring the composition of mixed-donor blood-derived

transfusion products.

Acknowledgements

This work was funded by the Unit of Animal Genomics and by the ERC DAMONA grant to

Michel Georges. We are grateful to Jean-Bernard Davière, Pierre Lenormand, Bonny Van

Ranst, Kristien Neyens and Miel Hostens for providing the samples and information needed

to conduct the experiments.

References

1. Hogeveen H, Huijps K, Lam TJGM. Economic aspects of mastitis: New developments. *N. Z.*

Vet. J. 59:16–23 (2011).

2. Viguier C, Arora S, Gilmartin N, Welbeck K, O'Kennedy R. Mastitis detection: current

trends and future perspectives. Trends Biotechnol 27: 486-493 (2009).

3. Blard G, Zhang Z, Coppieters W, Georges M. Identifying cows with subclinical mastitis by

bulk SNP genotyping of tank milk. *J Dairy Sci* 95:4109-4113 (2012).

4. Georges M, Charlier C, Hayes B. Genomics selection of livestock and beyond. *Nat Rev*

Genet 20:135-156 (2019).

5. Schukken YH, Wilson DJ, Welcome F, Garrison-Tikofsky L, Gonzales RN. Monitoring udder

health and milk quality using somatic cell counts. *Vet Res* 34: 579-596 (2003).

6. Marchini J, Howie B. Genotype imputation for genome-wide association studies. *Nat Rev*
Genet 11:499-511 (2010).

7. Daetwyler HD *et al.* Whole-genome sequencing of 234 bulls facilitates mapping of
monogenic and complex traits in cattle. *Nat Genet* 46: 858-865 (2014).

8. Wiggans GR, Cole JB, Hubbard SM, Sonstegard TS. Genomic selection in dairy cattle: the
USDA experience. *Annu Rev Anim Biosci* 5:309-327 (2017).

9. Charlier C. *et al.* Reverse genetic screen for embryonic lethal mutations comprising
fertility in cattle. *Genome Res* 26: 1-9 (2016).

10. Brooke-Taylor S, Dwyer K, Woodford K, Kost N. Systematic review of the gastrointestinal
effects of A1 compared with A2 β-casein. *Adv Nutr* 8:739-748 (2017).

Figure 1: Estimating Somatic Cell Counts (SCC) in the milk of individual cows by analyzing a
sample of milk from the farm's tank. Cows 1 to *n* contribute different amounts of milk
(buckets of various sizes in the figure) to the farm's tank. The milk contains somatic cells
(shown as small spheres in the milk colored by cow) whose numbers reflect the health status
of the cow's udder. Cow 1 has higher SCC, an indicator of subclinical mastitis. SCC are
unknown upon milking (indicated by the "?"). Cows are individually SNP genotyped once. In
scheme A this is done using SNP arrays (illustrated by the mesh) yielding genotype
information for the limited number of interrogated SNPs (high bars) that can be summarized
by the B-allele frequency as shown (white: 0, halve colored: 0.5, full colored: 1). SNP
genotypes of individual cows are coded in the same colors as the SCC. In scheme B, the
genotypes of the interrogated SNPs are augmented by imputation (illustrated by the
computer rack), yielding dosage information (B-allele frequency) for many more SNPs (small
bars). In scheme C, cows are genotyped individually by shallow whole genome sequencing
(SWGS) (illustrated by the sequencer). Sequence reads (gray lines) are aligned to the
reference genome and alternate alleles at SNP positions highlighted as color-coded tics. The
B-allele frequency at specific SNP positions is measured as the ratio of the number of reads
with alternate (B) vs the total number of reads. In scheme D, the genotype information from
SWGS is augmented by imputation improving the accuracy of the B-allele frequency estimates
for millions of SNPs (small bars). A small sample of milk (T(ank) M(ilk)) is periodically (f.i.
monthly or weekly) collected from the farm's tank. DNA is extracted from TM and genotyped
using SNP arrays (scheme A) or SWGS (schemes B, C and D). B-allele frequency for SNP *j* in
the milk ($\widehat{BAF_j}$) is estimated from the ratio of fluorescence intensities when using SNP arrays,
or from the proportion of reads with B allele in SWGS. The SCC of individual cows are
estimated from a set of linear equations modelling $\widehat{BAF_j}$ as the sum of B allele dosage (d_{ij})
multiplied by the proportion of the DNA in the tank contributed by cow *i* (f_i). The estimated
proportions of DNA contributed by each cow correspond to the values of f_i's that minimize
the sum of squared errors (ε_j) over all SNPs. The SSC for individual cows, *per se*, can be
estimated as $SCC_i = SCC_{tank} \times V_{tank} \times f_i/V_i$, where SCC_{tank} is the SCC measured in the
farm's tank, and V_i/V_{tank} is the proportion of the milk volume contributed by cow *i*.
230
231

232
233

$$\overline{BAF}_j = \sum_{j=1}^{m} f_i \times d_{ij} + \varepsilon_j$$

Estimates of individual SCC

Figure 2: Evaluating the efficiency of the proposed approach for the estimation of SCC in the
milk of individual cows by genotyping the tank milk, by simulation. **(A)** Reference scheme in
which individual cows and tank milk are genotyped with the same array interrogating 10K
(LD), 50K (MD) or 700F (HD) SNPs. **(B)** Scheme in which individual cows are genotyped with a
LD 10K SNP array and imputed to whole-genome (8 million SNPs), while the tank milk is
whole-genome sequenced at depth ranging from 0.25x to 5x. **(C)** Scheme in which individual
cows (0.25x and 1x) and tank milk (range: 0.25x to 5x) are genotyped by shallow whole-
genome sequencing (SWGS). **(D)** Scheme in which individual cows are genotyped by SWGS
(0.25x and 1x) followed by imputation to whole genome (8M SNPs), and tank milk is
genotyped by SWGS (range: 0.25x to 5x). In all graphs, the X axis corresponds to the number
of cows contributing milk to the tank. The dots mark parameter combinations that yield
satisfactory correlations ($r \geq 0.9$). Colored lines correspond to conditions that were used
with the real data as shown in Fig. 2.
248

249
250

Figure 3: Correlation between predicted and measured SCC in the milk of individual cows (left
column), as well as accuracies in classifying cows with SCC above and below a chosen
threshold value (right column), in farms with 133 (top row), 520 (middle row) and 120
(bottom row) cows, using scheme A (blue), scheme B (red), or scheme D (green). Scheme A:
cows and tank milk genotyped with LD SNP arrays (17K), no imputation. Scheme B: cows
genotyped with LD array and imputed to 13M SNPs, tank milk sequenced 3.5x (red) or 0.1x
(orange). Scheme D: cows genotyped by whole-genome sequencing (1x) and imputation to
HD, and tank milk sequenced at 3.5x (dark green) or 0.1x (light green).

259
260

261 **Figure 4: (A)** SCC predicted using scheme A for 21 tank milk samples collected over a 100-day
262 period from 138 cows total. Small grey circles: 20 predictions per cow. Large grey circles:
263 average of 21 measurements per cow. Red diamond: SCC measured on day 0. Green triangle:
264 SSC predictions on day 0. **(B)** Relationship between SCC values measured on day 0 and
265 average of 21 predictions sampled over a 100-day period (days -84 to +17). **(C)** Correlations
266 between measured (day 0) and predicted (day x) SCC as a function of the number of days from
267 day 0.

Methods

Simulated data. Reference scheme (A): We simulated farms with *n* (25, 50, 100, 250 and 500)

cows contributing milk to the tank. Cows were genotyped with SNP arrays for *m* (10K, 50K,

or 750K) markers without error. Minor Allele Frequencies (MAFs) were sampled from a

uniform $]0,0.5]$ distribution, and genotypes from the corresponding Hardy-Weinberg

distributions. SCS of individual cows (SCS_i) were simulated by sampling values from a

Weibull distribution with scale parameter α=1 and shape parameter β=2, and multiplying the

ensuing value by 200,000. Exact B-allele frequencies of individual SNPs (BAF_j) in the milk

were determined for each SNP *j* based on the combination of cellular contribution of the *n*

cows to the milk, and their genotype. It was assumed that B-allele frequencies were

estimated with a normally distributed error $N(0, 0.0025)$ (i.e. SE = 0.05), yielding m $\widehat{BAF_j}$.

Scheme B: Same setting as in the reference scheme with the following additions. For cows

genotyped for 10K or 50K SNPs, we simulated imputation by augmenting the data to 8 million

(M) genotypes using an error model mimicking real, MAF-dependent imputation accuracy.

The error model was constructed using a real data set for 800 unrelated Holstein-Friesian

individuals that were genotyped for the Illumina 777K array. This data set was split into a set

of 200 and a set of 600 individuals. The set of 200 was reduced first to the genotypes

interrogated by the Illumina 10K (LD) array and then to the genotypes interrogated by the

Illumina 50K SNP arrays. The reduced SNP sets were imputed back to the content of the

Illumina 777K (HD) SNP array using the 600 individuals as reference population. The

frequencies of imputing a given genotype depending on the real genotype, were scored for

MAF bins of 0.01 separately for the LD and 50K array data. We simulated genotyping-by-

sequencing of tank milk as follows. For each of the 8M SNP positions, we sampled local read

depth ($r \in$ integers) from a Poisson distribution with mean *C*, where *C* is the average genome-

wide coverage (0.25, 0.5, 1, 2 or 5). We then sampled *r* reads, each with a probability = BAF_j

(computed as above) of being the B-allele. Scheme C: Individual SNP genotypes and tank B-

allele frequencies (BAF_j) were generated as in scheme A (genotypes at 8 M SNP positions).

It was assumed that milk tank was genotyped by SWGS at average coverage of *C* (0.25, 0.5, 1,

2 or 5) and cows were genotyped by SWGS at average coverage of *C* (0.25, 0.5, or 1).

Genotyping-by-sequencing of individual cows was simulated by (i) sampling, for each of 8M

SNP positions, local read depth ($r \in$ integers) from a Poisson distribution with mean *C*, and

(ii) sampling *r* reads with probability 0, 0.5 or 1 to be the alternate allele (A) depending on the

genotype of the cow (RR, RA or AA). Genotyping-by-sequencing of the tank milk was done as

in Scheme A. <u>Scheme D:</u> Identical to scheme C except that cow genotypes were generated

at 8M SNP position using a MAF- and sequence-depth dependent imputation error model.

The error model was constructed using available SWGS data down sampled to 1x (176 cows)

or 0.25x coverage (192 cows). The cows were imputed to HD (777K SNPs) using a reference

population of 800 unrelated Holstein-Friesian individuals that were genotyped with the

Illumina 777K array. At each of the 777K SNP positions, the likelihood of the sequence data

under the three possible genotypes (RR, AR and AA), were computed following Chan et al.[3],

as:

310
$$L(nr_R, nr_A|\text{"RR"}, \varepsilon) = \binom{nr_R + nr_A}{nr_A} \times (1 - \varepsilon)^{nr_{AR}} \times \varepsilon^{nr_{RA}}$$

311
$$L(nr_R, nr_A|\text{"RA"}, \varepsilon) = \binom{nr_R + nr_A}{nr_A} \times 0.5^{(nr_R + nr_A)}$$

312
$$L(nr_R, nr_A|\text{"AA"}, \varepsilon) = \binom{nr_R + nr_A}{nr_A} \times (1 - \varepsilon)^{nr_A} \times \varepsilon^{nr_R}$$

where nr_R (respectively nr_A) is the number of R (respectively A reads) and ε is the sequencing

error rate set at 0.01. The corresponding $\log_{10} L$ were used as input for Beagle4[1]. Variant

positions without sequence coverage in any of the 176 (192) cows (hence not imputed by

Beagle4) were dealt with in a second round of imputation using Beagle5[2]. The imputation

accuracy was evaluated in 0.01 MAF-bins by comparing imputed and real genotypes at the

~17K variant positions interrogated by the Illumina LD array.

Real data. <u>Data set 1:</u> We obtained a sample of tank milk from a farm in France milking 133

Holstein-Friesian cows. All had been genotyped with an Illumina LD array interrogating 17K

SNPs using standard procedures. For all cows, genotypes were imputed to whole genome

using a reference population of 743 Holstein-Friesian animals sequenced at average depth of

15x (range: 4-48) and the Beagle software (v5.0)[1] yielding allelic dosages for a total of 13

million SNPs. Individual milk records, including volume and SCC (cells/ml) measured on the

day of the sample collection, were obtained for all cows that had contributed milk to the tank.

DNA was isolated from 1.5 ml tank milk using the NucleoMag kit (Macherey-Nagel). The tank

milk DNA was first genotyped using the Illumina LD array interrogating 17K SNPs. An Illumina

compatible NGS library was then prepared with 50ng of genomic DNA using the KAPA

HyperPlus kit (Roche). Sequencing was performed on a NextSeq500 instrument (Illumina),

yielding 63 million paired end reads of 2*75 bp, corresponding to a genome coverage of 3.5x.

Reads were mapped to the bosTau8 genome build using BWA mem. Reference (R) and

alternate (A) alleles were counted at 13M SNP positions of the HD array using the Bam-

ReadCount tool (https://github.com/genome/bam-readcount.git) for reads with a minimum

mapping quality of 30. Data set 2: We obtained samples of tank milk from a Belgian farm

including milk from 520 Holstein-Friesian cows. Milk volume and SCC (cells/ml) measured on

the same day, were obtained for all cows that had contributed milk to the tank. All cows were

genotyped with the Illumina LD array interrogating 17K SNPs using standard procedures, and

imputed to whole genome using whole genome sequence data (average depth: 15x; range:

4x-48x) from 743 Holstein-Friesian animals as reference (M. Georges, unpublished) and the

Beagle software (v5.0)[2] yielding allelic dosages for a total of 13 million SNPs. DNA extraction

from the tank milk samples and genotyping with the Illumina LD (17K) array were conducted

as for dataset 1. For sequencing of the tank milk, an illumina compatible sequencing library

was prepared using 12 ng of DNA and the Riptide High Throughput Rapid Library Prep

Kit (iGenomx). The library was sequenced on an Illumina NextSeq500 2*150 paired end flow

cell at 4X coverage. Data set 3: We obtained samples of tank milk from a Belgian farm

including milk from 120 Holstein-Friesian cows. Milk volume and SCC (cells/ml) measured on

the same day, were obtained for all cows that had contributed milk to the tank. All cows were

genotyped with the Illumina LD array interrogating 17K SNPs using standard procedures, and

imputed to whole genome using whole genome sequence data (average depth: 15x; range:

4x-48x) from 743 Holstein-Friesian animals as reference (M. Georges, unpublished) and the

Beagle software (v5.0)[2] yielding allelic dosages for a total of 13 million SNPs. We additionally

prepared Illumina compatible NGS library for each cow, using 12 ng of genomic DNA and the

Riptide High Throughput Rapid Library Prep Kit (iGenomx). Libraries were sequenced on an

Illumina Novaseq S4 2*150 paired end flow cell at average 1.08x depth (range: 0.26x-1.73x).

Cow genotype-by-sequencing data were imputed to HD (777K) density using a reference

population of 800 Holstein-Friesian animals genotyped with the bovine HD Illumina array

(777K SNPs) and the Beagle software (v5.0)[2] yielding allelic dosages for a total of 777K SNPs.

DNA extraction from the tank milk samples, genotyping with the Illumina LD (17K) array, and

sequencing (coverage 4x) were conducted as for datasets 1&2. Data set 4: In addition to

obtaining a sample of tank milk on the day of the milk recording (i.e. yielding the SCC

measured using with a cell counter) for the Belgian farm with 120 cows, we weekly collected

an additional 11 tank milk samples before and 9 samples after, spanning a total period of ~3

months. The corresponding DNA samples were genotyped using the Illumina LD (17K) array.

Statistical model. We defined a set of *m* linear equations of the form:

366
$$\widehat{BAF_j} = \sum_{j=1}^{m} f_i \times d_{ij} + \varepsilon_j$$

in which f_i is the proportion of the DNA in the tank milk contributed by cow *i*, d_{ij} is the

"dosage" of the alternate allele A for cow *i* and marker *j*, and ε_j is the error term for marker

j. When genotyping the tank milk with arrays, $\widehat{BAF_j}$ corresponds to the B-allele frequency

estimated by Genome Studio (Illumina). When genotyping the tank milk by SWGS, $\widehat{BAF_j}$

corresponds to the proportion of A reads at the corresponding genome position. For cow

genotypes obtained with arrays, d_{ij} corresponds to 0, 0.5 or 1 for genotypes RR, RA and AA,

respectively. For cow genotypes obtained by imputation, d_{ij} is the dosage of the A allele

estimated by Beagle. For cow genotypes obtained by SWGS, $d_{ij} = 0.5 \times$

$P(\text{"RA"}|nr_R, nr_A, q_j) + P(\text{"AA"}|nr_R, nr_A, q_j)$ where nr_R (respectively nr_A) is the number of R

(respectively A reads) for marker *j* and cow *i*, and q_j is the population frequency of the A allele

of marker j.

$P(\text{"RA"}|nr_R, nr_A, q_j) = \dfrac{2q_j(1-q_j) \times 0.5^{nr_R} \times 0.5^{nr_A} \times \frac{(nr_R + nr_A)!}{nr_R!}}{(1-q_j)^2 \times 1^{nr_R} \times 0^{nr_A} + 2q_j(1-q_j) \times 0.5^{nr_R} \times 0.5^{nr_A} \times \frac{(nr_R + nr_A)!}{nr_R!} + q_j^2 \times 0^{nr_R} \times 1^{nr_A}}$

$P(\text{"AA"}|nr_R, nr_A, q_j) = \dfrac{q_j^2 \times 0^{nr_R} \times 1^{nr_A}}{\left(1-q_j\right)^2 \times 1^{nr_R} \times 0^{nr_A} + 2q_j\left(1-q_j\right) \times 0.5^{nr_R} \times 0.5^{nr_A} \times \frac{(nr_R + nr_A)!}{nr_R!} + q_j^2 \times 0^{nr_R} \times 1^{nr_A}}$

For SNPs *j* without usable information for cow *i* (f.i. genotyping failure or no covering reads)

d_{ij} was set at $\widehat{BAF_j}$.

The f_i's were estimated by least square analysis, i.e. by minimizing $\sum_{j=1}^{m} \varepsilon_j^2$. When the tank

milk was genotyped by SWGS, we also performed a weighted least square analysis, i.e. we

estimated f_i's by minimizing $\sum_{j=1}^{m} w_j \varepsilon_j^2$, where w_j is the coverage $(nr_R + nr_A)$.

The SCC_i's were calculated from the f_i's

389
$$SCC_i = SCC_{tank} \times V_{tank} \times f_i/V_i$$

Where V_{tank} and V_i are the volumes of milk in the tank and contributed by cow i, respectively.

The accuracies of the predictions were measured by the (i) correlation (r) between real and

estimated SCC_i, and/or (ii) the ability to discriminate animals with SCC above versus below a

certain threshold value measured as $(T_P + T_N)/m$, where T_P stands for the number of true

positives, T_N for the number of true negatives, and m for the total number of cows.

To test the effect of sequence depth on accuracy we sampled reads overlapping SNP positions

with probability x, such that $E(C \times x) = D$, where D is the desired sequence depth.

References

1.   Browning BL, Browning SR. A unified approach to genotype imputation and haplotype

phase inference for large data sets of trios and unrelated individuals. *Am J Hum Genet*

84:210-223 (2009).

2.   Browning BL, Zhou Y, Browning SR. A one-penny imputed genome from next generation

reference panels. *Am J Hum Genet* 103:338-348 (2018).

3.   Chan AW, Hamblin MT, Jannink J-L. Evaluating imputation algorithms for low depth

genotyping-by-sequencing (GBS) data. *PLoS ONE* 11:e0160733 (2016).

407