Cryptococcus neoformans evades pulmonary immunity by modulating xylose precursor transport

Lucy X. Li[a,*,+], Camaron R. Hole[a,+], Javier Rangel-Moreno[b], Shabaana A. Khader[a], and Tamara L. Doering[a,#]

[a]Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, Missouri, USA

[b]Department of Medicine, Allergy/Immunology, and Rheumatology, University of Rochester School of Medicine, Rochester, New York, USA

Running Head: Xylose precursors and cryptococcal immunity

[#]Address correspondence to Tamara L. Doering, doering@wustl.edu.

[*]Present address: Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

[+]Co-first authors

ABSTRACT

Cryptococcus neoformans is a fungal pathogen that kills almost 200,000 people each

year and is distinguished by abundant and unique surface glycan structures that are rich

in xylose. A mutant strain of *C. neoformans* that cannot transport xylose precursors into

the secretory compartment is severely attenuated in virulence in mice, yet surprisingly is

not cleared. We found that this strain failed to induce the non-protective T helper cell type

2 (Th2) responses characteristic of wild-type infection, instead promoting sustained Inter-

leukin (IL)-12p40 induction and increased IL-17A (IL-17) production. It also stimulated

dendritic cells to release high levels of pro-inflammatory cytokines, a behavior we linked

to xylose expression. We further discovered that inducible bronchus associated lymphoid

tissue (iBALT) forms in response to infection with either wild-type cryptococci or the mu-

tant strain with reduced surface xylose; although iBALT formation is slowed in the latter

case, the tissue is better organized. Finally, our temporal studies suggest that lymphoid

structures in the lung restrict the spread of mutant fungi for at least 18 weeks after infec-

tion, in contrast to ineffective control of the pathogen after infection with wild-type cells.

These studies demonstrate the role of xylose in modulation of host response to a fungal

pathogen and show that cryptococcal infection triggers iBALT formation.

INTRODUCTION

Cryptococcus neoformans is a ubiquitous environmental fungus that causes pneumonia and meningitis. This opportunistic pathogen infects over a million individuals each year, with overall mortality exceeding 20% (1-3). Patient immune status is the main determinant of infection outcome, highlighting the importance of efficient host immune responses in the control of cryptococcosis.

C. neoformans infection begins when the organism is inhaled, which is followed by its proliferation in the lungs. This pulmonary infection may then disseminate to the brain, where it causes a frequently lethal meningoencephalitis. In the lungs, *C. neoformans* first interacts with host macrophages and dendritic cells (DC). These engulf the fungus and migrate to draining lymph nodes, where they present antigen to T cells to initiate the adaptive immune response. The protective immune response to this infection is primarily mediated by T cells: T helper cell type 1 (Th1) responses are protective against *C. neoformans* (4) and T helper type 17 (Th17) responses likely participate in pathogen clearance at mucosal surfaces (5-7). In contrast, T helper cell type 2 (Th2) responses are considered detrimental for protective immunity, leading to fungal growth and facilitating dissemination to the CNS (8, 9). The induction of these non-protective responses is not well understood, but cryptococcal glycans may contribute to this process (10).

Glycan structures participate in antigen recognition, immune activation, and immune regulation in multiple organisms (11). In *C. neoformans* the major virulence factor, a polysaccharide capsule, modulates the immune response via multiple mechanisms, including dampening immune cell activation and impeding phagocytosis (12, 13). The capsule is

primarily composed of two large polysaccharides: glucuronoxylomannan (GXM) and glu-

curonoxylomannogalactan (GXMGal) (14, 15). One major component of both polymers is

xylose (Xyl), which comprises almost one fourth of the polysaccharide capsule mass (16).

Xylose also occurs in cryptococcal glycolipids (17) and as both Xyl and Xyl-phosphate

modifications of protein-linked glycans (18-20). We demonstrate here that Xyl plays a

critical role in the immune recognition of, and response to, *C. neoformans.*

The incorporation of Xyl into cryptococcal glycan structures occurs in the lumen of the

secretory pathway, via enzymatic reactions that use the substrate molecule UDP-Xyl.

This xylose donor is imported into the synthetic compartment by two transporters, Uxt1

and Uxt2 (21). A mutant strain that lacks both transporters (*uxt1Δ uxt2Δ*) exhibits defects

in xylosylation, including incomplete synthesis of capsule polysaccharides, and consider-

able attenuation of virulence in mouse models of infection. Surprisingly, despite its com-

promised virulence, which contrasts with the rapid lethality of wild-type (WT) infection, the

mutant strain persists for months in the lungs of infected mice (21).

In pursuing the mechanism of *uxt1Δ uxt2Δ* persistence, we found that infection with the

mutant failed to induce the Th2 responses provoked by WT fungi, although it did promote

production of IL-12p40 and IL-17, likely via enhanced stimulation of DCs. The correspond-

ing inflammatory response was enriched in T and B cells, which were required for the

prolonged survival of the infected animals. Inflammation in the lung, although it increased

slowly compared to the response to WT infection, appeared to effectively contain the mu-

tant yeast cells for an extended period after infection. Finally, we observed the formation

of inducible bronchus associated lymphoid tissue (iBALT) in the lungs of both WT and

uxt1Δ uxt2Δ infected mice. iBALT resembles secondary lymphoid organs (lymph nodes)

in cellular composition and organization, with two distinctive zones: a central B cell follicle

and surrounding T cell zone (22). The T cell zone contains CD4 and CD8 T cells as well

as DCs (23); the B cell zone contains mainly follicular B cells and its organization is main-

tained by the continuous production of CXCL13 (24). DC activation and cytokine produc-

tion are also required for iBALT formation and maintenance (23, 25). In the lungs of mice

infected with the *uxt1Δ uxt2Δ* mutant, iBALT formation was significantly delayed. How-

ever, once formed, it was better organized than the iBALT produced when mice were

infected with fully xylosylated WT *C. neoformans.* Together, these results suggest that

luminal Xyl modifications of cryptococcal glycoconjugates alter immune recognition and

activation of immune cells, events that are required for efficient control of *C. neoformans*

infection.

RESULTS

UDP-Xyl transport is required for cryptococcal virulence and dissemination

We previously observed that a *C. neoformans* strain lacking UDP-Xyl transporters (*uxt1Δ*

uxt2Δ) was highly attenuated in virulence compared to the WT parental strain KN99α, yet

remained in the lungs of asymptomatic mice for at least 100 days after infection (21).

While defects in the mutant strain, such as altered capsule and a slightly decreased

growth rate, might explain its reduced virulence, this persistence was surprising. To in-

vestigate the development of this chronic infection, we first examined the kinetics of fungal

burden in mice after intranasal inoculation with WT and mutant strains. In A/JCr mice

infected intranasally with WT fungi, which typically succumb to infection roughly 18 days

post-infection (dpi), we observed a rapid and significant increase in pulmonary fungal

burden (Fig. 1A) along with significant dissemination to the spleen (Fig. 1B) and brain

(Fig. 1C). In contrast, the pulmonary burden in *uxt1Δ uxt2Δ*-infected mice increased

slowly from initial inoculum levels to a modest peak at 63 dpi (Fig. 1A). Colony-forming

units in the lung then gradually declined, although even after six months only 30% of mice

showed complete clearance (Fig. 1A, 189 dpi). Although *uxt1Δ uxt2Δ* infection was not

restricted to the lungs, only a fraction of the mice had measurable mutant cells in the

spleen (Fig. 1B) and brain (Fig. 1C) at 63 dpi. Furthermore, this limited dissemination

was only observed at the times of highest lung burden (63 and 126 dpi), with no fungi

detected at distal sites by 189 dpi (Fig. 1B, Fig. 1C).

Cryptococcal xylosylation influences production of pulmonary cytokines and cellular im-

mune response

We next sought to identify the mechanism(s) responsible for the unexpected persistence

of *uxt1Δ uxt2Δ* after intranasal infection. As summarized above, protection against *C.*

neoformans is generally associated with production of type 1 cytokines like interleukin

(IL)-2, IL-12, interferon gamma (IFN-γ), and tumor necrosis factor alpha (TNF-α). In con-

trast, a Th2 response to cryptococcal infection (characterized by the production of IL-4,

IL-5, and IL-13) compromises control of pulmonary fungal burden (26) and facilitates dis-

semination to the central nervous system (4, 26).

We hypothesized that the protracted host clearance of the *uxt1Δ uxt2Δ* mutant reflected

an altered host immune response to cells that display reduced xylose. To assess the host

response, we first analyzed cytokine levels over time in lung homogenates from mice

infected with WT or *uxt1Δ uxt2Δ* strains (Fig. 1). In mice infected with WT cryptococci, we

observed a strong Th2-type response, with significant induction of IL-4 and IL-5, over the

15 days prior to sacrifice (Fig. 1D, Fig. 1E). Other proinflammatory cytokines (IL-1α, IL-

1β, IL-6) were modestly increased (Supp. Fig. 1). In contrast, *uxt1Δ uxt2Δ* cells elicited a

more quiescent Th2 cytokine response at early stages of infection. Furthermore, at the

peak of fungal burden in *uxt1Δ uxt2Δ*-infected mice (day 63, Fig. 1A), there was a signif-

icant increase in the Th1 polarizing cytokine IL-12p40 compared to the levels observed

in naïve lungs (Fig. 1F); IL-12p40 levels then slowly declined, along with fungal burden.

Although the levels of IFNγ and TNFα in mutant-infected mice resembled those in the

lungs of naïve mice (Fig. 1G, Fig. 1H), IL-17 trended higher at the peak of infection (Fig.

1I). Overall, while WT infection rapidly induced non-protective Th2 responses, infection

with *uxt1Δ uxt2Δ* failed to do so (even at the peak of fungal burden) and instead promoted

the sustained induction of IL-12p40 and increased production of IL-17.

To measure the host adaptive immune response to *uxt1Δ uxt2Δ*, we used flow cytometry

to count lymphocytes in the lungs of mice infected with WT or *uxt1Δ uxt2Δ* cells. By 15

dpi, the B cell fraction of viable leukocytes in mutant-infected mice significantly exceeded

the value for WT-infected mice (Fig. 2A); this fraction further increased as mutant infection

progressed, to peak at over 30%. The same pattern was seen for CD4$^+$ and CD8$^+$ T cells

(Fig. 2B, Fig. 2C), initiating even earlier (at 12 dpi) and remaining high throughout the

time course. The total numbers of cells in all three populations remained elevated through

126 dpi (Fig. 2A-C, right), although they subsequently declined.

Survival of uxt1Δ uxt2Δ-infected mice is dependent on T cells

Because our results suggested that T and B cells play a protective role that contributes

to the prolonged survival of *uxt1Δ uxt2Δ*-infected mice, we next examined whether abla-

tion of these cell types altered the course of infection. Because the mouse strains used

for these studies were engineered in the C57BL/6 background, we included infections of

this strain as controls. C57BL/6 mice are less resistant to *C. neoformans* than the A/JCr

mice used above (27), as shown by their eventual susceptibility to *uxt1Δ uxt2Δ* infection

(Fig. 2D, solid black line). Notably, *Rag1-/-* mice in this background were significantly

more susceptible to the mutant fungi: all succumbed by 55 dpi (Fig 2D; solid blue line).

These animals still survived longer than WT-infected mice, possibly due to the slightly

slower growth of the mutant which we previously observed (21).

Since *Rag1-/-* mice lack both T and B cells, we also infected mice deficient in either T

cells (*TCRβ -/-*) or B cells (*μMT*) with *uxt1Δ uxt2Δ* cryptococci. B cell deficient mice ex-

hibited a more protracted course of disease than *Rag1-/-* animals, succumbing over an

85-day period (Fig. 2D, solid green line), while mice lacking only T cells succumbed to

infection with kinetics similar to those of the *Rag1-/-* mice (Fig. 2D, solid red line). In con-

trast, all mice inoculated with WT *C. neoformans* succumbed by 20 dpi (Fig. 2D; dashed

lines). Of this group, *Rag1-/-* and *TCRβ-/-* mice still succumbed slightly faster than

C57BL/6 and μMT mice (Fig. 2D). Together, these data show that T cells are the promi-

nent cell type responsible for increased survival of *uxt1Δ uxt2Δ*-infected mice, although

B cells may play a lesser role in protection.

Reduced surface xylose of C. neoformans stimulates DC activation

The antigen-driven activation of DCs is critical to induce protective immunity against *C.*

neoformans (28) and it is likely that their consequent cytokine and chemokine production

are responsible for T cell recruitment (23). We speculated that an enhanced DC response

to xylose-deficient fungi could explain the rise in T cells during *uxt1Δ uxt2Δ* infection (Fig.

2), even in the absence of expanded numbers of DCs (Fig. 3A). To test this idea, we grew

bone marrow derived DCs, stimulated them for 24 hours with heat-killed *C. neoformans*,

and measured the production of pro-inflammatory cytokines. In these studies we used

WT fungi, which display antigens modified with Xyl, or *uxt1Δ uxt2Δ* fungi, which are im-

paired in these modifications. We also tested strains with individual *uxt* mutations: *uxt1Δ*

shows an intermediate level of Xyl utilization while *uxt2Δ* resembles WT (21). Finally, we

assayed *uxs1Δ* cells, which lack all Xyl modification because they cannot synthesize

UDP-Xyl (29), and an acapsular control strain (*cap59Δ*) known to induce a potent DC

response (30).

We found that bone marrow derived DCs (BMDCs) cultured with *uxt1Δ uxt2Δ* cells re-

leased high levels of IL-1β and IL-6, comparable to the levels observed following expo-

sure to a completely acapsular control strain, *cap59Δ* (Fig. 3B-C). In contrast, individual

uxt mutants and *uxs1Δ* induced IL-1β and IL-6 levels similar to those induced by WT fungi

(close to background levels; Fig. 3B-C). TNF-α and IL-12p40 similarly showed the great-

est response upon challenge with *uxt1Δ uxt2Δ*, with levels like those induced by the acap-

sular strain, although a slight increase was also noted with other xylose-deficient strains

(*uxt1Δ* and *uxs1Δ*; Fig. 3D-E)). These results indicate that early interaction with DCs and

consequent induction of proinflammatory cytokines are influenced by the surface Xyl ex-

pression of *C. neoformans*.

The cellular response to cryptococcal infection includes the development of inducible

bronchus-associated lymphoid tissue (iBALT)

Our data suggested that *uxt1Δ uxt2Δ* mutant cells stimulate elevated production of pro-

inflammatory cytokines by DCs, which in turn significantly increases the T and B cells in

infected lungs (in both absolute numbers and as a fraction of total leukocytes). Flow cy-

tometry, however, provides no information as to the distribution of these cell populations

within the lung. To futher probe the cellular immune response, we therefore examined

lung histology, focusing on late time points of infection: 15 dpi for mice infected with WT

fungi and 126 and 189 dpi for *uxt1Δ uxt2Δ*-infected mice (also including 15 dpi to allow

direct comparison to WT). We used Movat's stain for these studies (Fig. 4), because the

Alcian Blue component stains cryptococcal cell walls and capsules turquoise, allowing

visualization of fungi as well as of host cells (31). In WT-infected mice at 15 dpi we saw

the typical pattern of overwhelming fungal burden throughout the lung, with little organized

inflammation (32). The tissue architecture was largely destroyed and there was a prepon-

derance of enlarged fungi with thick capsules. In contrast, mutant-infected mice at the

same time point showed relatively preserved tissue architecture, with limited regions of

organized inflammation. Most cryptococci were restricted to these areas and overall fun-

gal burden was noticeably lower, as expected for this mutant (Fig. 1A). By 126 dpi of

infection with mutant fungi, the regions of inflammation were larger and more highly con-

solidated; fungi were rarely observed in other areas of the lung. The last time point of

mutant infection (189 dpi) looked similar, although inflammation appeared slightly re-

duced, consistent with the decrease in B and T cells that we observed at that time point

(Fig. 2). Lymphoid structures were also less dense; fungi were less restricted to the cen-

ters of these regions and occasionally appeared in lung tissue beyond them.

In both groups of mice (WT- and mutant-infected), we noted distinctive lymphocytic accu-

mulations proximal to the basal side of the bronchial epithelium (Fig. 4). To probe these

structures, we stained lung sections for T cells (CD3), B cells (B220), and plasma cells

(IgG). In all samples we observed B cell follicles surrounded by CD3$^+$ T cell cuffs (Fig 5A),

suggestive of inducible bronchus-associated lymphoid tissue (iBALT) (24). Consistent

with identification as iBALT, these structures stained positive for CXCL13 (Fig. 5B).

To examine iBALT development, we performed blinded morphometric analysis of lung

tissue from mice infected with WT and mutant cryptococci. At 15 dpi with WT fungi, we

observed numerous iBALT structures, with compact B cell follicles surrounded by T cells

(Figs. 5C-E). In contrast, B cell follicles were considerably fewer and less organized at

the same time point in mice infected with the *uxt1Δ uxt2Δ* strain (Figs. 5C-E). The limited

formation of iBALT structures was also characterized by increased peri-vascular accumu-

lation of T cells (Fig. 5A) and decreased production of CXCL13 (Fig. 5B, Fig. 5F). By 126

dpi with *uxt1Δ uxt2Δ*, however, although iBALT structures were still less abundant than

those found in WT infection (Fig. 5C), their average B cell follicle size was significantly

larger (Fig. 5D). Additionally, the *uxt1Δ uxt2Δ* iBALT was more organized, with reduced

T cell cuffing (Fig. 5E), increased CXCL13 (Fig. 5F) and more IgG$^+$ plasma cells (Fig. 5A).

By 189 dpi, these structures regressed slightly, consistent with the reduced lymphocyte

infiltration, cytokine levels, and fungal burden at this time point. WT cryptococci thus in-

duce robust early formation of iBALT, while the *uxt1Δ uxt2Δ* mutant induces delayed but

ultimately larger and better organized iBALT.

DISCUSSION

We wished to investigate the unexpected observation that a mutant strain of *C. neofor-*

mans, despite significant defects in surface glycoconjugates, nevertheless persisted for

many months in infected mice. We found that *uxt1Δ uxt2Δ* cells showed an initial delay

after intranasal infection, but then began to accumulate slowly, eventually rising to a

modest peak, at which time (63 dpi) we also observed limited dissemination. The lung

burden then began to gradually decline, although by the end point of the study (189 dpi)

only three of ten animals had cleared the infection.

The *uxt1Δ uxt2Δ* mutant induced a generally subdued pulmonary immune response

compared to that induced by WT cryptococci. Notably, levels of IL-12p40, a cytokine

normally required for inducing a protective response against *C. neoformans* (33), were

elevated at 63 dpi and beyond (Fig. 1F); this may facilitate subsequent resolution of the

uxt1Δ uxt2Δ infection. The rise in IL-12p40 could reflect an increase in either IL-23 or IL-

12 (34), which stimulate Th17 and Th1 type responses, respectively. Since we observed

a trend towards increased IL-17 at 63 dpi, but no difference in IFN-γ levels, IL-12p40

was more likely associated with a Th17 response.

In terms of overall lung histology, the early host response of mutant-infected mice was

limited. By 128 dpi it had become more organized, with all fungi apparently contained

within inflammatory structures, although there was a slight degradation of fungal control

by the last time point imaged (189 dpi). We specifically quantitated the temporal pro-

gression for iBALT and found that B-cell follicles at the earliest time point of mutant in-

fections (15 dpi) were sparse and small compared to 15 dpi wild-type iBALT, with larger

T cell cuffs and smaller areas of CXCL13 staining. By 126 dpi, however, their follicle

size and CXCL3 staining had surpassed WT levels, although these parameters re-

gressed somewhat by 189 dpi. The development of iBALT, therefore, paralleled the abil-

ity of the host immune response to physically contain cryptococci within inflammatory

structures in the lung.

DCs reacted more strongly to *uxt1Δ uxt2Δ* than to WT *in vitro*, releasing greater amounts

of pro-inflammatory cytokines (Fig. 3B-E), even though the global cytokine response to

the mutant was less perturbed (Fig 1D-I). These cytokines may directly or indirectly lead

to the increased recruitment of T and B cells in *uxt1Δ uxt2Δ* infection (Fig. 2A-C) and

perhaps accelerate compartmentalization into compact B cell follicles and discrete T cell

areas in iBALT (Fig 5). Such differences in cellular composition and organization, coupled

with differences in the local cytokine production, may confer distinct properties on the

iBALT induced by WT and *uxt1Δ uxt2Δ*, influencing their ability to protect against *C.*

neoformans.

We expected that the highly immunostimulatory behavior of *uxt1Δ uxt2Δ in vitro* reflected

the lack of Xyl modifications on its surface glycoconjugates, a consequence of the ab-

sence of UDP-Xyl transport into the secretory compartment. We were surprised, there-

fore, that mutants unable to synthesize UDP-Xyl (*uxs1Δ*), which similarly lack Xyl modifi-

cations on secreted glycoconjugates, did not phenocopy this broad and robust DC re-

sponse. This suggests that Uxs1 itself is required for the strong response seen in the

absence of xylosylation in the secretory pathway. It may be that this protein plays addi-

tional roles unrelated to glycosylation (35), or that the stimulatory component is depend-

ent on cytosolic UDP-Xyl, rather than UDP-Xyl that has been transported into the secre-

tory pathway.

Unlike encapsulated strains, *C. neoformans* mutants lacking capsule induce the upregu-

lation of multiple genes involved in cytokine responses and enhance antigen processing

and presentation by DCs (36). Interestingly, the cytokine levels induced in DC by the

uxt1Δ uxt2Δ mutant were similar to those induced by the acapsular mutant *cap59Δ* (Fig

4A-C). Xyl may thus be critical for the immune suppressive effects normally exerted by

capsule material. The mechanisms behind the robust DC activation seen in these two

mutants, including the cellular receptors and fungal components responsible, remain to

be determined. It is possible that the *uxt1Δ uxt2Δ* mutant lacks immune suppressive

mechanisms that are present in WT *C. neoformans* infection, and therefore induces ro-

bust cytokine responses and consequent recruitment of T and B cells that stimulate iBALT

formation and limit early infection.

In this study, we report for the first time that *C. neoformans* infection triggers iBALT for-

mation. To our knowledge, the only other fungus reported to induce iBALT is *Pneumo-*

cystis (37). Notably, the iBALT detected in our experimental infections resembles sub-

pleural nodules that have been observed in cryptococcosis patients (38), although fur-

ther examination will be necessary to establish the similarities between histological fea-

tures of inflammatory cell infiltration in mice and humans. Although iBALT formation oc-

curred in mice infected with both mutant and WT fungi, it was inadequate to defend

against the overwhelming fungal burden that rapidly develops in a standard murine

model of cryptococcal infection. It may be that lower inocula would shift the balance in

favor of the host. In any event, the kinetics of iBALT formation and its organization are

dependent on virulence factors of the pathogen. Future studies will be needed to fully

delineate the molecular host-pathogen interactions that drive iBALT formation following

C. neoformans infection and the magnitude of the role that iBALT plays; in this regard

specific chemokine deficient mice, such as CXCL-13 deficient animals that lack the abil-

ity to induce iBALT structures (39), may be useful.

MATERIALS AND METHODS

Fungal strains

C. neoformans strains (Table S1) were grown at 30 °C in YPD medium (1% [wt/vol] yeast extract, 2% [wt/vol] peptone, 2% [wt/vol] dextrose) with shaking (230 rpm) or on YPD agar plates (YPD medium with 2% [wt/vol] agar) supplemented with the following antibiotics as appropriate: 100 μg/ml nourseothricin (NAT; Werner BioAgents) or Geneticin (G418; Invitrogen).

Mice

C57BL/6J (RRID: IMSR_JAX:000664) and A/JCr (RRID: IMSR_CRL:563) mice were obtained from Jackson Laboratory. *Rag1*-/- (RRID: IMSR_JAX:002216), *TCRβ*-/- (RRID: IMSR_JAX:002118), and μMT mice (RRID: IMSR_JAX:002288) (all on C57BL/6 background) were generously provided by Dr. Michael Diamond and Dr. Wayne Yokoyama (Washington University School of Medicine); breeders were originally purchased from Jackson Laboratory. All mice were 6 to 8 weeks old at the time of infection. A/JCr mice were female; male and female mice were used for the other strains with gender and age matched C57BL/6 controls.

Ethics statement

All animal protocols were reviewed and approved by the Animal Studies Committee of the Washington University School of Medicine and conducted according to National Institutes of Health guidelines for housing and care of laboratory animals.

C. neoformans infection

Fungal strains were cultured overnight (O/N) and diluted to 10^6 cells/mL in sterile PBS.

Mice were intranasally inoculated with a 50 μL aliquot, and then weighed and monitored

daily. Infected mice were sacrificed if they lost >20% relative to peak weight or at specified

time points. At the time of sacrifice, mice were perfused intracardially with 10 mL sterile

PBS, and organs were processed as described below for fungal burden, histology, flow

cytometric analysis, and cytokine measurements.

Organ burden and cytokines

Brain and spleen homogenates were harvested and plated for CFU at the specified time

points. 50 μL aliquots of the left lung homogenates (V_T = 1 mL) were similarly plated for

CFUs. The remaining sample was assayed for pulmonary cytokine levels using the Bio-

Plex Protein Array System (Bio-Rad Laboratories). Briefly, the lung homogenates were

mixed with an equal volume of PBS/0.1% Triton 100x/2x protease inhibitor (Pierce EDTA-

free protease inhibitor; Thermo Scientific), vortexed for 3 seconds, and clarified by cen-

trifugation (2500 x g, 10 min). Supernatant fractions were then assayed using the Bio-

Plex Pro Mouse Cytokine 23-Plex (Bio-Rad Laboratories) for the presence of IL-1α, IL-

1β, IL-2, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, IL-12 (p40), IL-12 (p70), IL-13, IL-17A, granulo-

cyte colony stimulating factor (G-CSF), granulocyte monocyte colony stimulating factor

(GM-CSF), interferon-γ (IFN-γ), CXCL1/keratinocyte-derived chemokine (KC),

CCL2/monocyte chemotactic protein-1 (MCP-1), CCL3/macrophage inflammatory pro-

tein-1α (MIP-1α), CCL4/MIP-1β, CCL5/regulated upon activation normal T cell expressed

and secreted (RANTES), and tumor necrosis factor-α (TNF-α).

Tissue staining and histologic analysis

Mice were perfused intracardially with sterile PBS, and the lungs inflated with 10% forma-
lin. Lung tissue was then fixed O/N in 10% formalin and submitted to the Washington
University Developmental Biology Histology Core for paraffin-embedding and sectioning.
Immunofluorescence staining for B cells (APC-conjugated rat anti-mouse CD45R/B220,
clone RA3-6B2, BD Biosciences, RRID: AB_627078), T cells (CD3-ε, clone M-20, Santa
Cruz Biotechnology, RRID: AB_631129), Ig G$^+$ plasma cells (FITC-donkey anti-mouse Ig
G, Jackson ImmunoResearch Laboratories, RRID: AB_2340796), and CXCL13-produc-
ing cells (goat α-mouse CXCL13, AF470, R&D Systems, RRID: AB_355378) was per-
formed as in Ref (37). Images were taken with a Zeiss Axioplan2 microscope and rec-
orded with a Hamamatsu camera. Blinded morphometric analysis of lung structures was
performed with the outline tool of Zeiss Axiovision software. Additional sections were
stained with Movat's pentachrome stain by the Washington University Pulmonary Mor-
phology Core.

Flow cytometric analysis

The right lung of individual mice was enzymatically digested (in 5 mL RPMI with 1 mg/mL
collagenase type IV) at 37 °C with shaking (230 rpm) for 30 min, and then sequentially
passed through sterile 70 and 40 μm pore nylon strainers (BD Biosciences, San Jose,
CA). Red blood cells in the samples were lysed by treatment for 3 min on ice with 5 mL
ammonium-chloride-potassium lysing buffer (8.024 g/L NH_4Cl, 1.001 g/L $KHCO_3$, 2.722
mg/L EDTA•Na_2 $2H_2O$), followed by the addition of 2 volumes of PBS. The remaining cells

were pelleted (1000 x g, 5 min, 4 °C), washed twice with PBS, diluted to 10^6 cells/mL in

PBS, and stained with LIVE/DEAD fixable blue dead cell stain (1:1000; Thermo Scien-

tific). Following incubation in the dark for 30 min at 4 °C, cells were washed with PBS and

FACS buffer (2% fetal bovine serum in PBS) before resuspension in FACS buffer. Sam-

ples were then blocked with CD16/CD32 (1:500; Fc Block™; BD Biosciences, RRID:

AB_394656) for 5 min and incubated for 30 min with optimal concentrations of fluoro-

chrome-conjugated antibodies (Table S2) diluted in Brilliant Stain Buffer (BD Biosci-

ences). After three washes with FACS buffer, the cells were fixed in 2% formalde-

hyde/FACS buffer. For data acquisition, >50,000 events were collected on a BD LSRFor-

tessa X-20 flow cytometer (BD Biosciences), and the data were analyzed with FlowJo

V10 (Fig. S1; TreeStar). The absolute number of cells in each leukocyte subset was de-

termined by multiplying the absolute number of CD45$^+$ cells by the percentage of cells

stained by fluorochrome-labeled antibodies for each cell population analyzed.

Isolation of bone marrow derived cells

Bone marrow was flushed from the femurs and tibiae of C57BL/6 mice using RPMI. Cells

were collected (1000 x g, 5 min, 4°C), resuspended in RPMI, and counted using a hemo-

cytometer. To prepare bone marrow derived dendritic cells (BMDCs), 2 x 10^6 bone mar-

row cells were plated in 10 mL R10 medium (10% FBS, 0.4% Penicillin-Streptomycin, 2

mM L-glutamate, 50 µM 2-β-mercaptoethanol in RPMI) supplemented with 1 ng/mL GM-

CSF, and incubated at 37 °C and 5% CO_2. Medium was changed 3 and 6 days after

plating, and cells were harvested on day 8. BMDCs were enriched by depletion of BMMs

using biotinylated α-F4/80 antibody (eBioscience, RRID: AB_466657) and anti-biotin con-

jugated magnetic beads (Miltenyi Biotec). The BMDCs in the flow through were positively

selected using α-CD11c magnetic beads according to the manufacturer's protocol (Mil-

tenyi Biotec).

Dendritic cell assays

To assay the ability of fungal strains to activate BMDCs, *C. neoformans* strains of interest

were grown O/N, washed in PBS, and incubated at 65 °C for 15 min to heat kill (HK) the

fungi. BMDCs and HK fungi (10^6 cells of each) were then co-incubated for 24 h, sedi-

mented, and the supernatant fractions transferred to 1.5 mL centrifuge tubes containing

10 μL of 100x protease inhibitor (ThermoScientific). IL-1β, IL-6, and TNF-α levels in su-

pernatants were determined by ELISA according to the manufacturer's protocol (R&D

systems).

Statistical analysis

Each experiment was performed a minimum of two times. Statistical analyses were con-

ducted using GraphPad Prism version 6.0f (GraphPad Software). All studies comparing

two groups were analyzed with a Student's t-test. Those with three or more groups were

compared using an ordinary one-way ANOVA with Tukey's *post-hoc* test. $p < 0.05$ was

considered statistically significant.

ACKNOWLEDGEMENTS

We thank members of the Doering laboratory for helpful discussions and assistance with

experiments. We also thank Dr. Thomas R. Kozel (University of Nevada School of Medi-

cine) for anti-GXM monoclonal antibody, and Dr. Michael Diamond and Dr. Wayne Yoko-

yama for immune deficient mouse lines.

This work was funded by National Institutes of Health grants R21 AI109623 and R01

AI102882 to TLD. LXL was partly supported by a National Research Science Award (T32

GM007200), a Sondra Schlesinger Graduate Fellowship (Washington University St. Louis

Microbiology Department), and a National Institute of Allergy and Infectious Diseases fel-

lowship (F30 AI120339). CRH was partly funded by a National Institute of Allergy and

Infectious Diseases training grant (T32 AI007172). SAK and JRM were supported by NIH

grant R01 AI111914. The funders had no role in study design, data collection and inter-

pretation, or the decision to submit the work for publication.

AUTHOR CONTRIBUTIONS

Conceptualization, LXL, CRH, SAK, and TLD; Methodology, LXL, CRH, JRM, SAK, TLD;

Investigation, LXL, CRH, and JRM; Writing – Original Draft, LXL and CRH; Writing – Re-

view & Editing, CRH, JRM, SAK, and TLD; Funding Acquisition, LXL, SAK and TLD; Re-

sources, SAK and TLD; Supervision, SAK and TLD.

DECLARATION OF INTERESTS

LXL, CRH, JRM, SAK and TLD have no interests to declare

References

References

1. Kwon-Chung KJ, Fraser JA, Doering TL, Wang Z, Janbon G, Idnurm A, Bahn YS.
2014. *Cryptococcus neoformans* and *Cryptococcus gattii,* the etiologic agents of
cryptococcosis. Cold Spring Harb Perspect Med 4:a019760.

2. Denning DW. 2016. Minimizing fungal disease deaths will allow the UNAIDS target
of reducing annual AIDS deaths below 500 000 by 2020 to be realized. Philos
Trans R Soc Lond B Biol Sci 371.

3. Rajasingham R, Smith RM, Park BJ, Jarvis JN, Govender NP, Chiller TM, Denning
DW, Loyse A, Boulware DR. 2017. Global burden of disease of HIV-associated
cryptococcal meningitis: an updated analysis. Lancet Infect Dis
doi:10.1016/S1473-3099(17)30243-8.

4. Wozniak KL, Levitz SM. 2011. T cell and dendritic cell immune responses to
Cryptococcus, p p 387-396. *In* Heitman J, Kozel TR, Kwon-Chung KJ, Perfect JR,
Casadeval A (ed), Cryptococcus. ASM Press, Washington, DC.

5. Zhang Y, Wang F, Tompkins KC, McNamara A, Jain AV, Moore BB, Toews GB,
Huffnagle GB, Olszewski MA. 2009. Robust Th1 and Th17 immunity supports
pulmonary clearance but cannot prevent systemic dissemination of highly virulent
Cryptococcus neoformans H99. Am J Pathol 175:2489-500.

6. Wozniak KL, Hardison SE, Kolls JK, Wormley FL. 2011. Role of IL-17A on
resolution of pulmonary *C. neoformans* infection. PLoS One 6:e17204.

7. Murdock BJ, Huffnagle GB, Olszewski MA, Osterholzer JJ. 2014. Interleukin-17A
enhances host defense against cryptococcal lung infection through effects

470   mediated by leukocyte recruitment, activation, and gamma interferon production.

471   Infect Immun 82:937-48.

472 8. Schulze B, Piehler D, Eschke M, von Buttlar H, Kohler G, Sparwasser T, Alber G.

473   2014. CD4(+) FoxP3(+) regulatory T cells suppress fatal T helper 2 cell immunity

474   during pulmonary fungal infection. Eur J Immunol 44:3596-604.

475 9. Wiesner DL, Smith KD, Kotov DI, Nielsen JN, Bohjanen PR, Nielsen K. 2016.

476   Regulatory T cell induction and retention in the lungs drives suppression of

477   detrimental type 2 Th cells during pulmonary cryptococcal infection. J Immunol

478   196:365-74.

479 10. Wiesner DL, Specht CA, Lee CK, Smith KD, Mukaremera L, Lee ST, Lee CG, Elias

480   JA, Nielsen JN, Boulware DR, Bohjanen PR, Jenkins MK, Levitz SM, Nielsen K.

481   2015. Chitin recognition via chitotriosidase promotes pathologic type-2 helper T

482   cell responses to cryptococcal infection. PLoS Pathog 11:e1004701.

483 11. Wolfert MA, Boons GJ. 2013. Adaptive immune activation: glycosylation does

484   matter. Nat Chem Biol 9:776-84.

485 12. Zaragoza O, Rodrigues ML, De Jesus M, Frases S, Dadachova E, Casadevall A.

486   2009. The capsule of the fungal pathogen *Cryptococcus neoformans*. Adv Appl

487   Microbiol 68:133-216.

488 13. Yauch LE, Lam JS, Levitz SM. 2006. Direct inhibition of T-cell responses by the

489   Cryptococcus capsular polysaccharide glucuronoxylomannan. PLoS Pathog

490   2:e120.

491 14. Agustinho DP, Miller LC, Li LX, Doering TL. 2018. Peeling the onion: the outer

492   layers of *Cryptococcus neoformans*. Mem Inst Oswaldo Cruz 113:e180040.

493   15.   Wang ZA, Li LX, Doering TL. 2018. Unraveling synthesis of the cryptococcal cell

494   wall and capsule. Glycobiology 28:719-730.

495   16.   Cherniak R, Valafar H, Morris LC, Valafar F. 1998. *Cryptococcus neoformans*

496   chemotyping by quantitative analysis of 1H nuclear magnetic resonance spectra

497   of glucuronoxylomannans with a computer-simulated artificial neural network. Clin

498   Diagn Lab Immunol 5:146-59.

499   17.   Castle SA, Owuor EA, Thompson SH, Garnsey MR, Klutts JS, Doering TL, Levery

500   SB. 2008. Beta1,2-xylosyltransferase Cxt1p is solely responsible for xylose

501   incorporation into *Cryptococcus neoformans* glycosphingolipids. Eukaryot Cell

502   7:1611-5.

503   18.   Klutts JS, Levery SB, Doering TL. 2007. A beta-1,2-xylosyltransferase from

504   *Cryptococcus neoformans* defines a new family of glycosyltransferases. J Biol

505   Chem 282:17890-9.

506   19.   Park JN, Lee DJ, Kwon O, Oh DB, Bahn YS, Kang HA. 2012. Unraveling unique

507   structure and biosynthesis pathway of N-linked glycans in human fungal pathogen

508   *Cryptococcus neoformans* by glycomics analysis. J Biol Chem 287:19501-15.

509   20.   Lee DJ, Bahn YS, Kim HJ, Chung SY, Kang HA. 2015. Unraveling the novel

510   structure and biosynthetic pathway of O-linked glycans in the Golgi apparatus of

511   the human pathogenic yeast *Cryptococcus neoformans*. J Biol Chem 290:1861-

512   73.

513   21.   Li LX, Rautengarten C, Heazlewood JL, Doering TL. 2018. Xylose donor transport

514   is critical for fungal virulence. PLoS Pathog 14:e1006765.

22. Randall TD, Carragher DM, Rangel-Moreno J. 2008. Development of secondary

lymphoid organs. Annu Rev Immunol 26:627-50.

23. Hwang JY, Randall TD, Silva-Sanchez A. 2016. Inducible bronchus-associated

lymphoid tissue: taming inflammation in the lung. Front Immunol 7:258.

24. Moyron-Quiroz JE, Rangel-Moreno J, Kusser K, Hartson L, Sprague F, Goodrich

S, Woodland DL, Lund FE, Randall TD. 2004. Role of inducible bronchus

associated lymphoid tissue (iBALT) in respiratory immunity. Nat Med 10:927-34.

25. Geurts van Kessel CH, Willart MA, Bergen IM, van Rijt LS, Muskens F, Elewaut D,

Osterhaus AD, Hendriks R, Rimmelzwaan GF, Lambrecht BN. 2009. Dendritic

cells are crucial for maintenance of tertiary lymphoid structures in the lung of

influenza virus-infected mice. J Exp Med 206:2339-49.

26. Koguchi Y, Kawakami K. 2002. Cryptococcal infection and Th1-Th2 cytokine

balance. Int Rev Immunol 21:423-38.

27. Huffnagle GB, Boyd MB, Street NE, Lipscomb MF. 1998. IL-5 is required for

eosinophil recruitment, crystal deposition, and mononuclear cell recruitment during

a pulmonary *Cryptococcus neoformans* infection in genetically susceptible mice

(C57BL/6). J Immunol 160:2393-400.

28. Leopold Wager CM, Hole CR, Wozniak KL, Wormley FL, Jr. 2016. Cryptococcus

and Phagocytes: Complex Interactions that Influence Disease Outcome. Front

Microbiol 7:105.

29. Bar-Peled M, Griffith CL, Doering TL. 2001. Functional cloning and

characterization of a UDP-glucuronic acid decarboxylase: the pathogenic fungus

Cryptococcus neoformans elucidates UDP-xylose synthesis. Proc Natl Acad Sci U S A 98:12003-12008.

30. Grijpstra J, Tefsen B, van Die I, de Cock H. 2009. The *Cryptococcus neoformans cap10* and *cap59* mutant strains, affected in glucuronoxylomannan synthesis, differentially activate human dendritic cells. FEMS Immunol Med Microbiol 57:142-50.

31. Rittershaus PC, Kechichian TB, Allegood JC, Merrill AH, Jr., Hennig M, Luberto C, Del Poeta M. 2006. Glucosylceramide synthase is an essential regulator of pathogenicity of *Cryptococcus neoformans*. J Clin Invest 116:1651-9.

32. Gish SR, Maier EJ, Haynes BC, Santiago-Tirado FH, Srikanta DL, Ma CZ, Li LX, Williams M, Crouch EC, Khader SA, Brent MR, Doering TL. 2016. Computational Analysis Reveals a Key Regulator of Cryptococcal Virulence and Determinant of Host Response. MBio 7.

33. Decken K, Kohler G, Palmer-Lehmann K, Wunderlin A, Mattner F, Magram J, Gately MK, Alber G. 1998. Interleukin-12 is essential for a protective Th1 response in mice infected with *Cryptococcus neoformans*. Infect Immun 66:4994-5000.

34. Oppmann B, Lesley R, Blom B, Timans JC, Xu Y, Hunte B, Vega F, Yu N, Wang J, Singh K, Zonin F, Vaisberg E, Churakova T, Liu M, Gorman D, Wagner J, Zurawski S, Liu Y, Abrams JS, Moore KW, Rennick D, de Waal-Malefyt R, Hannum C, Bazan JF, Kastelein RA. 2000. Novel p19 protein engages IL-12p40 to form a cytokine, IL-23, with biological activities similar as well as distinct from IL-12. Immunity 13:715-25.

35. Billmyre RB, Applen Clancey S, Li LX, Doering TL, Heitman J. 2020. 5-

fluorocytosine resistance is associated with hypermutation and alterations in

capsule biosynthesis in Cryptococcus. Nat Commun 11:127.

36. Lupo P, Chang YC, Kelsall BL, Farber JM, Pietrella D, Vecchiarelli A, Leon F,

Kwon-Chung KJ. 2008. The presence of capsule in *Cryptococcus neoformans*

influences the gene expression profile in dendritic cells during interaction with the

fungus. Infect Immun 76:1581-9.

37. Eddens T, Elsegeiny W, Garcia-Hernadez ML, Castillo P, Trevejo-Nunez G,

Serody K, Campfield BT, Khader SA, Chen K, Rangel-Moreno J, Kolls JK. 2017.

Pneumocystis-driven inducible bronchus-associated lymphoid tissue formation

requires Th2 and Th17 immunity. Cell Rep 18:3078-3090.

38. Haugen RK, Baker RD. 1954. The pulmonary lesions in cryptococcosis with

special reference to subpleural nodules. Am J Clin Pathol 24:1381-90.

39. Slight SR, Rangel-Moreno J, Gopal R, Lin Y, Fallert Junecko BA, Mehra S, Selman

M, Becerril-Villanueva E, Baquera-Heredia J, Pavon L, Kaushal D, Reinhart TA,

Randall TD, Khader SA. 2013. CXCR5(+) T helper cells mediate protective

immunity against tuberculosis. J Clin Invest 123:712-26.

FIGURES WITH LEGENDS

SUPPLEMENTARY FIGURES WITH LEGENDS

SUPPLEMENTARY TABLES AND REFERENCES

FIGURES WITH LEGENDS

Figure 1. Fungal burden and pulmonary cytokine production in WT and *uxt1Δ uxt2Δ* mice. All panels show the combined results of two independent experiments (5 mice per group per experiment). (A-C) Tissue homogenates of infected A/JCr mice were plated for CFUs at the indicated dpi (open symbols, WT; red symbols, *uxt1Δ uxt2Δ*; dashed line, initial inoculum). Values from individual mice are plotted along with the mean ± SEM. (D-I) Cytokine levels in lung homogenates at the indicated dpi (dashed line, naïve; gray bars, WT; red bars, *uxt1Δ uxt2Δ*). *, *p* < 0.05; **, *p* < 0.01; ***, *p* < 0.005 for Student's t-test comparing WT to mutant.

Figure 2. Accumulation of T and B cells in the lungs of mice infected with *uxt1Δ uxt2Δ* cells is required to prevent disease progression. CD19+ B cells (A), CD4+ T cells (B), and CD8+ T cells (C) were enumerated in the lungs of infected mice by flow cytometry (gray, WT; red, *uxt1Δ uxt2Δ*). Plotted are the combined mean ± SEM values from two independent experiments (5

mice per group per experiment). *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.005$ by Student's t-test. (D) Survival of mice after intranasal inoculation with 5×10^4 cells of WT (n = 5) or *uxt1Δ uxt2Δ* (n = 10). Results shown are combined from two independent experiments. **, $p < 0.01$; ***, $p < 0.005$ by the log-rank test.

Figure 3. Cytokine production by dendritic cells stimulated with cryptococcal antigens. (A) CD11c+ DC populations in the lungs of infected mice were quantified by flow analysis. (B-E) DCs were co-incubated for 24 h with heat-killed cells of the strains indicated or subjected to no treatment (no tx) and levels of IL-1β (B), IL-6 (C), TNF-α (D), and IL-12p40 (E) in the supernatant were quantified by ELISA. Shown are the mean ± SD (n = 3) values for one representative experiment from five independent experiments that yielded similar results. **, $p < 0.01$ by one-way ANOVA with Tukey's post hoc test.

Figure 4. Mutant cryptococci are primarily restricted to lymphoid structures. Micrographs show Movat-stained lung sections from A/JCr mice infected with WT or *uxt1Δuxt2Δ* as above. Scale bars, 100 µm (top row) and 10 µm (bottom row). White arrows highlight fungi (stained turquoise). Images shown are representative of two independent studies (4-5 mice per group per experiment).

Figure 5. iBALT complexity in the context of *C. neoformans* infection. (A-B) Representative immunofluorescent staining of B cells (B220⁺), T cells (CD3⁺), plasma cells (IgG⁺), and CXCL13 in the lungs of infected A/JCr mice. Scale bar = 100 μm. (C-E) Morphometric analysis of iBALT structures in A/JCr mice infected with WT (gray bars) and *uxt1Δ uxt2Δ* (red bars). (F) Quantification of immunofluorescent staining of CXCL13. Plots show the combined mean ± SEM of two independent studies (4-5 mice per group per experiment); colors as in panels C-E. *, $p < 0.05$; ***, $p < 0.005$ by Student's t-test.

SUPPLEMENTARY MATERIAL

Supplemental Figure 1. Pro-inflammatory cytokines in lung homogenates at the indicated dpi (grey, naïve; white, WT; black, *uxt1Δ uxt2Δ*). Results shown are for individual mice from two independent experiments (n = 5 per group per experiment) and are plotted with the mean ± SEM. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.005$ by Student's t-test.

Supplemental Figure 2. Example of the multi-color flow cytometry gating strategy used to enumerate lung infiltrating immune cells.

Table S1. *C. neoformans* strains utilized in these studies.

C. neoformans strain[a]	Origin
KN99α	(Nielsen et al., 2003)
uxt1Δ	(Li et al., 2018)
UXT1	(Li et al., 2018)
uxt2Δ	(Li et al., 2018)
UXT2	(Li et al., 2018)
uxt1Δ uxt2Δ	(Li et al., 2018)
uxs1Δ	(Gish et al., 2016)
cap59Δ	(Moyrand and Janbon, 2004)

[a] All mutant strains are derived from KN99α

Table S2. Antibodies for flow analysis.

Antigen	Clone	Fluorophore	Dilution	Company
CD3	17-A2	APC-Cy7	1:250	BioLegend
CD4	GK1.5	BUV737	1:500	BD Biosciences
CD8a	53-6.7	BV650	1:125	BD Biosciences
CD11b	M1/70	BV510	1:250	BD Biosciences
CD11c	N418	PE	1:125	BD Biosciences
CD1/CD32 Fc Block	2.4G2	(not applicable)	1:500	BD Biosciences
CD19	1D3	BV786	1:125	BD Biosciences
CD45	30-F11	Pacific Blue	1:250	BioLegend
F4/80	BM8	APC	1:250	Thermo

REFERENCES FOR SUPPLEMENTAL MATERIAL

Gish, S.R., Maier, E.J., Haynes, B.C., Santiago-Tirado, F.H., Srikanta, D.L., Ma, C.Z., Li, L.X., Williams, M., Crouch, E.C., Khader, S.A.*, et al.* (2016). Computational analysis reveals a key regulator of Cryptococcal virulence and determinant of host response. mBio *7*, e00313-00316.

Li, L.X., Rautengarten, C., Heazlewood, J.L., and Doering, T.L. (2018). Xylose donor transport is critical for fungal virulence. PLoS Pathog *14*, e1006765.

Moyrand, F., and Janbon, G. (2004). *UGD1*, encoding the *Cryptococcus neoformans* UDP-glucose dehydrogenase, is essential for growth at 37 degrees C and for capsule biosynthesis. Eukaryot Cell *3*, 1601-1608.

Nielsen, K., Cox, G.M., Wang, P., Toffaletti, D.L., Perfect, J.R., and Heitman, J. (2003). Sexual cycle of *Cryptococcus neoformans var. grubii* and virulence of congenic a and alpha isolates. Infection and immunity *71*, 4831-4841.