Cryptococcus neoformans evades pulmonary immunity by modulating xylose transport

Lucy X. Li[a*], Camaron R. Hole[a], Javier Rangel-Moreno[b], Shabaana A. Khader[a], and Tamara L. Doering[a,#]

[a]Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, Missouri, USA

[b]Department of Medicine, Allergy/Immunology, and Rheumatology, University of Rochester School of Medicine, Rochester, New York, USA

Running Head: Xylose and cryptococcal immunity

[#]Address correspondence to Tamara L. Doering, doering@wustl.edu.

*Present address: Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

1 ABSTRACT

Cryptococcus neoformans is a fungal pathogen that kills almost 200,000 people each

year and is distinguished by abundant and unique surface glycan structures that are rich

in xylose. A mutant strain of *C. neoformans* that cannot transport xylose precursors into

the secretory compartment is severely attenuated in virulence in mice, yet surprisingly is

not cleared. We found that this strain failed to induce the non-protective T helper cell type

2 (Th2) responses characteristic of wild type infection, instead promoting sustained Inter-

leukin (IL)-12p40 induction and increased IL-17A (IL-17) production. It also stimulated

dendritic cells to release high levels of pro-inflammatory cytokines, a behavior we linked

to xylose expression. Finally, we discovered that inducible bronchus associated lymphoid

tissue (iBALT) forms in response to cryptococcal infection. Although iBALT formation is

delayed upon infection with xylose-deficient cryptococci, it subsequently restricts infec-

tion. These studies demonstrate that cryptococcal infection triggers iBALT formation and

elucidate the role of xylose in the modulation of host response to this fungal pathogen.

INTRODUCTION

Cryptococcus neoformans is a ubiquitous environmental fungus that causes pneumonia and meningitis. This opportunistic pathogen infects over a million individuals each year, with overall mortality exceeding 20% (1-3). Patient immune status is the main determinant of infection outcome, highlighting the importance of efficient host immune responses in the control of cryptococcosis.

C. neoformans infection begins when the organism is inhaled, followed by its proliferation in the lungs. This pulmonary infection may then disseminate to the brain, where it causes a frequently lethal meningoencephalitis. In the lungs, *C. neoformans* first interacts with host macrophages and dendritic cells (DC). These can engulf the fungus and migrate to draining lymph nodes, where they present antigen to T cells to initiate the adaptive immune response. The protective immune response to *C. neoformans* is primarily mediated by T cells. Although T helper cell type 1 (Th1) responses are protective against *C. neoformans* (4), it is likely that T helper type 17 (Th17) responses participate in pathogen clearance at mucosal surfaces (5-7). In contrast, T helper cell type 2 (Th2) responses are considered detrimental for protective immunity, leading to fungal growth and facilitating dissemination to the CNS (8, 9). The induction of these non-protective responses is not well understood, but cryptococcal glycans may contribute to this process (10).

Glycan structures participate in antigen recognition, immune activation, or immune regulation in multiple organisms (11). In *C. neoformans* the major virulence factor, a polysaccharide capsule, modulates the immune response via multiple mechanisms (12, 13). The capsule is primarily composed of two large polysaccharides: glucuronoxylomannan

(GXM) and glucuronoxylomannogalactan (GXMGal) (14, 15). One major component of

both polymers is xylose (Xyl), which comprises almost one fourth of the polysaccharide

capsule mass (16). Xylose also occurs in cryptococcal glycolipids (17) and as both Xyl

and Xyl-phosphate modifications of protein-linked glycans (18-20). We demonstrate here

that Xyl plays a critical role in the immune recognition of, and response to, *C. neoformans.*

The incorporation of Xyl into cryptococcal glycan structures occurs in the lumen of the

secretory pathway, via enzymatic reactions that use the substrate molecule UDP-Xyl.

This xylose donor is imported into the synthetic compartment by two transporters, Uxt1

and Uxt2 (21). A mutant strain that lacks both transporters (*uxt1Δ uxt2Δ*) exhibits defects

in glycosylation, including incomplete synthesis of capsule polysaccharides, and consid-

erable attenuation of virulence in mouse models of infection. Surprisingly, despite its com-

promised virulence, this attenuated strain persists in the lungs of infected mice. In pursu-

ing the mechanism of this persistence, we observed the formation of inducible bronchus

associated lymphoid tissue (iBALT) in the lungs of both wild-type (WT) and *uxt1Δ uxt2Δ*

infected mice. In the lungs of mice infected with the mutant strain, iBALT formation was

significantly delayed. However, once formed, it was better organized than the iBALT

formed when mice were infected with fully xylosylated WT *C. neoformans.* Furthermore,

animals infected with the mutant experienced a long-term asymptomatic infection rather

than the rapid and lethal disease observed upon WT infection. Together, these results

suggest that luminal Xyl modifications of cryptococcal glycoconjugates interfere with im-

mune recognition and activation of immune cells, events that are required for efficient

control of the pulmonary infection.

RESULTS

UDP-Xyl transport is required for cryptococcal virulence and dissemination

We previously observed that a *C. neoformans* strain lacking UDP-Xyl transporters (*uxt1Δ uxt2Δ*) was highly attenuated in virulence compared to the WT parental strain KN99α, yet persisted in the lungs of asymptomatic mice for at least 100 days after infection (21). To investigate the development of this chronic infection, we first examined the kinetics of fungal burden in mice after intranasal inoculation with WT and mutant strains. In A/JCr mice infected intranasally with WT fungi, which typically succumb to infection roughly 18 days post-infection (dpi), we observed a rapid and significant increase in pulmonary fungal burden (Fig 1A) along with significant dissemination to the spleen (Fig 1B) and brain (Fig 1C). In contrast, the pulmonary burden in *uxt1Δ uxt2Δ*-infected mice increased only gradually from initial inoculum levels to a modest peak at 63 dpi (Fig 1A). Colony-forming units in the lung then gradually declined, with complete clearance occurring in 30% of the mice by 189 dpi (Fig 1A). Although this infection was not limited to the lungs, only a fraction of the mice had measurable *uxt1Δ uxt2Δ* cells in the spleen (Fig 1B) and brain (Fig 1C) at 63 dpi. Furthermore, this limited dissemination was only observed at the times of highest lung burden (63 and 126 dpi), with no fungi detected at distal sites by 189 dpi (Fig 1B, Fig 1C). These results suggest that the A/JCr mice would eventually clear the *uxt1Δ uxt2Δ* infection.

Cryptococcal xylosylation influences differential production of pulmonary cytokines.

We next sought to identify the mechanism(s) responsible for the protracted host clearance

of *uxt1Δ uxt2Δ* infection. As summarized above, protection against *C. neoformans* is gen-

erally associated with production of type 1 cytokines like interleukin (IL)-2, IL-12, inter-

feron gamma (IFN-γ), and tumor necrosis factor alpha (TNF-α). In contrast, a Th2 re-

sponse to cryptococcal infection (characterized by the production of IL-4, IL-5, IL-10, and

IL-13) compromises control of pulmonary fungal burden (22) and facilitates dissemination

to the central nervous system (4, 22).

We hypothesized that the increased survival of mice infected with the mutant cryptococci

reflected an alteration in the immune response elicited by pulmonary infection. To test

this hypothesis, we analyzed cytokine levels over time in lung homogenates from mice

infected with WT or *uxt1Δ uxt2Δ* strains (Fig 1). A strong Th2-type response with signifi-

cant induction of IL-4 and IL-5 over the 15 days prior to sacrifice was detected in mice

infected with WT cryptococci (Fig 1D, Fig 1E). In contrast, *uxt1Δ uxt2Δ* cells elicited a

more quiescent cytokine response at early stages of infection. Furthermore, at the peak

of fungal burden in *uxt1Δ uxt2Δ*-infected mice (day 63, Fig. 1A), there was a significant

increase in the Th1 polarizing cytokine IL-12p40 compared to the levels observed in naïve

lungs (Fig 1F). Subsequently, IL-12p40 levels slowly declined, along with fungal burden

(Fig 1F). Although levels of IFNγ and TNFα (Fig 1G, Fig 1H) were similar to those in the

lungs of naïve mice, IL-17 trended higher at the peak of infection (Fig 1I). Overall, while

WT infection rapidly induced non-protective Th2 responses, infection with *uxt1Δ uxt2Δ*

failed to do so (even at the peak of fungal burden) and instead promoted the sustained

induction of IL-12p40 and increased production of IL-17.

Inducible bronchus-associated lymphoid tissue develops following cryptococcal infection

The differences that we observed in fungal burden and cytokine levels between mutant and WT infection suggested alterations in the immune response at the cellular level. To test this, we performed blinded morphometric analysis of inflammatory cell infiltration in the lungs at 15 dpi in mice infected with WT fungi and at 15 and 189 dpi in *uxt1Δ uxt2Δ*-infected mice. The animals infected with the mutant strain showed significantly reduced lung inflammation at both time points (Fig 2A), consistent with the more quiescent cytokine response reported above (Fig 1). We also noted distinctive lymphocytic accumulations proximal to the basal side of the bronchial epithelium in both groups of mice (Fig 2B). The lymphoid populations within these structures were organized into B cell follicles surrounded by CD3⁺ T cell cuffs (Fig 2C). The cellular composition of these inflammatory foci suggested inducible bronchus-associated lymphoid tissue (iBALT) (23).

iBALT is a tertiary lymphoid tissue that forms in the lung. It resembles secondary lymphoid organs (lymph nodes) in cellular composition and organization, with two distinctive zones: the B cell follicle and the T cell zone (24). The B cell follicle contains mainly follicular B cells and its organization is maintained by the continuous production of CXCL13. It is surrounded by a T cell zone that contains CD4 and CD8 T cells as well as dendritic cells (DCs) (25). At 15 dpi with WT cryptococci, we observed numerous iBALT structures, with compact B cell follicles surrounded by T cells (Fig 2C, Fig 2D). In contrast, we found considerably fewer and less organized B cell follicles at 15 dpi in mice infected with the *uxt1Δ uxt2Δ* strain (Fig 2C, Fig 2D). By late in infection (189 dpi for *uxt1Δ uxt2Δ* and 15

dpi for WT), although iBALT structures in mice infected with the mutant strain were still

less abundant than those found in lungs of WT (Fig. 2D), their average size was signifi-

cantly larger (Fig. 2E). Additionally, the iBALT was more organized and even contained

IgG$^+$ plasma cells (Fig 2C). Early in *uxt1Δ uxt2Δ* infection, the compromised formation of

iBALT structures was accompanied by increased peri-vascular accumulation of T cells

(Fig. 2F) and a significantly decreased production of CXCL13 (Fig 2G, Fig2H), a chemo-

kine essential for iBALT formation and organization (23). In contrast, at late time points

in this protracted developmental process, T cell cuffing was significantly reduced (Fig 2F)

and CXCL13 expression was increased (Fig 2G, Fig 1H). While infection with WT crypto-

cocci induced early and robust formation of iBALT, *uxt1Δ uxt2Δ* infection induced delayed

but ultimately larger iBALT.

To quantify the host adaptive immune response to *uxt1Δ uxt2Δ*, we used flow cytometry

to enumerate lymphocytes in the lungs of mice infected with WT or *uxt1Δ uxt2Δ* cells. By

12-15 dpi, the fraction of CD19$^+$ B cells (Fig 3A) or T cells (CD4$^+$ and CD8$^+$; Fig 3B, Fig

3C) in *uxt1Δ uxt2Δ*-infected mice significantly exceeded those of WT-infected mice, and

continued to accumulate as infection progressed. The total numbers of cells in these pop-

ulations remained elevated through 126 dpi (Fig. 3A-C, right), although they subsequently

declined, consistent with the decreased average size in T cell cuffing observed at 189

versus 15 dpi (Fig 2F).

Survival of uxt1Δ uxt2Δ-infected mice is dependent on T cells

To test the protective role of T and B cells in the prolonged survival of *uxt1Δ uxt2Δ*-in-

fected mice, we examined whether ablation of these cell types altered the course of in-

fection. Because the mouse strains used for these studies were engineered in the

C57BL/6 background, we included infections of this strain as controls. C57BL/6 mice are

less resistant to *C. neoformans* than A/JCr mice used above (26), as shown by their even-

tual susceptibility to *uxt1Δ uxt2Δ* infection (Fig. 3D, solid black line). In contrast, *Rag1-/-*

mice were significantly more susceptible to this strain, and succumbed by 55 dpi (Fig 3D;

solid blue line).

Since *Rag1-/-* mice lack both T and B cells, we also infected mice deficient in either T

cells (*TCRβ -/-*) or B cells (*µMT*) with *uxt1Δ uxt2Δ* cryptococci. B cell deficient mice ex-

hibited a more protracted course of disease than *Rag1-/-*, succumbing over an 85-day

period (Fig 3D, solid green line), while mice lacking only T cells succumbed to infection

with kinetics similar to those of the *Rag1-/-* mice (Fig 3D, solid red line). In contrast, all

mice inoculated with WT *C. neoformans* succumbed by 20 dpi (Fig 3D; dashed lines). Of

this group, *Rag1-/-* and *TCRβ-/-* mice still succumbed slightly faster than C57BL/6 and

µMT mice (Fig 3D). Together, these data show that T cells are the prominent cell type

responsible for increased survival of *uxt1Δ uxt2Δ*-infected mice, although B cells may also

play a role in protection.

Lack of luminal xylose modification in C. neoformans stimulates DC activation

DCs are critical for induction of protective immunity against *C. neoformans* (27), and their

activation and cytokine production are required for iBALT formation and maintenance (25,

28). During iBALT formation, antigen-driven activation of DCs and their consequent cyto-

kine and chemokine production are likely responsible for T and B cell recruitment (25),

which could potentially both induce the observed increase in these cell populations and

contribute to the protective effect of T cells during cryptococcal infection. For these rea-

sons, we wished to specifically measure DC responses. To do this we tested DC interac-

tions with WT cryptococcal cells, which exhibit antigens modified with Xyl, and *uxt1Δ*

uxt2Δ cryptococci, which are impaired in these modifications. We also tested single *uxtΔ*

mutants: *uxt1Δ* shows an intermediate level of Xyl utilization while *uxt2Δ* is similar to WT

(21). In these assays, we measured the ability of heat-killed fungi to stimulate the produc-

tion of pro-inflammatory cytokines by DC, comparing WT, *uxt1Δ uxt2Δ*, single *uxt* mu-

tants, *uxs1Δ* (which lacks all Xyl modification because it cannot synthesize UDP-Xyl), and

an acapsular control strain (*cap59Δ*) that induces a potent DC response (29). We found

that bone marrow derived DCs (BMDCs) cultured with *uxt1Δ uxt2Δ* cells released high

levels of IL-1β, IL-6, and TNF-α, similar to the levels observed following exposure to a

completely acapsular control strain, *cap59Δ* (Fig 4A-C). In contrast, individual *uxt* mutants

and *uxs1Δ* induced IL-1β and IL-6 levels similar to those induced by WT fungi (close to

background levels; Fig 4A, Fig 4B). TNF-α similarly showed the greatest response upon

challenge with *uxt1Δ uxt2Δ*, with levels like those induced by an acapsular strain, alt-

hough some increase was also noted with other xylose-deficient strains (*uxt1Δ* and

uxs1Δ). These results suggest that early interaction with DCs and consequent induction

of proinflammatory cytokines are regulated by Xyl expression in Cryptococcus.

DISCUSSION

In this study, we report for the first time that *C. neoformans* infection triggers iBALT for-

mation. Notably, the iBALT detected in our experimental infections resembles subpleural

nodules that have been observed in cryptococcosis patients (30). To our knowledge, the

only other fungus reported to induce iBALT is *Pneumocystis* (31). Further examination

will be necessary to establish the similarities between histological features of inflamma-

tory cell infiltration in mice and humans infected with *C. neoformans* and to delineate the

molecular host-pathogen interactions that drive iBALT formation following *C. neoformans*

infection.

When we infected mice with mutant fungi unable to use Xyl for luminal glycoconjugate

synthesis (*uxt1Δ uxt2Δ*), we observed a delay in fungal accumulation and in the organi-

zation of the iBALT structures. When *uxt1Δ uxt2Δ* burden did peak, it was accompanied

by a rise in IL-12p40 (Fig 1F), a cytokine normally required for inducing a protective re-

sponse against *C. neoformans* (32), which may have facilitated subsequent resolution

of the *uxt1Δ uxt2Δ* infection. A change in IL-12p40 level might reflect an increase in ei-

ther IL-23 or IL-12 (33), which stimulate a Th17 or Th1 type response, respectively; the

former is required for iBALT formation(31). Since we saw no difference in IFN-γ levels,

IL-12p40 is more likely associated with a Th17 response. We did observe a trend to-

wards increased IL-17 levels when *uxt1Δ uxt2Δ* burden was the highest, which was ab-

sent in WT infection. Our results show that cryptococcal infection can induce iBALT

structures, although this does not prevent mice from succumbing to the disease. We fur-

ther show that the kinetics of iBALT formation and its organization are dependent on vir-

ulence factors of the pathogen. Future studies should futher define the functional role of

iBALT in mediating immunity to Cryptococcus and determine whether the iBALT is me-

diating early and local immune protection in the lung.

DCs reacted more strongly to *uxt1Δ uxt2Δ* than to WT *in vitro*, releasing greater amounts

of pro-inflammatory cytokines (Fig 4A-C), even though the global cytokine response to

the mutant is less perturbed (Fig 1D-I). These cytokines may directly or indirectly lead to

the increased recruitment of T and B cells in *uxt1Δ uxt2Δ* infection (Fig 3A-C) and perhaps

accelerate compartmentalization into compact B cell follicles and discrete T cell areas in

iBALT (Fig 2). Such differences in cellular composition and organization, coupled with the

differences in the local cytokine production, may confer distinct properties on the iBALT

induced by WT and *uxt1Δ uxt2Δ*, influencing their ability to protect against *C. neoformans*.

We expected that the highly immunostimulatory behavior of *uxt1Δ uxt2Δ in vitro* reflected

the lack of Xyl modifications on its surface glycoconjugates, a consequence of the ab-

sence of UDP-Xyl transport into the secretory compartment. We were surprised, there-

fore, that mutants unable to synthesize UDP-Xyl (*uxs1Δ*), which similarly lack Xyl modifi-

cations on secreted glycoconjugates, did not phenocopy this broad and robust DC re-

sponse. This suggests that Uxs1 itself is required for the strong response seen in the

absence of xylosylation in the secretory pathway. It may be that this protein plays addi-

tional roles unrelated to glycosylation, or that the stimulatory component is dependent on

cytosolic UDP-Xyl, rather than UDP-Xyl that has been transported into the secretory path-

way.

Unlike encapsulated strains, *C. neoformans* mutants lacking capsule induce the upregu-

lation of multiple genes involved in cytokine responses and enhance antigen processing

and presentation by DCs (34). Interestingly, the cytokine levels induced in DC by the

uxt1Δ uxt2Δ mutant were similar to those induced by the acapsular mutant *cap59Δ* (Fig

4A-C). Xyl may thus be critical for the immune suppressive effects normally exerted by

capsule material. The mechanisms behind the robust DC activation seen in these two

mutants, including the cellular receptors and fungal components responsible, remain to

be determined. It is possible that the *uxt1Δ uxt2Δ* mutant lacks immune suppressive

mechanisms that are present in WT *C. neoformans* infection, and therefore induces ro-

bust cytokine responses and consequent recruitment of T and B cells to limit early infec-

tion. Further studies addressing the nature, type, and extent of the T and B cell responses

in the context of WT and *uxt1Δ uxt2Δ* infection should allow us to fully understand the the

initiation, formation, and organization of iBALT structures in the lung during *C. neofor-*

mans infection. Further, as iBALT is induced in WT *C. neoformans* infection where the

mice succumb to the disease, it is unclear if iBALT structures play a protective or patho-

logical role in immunity. Specific chemokine deficient mice, such as CXCL-13 deficient

animals that lack the ability to induce iBALT structures (35), will be useful in future studies

to define the function of iBALT in fungal infections.

Control of *uxt1Δ uxt2Δ* infection requires both T and B cells (Fig 3D), which suggests that

iBALT formation may play a key role in restricting disease. Even in the absence of T and

B cells, however, mutant-infected mice still survived more than twice as long as WT-in-

fected animals (Fig 3D). This may be due to the slightly slower growth of the mutant,

which we have previously observed (21).

We have found that *C. neoformans* infection induces iBALT formation, although the for-

mation of iBALT by itself is insufficient to protect against wild-type fungal infection. Inoc-

ulation with a *uxt1Δ uxt2Δ* strain, however, induced similar lymphoid structures, which do

appear to restrict infection despite their delayed organization. *In vitro*, this mutant induced

DCs to release much higher levels of proinflammatory cytokines than WT fungi. This is

presumably a consequence of the lack of surface Xyl modification, suggesting a role for

this structure in the immunosuppressive features of capsule polysaccharides or other sur-

face exposed or secreted glycoconjugates, although the full mechanism remains to be

defined. By influencing the dynamics of the inflammatory process, we may be able to

improve pulmonary responses to help control infection and use this information to help

guide novel therapies that stimulate the induction of protective lung fungal immunity.

MATERIALS AND METHODS

Fungal strains

C. neoformans strains (Table S1) were grown at 30 °C in YPD medium (1% [wt/vol] yeast

extract, 2% [wt/vol] peptone, 2% [wt/vol] dextrose) with shaking (230 rpm) or on YPD agar

plates (YPD medium with 2% [wt/vol] agar) supplemented with the following antibiotics as

appropriate: 100 μg/ml nourseothricin (NAT; Werner BioAgents) or Geneticin (G418; Invi-

trogen).

Mice

C57BL/6J (RRID: IMSR_JAX:000664) and A/JCr (RRID: IMSR_CRL:563) mice were ob-

tained from Jackson Laboratory. *Rag1-/-* (RRID: IMSR_JAX:002216), *TCRβ-/-* (RRID:

IMSR_JAX:002118), and μMT mice (RRID: IMSR_JAX:002288) (all on C57BL/6 back-

ground) were generously provided by Dr. Michael Diamond and Dr. Wayne Yokoyama

(Washington University School of Medicine); breeders were originally purchased from

Jackson Laboratory. All mice were 6 to 8 weeks old at the time of infection. A/JCr mice

were female; male and female mice were used for the other strains with gender and age

matched C57BL/6 controls.

Ethics statement

298   All animal protocols were reviewed and approved by the Animal Studies Committee of

299   the Washington University School of Medicine and conducted according to National Insti-

300   tutes of Health guidelines for housing and care of laboratory animals.

C. neoformans infection

Fungal strains were cultured overnight (O/N) and diluted to 10^6 cells/mL in sterile PBS.

Mice were intranasally inoculated with a 50 μL aliquot, and then weighed and monitored

daily. Infected mice were sacrificed if they lost >20% relative to peak weight or at specified

time points. At the time of sacrifice, mice were perfused intracardially with 10 mL sterile

PBS, and organs were processed as described below for fungal burden, histology, flow

cytometric analysis, and cytokine measurements.

Organ burden and cytokines

Brain and spleen homogenates were harvested and plated for CFU at the specified time points. 50 μL aliquots of the left lung homogenates (V_T = 1 mL) were similarly plated for CFUs. The remaining sample was assayed for pulmonary cytokine levels using the Bio-Plex Protein Array System (Bio-Rad Laboratories). Briefly, the lung homogenates were mixed with an equal volume of PBS/0.1% Triton 100x/2x protease inhibitor (Pierce EDTA-free protease inhibitor; Thermo Scientific), vortexed for 3 seconds, and clarified by centrifugation (2500 x g, 10 min). Supernatant fractions were then assayed using the Bio-Plex Pro Mouse Cytokine 23-Plex (Bio-Rad Laboratories) for the presence of IL-1α, IL-1β, IL-2, IL-3, IL-4, IL-5, IL-6, IL-9, IL-10, IL-12 (p40), IL-12 (p70), IL-13, IL-17A, granulocyte colony stimulating factor (G-CSF), granulocyte monocyte colony stimulating factor (GM-CSF), interferon-γ (IFN-γ), CXCL1/keratinocyte-derived chemokine (KC), CCL2/monocyte chemotactic protein-1 (MCP-1), CCL3/macrophage inflammatory protein-1α (MIP-1α), CCL4/MIP-1β, CCL5/regulated upon activation normal T cell expressed and secreted (RANTES), and tumor necrosis factor-α (TNF-α).

Immunofluorescence staining and histologic analysis

Mice were perfused intracardially with sterile PBS, and the lungs inflated with 10% formalin. Lung tissue was then fixed O/N in 10% formalin, and submitted to the Washington University Developmental Biology Histology Core for paraffin-embedding, sectioning, and staining with H & E. Immunofluorescence staining for B cells (APC-conjugated rat anti-mouse CD45R/B220, clone RA3-6B2, BD Biosciences, RRID: AB_627078), T cell cuffing

(CD3-ε, clone M-20, Santa Cruz Biotechnology, RRID: AB_631129), and CXCL13-pro-

ducing cells (goat α-mouse CXCL13, AF470, R&D Systems, RRID: AB_355378) was per-

formed as in Ref (31). Images were taken with a Zeiss Axioplan2 microscope and rec-

orded with a Hamamatsu camera. Blinded morphometric analysis of lung structures was

performed with the outline tool of Zeiss Axiovision software.

Flow cytometric analysis

The right lung of individual mice was enzymatically digested (in 5 mL RPMI with 1 mg/mL

collagenase type IV) at 37 °C with shaking (230 rpm) for 30 min, and then sequentially

passed through sterile 70 and 40 μm pore nylon strainers (BD Biosciences, San Jose,

CA). Red blood cells in the samples were lysed by treatment for 3 min on ice with 5 mL

ammonium-chloride-potassium lysing buffer (8.024 g/L NH_4Cl, 1.001 g/L $KHCO_3$, 2.722

mg/L EDTA•Na_2 $2H_2O$), followed by the addition of 2 volumes of PBS. The remaining cells

were pelleted (1000 x g, 5 min, 4 °C), washed twice with PBS, diluted to 10^6 cells/mL in

PBS, and stained with LIVE/DEAD fixable blue dead cell stain (1:1000; Thermo Scien-

tific). Following incubation in the dark for 30 min at 4 °C, cells were washed with PBS and

FACS buffer (2% fetal bovine serum in PBS) before resuspension in FACS buffer. Sam-

ples were then blocked with CD16/CD32 (1:500; Fc Block™; BD Biosciences, RRID:

AB_394656) for 5 min and incubated for 30 min with optimal concentrations of fluoro-

chrome-conjugated antibodies (Table S2) diluted in Brilliant Stain Buffer (BD Biosci-

ences). After three washes with FACS buffer, the cells were fixed in 2% formalde-

hyde/FACS buffer. For data acquisition, >50,000 events were collected on a BD LSRFor-

tessa X-20 flow cytometer (BD Biosciences), and the data were analyzed with FlowJo

V10 (Fig. S1; TreeStar). The absolute number of cells in each leukocyte subset was de-

termined by multiplying the absolute number of CD45$^+$ cells by the percentage of cells

stained by fluorochrome-labeled antibodies for each cell population analyzed.

Isolation of bone marrow derived cells

Bone marrow was flushed from the femurs and tibiae of C57BL/6 mice using RPMI. Cells

were collected (1000 x g, 5 min, 4°C), resuspended in RPMI, and counted using a hemo-

cytometer. To prepare bone marrow derived dendritic cells (BMDCs), 2 x 10^6 bone mar-

row cells were plated in 10 mL R10 medium (10% FBS, 0.4% Penicillin-Streptomycin, 2

mM L-glutamate, 50 µM 2-β-mercaptoethanol in RPMI) supplemented with 1 ng/mL GM-

CSF, and incubated at 37 °C and 5% CO_2. Medium was changed 3 and 6 days after

plating, and cells were harvested on day 8. BMDCs were enriched by depletion of BMMs

using biotinylated α-F4/80 antibody (eBioscience, RRID: AB_466657) and anti-biotin con-

jugated magnetic beads (Miltenyi Biotec). The BMDCs in the flow through were positively

selected using α-CD11c magnetic beads according to the manufacturer's protocol (Mil-

tenyi Biotec).

Dendritic cell assays

To assay the ability of fungal strains to activate BMDCs, *C. neoformans* strains of interest

were grown O/N, washed in PBS, and incubated at 65 °C for 15 min to heat kill (HK) the

fungi. BMDCs and HK fungi (10^6 cells of each) were then co-incubated for 24 h, sedi-

mented, and the supernatant fractions transferred to 1.5 mL centrifuge tubes containing

10 μL of 100x protease inhibitor (ThermoScientific). IL-1β, IL-6, and TNF-α levels in su-

pernatants were determined by ELISA according to the manufacturer's protocol (R&D

systems).

Statistical analysis

Each experiment was performed a minimum of two times. Statistical analyses were con-

ducted using GraphPad Prism version 6.0f (GraphPad Software). All studies comparing

two groups were analyzed with a Student's t-test. Those with three or more groups were

compared using an ordinary one-way ANOVA with Tukey's *post-hoc* test. $p < 0.05$ was

considered statistically significant.

ACKNOWLEDGEMENTS

We thank members of the Doering laboratory for helpful discussions and assistance with

experiments. We also thank Dr. Thomas R. Kozel (University of Nevada School of Medi-

cine) for the anti-GXM monoclonal antibody, and Dr. Michael Diamond and Dr. Wayne

Yokoyama for immune deficient mouse lines.

This work was funded by National Institutes of Health grants R21 AI109623 and R01

AI102882 to TLD. LXL was partly supported by a National Research Science Award (T32

GM007200), a Sondra Schlesinger Graduate Fellowship (Washington University St. Louis

Microbiology Department), and a National Institute of Allergy and Infectious Diseases fel-

lowship (F30 AI120339). CRH was partly funded by a National Institute of Allergy and

Infectious Diseases training grant (T32 AI007172). SAK and JRM were supported by NIH

grant R01 AI111914. The funders had no role in study design, data collection and inter-

pretation, or the decision to submit the work for publication.

403   AUTHOR CONTRIBUTIONS

404   Conceptualization, LXL, CRH, SAK, and TLD; Methodology, LXL, CRH, JRM, SAK, TLD;

405   Investigation, LXL, CRH, and JRM; Writing – Original Draft, LXL and CRH; Writing – Re-

406   view & Editing, LXL, CRH, JRM, SAK, and TLD; Funding Acquisition, LXL, SAK and TLD;

407   Resources, SAK and TLD; Supervision, SAK and TLD.

409   DECLARATION OF INTERESTS

References

1.   Kwon-Chung KJ, Fraser JA, Doering TL, Wang Z, Janbon G, Idnurm A, Bahn YS.

2014. *Cryptococcus neoformans* and *Cryptococcus gattii,* the etiologic agents of

cryptococcosis. Cold Spring Harb Perspect Med 4:a019760.

2.   Denning DW. 2016. Minimizing fungal disease deaths will allow the UNAIDS target

of reducing annual AIDS deaths below 500 000 by 2020 to be realized. Philos

Trans R Soc Lond B Biol Sci 371.

3.   Rajasingham R, Smith RM, Park BJ, Jarvis JN, Govender NP, Chiller TM, Denning

DW, Loyse A, Boulware DR. 2017. Global burden of disease of HIV-associated

cryptococcal   meningitis: an   updated   analysis.   Lancet   Infect   Dis

doi:10.1016/S1473-3099(17)30243-8.

4.   Wozniak KL, Levitz SM. 2011. T cell and dendritic cell immune responses to

Cryptococcus, p p 387-396. *In* Heitman J, Kozel TR, Kwon-Chung KJ, Perfect JR,

Casadeval A (ed), Cryptococcus. ASM Press, Washington, DC.

5.   Zhang Y, Wang F, Tompkins KC, McNamara A, Jain AV, Moore BB, Toews GB,

Huffnagle GB, Olszewski MA. 2009. Robust Th1 and Th17 immunity supports

pulmonary clearance but cannot prevent systemic dissemination of highly virulent

Cryptococcus neoformans H99. Am J Pathol 175:2489-500.

6.   Wozniak KL, Hardison SE, Kolls JK, Wormley FL. 2011. Role of IL-17A on

resolution of pulmonary *C. neoformans* infection. PLoS One 6:e17204.

7.   Murdock BJ, Huffnagle GB, Olszewski MA, Osterholzer JJ. 2014. Interleukin-17A

enhances host defense against cryptococcal lung infection through effects

mediated by leukocyte recruitment, activation, and gamma interferon production.

Infect Immun 82:937-48.

8.   Schulze B, Piehler D, Eschke M, von Buttlar H, Kohler G, Sparwasser T, Alber G.

2014. CD4(+) FoxP3(+) regulatory T cells suppress fatal T helper 2 cell immunity

during pulmonary fungal infection. Eur J Immunol 44:3596-604.

9.   Wiesner DL, Smith KD, Kotov DI, Nielsen JN, Bohjanen PR, Nielsen K. 2016.

Regulatory T cell induction and retention in the lungs drives suppression of

detrimental type 2 Th cells during pulmonary cryptococcal infection. J Immunol

196:365-74.

10. Wiesner DL, Specht CA, Lee CK, Smith KD, Mukaremera L, Lee ST, Lee CG, Elias

JA, Nielsen JN, Boulware DR, Bohjanen PR, Jenkins MK, Levitz SM, Nielsen K.

2015. Chitin recognition via chitotriosidase promotes pathologic type-2 helper T

cell responses to cryptococcal infection. PLoS Pathog 11:e1004701.

11. Wolfert MA, Boons GJ. 2013. Adaptive immune activation: glycosylation does

matter. Nat Chem Biol 9:776-84.

12. Zaragoza O, Rodrigues ML, De Jesus M, Frases S, Dadachova E, Casadevall A.

2009. The capsule of the fungal pathogen *Cryptococcus neoformans*. Adv Appl

Microbiol 68:133-216.

13. Yauch LE, Lam JS, Levitz SM. 2006. Direct inhibition of T-cell responses by the

Cryptococcus capsular polysaccharide glucuronoxylomannan. PLoS Pathog

2:e120.

14. Agustinho DP, Miller LC, Li LX, Doering TL. 2018. Peeling the onion: the outer

layers of *Cryptococcus neoformans*. Mem Inst Oswaldo Cruz 113:e180040.

15.   Wang ZA, Li LX, Doering TL. 2018. Unraveling synthesis of the cryptococcal cell

460   wall and capsule. Glycobiology 28:719-730.

461   16.   Cherniak R, Valafar H, Morris LC, Valafar F. 1998. *Cryptococcus neoformans*

462   chemotyping by quantitative analysis of 1H nuclear magnetic resonance spectra

463   of glucuronoxylomannans with a computer-simulated artificial neural network. Clin

464   Diagn Lab Immunol 5:146-59.

465   17.   Castle SA, Owuor EA, Thompson SH, Garnsey MR, Klutts JS, Doering TL, Levery

466   SB. 2008. Beta1,2-xylosyltransferase Cxt1p is solely responsible for xylose

467   incorporation into *Cryptococcus neoformans* glycosphingolipids. Eukaryot Cell

468   7:1611-5.

469   18.   Klutts JS, Levery SB, Doering TL. 2007. A beta-1,2-xylosyltransferase from

470   *Cryptococcus neoformans* defines a new family of glycosyltransferases. J Biol

471   Chem 282:17890-9.

472   19.   Park JN, Lee DJ, Kwon O, Oh DB, Bahn YS, Kang HA. 2012. Unraveling unique

473   structure and biosynthesis pathway of N-linked glycans in human fungal pathogen

474   *Cryptococcus neoformans* by glycomics analysis. J Biol Chem 287:19501-15.

475   20.   Lee DJ, Bahn YS, Kim HJ, Chung SY, Kang HA. 2015. Unraveling the novel

476   structure and biosynthetic pathway of O-linked glycans in the Golgi apparatus of

477   the human pathogenic yeast *Cryptococcus neoformans*. J Biol Chem 290:1861-

478   73.

479   21.   Li LX, Rautengarten C, Heazlewood JL, Doering TL. 2018. Xylose donor transport

480   is critical for fungal virulence. PLoS Pathog 14:e1006765.

22.   Koguchi Y, Kawakami K. 2002. Cryptococcal infection and Th1-Th2 cytokine

balance. Int Rev Immunol 21:423-38.

23.   Moyron-Quiroz JE, Rangel-Moreno J, Kusser K, Hartson L, Sprague F, Goodrich

S, Woodland DL, Lund FE, Randall TD. 2004. Role of inducible bronchus

associated lymphoid tissue (iBALT) in respiratory immunity. Nat Med 10:927-34.

24.   Randall TD, Carragher DM, Rangel-Moreno J. 2008. Development of secondary

lymphoid organs. Annu Rev Immunol 26:627-50.

25.   Hwang JY, Randall TD, Silva-Sanchez A. 2016. Inducible bronchus-associated

lymphoid tissue: taming inflammation in the lung. Front Immunol 7:258.

26.   Huffnagle GB, Boyd MB, Street NE, Lipscomb MF. 1998. IL-5 is required for

eosinophil recruitment, crystal deposition, and mononuclear cell recruitment during

a pulmonary *Cryptococcus neoformans* infection in genetically susceptible mice

(C57BL/6). J Immunol 160:2393-400.

27.   Leopold Wager CM, Hole CR, Wozniak KL, Wormley FL, Jr. 2016. Cryptococcus

and Phagocytes: Complex Interactions that Influence Disease Outcome. Front

Microbiol 7:105.

28.   Geurts van Kessel CH, Willart MA, Bergen IM, van Rijt LS, Muskens F, Elewaut D,

Osterhaus AD, Hendriks R, Rimmelzwaan GF, Lambrecht BN. 2009. Dendritic

cells are crucial for maintenance of tertiary lymphoid structures in the lung of

influenza virus-infected mice. J Exp Med 206:2339-49.

29.   Grijpstra J, Tefsen B, van Die I, de Cock H. 2009. The *Cryptococcus neoformans*

cap10 and *cap59* mutant strains, affected in glucuronoxylomannan synthesis,

differentially activate human dendritic cells. FEMS Immunol Med Microbiol 57:142-

50.

30. Haugen RK, Baker RD. 1954. The pulmonary lesions in cryptococcosis with

special reference to subpleural nodules. Am J Clin Pathol 24:1381-90.

31. Eddens T, Elsegeiny W, Garcia-Hernadez ML, Castillo P, Trevejo-Nunez G,

Serody K, Campfield BT, Khader SA, Chen K, Rangel-Moreno J, Kolls JK. 2017.

Pneumocystis-driven inducible bronchus-associated lymphoid tissue formation

requires Th2 and Th17 immunity. Cell Rep 18:3078-3090.

32. Decken K, Kohler G, Palmer-Lehmann K, Wunderlin A, Mattner F, Magram J,

Gately MK, Alber G. 1998. Interleukin-12 is essential for a protective Th1 response

in mice infected with *Cryptococcus neoformans*. Infect Immun 66:4994-5000.

33. Oppmann B, Lesley R, Blom B, Timans JC, Xu Y, Hunte B, Vega F, Yu N, Wang

J, Singh K, Zonin F, Vaisberg E, Churakova T, Liu M, Gorman D, Wagner J,

Zurawski S, Liu Y, Abrams JS, Moore KW, Rennick D, de Waal-Malefyt R, Hannum

C, Bazan JF, Kastelein RA. 2000. Novel p19 protein engages IL-12p40 to form a

cytokine, IL-23, with biological activities similar as well as distinct from IL-12.

Immunity 13:715-25.

34. Lupo P, Chang YC, Kelsall BL, Farber JM, Pietrella D, Vecchiarelli A, Leon F,

Kwon-Chung KJ. 2008. The presence of capsule in *Cryptococcus neoformans*

influences the gene expression profile in dendritic cells during interaction with the

fungus. Infect Immun 76:1581-9.

35. Slight SR, Rangel-Moreno J, Gopal R, Lin Y, Fallert Junecko BA, Mehra S, Selman

M, Becerril-Villanueva E, Baquera-Heredia J, Pavon L, Kaushal D, Reinhart TA,

Randall TD, Khader SA. 2013. CXCR5(+) T helper cells mediate protective

immunity against tuberculosis. J Clin Invest 123:712-26.

FIGURES WITH LEGENDS

Figure 1. Fungal burden and pulmonary cytokine production in WT and *uxt1Δ uxt2Δ* mice. All panels show the combined results of two independent experiments (5 mice per group per experiment). (A-C) Tissue homogenates of infected A/JCr mice were plated for CFUs at the indicated dpi (open circles, WT; red circles, *uxt1Δ uxt2Δ*; dashed line, initial inoculum). Values from individual mice are plotted along with the mean ± SEM. (D-I) Levels of the cytokines indicated in lung homogenates over time after infection (gray bars, WT; red bars, *uxt1Δ uxt2Δ*; dashed line, naïve). *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.005$ for Student's t-test comparing wild type to mutant.

Figure 2. Inflammation and iBALT complexity in the context of *C. neoformans* infection. (A) Percent of area covered by inflammatory cells in individual lungs and (B) representative H&E micrographs from lungs collected at 15 and 189 dpi are shown. Scale bar = 100 μm. (C) Representative immunofluorescent staining of B cells (B220⁺), T cells (CD3⁺), and plasma cells (IgG⁺) in the lungs of infected A/JCr mice. Scale bar = 100 μm. (D-F) Morphometric analysis of iBALT structures in A/JCr mice infected with WT and *uxt1Δ uxt2Δ*. (G,H) Quantification and examples of immunofluorescent staining of CXCL13 (red). Nuclei are stained with DAPI (blue) and scale bar = 100 μm. Plots show the combined mean ± SEM of two independent studies (4-5 mice per group per experiment). *, $p < 0.05$; ***, $p < 0.005$ by Student's t-test.

Figure 3. Accumulation of T and B cells in the lungs of mice infected with *uxt1Δ uxt2Δ* cells is required to prevent disease progression. (A) CD19+ B cells, (B) CD4+ T cells, and (C) CD8+ T cells were enumerated in the lungs of infected mice by flow cytometry (gray, WT; red, *uxt1Δ uxt2Δ*). Plotted are the combined mean ± SEM values from two independent experiments (5

mice per group per experiment). *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.005$ by Student's t-test. (D) Survival of mice after intranasal inoculation with 5×10^4 cells of WT (n = 5) or *uxt1Δ uxt2Δ* (n = 10). Results shown are combined from two independent experiments. **, $p < 0.01$; ***, $p < 0.005$ by the log-rank test.

Figure 4. Cytokine production by dendritic cells stimulated with cryptococcal antigens. DCs were co-incubated for 24 h with heat-killed cells of the strains indicated or subjected to no treatment (no tx). Levels of (B) IL-1β, (C) IL-6, and (D) TNF-α in the supernatant were quantified by ELISA. Data shown is the mean ± SD (n = 3) of one representative experiment from five independent experiments that yielded similar results. **, $p < 0.01$ by one-way ANOVA with Tukey's post hoc test.

SUPPLEMENTARY MATERIAL

Supplemental Figure 1. Example of the multi-color flow cytometry gating strategy used to enumerate lung infiltrating immune cells.

Table S1. *C. neoformans* strains utilized in these studies.

C. neoformans strain[a]	Origin
KN99α	(Nielsen et al., 2003)
uxt1Δ	(Li et al., 2018)
UXT1	(Li et al., 2018)
uxt2Δ	(Li et al., 2018)
UXT2	(Li et al., 2018)
uxt1Δ uxt2Δ	(Li et al., 2018)
uxs1Δ	(Gish et al., 2016)
cap59Δ	(Moyrand and Janbon, 2004)

[a] All mutant strains are derived from KN99α

Table S2. Antibodies for flow analysis.

Antigen	Clone	Fluorophore	Dilution	Company	RRID
CD3	17-A2	APC-Cy7	1:250	BioLegend	AB_2242784
CD4	GK1.5	BUV737	1:500	BD Biosciences	AB_2738734
CD8a	53-6.7	BV650	1:125	BD Biosciences	AB_2738084
CD16/CD32 Fc Block	2.4G2	(not applicable)	1:500	BD Biosciences	AB_394656
CD19	1D3	BV786	1:125	BD Biosciences	AB_2738141
CD45	30-F11	Pacific Blue	1:250	BioLegend	AB_493535

REFERENCES FOR SUPPLEMENTAL MATERIAL

Gish, S.R., Maier, E.J., Haynes, B.C., Santiago-Tirado, F.H., Srikanta, D.L., Ma, C.Z., Li, L.X., Williams, M., Crouch, E.C., Khader, S.A., *et al.* (2016). Computational analysis reveals a key regulator of Cryptococcal virulence and determinant of host response. mBio *7*, e00313-00316.

Li, L.X., Rautengarten, C., Heazlewood, J.L., and Doering, T.L. (2018). Xylose donor transport is critical for fungal virulence. PLoS Pathog *14*, e1006765.

Moyrand, F., and Janbon, G. (2004). UGD1, encoding the *Cryptococcus neoformans* UDP-glucose dehydrogenase, is essential for growth at 37 degrees C and for capsule biosynthesis. Eukaryot Cell *3*, 1601-1608.

Nielsen, K., Cox, G.M., Wang, P., Toffaletti, D.L., Perfect, J.R., and Heitman, J. (2003). Sexual cycle of *Cryptococcus neoformans var. grubii* and virulence of congenic a and alpha isolates. Infection and immunity *71*, 4831-4841.