A systemic approach provides insights into the salt stress adaptation mechanisms of

contrasting bread wheat genotypes

Running title: Salt stress adaptation mechanisms of bread wheat studied by a systemic

approach

Diana Duarte-Delgado[1], Said Dadshani[1], Heiko Schoof[2], Benedict C. Oyiga[1], Michael

Schneider[1], Boby Mathew[1], Jens Léon[1], Agim Ballvora[1*]

[1] INRES-Plant Breeding, University of Bonn, Bonn, Germany

[2] INRES-Crop Bioinformatics, University of Bonn, Bonn, Germany.

*To whom correspondence should be addressed. Katzenburgweg 5, 53115 Bonn, Germany. E

mail: ballvora@uni-bonn.de

Highlight: The implementation of a systemic approach provided insights into salt stress

adaptation response mechanisms of contrasting bread wheat genotypes from two mapping

populations at both osmotic and ionic phases.

Abstract

Bread wheat is one of the most important crops for human diet but the increasing soil

salinization is causing yield reductions worldwide. Physiological, genetic, transcriptomics and

bioinformatics analyses were integrated to study the salt stress adaptation response in bread

wheat. A comparative analysis to uncover the dynamic transcriptomic response of contrasting

genotypes from two wheat populations was performed at both osmotic and ionic phases in

time points defined by physiologic measurements. The differential stress effect on the

expression of photosynthesis, calcium binding and oxidative stress response genes in the

contrasting genotypes supported the greater photosynthesis inhibition observed in the

susceptible genotype at the osmotic phase. At the ionic phase genes involved in metal ion

binding and transporter activity were up-regulated and down-regulated in the tolerant and

susceptible genotypes, respectively. The stress effect on mechanisms related with protein

synthesis and breakdown was identified at both stress phases. Based on the linkage

disequilibrium blocks it was possible to select salt-responsive genes as potential components

operating in the salt stress response pathways leading to salt stress resilience specific traits.

Therefore, the implementation of a systemic approach provided insights into the adaptation

response mechanisms of contrasting bread wheat genotypes at both salt stress phases.

Keywords: Bread wheat, comparative transcriptomics, ionic stress, osmotic stress, QTL

dissection, salt stress, systemic approach.

Introduction

Bread wheat (*Triticum aestivum* L.) is a key staple crop for global food security and to feed

the world population by 2050 its production needs to be increased substantially (Curtis and

Halford, 2014). Therefore, breeding programs should emphasize on the genetic improvement

of complex traits to increase yield potential under growth limiting conditions (Hawkesford *et*

al., 2013). The genetic studies of complex traits in wheat are challenging because it is an

allohexapolyploid species containing three sub genomes (AABBDD) with highly repetitive

DNA sequences (85%) and a total genome size of 16 Gb (Loginova and Silkova, 2018;

IWGSC, 2018). Efforts from the IWGSC (International Wheat Genome Sequencing

Consortium) have resulted in the release of an annotated and highly contiguous chromosome-

level assembly sequence draft of the Chinese Spring cultivar representing 94% of the whole

genome (Shi and Ling, 2018; IWGSC; Borrill *et al.*, 2019).

The world is losing 2000 hectares of arable soil daily due to salt-induced degradation which is

a serious threat for global food security (Zaman *et al.*, 2016). Among all the abiotic stress

factors, soil salinity can cause significant yield reductions and decreased grain quality in

wheat (Zheng *et al.*, 2009). The salt stress adaptation response is a complex trait since the

changes caused in key physiological processes are the effect of a coordinated action of gene

networks in several metabolic pathways (Tuteja, 2007; Gupta and Huang, 2014). Therefore,

the development of cultivars with increased salinity tolerance can be facilitated by a better

understanding of the mechanistic basis underlying the stress adaptation response. This

objective can be achieved through the integration of a range of experimental and statistical

methods from various disciplines such as genetics, transcriptomics, bioinformatics and stress

physiology to develop systemic approaches to gain a comprehensive understanding of

relevant agronomic complex characters as the tolerance to saline conditions (Civelek and

Lusis, 2014).

High salinity leads to physiological drought conditions, causes ion toxicity and cell oxidative

damage that affect the plant growth (Tuteja, 2007; Gupta and Huang, 2014). Thus, plant

growth response to salinity comprises two phases. The first corresponds to the osmotic phase

which is independent of the sodium accumulation in tissues. The rapid and often transient

impact on plant growth in this phase is attributed to the osmotic effect of the salts in the

rhizosphere because of a reduced water potential (Ismail *et al.*, 2014; Parihar *et al.*, 2015;

Julkowska and Testerink, 2015). Consequently, the osmotic tolerant plants have the ability to

adapt to the drought aspect of the stress through the maintenance of the stomatal conductance

and the leaf turgor (Carillo *et al.*, 2011). The early signaling events in the osmotic phase that

occur within seconds to hours of salt stress exposure are crucial for the acclimation response

of the plants (Julkowska and Testerink, 2015). A model proposes that in the osmotic phase the

root senses salt stress and second messengers as ROS (Reactive Oxygen Species) and Ca^{2+} are

spread as signals to the aerial parts to trigger adaptive responses to cope with the Na^+ ions that

reach photosynthetic tissues and cause toxic effects in the following stress phase (Köster *et*

al., 2018). Second, the ionic phase continues as a result of salt accumulation in leaves and

takes days or even weeks to be manifested. In this phase the senescence of older leaves is

caused by the inability of the plant to tolerate the toxic concentrations of salts in the tissues

(Roy *et al.*, 2014; Ismail *et al.*, 2014). To reduce the toxicity effects, tolerant plants can

excrete Na^+ from leaves through the roots or compartmentalize Na^+ and Cl^- in vacuoles to

avoid toxic concentrations in the cytoplasm (Munns and Tester, 2008; James *et al.*, 2012).

Hence, the limiting effect of salinity on crop productivity in both stress phases is mainly due

to its effect on the photosynthetic process which results in a substantial decrease in the

biomass accumulation (Demetriou *et al.*, 2007; Silveira and Carvalho, 2016).

The genetic mapping studies have allowed the detection of statistical associations of

molecular markers with phenotypic values of a quantitative trait to find locations of

quantitative trait loci (QTL) in the genome of a given species (Kang *et al.*, 2016; Ishikawa,

2017). Several mapping analyses in bread wheat have identified QTL with effect on salt

stress-related traits (Xu *et al.*, 2013; Hussain *et al.*, 2017; Dadshani, 2018; Oyiga *et al.*, 2018).

The recent studies by Oyiga *et al*. (2018) and Dadshani (2018) identified QTL for salt stress-

related traits at germination, seedling and adult stages in association and advanced backcross-

QTL (AB-QTL) mapping populations, respectively. Because of the limited mapping

resolution of these studies, the linkage disequilibrium (LD) blocks where the QTL are

localized contain many genes that are potential causal quantitative trait genes (QTGs)

influencing the trait variation. A combination of genetic mapping analyses with

transcriptomics can greatly reduce the number of these genes and provide with strong

candidate QTGs for the identified QTL (González-Prendes *et al.*, 2017; Ishikawa, 2017). In

addition, transcriptomics analyses allow to identify the pathways that are regulated in the

osmotic and ionic phases and give insights into the salt stress regulation of the photosynthesis

from contrasting genotypes (Chaves *et al.*, 2009; Liu *et al.*, 2019). A time course

transcriptomic analysis is therefore appropriate to identify key events during the salt stress

response (Zhang *et al.*, 2016; Liu *et al.*, 2019).

The Next-Generation Sequencing platforms comprise rapidly evolving high throughput

methods which are becoming the election to provide new insights into the whole genome,

transcriptome, and epigenome of plants to assist the breeders to understand the biological

function of the genes (Yadav *et al.*, 2018). In contrast to regular RNA-seq, the Massive

Analysis of cDNA 3'-ends (MACE) sequencing protocol is an alternative to perform

transcriptomic analyses where a single sequence fragment represents one transcript

(Hrdlickova *et al.*, 2017; Tzfadia *et al.*, 2018). Therefore, the output from this protocol is

digital, strand-specific, and the quantification of the expression is simpler and more precise in

each library (Asmann *et al.*, 2009; Hrdlickova *et al.*, 2017). This sequencing strategy can also

contribute to better define gene models towards the 3'-ends (Tzfadia *et al.*, 2018).

This study presents a systemic approach that integrates the dynamic transcriptomic responses

of salt-tolerant and salt-sensitive wheat genotypes with photosynthetic activity and genetic

mapping studies. The main goal of our study was to provide insights into the molecular

mechanisms of the salt stress response in bread wheat through a comparative transcriptomic

analysis and the identification of potential candidate genes operating in salt stress response

pathways.

Methods

Contrasting genotypes from the mapping populations and tissue sampling

The elite German winter wheat cultivar Zentos (salt-tolerant; Belderok *et al.*, 2000) and the

synthetic genotype Syn86 (salt-susceptible; Lange and Jochemsen, 1992) were the contrasting

parents used to study the foliar transcriptome during the osmotic stress response (Kunert *et*

al., 2007; Dadshani, 2018). Altay2000 (salt-tolerant; Altay, 2012) and Bobur (salt-

susceptible;   Amanov,   2017;   available   at

http://www.iwwip.org/images/dosya/636220600428830708.pdf) were the contrasting Turkish

and Uzbeq winter wheat cultivars used to study the transcriptome during the ionic stress

response, respectively (Oyiga *et al.*, 2018). A differential photosynthesis response of Syn86

and Zentos was revealed by the sensor-based measurement of the time course of the

photosynthesis rate performed under hydroponics conditions with 100 mol m^{-3} NaCl

(Dadshani, 2018). This study allowed the identification of the "turning points" of the

photosynthesis rate from 0 to 45 min after stress exposure (ASE). As "turning points" were

considered the time points with maximum variation response, as revealed by the change of

direction from the curve slope (Fig. 1A).

Homogeneous seedlings adapted to hydroponics conditions were used for sampling leaves

under control conditions and a salt stress treatment of 100 mol m^{-3} NaCl. The detailed

procedures of the hydroponic systems are described in Dadshani (2018) and Oyiga *et al*.

(2016) for osmotic and ionic phase experiments, respectively. Osmotic stress conditions were

sampled in the photosynthesis turning time points identified at 8, 15 and 30 min (Fig. 1A) and

at 4 h ASE whereas control conditions were sampled at 0, 30 min and 4 h in plants grown in

hydroponic boxes without NaCl. Samples for the ionic stress conditions were collected at 11

days and 24 days in both salt stressed and unstressed plants. A biologically averaged

experiment was conducted where each sampled condition consisted on a pool of five leaves

from four plants that were harvested and immediately stored in liquid nitrogen for posterior

homogenization. This sampling strategy allowed an exploratory analysis with less amount of

sequencing data assuming that most expression measurements from RNA pools are similar to

the averages from the individuals that are included in the corresponding pools (Kendziorski *et*

al., 2005; Biswas *et al.*, 2013)*.*

MACE reads processing and mapping to the reference genome

The total RNA isolation and the MACE library construction were performed at GenXPro

GmbH (Frankfurt, Germany). An Illumina NextSeq 500 system was used to sequence

fourteen and eight libraries from the osmotic and ionic stress experiments, respectively. The

adapters from the reads were trimmed with Cutadapt (Martin, 2011). The quality control of

the prepared libraries was carried out using FastQC (Andrews, 2010) and the short reads with

less than 35 bp were removed with Trimmomatic (Bolger *et al.*, 2014). The retained reads

were aligned to the reference wheat genome assembly version "RefSeq v1.0" (Alaux *et al.*,

2018) using Tophat (Trapnell *et al.*, 2009). Assemblies of novel transcripts were produced

with the prediction tool of Cufflinks (Roberts *et al.*, 2011; Trapnell *et al.*, 2012). The markdup

tool from SAMtools was employed to produce deduplicated alignment files and estimate the

amount of read duplication (Li *et al.*, 2009).

A new annotation file was elaborated to count reads beyond the predicted 3'-ends of high

confidence (HC) and low confidence (LC) gene models (IWGSC, 2018) with the purpose to

contribute to gene model improvement and to better estimate gene expression levels. For that,

the annotated gene models were extended by 40 % downstream of the predicted 3'-end in the

case of intergenic regions greater than 1000 bp but smaller than three times the gene size.

When the intergenic distance was larger, the elongated target sequence corresponded to the

size of the gene. Then, the stranded option from the featureCounts tool (Liao *et al.*, 2014) of

the Subread software (Liao *et al.*, 2013) was used to count the unique mapped reads assigned

to the elongated HC and LC gene models and to novel predicted transcripts. The read count

data was normalized to counts per million. An average normalized value of 2.5 across

libraries from the same genotype was selected as threshold to define a transcriptomic

background aiming to select the genes adequately represented and to reduce the number of

low expressed transcripts that might cause sampling noise (Sha *et al.*, 2015; Lin *et al.*, 2016)

A merged alignment file with all reads from the libraries was generated to compare the

number of reads counted using the extended gene models with those counted employing the

reference annotation. To exemplify the improvement in transcript quantification with the

extended gene models, two windows of 5 Mbp (23 to 28 Mbp coordinates of the

chromosomes 5B and 7D) were inspected in the alignment files from two libraries. The online

web tool GenomeView (Abeel *et al.*, 2012) allowed the visualization of the alignments in the

reference genome to observe the coordinates of the reads scored beyond the 3'-end gene

boundaries.

Identification of salt-responsive genes and gene ontology (GO) enrichment analysis

After filtering the low expressed transcripts, salt-responsive genes were identified using the

raw count data of fragments as input in the GFOLD (generalized fold change) software (Feng

et al., 2012). The GFOLD value is a reliable estimator of the relative difference of gene

expression which allows the generation of gene rankings that are particularly useful when

biological averaging is available (Feng *et al.*, 2012). Density plots with $\log_{10}$ normalized

expression values were generated to assess whether count normalization was appropriate to

allow the comparison of the libraries from the same genotype (Klaus and Huber, 2016;

available at https://www.huber.embl.de/users/klaus/Teaching/DESeq2Predoc_2014.html). The

0 min condition was used as control for both 8 and 15 min ASE, with the assumption that no

physiological changes occur in this short period of time under normal conditions. A high

absolute GFOLD value indicated greater up- or down-regulation of the genes. Genes with

GFOLD values >1 or $<$ -1 were considered for further analyses. Venn diagrams were used

to summarize the comparisons of the salt-responsive genes across genotypes and time points.

The GO enrichment tool from the STEM software was implemented to distinguish the

categories of genes over-represented in the contrasting genotypes (Ernst *et al.*, 2005; Ernst

and Bar-Joseph, 2006) using the available GO category assignation in the annotation version

RefSeq v1.0 (Alaux *et al.*, 2018). Only the gene categories from the transcriptomic

background of each genotype were retained in the analysis (Timmons *et al.*, 2015). A

Bonferroni multiple hypothesis correction test was implemented, thus GO terms with a

corrected p-value < 0.001 and < 0.005 were considered as over-represented during the

osmotic and the ionic phases, respectively. Over-represented categories during the osmotic

phase were selected to graphic the time course expression profiles of the corresponding genes.

A LOESS (locally estimated scatterplot smoothing) curve was used to represent the

expression tendency of the clusters of genes.

Identification of potential QTGs

The identification of QTGs was done by localizing salt-responsive genes revealed with the

transcriptomic analysis within the LD blocks of the corresponding QTL. Thus, adjacent

markers in strong LD ($R^2 \geq 0.8$) with the significant single nucleotide polymorphism (SNP)

were assigned to one LD block (Cirilli *et al.*, 2018). Polymorphic markers in the contrasting

genotypes Altay2000 and Bobur were chosen from the association mapping analysis.

Afterwards, the positions of the LD blocks in the reference genome sequence RefSeq v1.0

were determined according to the IWGSC RefSeq v1.0 BLAST results of the SNPs-flanking

sequences (Alaux *et al.*, 2018). Consequently, the LD block coordinates enabled the

identification of genes whose transcripts showed up- or down- regulation upon salt treatment

and were recognized as potential candidate genes operating within the QTL. The wheat RNA-

seq atlas expVIP was used to compare the expression of the genes from the LD blocks with

the expression determined in other abiotic stress experiments in the species (Borrill *et al.*,

2016).

Results

Sequence processing and reference genome mapping

To identify up- or down-regulated transcripts during the salt stress response the samples

collected were sequenced using the MACE approach. An overview of the sequence

processing and reference genome mapping across both osmotic and ionic stress experiments is

presented in the Table 1. The sequencing process yielded a higher number of total and

duplicated reads in the eight libraries from Altay2000 and Bobur than in the 14 libraries from

the osmotic stress genotypes. The exclusion of less amount of reads after the quality control

filtering and a greater average mapping efficiency were observed in the ionic stress libraries

compared to those from the osmotic stress experiment (Table 1; see details in Supplementary

Table S1).

The use of the reference annotation file scored 83% of the total number of unique mapped

reads while with the extended gene models 88% of the reads were counted. Therefore, with

the extended annotation an additional amount of ca. 5 million reads were detected in 12019

genes (see Supplementary Table S2) which accounted for 4.5 % of the gene models predicted

in the RefSeq v1.1 genome annotation (IWGSC, 2018). The observation of 10 Mbp from two

alignment files allowed the identification of five genes extended from 186 to 470 bp (Table

2). Two of these genes also show a prolonged 3'-end according to RNA-seq data from

Pingault *et al*. (2015) visualized in the RefSeq v1.0 genome browser (Alaux *et al*., 2018)

(Table 2).

Identification of salt-responsive genes

To compare the level of expression of genes in response to salt stress in the osmotic and ionic

phases, the GFOLD tool was used to identify salt-responsive genes in the two tolerant and

two susceptible genotypes studied. The overlapping densities from Syn86 and Zentos

indicated an appropriate homogeneity of the sequencing depth from the libraries and an

adequate expression normalization (Supplementary Fig. S1). Differently, a greater mean of

the expression values was observed at 24 days ASE when compared to the mean in the other

time points (Supplementary Fig. S1). This type of distribution of the expression is an indicator

of high levels of PCR duplication of reads and therefore the deduplicated alignment files were

used for the differential expression analysis at the ionic phase. The density plots produced

after deduplication revealed a better homogeneity among samples from the same genotype

(Supplementary Fig. S1). The removal of low expressed transcripts leaded to the reduction of

the number reads for the differential expression analysis by $3.1\% \pm 0$ $3.1\% \pm 0.4$, $3.2\% \pm 0.2$,

$3.2\% \pm 0.6$ and $3.9\% \pm 1.1$ for the genotypes Syn86, Zentos, Bobur and Altay2000,

respectively.

The differential expression analysis showed a reduced variability among genotypes (mean $\pm$

standard deviation) concerning the percentage of identified novel transcripts ($4.5 \pm 0.4\%$) (see

genome coordinates in Supplementary Table S3), LC ($4.6 \pm 0.6\%$) and HC ($90.9 \pm 1.0\%$)

gene models. The D subgenome contained the greatest percentage of salt-responsive genes

(35.8 ± 1.7%) followed by subgenomes B (31.4 ± 1.1%) and A (31.3 ± 1.9%) and unplaced

superscafolds (1.5 ± 0.4%) (see all salt-responsive genes in Supplementary Table S4).

Comparative analysis of the osmotic stress response

To better understand the early plant reaction to salt exposure, the comparative transcription

profiling at osmotic stress phase was performed. This analysis revealed up- and down-

regulated salt-responsive genes across the four time points studied. The greatest amount of

them was observed at 15 min ASE in Zentos and at 30 min in Syn86, whereas the lowest

number of salt-responsive genes was identified at 8 min ASE in both genotypes (Fig. 2B,C).

Thirty eight and fourteen genes were differentially expressed simultaneously across all the

time points in Syn86 and Zentos, respectively (Fig. 2B,C). The distribution of up and down-

regulated genes across the time points of the osmotic stress experiment are presented in the

Fig. 3A. Zentos had the highest number of up- and down-regulated genes at 15 min and 4 h

ASE, respectively. In contrast, Syn86 had the highest number of up-regulated genes at 4 h and

of down-regulated at 30 min. In total, Zentos showed 75% of up-regulated genes while Syn86

had 60%.

The GO enrichment analysis allowed to compare the over-represented gene categories among

the up- and down-regulated genes identified in each genotype and time point. Highlighted in

the heatmaps are the 24 and 18 ontology terms that were exclusively up- and down-regulated,

respectively (Fig. 4A,B). Among the up-regulated categories, the over-representation of

response to wounding genes and tryptophan synthase activity were recognized in the

susceptible genotype, whereas in Zentos the calcium binding category was identified. Defense

response to fungus and bacterium, transcription factor activity and protein kinase coding

genes were over-represented and up-regulated in both genotypes (Fig. 4A). The down-

regulated categories revealed the over-representation of spermine and spermidine biosynthesis

and antioxidant activity genes in Syn86, while the lipid binding category was observed in

both genotypes (Fig. 4B).

Some gene ontologies such as xyloglucan:xyloglucosyl transferase activity and oxidative

stress were over-represented in both genotypes and revealed differential expression profiles.

The calcium binding category also showed differential expression profiles in the genotypes,

but unlike Zentos, this term was not over-represented in Syn86 (p-value > 0.05) even though

more genes showed differential expression (129 vs 50 genes). Additionally, ontology terms

related to photosynthesis were over-represented in both up- and down-regulated genes from

Syn86 (Fig. 4A,B). Because of the relevance of these categories in the osmotic stress response

and the link of oxidative stress, calcium binding and photosynthesis categories with the

photosynthetic response under salt stress, a comparison of the expression profiles in the

contrasting genotypes will be presented in the next section.

Time course of gene expression and photosynthesis rate during the osmotic phase

A comparison of the expression profiles of the salt-responsive calcium binding, oxidative

stress response and xyloglucan:xyloglucosyl transferase activity genes was performed to

understand the time course of gene expression differences of these categories in the

contrasting genotypes (Fig. 1). The expression of 101 photosynthesis-related genes is only

shown in the susceptible genotype (Fig. 1B) since the related categories were not over-

represented in the tolerant genotype. The up-regulation of eight electron transport in PSII

(photosystem II) genes was observed at 8 min ASE when the photosynthesis rate starts to

decrease in Syn86 (Fig. 1A). When the photosynthesis rate showed recovery but was still

inhibited (Fig. 1A), 91 transcripts from both photosystems I and II were down-regulated at 30

min with relative expression values ranging from -1.1 to -3.4 (Fig. 1B).

The LOESS curve from the 50 salt-responsive calcium binding genes of the tolerant genotype

revealed a gene up-regulation tendency at 15 min. Thirty-four transcripts were identified in

this time point with relative expression values ranging from 1.0 to 3.4 (Fig. 1C). From these

genes, 32 contained an EF hand calcium binding domain. The susceptible genotype presented

heterogeneous expression patterns for 129 salt-responsive genes of this category. In this

genotype, the greatest amount of calcium binding genes (40) were down-regulated at 30 min

with GFOLD values ranging from -1.0 to -3.1 (Fig. 1D). The majority from these genes (26)

were components of the oxygen-evolving complex from the PSII (Wang *et al.*, 2019). This

result was also in line with the suppressed photosynthesis rate of Syn86 at this time point

(Fig. 1A). Furthermore, 35 and 21 additional genes from this category were up-regulated at 30

and 15 min, respectively.

On the other hand, 33 salt-responsive genes from the oxidative stress response category were

identified in Zentos. Eight and 10 of them were up-regulated and showed relative expression

values lower than 2.5 at 15 and 30 min ASE, respectively. The down-regulation of eight genes

was observed at 4 h with expression values ranging from -1.0 to -2.4 (Fig. 1E). In contrast,

Syn86 contained 59 genes displaying heterogeneous expression patterns with greater relative

expression values than Zentos (Fig. 1F). Thus, 11, 17 and 22 genes were down-regulated at 8,

30 min and 4 h, respectively, while 21 transcripts at 30 min and 11 genes at 4 h were up-

regulated. The expression values from the down-regulated transcripts fluctuated from -1.0 to -

3.5 and the up-regulated genes revealed a GFOLD value range from 1.0 to 4.2. The greatest

number of salt-responsive oxidative stress genes observed at 30 min (38 transcripts), which

included both up- and down-regulated transcripts, agrees as well with the inhibited

photosynthetic activity observed in Syn86 in this time point (Fig. 1A).

Finally, all the salt-responsive cell wall genes corresponded to the xyloglucan:xyloglucosyl

transferase activity category. Eighteen genes were identified in Zentos from which 14 showed

up-regulation both at 8 and 15 min with GFOLD values ranging from 1.1 to 4.0 (Fig. 1G).

Otherwise, 24 genes from Syn86 were observed in this category where the LOESS curve

highlighted the down-regulation of 16 transcripts at 30 min (Fig. 1H). The relative expression

values of the down-regulated transcripts ranged from -1.0 to -4.3.

Comparative analysis of the ionic stress response

To better understand the later phase of plant reaction to salt exposure, the comparative

transcription profiling at ionic stress phase was performed with the expression values

calculated after deduplication. This analysis revealed the fewest amount of salt-responsive

genes in Altay at 11 days (Fig. 2D). The simultaneous differential expression of nine genes

was identified across genotypes and time points (Fig. 2D). The Fig. 3B shows the distribution

of up and down-regulated genes. At 24 days ASE Bobur showed a greater proportion of up-

regulated than of down-regulated ones, whereas the opposite pattern with greater amount of

down-regulated transcripts was found in Bobur at 11 days ASE and in Altay2000 at both time

points (Fig. 3B). In total, Altay2000 and Bobur contained 61% and 53% of down-regulated

genes, respectively.

The Fig. 4C summarizes the GO enrichment analysis at the ionic stress phase which is

separated by the up- and down-regulated genes in the two genotypes and the two time points.

Three GO terms specific for this stress phase were identified and half of the enriched

categories shared the same stress effect in the two genotypes (Fig. 4C). For instance,

transferase activity, chitinase activity and response to oxidative stress were down-regulated in

both genotypes (Fig. 4C). On the other hand, translation and transcription factor activity terms

were down-regulated in the tolerant genotype while metal ion binding was up-regulated. The

up-regulation of the response to auxin category and the down-regulation of transporter

activity were observed in the susceptible genotype (Fig. 4C).

Comparative analysis of osmotic and ionic stress responses

The implementation of a transcriptomic approach allowed the comparison of the salt stress

response during the osmotic and ionic phases in the analyzed genotypes. Syn86 was the

genotype presenting the highest amount of salt-responsive genes, from three to five times

more genes than the three cultivars. From all the differentially expressed genes, 79 were stress

responsive in the four genotypes while 46 and 229 transcripts were expressed in both tolerant

and both sensitive genotypes, respectively (Fig. 2A).

A total of 17 GO terms were over-represented in both the osmotic and ionic phases of salt

stress (Fig. 4). The translation category appeared down-regulated in the salt-sensitive Syn86

at the osmotic phase and the tolerant Altay2000 at the ionic phase. The serine-type

endopeptidase inhibitor activity term presented opposite relative expression values in the

tolerant genotypes of both salt stress phases. These genes were down-regulated in the tolerant

genotype and up-regulated in the salt-sensitive one during the osmotic phase. On the contrary,

this category showed up-regulation in the salt-tolerant genotype at the ionic stress phase. The

response to oxidative stress category was both up- and down-regulated in the contrasting

genotypes from at osmotic stress phase while it was only down-regulated in both genotypes

studied during the ionic phase.

 Identification of candidate QTGs

To unravel candidate QTGs that might contain alleles controlling salt stress-related traits, salt-

responsive transcripts within the LD blocks harboring markers with significant phenotypic

effect were identified and analyzed. The Table 3 presents the candidate QTGs from two QTL

identified in an association panel (Oyiga *et al.*, 2018) and an AB-QTL mapping population

(Dadshani, 2018) on the chromosome 2A. A 36 Mbp LD block was covered by the QTL

interval detected by the marker RAC875_c38018_278. This marker showed an effect on shoot

fresh weight after salt stress in the association mapping panel. Two differentially expressed

genes were found in this region, one salt-responsive in the sensitive genotype and one in the

tolerant (Table 3). Among them, TraesCS2A02G395000 showed the strongest stress response

since the gene coding an oxoglutarate/iron-dependent dioxygenase was suppressed in the salt-

susceptible genotype with an expression value of -2.4. On the AB-QTL mapping population, a

9 Mbp LD block constituted the QTL interval from the marker BS00041707_51. A marker-

trait association with kernel weight under stress was discovered with this SNP. This region

included two up-regulated genes in Syn86 with similar relative expression values:

TraesCS2A02G327600 which codes for a copper amine oxidase and TraesCS2A02G331100

which codes for an amino acid transporter. The stress effect on the expression was confirmed

in other abiotic stress studies for the four candidate QTGs determined in both intervals (Table

3).

Discussion

Novel regions with transcription in wheat identified with MACE sequencing

The bioinformatic pipeline implemented detected salt-responsive genes in the contrasting

wheat genotypes studied. This pipeline has also been applied in recent transcriptomics studies

with RNA pools from plants and the real-time PCR validations analyses have confirmed its

high accuracy (Qiao *et al.*, 2017; Vidya *et al.*, 2018). Besides uncovering the dynamic

transcriptomic response during salt stress, the MACE-derived sequence analysis conferred

evidence of two type of novel regions with transcription in wheat. Firstly, the differential

expression analysis assigned a putative role in the salt stress response to *in silico* predicted

novel transcripts. These novel salt-responsive transcripts might enrich the wheat variable

pangenome that represents 39% of the pangenome according to the analysis of the whole

genome of 18 cultivars (Montenegro *et al.*, 2017). Secondly, the detection of unpredicted 3'-

ends of genes gave indication of longer gene transcription contributing to the improvement of

the current gene models (IWGSC, 2018). These 3'-end-extended transcripts might reveal that

some current gene model predictions were based on transcripts with incomplete read coverage

in the region possibly due to low expression levels in previous transcriptomic experiments

(Roberts *et al.*, 2011). The genes identified with extended transcription can be included in

computational prediction approaches to better define gene structures (Inatsuki *et al.*, 2016;

Tzfadia *et al.*, 2018). The real-time PCR validation of both novel transcripts and 3'-ends is

necessary to confirm the transcription of these regions.

Comparative transcriptomic and time course of photosynthesis rate during the osmotic phase

of stress

The adequate expression normalization of the libraries from Syn86 and Zentos allowed to

compare the gene expression values of the same genotype at the different conditions (Klaus

and Huber, 2016; available at https://www.huber.embl.de/users/klaus/Teaching/

DESeq2Predoc2014.html). The osmotic stress experiment revealed the early up-regulation

and the posterior down-regulation of photosynthesis-related transcripts in the susceptible

genotype. The up-regulation at 8 min of the electron transport in PSII category can be linked

to the over-excitation of this system which leads to an increase in the generation of ROS

(Parihar *et al.*, 2015; Foyer, 2018). The down-regulation of photosynthesis-related genes at 30

min ASE might be a consequence of excessive ROS accumulation that inhibits the repair of

photodamaged PSII at both transcriptional and translational levels (Allakhverdiev *et al.*, 2002;

Murata *et al.*, 2007; Saibo *et al.*, 2009; Queval and Foyer, 2012). Nevertheless, the

transcriptional suppression of photosynthesis is not observed at 4 h of stress (Fig. 1A), which

indicates that there are mechanisms that allow the plant to recover the normal expression level

of photosynthesis-related genes.

The reduced oxidative stress response of Zentos can be attributed to a restrained ROS

production in the tolerant genotype which might have a stimulating effect on the growth under

stressful conditions (Queval and Foyer, 2012). The reduced photosynthesis inhibition of this

genotype can be therefore linked to a lower oxidative damage of the photosynthetic apparatus.

Differently, the susceptible genotype revealed both the down- and up-regulation of genes

implicated in oxidative damage protection with greater relative expression values than the

tolerant genotype (Fig. 1E,F). These results indicate that salt stress exerted a stronger effect

on the oxidative damage protection system of Syn86 at the transcriptional level which

supports its greater photosynthesis inhibition.

The over-representation of genes coding for calcium binding proteins at 15 min in Zentos

agrees with an earlier timing of calcium and ROS signaling proposed for salt-tolerant

genotypes (Ismail *et al.*, 2014). These molecules interact to regulate salt stress response

(Ismail *et al.*, 2014; Parihar *et al.*, 2015; Köster *et al.*, 2018). A delayed Ca^{2+}/ROS signaling

will lead to the activation of jasmonic acid signaling pathway that will culminate in cell death.

Differently, an earlier activation of calcium- and ROS-dependent signaling induces the

constraint of jasmonic acid signaling through the activation of the abscisic acid signaling

pathway (Ismail *et al.*, 2014). We can infer that the calcium binding genes that exhibited an

early up-regulation in Zentos may play a role in activating the signaling pathway leading to

abscisic acid accumulation to stimulate growth under stress conditions (Ergen *et al.*, 2009). In

contrast, in the susceptible genotype the calcium binding up-regulation was delayed and

occurred at 30 min ASE which is consistent with the described model (Ismail *et al.*, 2014). In

addition to a delayed calcium binding up-regulation, the salt-driven suppression of calcium

binding genes related to photosynthesis was observed as well. This result is also linked to the

photosynthesis inhibition observed in Syn86.

The present study revealed as well the differential response of the xyloglucan:xyloglucosyl

transferase activity term in the contrasting genotypes. The greater transcription observed in

the tolerant genotype might enable plant growth under stress and might be beneficial for cell

wall strengthening, the prevention of excessive water loss and to maintain turgor pressure

because of the biosynthesis of xyloglucan in the cell wall (Eckardt, 2008; Le Gall *et al.*,

2015). On the other hand, the down-regulation of these genes in the salt-sensitive genotype

might be linked to the inhibition of cell expansion and cell wall synthesis that limit plant

growth under stress conditions (Wang *et al.*, 2018). The synergy of the described

transcriptional events might be crucial for the contrasting salt stress response of Syn86 and

Zentos during the osmotic phase.

Transcriptomic response during the ionic phase

The distribution of the expression values at the ionic phase insinuated a high amount of PCR

duplication in the libraries at 24 days ASE which coincided with the higher amount of reads

sequenced in this time point in both genotypes. The use of deduplicated read alignments was

therefore chosen to reduce false positives in the identification of differentially expressed

genes and to allow a better comparability among libraries. PCR duplication leads to the

overestimation of transcript abundance because the number of reads is biased in the libraries

(Klepikova *et al.*, 2017).

Among the over-represented transcripts identified, the ionic stress effect on the expression of

metal ion binding genes was in accordance with their role in oxidative stress response and

antioxidant activity in the photosynthetic electron transport system because of their redox

potentials (Palma *et al.*, 2013; Yruela, 2013). Therefore, the up-regulation of this mechanism

might counteract the oxidative damage in the tolerant genotype at 24 ASE since at 11 days

oxidative stress response activity was down-regulated. The observed down-regulation of

transporter activity in Bobur suggests two possibilities. First, it might be a strategy of the

susceptible genotype to avoid Na^+ transport and accumulation in the photosynthetic tissues

but also may point out at a potential harmful effect of this mechanism for the uptake of non-

toxic elements relevant for plant growth when the expression of non-selective cation channels

is affected (Assaha *et al.*, 2017). A second possibility might be the down-regulation of

transporters responsible for Na^+ exclusion from leaves that will lead to genotype susceptibility

(Cotsaftis *et al.*, 2012; Wu, 2018).

The similar stress response of some GO terms observed in the contrasting genotypes at the

ionic stress phase supports the finding that some earlier transcriptional responses might

present stronger differences and might cause a greater impact in the contrasting acclimation

response of the genotypes to long-term salt stress (Julkowska and Testerink, 2015).

Nevertheless, it is also possible that when similar pathways are salt-responsive in both

genotypes the difference might lie in the magnitude of the expression to affect the differential

response.

Comparative transcriptomic analysis of osmotic and ionic phases of salt stress

The stress effect on mechanisms related with protein synthesis and breakdown was identified

in the comparative analysis of both stress phases. An accumulation of aberrant proteins in

cells can result from stress-related ROS damage which can lead to the transient suppression of

the *de novo* synthesis of proteins and the intracellular protein degradation by proteases (Kidrič

et al., 2014; Zhu, 2016; Robles and Quesada, 2019). The differential response of the

translation and the serine-type endopeptidase inhibitor categories across the stress phases

suggests that the regulation of these mechanisms are stress stage specific.

Identification of candidate QTGs

Among the two salt-responsive genes observed in the QTL interval from the association

mapping analysis, the oxoglutarate/iron-dependent dioxygenase showed the strongest down-

regulation. This gene superfamily with a wide functional diversification is involved in the

biosynthesis of several specialized secondary metabolites responsive to biotic and abiotic

stresses (Farrow and Facchini, 2014; Xu and Song, 2017). Therefore, this gene is a strong

candidate that can be prioritized for further validation analyses. The AB-QTL mapping

interval contained as well two salt-responsive genes including a copper amine oxidase and an

amino acid transporter with similar magnitudes of relative expression. The up-regulation of

both genes in the sensitive genotype can be linked to the positive phenotypic effect of the

allele from Syn86 in the variation of kernel weight under salt stress (Dadshani, 2018). Studies

in *Arabidopsis* have shown the involvement of copper amine oxidases in the biosynthesis of

nitric oxide which is a signaling molecule that participates in adaptive responses to biotic or

abiotic stresses (Neill *et al.*, 2002; Wimalasekera *et al.*, 2011; Groß *et al.*, 2017). On the other

hand, there are amino acid transporters up-regulated by salt stress that are involved in the

transport of amino acids as proline which accumulates under stress to act as an osmolyte for

osmotic adjustment (Hayat *et al.*, 2012; Wan *et al.*, 2017). The differential expression of the

genes in the interval may contribute concomitantly to the phenotypic variation (González-

Prendes *et al.*, 2017). Only the implementation of a higher resolution mapping approach and

functional studies could help to confirm the causality of the selected candidate genes on the

trait of interest.

The above mentioned results validate the use of the recent genome assembly to facilitate the

integration of genomic and transcriptomic resources to resolve QTL and advance in the

understanding of the genetic variation underlying agronomic complex traits in bread wheat

(Adamski *et al.*, 2018; IWGSC, 2018). This screening approach to target potential functional

candidate genes is relevant since the mapping resolution of the studies is limited and the

average gene density considering solely HC models is of approximately seven genes per Mbp

(IWGSC, 2018). Nevertheless, this strategy can be more robust when expression data of other

tissues under salt stress and different time points can be also included.

Conclusions and future perspectives

To our knowledge this is one of the first studies to combine transcriptomics, bioinformatics,

genetics and stress physiology analyses in a systemic approach to obtain a more

comprehensive understanding of the salt stress adaptation response in bread wheat at both

osmotic and ionic phases. We expect that our results will encourage the wheat research

community to perform functional analysis of some prioritized genes. This will lead to a better

QTL dissection to finally shed light on novel genes controlling regulatory pathways for salt

stress-related traits.

Supplementary data

Table S1. Overview of reads processing and reference genome mapping from all libraries.

Table S2. Genes with extended 3'-end and position in the reference genome of the clusters of

additional reads scored.

Table S3. Reference genome coordinates from the novel salt-responsive transcripts identified

in the four genotypes.

Table S4. GFOLD values of the salt-responsive genes identified in the time points studied in

the four genotypes.

Fig. S1. Density plots with the $\log_{10}$ normalized expression values of the libraries from the

four genotypes studied.

Data availability

Acknowledgements

This study was financed by the "Bundesministerium für wirtschaftliche Zusammenarbeit und

Entwicklung" via "German Agency for International Cooperation", Germany (Project

09.7860.1-001.00). DDD was funded by the PhD scholarship call 679 from Colciencias

(Science, Technology and Innovation Department from Colombia). The authors would like to

acknowledge A. Shresta and S. Parra for the critical reading of the manuscript.

References

Abeel T, Van Parys T, Saeys Y, Galagan J, Van de Peer Y. 2012. GenomeView: a next-generation genome browser. Nucleic Acids Research **40**, e12.

Adamski NM, Borrill P, Brinton J, Harrington S, Marchal C, Uauy C. 2018. A roadmap for gene functional characterisation in wheat. PeerJ Preprints **6**, e26877v1.

Alaux M, Rogers J, Letellier T, *et al.* 2018. Linking the International Wheat Genome Sequencing Consortium bread wheat reference genome sequence to wheat genetic and phenomic data. Genome Biology **19**, 111.

Allakhverdiev SI, Nishiyama Y, Miyairi S, Yamamoto H, Inagaki N, Kanesaki Y, Murata N. 2002. Salt stress inhibits the repair of photodamaged photosystem II by suppressing the transcription and translation of *psbA* genes in *S*ynechocystis. Plant Physiology **130**, 1443–1453.

Altay F. 2012. Yield stability of some Turkish winter wheat (*Triticum aestivum* L.) genotypes in the western transitional zone of Turkey. Turkish Journal of Field Crops **17**, 129–134.

Asmann YW, Klee EW, Thompson EA, *et al.* 2009. 3' tag digital gene expression profiling of human brain and universal reference RNA using Illumina Genome Analyzer. BMC Genomics **10**, 531.

Assaha DVM, Ueda A, Saneoka H, Al-Yahyai R, Yaish MW. 2017. The role of Na^+ and K^+ transporters in salt stress adaptation in glycophytes. Frontiers in Physiology **8**, 509.

Belderok B, Mesdag H, Donner DA. 2000. *Bread-making quality of wheat: A century of breeding in Europe*. Dordrecht: Kluwer Academic Publisher.

Biswas S, Agrawal YN, Mucyn TS, Dangl JL, Jones CD. 2013. Biological averaging in RNA-Seq. arXiv, 1309.0670v2 [q-bio.QM].

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics (Oxford, England) **30**, 2114–2120.

Borrill P, Harrington SA, Uauy C. 2019. Applying the latest advances in genomics and phenomics for trait discovery in polyploid wheat. The Plant Journal **97**, 56–72.

Borrill P, Ramirez-Gonzalez R, Uauy C. 2016. expVIP: a customizable RNA-seq data analysis and visualization platform. Plant Physiology **170**, 2172–2186.

Carillo P, Annunziata MG, Pontecorvo G, Fuggi A, Woodrow P. 2011. Salinity stress and salt tolerance. In: Shanker A, Venkateswarlu B, eds. *Abiotic stress in plants - Mechanisms and adaptations*. IntechOpen, doi: 10.5772/22331.

Chaves MM, Flexas J, Pinheiro C. 2009. Photosynthesis under drought and salt stress: regulation mechanisms from whole plant to cell. Annals of Botany **103**, 551–560.

Cirilli M, Giovannini D, Ciacciulli A, Chiozzotto R, Gattolin S, Rossini L, Liverani A, Bassi D. 2018. Integrative genomics approaches validate *PpYUC11*-like as candidate gene for the stony hard trait in peach (*P. persica* L. Batsch). BMC Plant Biology **18**, 88.

Civelek M, Lusis AJ. 2014. Systems genetics approaches to understand complex traits. Nature reviews. Genetics **15**, 34–48.

Cotsaftis O, Plett D, Shirley N, Tester M, Hrmova M. 2012. A two-staged model of Na$^+$ exclusion in rice explained by 3D modeling of *HKT* transporters and alternative splicing. PLOS ONE **7**, e39865.

Curtis T, Halford NG. 2014. Food security: the challenge of increasing wheat yield and the importance of not compromising food safety. The Annals of Applied Biology **164**, 354–372.

Dadshani S. 2018. Genetic and physiological characterization of traits related to salinity tolerance in an advanced backcross population of wheat. PhD thesis, University of Bonn.

Demetriou G, Neonaki C, Navakoudis E, Kotzabasis K. 2007. Salt stress impact on the molecular structure and function of the photosynthetic apparatus - The protective role of polyamines. Biochimica Et Biophysica Acta **1767**, 272–280.

Eckardt NA. 2008. Role of xyloglucan in primary cell walls. The Plant Cell **20**, 1421–1422.

Ergen NZ, Thimmapuram J, Bohnert HJ, Budak H. 2009. Transcriptome pathways unique to dehydration tolerant relatives of modern wheat. Functional & Integrative Genomics **9**, 377–396.

Ernst J, Bar-Joseph Z. 2006. STEM: a tool for the analysis of short time series gene expression data. BMC Bioinformatics **7**, 191.

Ernst J, Nau GJ, Bar-Joseph Z. 2005. Clustering short time series gene expression data. Bioinformatics (Oxford, England) **21 Suppl 1**, i159-168.

Farrow SC, Facchini PJ. 2014. Functional diversity of 2-oxoglutarate/Fe(II)-dependent dioxygenases in plant metabolism. Frontiers in Plant Science **5**, 524.

Feng J, Meyer CA, Wang Q, Liu JS, Shirley Liu X, Zhang Y. 2012. GFOLD: a generalized fold change for ranking differentially expressed genes from RNA-seq data. Bioinformatics (Oxford, England) **28**, 2782–2788.

Foyer CH. 2018. Reactive oxygen species, oxidative signaling and the regulation of photosynthesis. Environmental and Experimental Botany **154**, 134–142.

González-Prendes R, Quintanilla R, Cánovas A, Manunza A, Figueiredo Cardoso T, Jordana J, Noguera JL, Pena RN, Amills M. 2017. Joint QTL mapping and gene expression analysis identify positional candidate genes influencing pork quality traits. Scientific Reports **7**, 39830.

Groß F, Rudolf E-E, Thiele B, Durner J, Astier J. 2017. Copper amine oxidase 8 regulates arginine-dependent nitric oxide production in *Arabidopsis thaliana*. Journal of Experimental Botany **68**, 2149–2162.

Gupta B, Huang B. 2014. Mechanism of salinity tolerance in plants: Physiological, biochemical, and molecular characterization. International Journal of Genomics **2014**, 701596.

Hawkesford MJ, Araus J-L, Park R, Calderini D, Miralles D, Shen T, Zhang J, Parry MAJ. 2013. Prospects of doubling global wheat yields. Food and Energy Security **2**, 34–48.

Hayat S, Hayat Q, Alyemeni MN, Wani AS, Pichtel J, Ahmad A. 2012. Role of proline under changing environments. Plant Signaling & Behavior **7**, 1456–1466.

Hrdlickova R, Toloue M, Tian B. 2017. RNA-Seq methods for transcriptome analysis. Wiley Interdisciplinary Reviews. RNA **8**, e1364.

Hussain B, Lucas SJ, Ozturk L, Budak H. 2017. Mapping QTLs conferring salt tolerance and micronutrient concentrations at seedling stage in wheat. Scientific Reports **7**, 15662.

Inatsuki T, Sato K, Sakakibara Y. 2016. Prediction of gene structures from RNA-seq data using dual decomposition. IPSJ Transactions on Bioinformatics **9**, 1–6.

Ishikawa A. 2017. A strategy for identifying quantitative trait genes using gene expression analysis and causal analysis. Genes **8**, 347.

Ismail A, Takeda S, Nick P. 2014. Life and death under salt stress: Same players, different timing? Journal of Experimental Botany **65**, 2963–2979.

IWGSC. 2018. Shifting the limits in wheat research and breeding using a fully annotated reference genome. Science **361**, eaar7191.

James RA, Blake C, Zwart AB, Hare RA, Rathjen AJ, Munns R. 2012. Impact of ancestral wheat sodium exclusion genes *Nax1* and *Nax2* on grain yield of durum wheat on saline soils. Functional Plant Biology **39**, 609–618.

Julkowska MM, Testerink C. 2015. Tuning plant signaling and growth to survive salt. Trends in Plant Science **20**, 586–594.

Kang YJ, Lee T, Lee J, Shim S, Jeong H, Satyawan D, Kim MY, Lee S-H. 2016. Translational genomics for plant breeding with the genome sequence explosion. Plant Biotechnology Journal **14**, 1057–1069.

Kendziorski C, Irizarry RA, Chen K-S, Haag JD, Gould MN. 2005. On the utility of pooling biological samples in microarray experiments. Proceedings of the National Academy of Sciences of the United States of America **102**, 4252–4257.

Kidrič M, Kos J, Sabotič J. 2014. Proteases and their endogenous inhibitors in the plant response to abiotic stress. Botanica Serbica **38**, 139–158.

Klepikova AV, Kasianov AS, Chesnokov MS, Lazarevich NL, Penin AA, Logacheva M. 2017. Effect of method of deduplication on estimation of differential gene expression using RNA-seq. PeerJ **5**, e3091.

Köster P, Wallrad L, Edel KH, Faisal M, Alatar AA, Kudla J. 2018. The battle of two ions: $Ca2^+$ signalling against Na^+ stress. Plant Biology (Stuttgart, Germany) **21**, 39–48.

Kunert A, Naz AA, Dedeck O, Pillen K, Léon J. 2007. AB-QTL analysis in winter wheat: I. Synthetic hexaploid wheat (T. turgidum ssp. dicoccoides x T. tauschii) as a source of favourable alleles for milling and baking quality traits. Theoretical and Applied Genetics **115**, 683–695.

Lange W, Jochemsen G. 1992. Use of the gene pools of *Triticum turgidum* ssp. *dicoccoides* and *Aegilops squarrosa* for the breeding of common wheat (*T. aestivum*), through chromosome-doubled hybrids. Euphytica **59**, 213–220.

Le Gall H, Philippe F, Domon J-M, Gillet F, Pelloux J, Rayon C. 2015. Cell wall metabolism in response to abiotic stress. Plants (Basel, Switzerland) **4**, 112–166.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics (Oxford, England) **25**, 2078–2079.

Li Q, Zheng Q, Shen W, Cram D, Fowler DB, Wei Y, Zou J. 2015. Understanding the biochemical basis of temperature-induced lipid pathway adjustments in plants. The Plant Cell **27**, 86–103.

Liao Y, Smyth GK, Shi W. 2013. The Subread aligner: fast, accurate and scalable read mapping by seed-and-vote. Nucleic Acids Research **41**, e108.

Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics (Oxford, England) **30**, 923–930.

Lin Y, Golovnina K, Chen Z-X, Lee HN, Negron YLS, Sultana H, Oliver B, Harbison ST. 2016. Comparison of normalization and differential expression analyses using RNA-Seq data from 726 individual Drosophila melanogaster. BMC Genomics **17**, 28.

Liu A, Xiao Z, Li M-W, *et al. 2019. Transcriptomic reprogramming in soybean seedlings under salt stress. Plant, Cell & Environment **42**, 98–114.

Liu Z, Xin M, Qin J, Peng H, Ni Z, Yao Y, Sun Q. 2015. Temporal transcriptome profiling reveals expression partitioning of homeologous genes contributing to heat and drought acclimation in wheat (*Triticum aestivum* L.). BMC Plant Biology **15**, 152.

Loginova DB, Silkova OG. 2018. The genome of bread wheat *Triticum aestivum* L.: Unique structural and functional properties. Russian Journal of Genetics **54**, 403–414.

Martin M. 2011. CUTADAPT removes adapter sequences from high-throughput sequencing reads. EMBnet.journal **17**, 10–12.

Montenegro JD, Golicz AA, Bayer PE, *et al. 2017. The pangenome of hexaploid bread wheat. The Plant Journal **90**, 1007–1013.

Munns R, Tester M. 2008. Mechanisms of salinity tolerance. Annual Review of Plant Biology **59**, 651–681.

Murata N, Takahashi S, Nishiyama Y, Allakhverdiev SI. 2007. Photoinhibition of photosystem II under environmental stress. Biochimica Et Biophysica Acta **1767**, 414–421.

Neill SJ, Desikan R, Clarke A, Hurst RD, Hancock JT. 2002. Hydrogen peroxide and nitric oxide as signalling molecules in plants. Journal of Experimental Botany **53**, 1237–1247.

Oyiga BC, Sharma RC, Baum M, Ogbonnaya FC, Léon J, Ballvora A. 2018. Allelic variations and differential expressions detected at quantitative trait loci for salt stress tolerance in wheat. Plant, Cell & Environment **41**, 919–935.

Oyiga BC, Sharma RC, Shen J, Baum M, Ogbonnaya FC, Léon J, Ballvora A. 2016. Identification and characterization of salt tolerance of wheat germplasm using a multivariable screening approach. Journal of Agronomy and Crop Science **202**, 472–485.

Palma JM, Gupta DK, Corpas FJ. 2013. Metalloenzymes involved in the metabolism of reactive oxygen species and heavy metal stress. In: Gupta DK, Corpas FJ, Palma JM, eds. *Heavy Metal Stress in Plants*. Berlin, Heidelberg: Springer, 1–17.

Parihar P, Singh S, Singh R, Singh VP, Prasad SM. 2015. Effect of salinity stress on plants and its tolerance strategies: a review. Environmental Science and Pollution Research International **22**, 4056–4075.

Pingault L, Choulet F, Alberti A, Glover N, Wincker P, Feuillet C, Paux E. 2015. Deep transcriptome sequencing provides new insights into the structural and functional organization of the wheat genome. Genome Biology **16**, 29.

Qiao Y, Zhang J, Zhang J, Wang Z, Ran A, Guo H, Wang D, Zhang J. 2017. Integrated RNA-seq and sRNA-seq analysis reveals miRNA effects on secondary metabolism in *Solanum tuberosum* L. Molecular Genetics and Genomics **292**, 37–52.

Queval G, Foyer CH. 2012. Redox regulation of photosynthetic gene expression. Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences **367**, 3475–3485.

Roberts A, Pimentel H, Trapnell C, Pachter L. 2011. Identification of novel transcripts in annotated genomes using RNA-Seq. Bioinformatics (Oxford, England) **27**, 2325–2329.

Robles P, Quesada V. 2019. Transcriptional and post-transcriptional regulation of organellar gene expression (OGE) and its roles in plant salt tolerance. International Journal of Molecular Sciences **20**, 1056.

Roy SJ, Negrão S, Tester M. 2014. Salt resistant crop plants. Current Opinion in Biotechnology **26**, 115–124.

Saibo NJM, Lourenço T, Oliveira MM. 2009. Transcription factors and regulation of photosynthetic and related metabolism under environmental stresses. Annals of Botany **103**, 609–623.

Sha Y, Phan JH, Wang MD. 2015. Effect of low-expression gene filtering on detection of differentially expressed genes in RNA-seq data. Conference Proceedings IEEE Engineering in Medicine and Biology Society **2015**, 6461–6464.

Shi X, Ling H-Q. 2018. Current advances in genome sequencing of common wheat and its ancestral species. The Crop Journal **6**, 15–21.

Silveira JAG, Carvalho FEL. 2016. Proteomics, photosynthesis and salt resistance in crops: An integrative view. Journal of Proteomics **143**, 24–35.

Timmons JA, Szkop KJ, Gallagher IJ. 2015. Multiple sources of bias confound functional enrichment analysis of global-omics data. Genome Biology **16**, 186.

Trapnell C, Pachter L, Salzberg SL. 2009. TopHat: discovering splice junctions with RNA-Seq. Bioinformatics (Oxford, England) **25**, 1105–1111.

Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL, Pachter L. 2012. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. Nature Protocols **7**, 562–578.

Tuteja N. 2007. Mechanisms of high salinity tolerance in plants. Methods in Enzymology **428**, 419–438.

Tzfadia O, Bocobza S, Defoort J, *et al.* 2018. The 'TranSeq' 3'-end sequencing method for high-throughput transcriptomics and gene space refinement in plant genomes. The Plant Journal **96**, 223–232.

Vidya SM, Kumar HSV, Bhatt RM, Laxman RH, Ravishankar KV. 2018. Transcriptional profiling and genes involved in acquired thermotolerance in banana: a non-model crop. Scientific Reports **8**, 10683.

Wan Y, King R, Mitchell RAC, Hassani-Pak K, Hawkesford MJ. 2017. Spatiotemporal expression patterns of wheat amino acid transporters reveal their putative roles in nitrogen transport and responses to abiotic stress. Scientific Reports **7**, 5461.

Wang S, Wong D, Forrest K, *et al.* 2014. Characterization of polyploid wheat genomic diversity using a high-density 90,000 single nucleotide polymorphism array. Plant Biotechnology Journal **12**, 787–796.

Wang Q, Yang S, Wan S, Li X. 2019. The significance of calcium in photosynthesis. International Journal of Molecular Sciences **20**, 1353.

Wang J, Zhu J, Zhang Y, Fan F, Li W, Wang F, Zhong W, Wang C, Yang J. 2018. Comparative transcriptome analysis reveals molecular response to salinity stress of salt-tolerant and sensitive genotypes of indica rice at seedling stage. Scientific Reports **8**, 2085.

Wimalasekera R, Villar C, Begum T, Scherer GFE. 2011. *COPPER AMINE OXIDASE1* (*CuAO1*) of *Arabidopsis thaliana* contributes to abscisic acid- and polyamine-induced nitric oxide biosynthesis and abscisic acid signal transduction. Molecular Plant **4**, 663–678.

Wu H. 2018. Plant salt tolerance and Na^+ sensing and transport. The Crop Journal **6**, 215 – 225.

Xu Y, Li S, Li L, Zhang X, Xu H, An D. 2013. Mapping QTLs for salt tolerance with additive, epistatic and QTL $\times$ treatment interaction effects at seedling stage in wheat. Plant Breeding **132**, 276–283.

Xu Z, Song J. 2017. The 2-oxoglutarate-dependent dioxygenase superfamily participates in tanshinone production in *Salvia miltiorrhiza*. Journal of Experimental Botany **68**, 2299–2308.

Yadav P, Vaidya E, Rani R, Yadav NK, Singh BK, Rai PK, Singh D. 2018. Recent perspective of next generation sequencing: Applications in molecular plant biology and crop improvement. Proceedings of the National Academy of Sciences, India Section B: Biological Sciences **88**, 435–449.

Yruela I. 2013. Transition metals in plant photosynthesis. Metallomics: Integrated Biometal Science **5**, 1090–1109.

Zaman M, Shahid SA, Pharis RP. 2016. Salinity a serious threat to food security – Where do we stand? Soils Newsletter **39**, 9–10.

Zhang Y, Liu Z, Khan AA, *et al.* 2016. Expression partitioning of homeologs and tandem duplications contribute to salt tolerance in wheat (*Triticum aestivum* L.). Scientific Reports **6**, 21476.

Zheng Y, Xu X, Li Z, Yang X, Zhang C, Li F, Jiang G. 2009. Differential responses of grain yield and quality to salinity between contrasting winter wheat cultivars. Seed Science and Biotechnology **3**, 40–43.

Zhu J-K. 2016. Abiotic stress signaling and responses in plants. Cell **167**, 313–324.

Table 1. Summary of MACE libraries processing and reference genome mapping (mean ± standard deviation).

	Osmotic phase	Ionic phase
Libraries	14	8
Total millions of reads	5.0 ± 0.7	9.0 ± 2.4
Reads excluded after QC[a] (%)	7.8 ± 1.6	2.6 ± 0.3
Mapping efficiency (%)	83.6 ± 1.2	93.0 ± 0.3
Millions of mapped reads	3.8 ± 0.5	8.2 ± 2.2
Multiple aligned reads (%)	21.7 ± 3.3	19.7 ± 1.7
Millions of unique mapped reads	3.0 ± 0.4	6.6 ± 1.8
Reads after deduplication (%)	62.2 ± 3.7	53.9 ± 3.2

[a]QC: Quality control

Table 2. Comparison of the reads scored with the reference annotation (R) and with extended gene models (E).

Gene[a]	Reads scored (R/E)		Extension size (bp)
	Library 1[b]	Library 2	
TraesCS5B02G027100	1/9	2/3	242
TraesCS5B02G028200	27/28	23/27	196
TraesCS7D02G049200	1/2	5/16	186
TraesCS7D02G062000LC	5/17	4/9	368
TraesCS7D02G051200	4/107	8/268	470

[a] Genes in bold letters also show a prolonged 3'-end according to RNA-seq data from Pingault et al. (2015)
[b] Library 1 is from Altay2000 at 11 days under control and Library 2 is from Altay2000 at 24 days of stress without deduplication.

Table 3. Differentially expressed genes in LD blocks of markers with effect on salt stress-related traits in the Chr 2A.

Marker[a]	LD interval and size (cM / Mbp)	No. SNPs in interval (No. in LD)	Gene relative expression[d]	Annotation[e]	Abiotic stress effect[f]
RAC875_c38018_278[b]	109.51-112.14= 2.63/ 612-648=36	7 (5)	TraesCS2A02G389400: **1.5**	Leucine zipper, homeobox-associated	1) ↑, 2) ↑
			TraesCS2A02G395000: *-2.4*	Oxoglutarate/iron-dependent dioxygenase	1) ↓
BS00041707_51[c]	105.53-108.46=2.93/ 556-565=9	34(5)	TraesCS2A02G327600: *2.1*	Copper amine oxidase	1) ↑, 2) ↑
			TraesCS2A02G331100: *1.8*	Amino acid transporter, transmembrane domain	1) ↑

[a] Marker names according to Wang et al. (2014)
[b] From mapping study (Oyiga *et al.*, 2018)
[c] From mapping study (Dadshani, 2018)
[d] GFOLD values from tolerant genotypes are bold and from the susceptible are in italics.
[e] Based on the Interpro results from the RefSeqv1.0 annotation (Alaux *et al.*, 2018).
[f] Abiotic stress response based on studies deposited in the wheat expression atlas expVIP (Borrill *et al.*, 2016). 1) Drought and heat (Liu *et al.*, 2015) and 2) cold (Li *et al.*, 2015). The direction of the arrows indicates the stress effect on expression, ↑ when the gene is up-regulated and ↓ when is down-regulated.

Fig. 1. Photosynthesis rate curve (A) and GFOLD values of time course relative expression of some selected gene ontologies from the contrasting genotypes studied during the osmotic stress phase: photosynthesis-related (B), calcium binding (C, D), oxidative stress response (E, F) and xyloglucan:xyloglucosyl transferase activity transcripts (G, H) are shown in Syn86 (B, D, F and H) and Zentos (C, E and G). In each expression profile frame, the gray lines show the time course expression pattern of each gene and the red line is a LOESS (locally estimated scatterplot smoothing) curve that represents the expression tendency of the cluster of genes. The photosynthesis rate curve was adapted from Dadshani (2018), where the shadows represent the standard deviation of the measurements and the time points selected for the transcriptomic analysis are highlighted with blue arrows.

Fig. 2. Venn diagrams of the salt-responsive genes in the contrasting genotypes studied. The total amount of genes in each genotype and/or time point are shown above each diagram. (A) Diagram with the four genotypes. The blue number represents the genes shared by the tolerant genotypes while the red number indicates the genes shared by the salt-susceptible; (B) diagram of the salt-responsive genes in Syn86 by time point; (C) diagram of the salt-responsive genes in Zentos by time point; (D) diagram of the salt-responsive genes in the two sampled days from the ionic stress phase.

Fig. 3. Distribution of up- and down-regulated salt-responsive genes across stress time points. (A) Osmotic phase and (B) ionic phase time points.

Fig. 4. GO terms over-represented during the salt stress response. (A) Up-regulated and (B) down-regulated categories identified in the four stress time points sampled during the osmotic phase; (C) up- and down-regulated categories observed in the two stress time points from the ionic phase. Bold ontologies are categories specific for each heatmap. The $-\log_{10}$ transformation of the corrected p-values highlights the categories with greater significance that are therefore better over-represented.

Fig. 1

Fig. 2

Fig. 3

A

B

Fig. 4

A

B

C