**Working title:**

# Detection of H3N8 influenza A virus with multiple mammalian-

# adaptive mutations in a rescued Grey seal (*Halichoerus grypus*) pup

Divya Venkatesh[1], Carlo Bianco[2, 3], Alejandro Núñez[2], Rachael Collins[4], Darryl Thorpe[5], Scott M.

Reid[6], Steve Essen[6, 7], Natalie McGinn[6, 7],  James Seekings[6, 7],  Jayne Cooper[6], Ian H. Brown[6, 7],

Nicola S. Lewis[1, 7]

1.  Department of Pathobiology and Population Sciences, Royal Veterinary College, Hatfield,

Hertfordshire, United Kingdom

2.  Pathology Department, Animal and Plant Health Agency (APHA-Weybridge), Woodham

Lane, New Haw, Addlestone, KT15 3NB, United Kingdom

3.  Diagnostic & Consultant Avian Pathology,  Pathology Department, Animal and Plant

Health Agency (APHA- Lasswade) Pentlands Science Park, Bush Loan, Penicuik,

Midlothian, EH26 0PZ, United Kingdom.

4.  Animal and Plant Health Agency, Starcross Veterinary Investigation Centre, Staplake

Mount, Starcross, Devon, EX6 8PE, United Kingdom

5.  British Divers Marine Life Rescue, Lime House, Regency Close, Uckfield, East Sussex,

TN22 1DS, United Kingdom

6.  Virology Department, Animal and Plant Health Agency (APHA-Weybridge), Woodham

Lane, New Haw, Addlestone, KT15 3NB, United Kingdom

7.  OIE/FAO/EURL International Reference Laboratory for avian influenza, swine influenza

and Newcastle Disease, Animal and Plant Health Agency (APHA) - Weybridge,

Addlestone, Surrey, United Kingdom

<u>ABSTRACT</u>

Avian Influenza A Viruses (IAV) in different species of seals display a spectrum of pathogenicity, from

subclinical infection to mass mortality events. Here we present an investigation of avian IAV infection

in a 3-4 month old Grey seal (*Halichoerus grypus*) pup, rescued from St Michael's Mount, Cornwall in

2017. The pup underwent medical treatment, but died after two weeks; post-mortem examination

and histology indicated sepsis as the cause of death. IAV NP antigen was detected by

immunohistochemistry in the nasal mucosa, and sensitive real-time reverse transcription polymerase

chain reaction assays detected trace amounts of viral RNA within the lower respiratory tract,

suggesting that the infection may have been cleared naturally. IAV prevalence among Grey seals

may therefore be underestimated. Moreover, contact with humans during the rescue raised concerns

about potential zoonotic risk. Nucleotide sequencing revealed the virus to be of subtype H3N8.

Combining a GISAID database BLAST search and time-scaled phylogenetic analyses, we inferred

that the seal virus originated from an unsampled, locally circulating (in Northern Europe) viruses,

likely from wild Anseriformes. From examining the protein alignments, we found several residue

changes in the seal virus that did not occur in the bird viruses, including D701N in the PB2 segment,

a rare mutation, and a hallmark of mammalian adaptation of bird viruses. IAVs of H3N8 subtype have

been noted for their particular ability to cross the species barrier and cause productive infections,

including historical records suggesting that they may have caused the 1889 pandemic. Therefore,

infections such as the one we report here may be of interest to pandemic surveillance and risk, and

may help us better understand the determinants and drivers of mammalian adaptation in influenza.

48
49

51                                                                                                   (Word count: 277)

## INTRODUCTION

Influenza A viruses (IAVs) are important pathogens for humans and livestock including pigs and poultry. They are segmented RNA viruses, whose genomes consist of eight segments of RNA, which code for ~11 proteins/polypeptides. IAVs are classified into several subtypes based on the antigenic properties of two surface glycoproteins haemagglutinin (HA, avian subtypes H1-H16) and neuraminidase (NA, avian subtypes N1-N9). Viruses of most subtypes can be found in wild waterfowl and shorebirds which are their natural reservoir (Alexander, 2007; Easterday et al., 1968) and can infect both domestic birds and mammalian species in spill-over infections. A few IAV lineages have become established in mammals such as humans, pigs, horses, and dogs though only of subtypes H1, H2 and H3 in combination with N1, N2 or N8 (Reperant et al., 2009; Webster et al., 1992).

IAV in birds replicates mainly in the intestine and is transmitted through the faecal-oral route, although respiratory tropism and oropharyngeal shedding has been noted (Webster et al., 1978; Daoust et al., 2011; Höfle et al., 2012; França et al., 2012; Daoust et al., 2013), and the virus can survive in the environment for fairly long periods (Brown et al., 2009; Stallknecht et al., 1990). This creates conditions conducive to viral exchange with marine mammals such as pinnipeds (seals), whose habitats and prey intersect with those of waterfowl and shorebirds. Several cases of marine mammal infection with IAV of many subtypes including H1N1, H3N3, H3N8, H4N5, H4N6, H7N7 and H10N7 have been documented with a spectrum of effects ranging from mass die-offs to asymptomatic; in a majority of these cases, the source is implicated to be avian (Fereidouni et al., 2016; White, 2013). A few studies have also suggested that Grey seals may act as an endemically infected reservoir which may disseminate viruses in coastal ecosystems to other mammals, coastal birds, and potentially humans (Duignan et al., 1995, 1997; Puryear et al., 2016). Indeed, seroprevalence levels in live-captured healthy Grey seal populations (20 – 26%) (Bodewes et al., 2015; Puryear et al., 2016) is comparable with levels found in wild birds (31% - 60%) depending on species, geography, seasonality and other factors (Curran et al., 2015; Fereidouni et al., 2010; Wilson et al., 2013).

In seals, IAV binds the same type of sialyloligosaccharide receptors as birds (SAα2,3Gal), but the

receptors are located in the respiratory tract and lungs instead of the intestinal tract, suggesting that

inhalation is the most likely route of transmission (Ito et al., 1999; Ramis et al., 2012; White, 2013).

Seals may also be infected by IAV from non-avian sources; there is serological evidence for infection

of Baikal and ringed seals in Russia with human H3N2 strains A/Aichi/2/68 and A/Bangkok/1/79

(Ohishi et al., 2004) and a human pandemic-2009 H1N1 virus was isolated from Elephant seals in

2010 on the coast of California, USA.  Furthermore, it has been shown that IAV from seals can

replicate in human tissue and that seal IAV can be systemically virulent in primates (Murphy et al.,

1983; Webster et al., 1981; White, 2013). Seals may therefore be potential sources of pandemic

influenza.

In this paper, we report an H3N8 IAV infection of a rescued Grey seal pup in coastal England. We

provide molecular and histological evidence for presence of IAV in the seal's respiratory tissues, and

conclude from the post-mortem that the clinical presentation was not caused by IAV. To our

knowledge, we provide the first whole genome sequence for an IAV isolated from a Grey seal, as

attempts so far have been unsuccessful. We use phylogenetic analyses to find the putative sources

of this virus and look for adaptive changes in its sequence. We compare this case with previously

described seal infections to identify unique or parallel elements which may have implications for

animal and human health.

## METHODS

*Clinical history and medical interventions*

In February 2017, a female Grey seal (*Halichoerus grypus*) pup was rescued from St Michael's

Mount, Cornwall. Physical examination of the animal on admittance to the rehabilitation centre

revealed an emaciated and dehydrated subject (body weight = 20 kg). The estimated age was 3 to 4

months. The animal exhibited a mucoid nasal discharge and a 1 cm wound on the ventral thorax. Its

temperature was 39.9 °C and breathing rate was abnormal (continuous breathing pattern, 12 breaths

per minute). The pup was monitored and provided nutritional support, fluid therapy and antibiotic

treatment, but died suddenly 14 days after admittance. The carcass and a nasal swab were

submitted to the Animal and Plant Health Agency (APHA) for virological investigation.

*Pathology, histopathology and immunohistochemistry*

The carcass underwent a full post-mortem examination. A set of tissues was sampled and fixed in

buffered formalin (nasal turbinates, trachea, lung, kidney, soft tissues adjacent to the

cutaneous/subcutaneous lesion, and lymph node). A standard histopathological examination was

carried out on the tissues (Hematoxylin & eosin), followed by an immunohistochemical investigation

targeting nucleoprotein to detect intra-lesional IAV along the respiratory tract (Brookes et al., 2010).

*Real-time reverse transcription polymerase chain reaction (RRT-PCR)*

RNA was extracted from the nasal swab and tissue suspensions using the QIAmp viral RNA

BioRobot kit customised for APHA in conjunction with a Universal BioRobot (Qiagen, Manchester,

UK) (Slomka et al., 2009). RRT-PCR testing of the RNA extracts comprised (i) the Matrix (M)-gene

assay for generic IAV detection using the primers and probes of (Nagy et al., 2010) and (ii) H5 and H7

IAV RRT-PCR assays to test for notifiable avian influenza (Slomka et al., 2007, 2009). For each RRT-

PCR assay, samples producing a threshold cycle (CT) value <36.0 were considered positive (Slomka

et al., 2010). The RNA was also tested by an IAV N1-specific RRT-PCR according to the procedure

described by (Payungporn et al., 2006), an IAV N5-specific RRT-PCR (James et al. 2018) and two

IAV subtype N8-specific RRT-PCRs (James et al., 2018) with the same positive/negative acceptance

criteria. All amplifications were carried out in an MX3000P qPCR System (Agilent).

*Virus isolation and whole genome sequencing*

Attempted virus isolation in 9- to 11-day old SPF embryonated fowls' eggs was performed on the

nasal swab sample according to the internationally-recognised European Union (EU) and OIE

methods (EU, 2006; OIE, 2015), but was unsuccessful. RNA was sequenced using the MiSeq

platform. Briefly, viral RNA was extracted from the egg-isolate of the virus from the nasal swab using

the QIAmp viral RNA mini-kit without the addition of carrier RNA (Qiagen, Manchester, UK). cDNA

was synthesized from RNA using a random hexamer primer mix and cDNA Synthesis System

(Roche, UK). The Sequence library was prepared using a NexteraXT kit (Illumina, Cambridge, UK).

Quality control and quantification of the cDNA and Sequence Library was performed using

Quantifluor dsDNA System (Promega, UK). Sequence libraries were run on a Miseq using MiSeq V2

300 cycle kit (Illumina, Cambridge, UK) with 2x150 base paired-end reads. The raw sequence reads

were analysed using publicly available bioinformatics software, following an in house pipeline,

available on github ([https://github.com/ellisrichardj/FluSeqID/blob/master/FluSeqID.sh](https://github.com/ellisrichardj/FluSeqID/blob/master/FluSeqID.sh)). This pipeline

de-novo assembles the raw data using the Velvet assembler (Zerbino and Birney, 2008), blasts the

resulting contigs against a local database of influenza genes using Blast+ (Camacho et al., 2009),

then maps the raw data against the highest scoring blast hit using the Burrows-Wheeler Aligner (Li,

2013).  The consensus sequence was extracted from the resultant bam file using a modified

SAMtools software package (Li et al., 2009), script (vcf2consensus.pl) available at:

[https://github.com/ellisrichardj/csu_scripts/blob/master/vcf2consensus.pl](https://github.com/ellisrichardj/csu_scripts/blob/master/vcf2consensus.pl)).

*Phylogenetic analysis of the seal virus*

BLAST (Basic Local Alignment Search Tool) was used on GISAID (Elbe and Buckland-Merrett, 2017;

Shu and McCauley, 2017) to find the top 50 closest related viral segments for each segment of the

seal virus. We combined the seal virus sequence (query) along with the BLAST-hits sequences

(blasthit) for each segment for phylogenetic analysis. We removed sequences containing duplicate

strain names and aligned with MAFFT (Katoh and Standley, 2013) using automatic settings.

Alignments for each segment were inspected manually on AliView (Larsson, 2014) and the ends

trimmed to the starting ATG and end STOP codon. Exploratory trees were run using FastTree (Price

et al., 2009), after which we used IQ-TREE (Nguyen et al., 2015) to make the final maximum-

likelihood tree with 1000 iterations of alrt (approximate likelihood ratio test) for branch support.

Tempest v1.5 (Rambaut et al., 2016) was used to test ML trees for clock-like behaviour. Trees for all

segments except MP showed clock-like behaviour (**Figure S1**), so results from the MP dataset were

excluded. BEAST v1.10.1 (Bayesian Evolutionary Analysis Sampling Trees) (Suchard et al., 2018) was

used to determine the putative time and source of emergence of the different segments of the seal

virus. BEAST performs Bayesian analysis of molecular sequences using MCMC (Markov Chain

Monte Carlo) methods. For all segments other than MP and NS which have multiple reading frames,

we used the SRD06 site model which partitions the codons into 1+2 and 3. For NS we used a GTR

model with no codon partitioning. The rest of the priors were kept identical for all segments: an

uncorrelated relaxed lognormal clock, GMRF (Gaussian Markov random field) Bayesian skyride tree

(Minin et al., 2008), 7 million MCMC generations with sampling every 7000. Two separate runs were

performed for each segment, which were combined after logs were inspected in Tracer

v1.7.1(Rambaut et al., 2018) for appropriate mixing and ESS (effective sample size) values > 200.

Trees were summarised into median clade credibility trees (MCC) and plotted in R v3.5 (using the

ggtree package (Yu et al., 2017).

*Amino acid substitutions*

Trimmed alignments of each segment were manually inspected in AliView software (Larsson, 2014),

translated into amino-acids, and checked for amino acid changes across each dataset. We identified

several substitutions in the seal virus that did not occur in any of the bird virus sequences. We

recorded these substitutions, and used the H3-numbering of the sequence using the HA subtype

numbering conversion tool available from FluDB ((Burke and Smith, 2014),

https://tinyurl.com/HAnumbering) for the HA protein. We also looked for differences in glycosylation

patterns between the seal and the related wild bird HA and NA glycoproteins using a program to

detect Asn-X-Ser or Asn-X-Thr (where X is any amino acid other than Proline) patterns in the amino

acid sequences (personal communication from Todd Davis, CDC, USA)

RESULTS

*Pathology and immunohistochemistry*

The post-mortem revealed abundant purulent fluid (approximately 1 litre) in the subcutaneous

tissues of the right side of the body extending from the neck to the thoracic region (**Figure 1A**). All

other organs were macroscopically unremarkable. The blubber sternal thickness was 0.4 cm.

Histology identified a severe, chronic-active, fibrino-purulent cellulitis and fasciitis and a severe,

acute renal infarct with thrombosis. In the respiratory system a diffuse mild to moderate rhinitis with

epithelial hyperplasia and presence of mites was observed in the nasal cavity and a focal broncho-

interstitial pneumonia with thrombosis and pulmonary nematodes were seen in the lungs, with no

changes observed in trachea and bronchi. Influenza A virus antigen was detected by

immunohistochemistry only in the nasal mucosa, in the nuclei of scattered isolated epithelial cells

(**Figure 1B**). The pathological findings were suggestive of a thromboembolic event and sepsis

caused by the cellulitis as the cause of the death of the animal.

*Detection and subtyping of IAV by RRT-PCR*

The nasal swab was positive for the M-gene (CT value of 22.36), signifying the presence of IAV RNA

but was negative for NAI haemagglutinin (HA) subtypes H5 and H7 as well as the IAV subtypes N1

and N5. However, IAV subtype N8 was detected by one of the specific RRT-PCR assays employed

(CT value of 24.38). In addition, trace amounts of IAV RNA were detected by the M-gene RRT-PCR

assay in tracheal bronchi (CT value of 35.88), the left lung (CT value of 37.73) and the right cranial

lung (CT value of 38.67). Due to low levels of virus detected in the lower respiratory tract, and the

presence of clinical signs not consistent with influenza infection, we concluded that the virus was of

no or low pathogenicity. It is possible that at the point the seal was rescued, it had already cleared

the infection in the lungs naturally, which would explain trace amounts of detection in these tissues.

This is consistent with previous studies which found that Grey seals were likely to remain

asymptomatic with IAV infection while a proportion of the surveyed adults and juveniles were

seropositive (Bodewes et al., 2015; Puryear et al., 2016).

Sequencing results from the nasal swab egg-isolate indicated no mixed infection, just a single virus,

from which it was possible to sequence all eight gene segments. The virus-derived sequence was

named A/Grey seal/England/027661/2017 and sequences were deposited in GISAID database

(Global Initiative on Sharing All Influenza Data) with isolate ID: EPI_ISL_381748.

*Source of viral segments*

Segment gene sequences which were found to be most closely related to the genes of the seal virus

by BLAST were of avian origin. Overall, the strains in the set of BLAST-hits for each segment gene,

all came from different avian viruses, isolated from different bird-types, and of different subtypes.

They were from strains that were mostly isolated from wild birds such as mallard, Eurasian teal,

White-fronted goose, Black-headed gull (*Anas platyrhynchos*, *Anas crecca*, *Anser albifrons*,

*Chroicocephalus ridibundus*) and others, along with a few from domestic birds such as chickens

(*Gallus gallus domesticus*). This indicates a wild bird origin for the seal virus, which is consistent with

previously detected seal infections (Fereidouni et al., 2016; White, 2013), and the hypothesis that

overlapping habitats for wild birds and seals make transmission from wild birds possible, but less

likely from poultry, pigs or humans.

Also similar to previously reported infections (Anthony et al., 2012; Zohari et al., 2014), the set of

closest related strains were isolated from within the local region, in this case, Northern Europe (The

Netherlands, Germany, UK etc.) with the exception of three strains from China being among the

close relatives of the PB1 sequence. The years of detection of the closest-related avian viruses for

each segment ranged from the 2007 (NA) to 2017 (NS).

**Table S1** shows a summary for how often the same strain appears as a BLAST-hit for each segment

of the seal virus. A large proportion of these occur singly (250 strains) or for a maximum of two

segments (40 strains). However, some strains were found to map for 3, 4 or 5 segments (19, 2 and 1

strains(s) respectively). Of these, all the strains that mapped for at least 3 segments were isolated

from birds in the Netherlands between 2011-15, except for one chicken virus from France in 2016.

The subtypes varied greatly, but they were usually isolated from wild birds. In the next section, we

further examine the closest available sequence(s) for each seal gene segment to try and understand

the emergence and further propagation of this virus.

*Emergence of viral segments*

There is no dedicated surveillance program for seals in England so infection status in seals as

endemic or one-off spill over event is uncertain. Previous studies in the US have indicated that Grey

seals are possibly a reservoir for IAV and other viruses (Duignan et al., 1995, 1997; Puryear et al.,

2016), We performed a time-scaled analysis with the set of closely related avian virus segment

sequences, to test if all segments from seal virus had a similar point of introduction, i.e., similar time

to most recent common ancestor (TMRCA) of the seal and wild bird viruses for all segments, or the

extent to which this varied segment to segment. We used BEAST to reconstruct time-scaled

phylogenies for each of our segment datasets.

Maximum clade credibility trees for each segment are presented in **Figure 2 (A-G)**; a maximum-

likelihood tree is presented for the MP gene segment in **Figure 2H** as it was excluded from BEAST

analysis due to lack of clock-like behaviour in the dataset. Time to the putative ancestor strain of the

segments from the seal virus and its closest related segment is inferred as ranging from 1999 (NA) to

between 2011 and 2015 (all other gene segments), summarised in **Figure 3** and **Table 1**. Gaps in

surveillance, and the availability of just one seal strain will likely affect the inference of TMRCAs, but

the variation between segments is also likely a testament to the high levels of reassortment seen in

wild bird IAVs (Lu et al., 2014). Indeed, the closest associated virus varies in host, subtype and

geography of isolation for each segment, as can be seen in the highlighted clades in **Figure 2 (A-H),**

and summarised in **Table 1**. Consistent with data of BLAST hits (Table S1), many of the closest

related segments are from strains isolated in the Netherlands (HA, MP, NP, NS, PA, PB1, PB2).

Where multiple strains are equally closely related to the seal virus, e.g. for PB1 and PB2, the avian

influenza strains come from the Netherlands and France. The closest strain to the seal NA gene that

has been sampled is from 2009 at the latest, from Norway. For the NS gene, the closest segments

appear to be from Russian strains in chickens as recent as December 2017, which may indicate

onward transmission via unsampled intermediaries; however the posterior probability support value

for the node is low (0.36).

From the above observations, we conclude that most of the Grey seal virus segments are derived

from viruses in wild birds in Northern Europe, possibly via unsampled intermediaries – but not in a

single transmission event. In contrast, the 2011 H3N8 Harbour seal virus appears to have been

derived from a single source. The closest related strain for all internal gene segments as identified by

BLAST and maximum-likelihood trees is the  A/American_black_duck/New_Brunswick/03552/2009

(an H4N6 isolated on 2009-09-11). The closest related strains for the glycoproteins are different but

this likely indicates a bias for sequencing glycoproteins over the internal genes. They are both H3N8

viruses from the same area (North-eastern US) isolated closer in time to the Harbour seal virus:

A/blue-winged_teal/New_Brunswick/00291/2010 isolated 2010-09-14 (for HA) and

A/northern_pintail/Minnesota/AI09-4322/2009 isolated 2009-09-12 (for NA). See **Table S2A** and **B**

for comparison.

*Substitutions for mammalian adaptation*

Previous analyses have indicated that mammalian adaptive mutations can occur in avian viruses

when they are transmitted into seals. A study of H10N7 viruses in Harbour seals in Northern Europe,

which unlike the present case was demonstrated to have transmitted to other seals and caused an

outbreak, showed that mutations were likely to occur early on after transmission to seals and then

plateau (Bodewes et al., 2016). We looked for differences between the seal virus and wild bird

viruses in the segment amino acid alignments of our datasets, to check if they any had putative

adaptive implications.

We found several amino acid substitutions in the seal virus that did not occur in any of the related

bird viruses. These substitutions are summarised in **Table 2**, along with references for those that

have been identified in previous studies of mammalian adaptation. Many of these changes occur in

the polymerase complex genes (Mänz et al., 2013): D701N in the PB2 segment is a rare mutation,

and a hallmark of mammalian adaptation of bird viruses, regardless of genetic background (Liu et al.,

2018; Steel et al., 2009)). Liu and Steel et. al. have elucidated that the basis of this adaptation is that

it allows for better replication in mammalian cells and showed that it has been associated with

increased transmission in ferret experiments. Another mutation in the PB2 gene, at residue 105 in its

NP-binding region (Poole et al., 2004) was also found in studies that used phylogenetic modelling

(Tamuri et al., 2009) and mutual information statistics (Miotto et al., 2008, 2010). Both the mutations

found in the PB1 gene are also of interest. PB1-I517V was found by Tamuri et al. and S678N found

in the seal PB1 gene has been associated with increased polymerase activity and virulence in mice

(Gabriel et al., 2005).

Changes were also found in the HA gene (**Table 2**), but the implications are less clear. There appear

to be no changes in the glycosylation patterns between the HA and NA of the Grey seal virus in

comparison to related wild bird viruses (**Table S3A and B**).

We compared all the substitutions with previously described mutations in seal infections, and found

that apart from D701N, which was also found in the H3N8 seal virus infection in Massachusetts in

2011 (Anthony et al., 2012), there were no convergent amino acid changes. In the H3 HA gene, we

found substitutions in residue 81 (reference H3 numbering - 64), and residue 176 (reference H3

numbering - 160), and these are also changed in the 2011 Massachusetts virus (Anthony et al., 2012)

but to different amino acids. The latter mutation was not implicated in receptor binding for the seal

viruses as was the case with H5N1 and some human H3N2 viruses, because the glycosylation site is

absent in both seal and wild bird viruses (**Table S3A**). The HA of the 2011 Massachusetts virus  had

an F110S mutation, where the 110 residue has been previously found to be a critical component of

the influenza fusion peptide, which may impact replication in mammalian cells (Anthony et al., 2012;

Liu et al., 2011). Our reported seal virus retains F at position 110, but whether the mutation in the

adjoining residue at S111G (reference H3 numbering 95) has any effect on HA fusion properties is

unknown. The presence of Serine at position 66 in the PB1 sequence, which enables production of

PB1-F2 (Conenello et al., 2007) was found in the 2011 H3N8 Harbour seal virus but was not seen in

this 2017 H3N8 Grey seal virus. Changes at positions 226 and 228 in HA (reference H3 numbering)

which can change receptor-binding preferences between avian and mammalian hosts (Connor et al.,

1994; Matrosovich et al., 2000), were not found in either of the H3N8 seal viruses, but the H10

equivalent of H3-Q226L was identified in viruses from the 2014-15 H10N7 outbreak in Europe

(Dittrich et al., 2018).

<u>DISCUSSION</u>

In the seal infection case reported here, the animal was referred to a rescue centre because it was

stranded and the IAV was detected only incidentally. The vast majority of Grey seals admitted to

rehabilitation centres in the UK are pups within the first year of life. Malnutrition is the single most

common reason for pinnipeds to be taken into rehabilitation centres (Barnett et al., 2000; Van Bonn,

2015). Wounds are often recorded and hold clinical significance, as in this case, and have been

considered predisposing factors for fatal non-specific septicaemia (Baily, 2014). The cause of trauma

and wounds may be anthropic (entanglement in fishing nets or gears) or biologic (conspecific

aggressiveness or hierarchical to cannibalistic behaviours). Wounds caused by bites of other seals or

predators were most often seen (Barnett et al., 2000; Van Bonn, 2015). Pulmonary and nasal

parasites are also very well documented clinico-pathological conditions among rescued seal pups,

and likely caused the hyperplastic rhinitis identified in this study seal. No histopathological findings

consistent with IAV derived damage were detected along the respiratory tract. The

immunohistochemistry demonstrated productive viral replication only in nasal mucosa, but not in

lower respiratory segments, although the more sensitive RRT-PCR revealed presence of trace

amounts of viral RNA within the lower respiratory tract, suggesting that the infection may have been

cleared naturally or indicating a passive translocation of antigen from upper respiratory segments.

Efficient clearance in healthy Grey seals may explain why previous attempts at sequencing IAV from

identified during surveillance were unsuccessful (Puryear et al., 2016).

Puryear et al. discuss the possibility that Grey seals may be more prone to infection due to gain in

population density since the marine mammal protection act in the US in 1972, along with more

socially gregarious and aggressive behaviour in comparison to Harbour seals, all of which contribute

to high pathogen transmissibility. Why we see differences between Grey and Harbour seals in their

resistance to diseases caused by viral agents is however unclear. Phylogenetic analyses have been

unable to resolve the relationships between different seal species with sufficient support but Grey

seals are either placed as a sister group to Caspian seals (*Pusa Caspica*) and separate from Harbour

seals, or in a basal position to both *Phoca* and *Pusa* genus (Berta et al., 2018; Fulton and Strobeck,

2006). It might be useful to explore long-term evolutionary history of host-pathogen relationships

along with host physiology and immune response differences between different seal species to

understand differences in viral pathogenicity.

In our dataset of closest related sequences to the seal virus, we find that while there are several

substitutions found in the seal virus which do not occur in the bird viruses; we generally do not find

substitutions in any of the bird virus strains that do not occur in other bird viruses too. This likely

indicates adaptation to the seal environment, a hypothesis supported by the occurrence of the

D701N mutation, a known rare marker of mammalian adaptation (Liu et al., 2018; Steel et al., 2009).

D701N is common in canine and horse H3N8 IAV (see **Table S4**), but does not occur in birds, and

was found associated with highly-pathogenic H5N1 viruses which infected humans (Gabriel et al.,

2005; Jong et al., 2006; Li et al., 2005). We found one reported instance of the D701N mutation in

the other avian H3N8 virus that infected pinnipeds, but not in the H10N7 outbreak, despite sustained

transmission in seals for several months, nor in any other previously sequenced seal PB2. H3N8 IAV

are noted for the ability cross species barriers, so it would be relevant to consider if the subtype of

the virus has any influence on the kind of adaptive mutations that occur in the polymerase genes,

and if so what sort of mechanisms this might involve. We also note that many of the putative

adaptive substitutions occur in the polymerase complex genes, which are increasingly being

recognised for their vital role in mediating viral host range (in addition to receptor compatibility with

glycoproteins).

The occurrence of adaptive substitutions in the viral sequence could mean that it has been

circulating in seals for a certain amount of time during which these substitutions have accumulated.

Alternatively, this event might represent an early rapidly-adapting virus from a spill-over event as

seen in the H10N7 outbreak (Bodewes et al., 2016). Given the subclinical nature of the infection, we

propose the former explanation is more likely to be correct. However, since the case of the seal virus

in this study appears to be a singular infection as far as we know, it is difficult to know if and how

long the virus has been in seals.  Both this study and Bodewes et al. report that despite mammalian

adaptation of the virus, continued contact with bird reservoirs allows for further exchange of viral

segments between the two hosts. It should be noted that the segment for which this study has

tentative evidence of onward transmission (NS) does not appear to have acquired any putative

mammalian-adaptive mutations. The nature of the Grey seal reservoir, if it exists, and its relationship

with the avian reservoir is currently unknown.

The 2011 North American H3N8 and 2014 European H10N7 viruses which caused outbreaks in

Harbour seals were found to have acquired mutations to enable recognition of sialyloligosaccharide

receptors found more abundantly in mammalian tissues (SAα2,6Gal) but which retained the ability to

interact with avian receptors (SAα2,3Gal). A later detailed structural and functional analysis of the

2011 H3N8 seal HA indicated a true avian receptor binding preference (Yang et al., 2015), as did a

mutational analysis of the H10N7 viruses (Dittrich et al., 2018).  Although there are some common

HA residues that are changed in both the seal virus reported here and the 2011 H3N8 virus, we

found limited convergence in the substitutions and residues involved between the different seal

viruses, likely attributable to the different genetic backgrounds from different avian sources and

possibly differences in host environment of reservoir vs susceptible species. In addition, to the best

of our knowledge, ours is the first Grey seal virus sequence that is being made publicly available (via

GISAID). It is therefore uncertain whether the type of mutations occurring in a putative reservoir host

vis-à-vis avian sequence might be different from those occurring in related hosts with pathogenic

outcomes.

Our phylogeny and molecular clock analyses suggest different lineages of source viruses, and time

of introduction of different segments (with the caveat of being inferred for a single sequence). It is not

surprising to find that the closest sampled and sequenced viruses for each segment are of different

subtypes, hosts and times of isolation, given the relatively low surveillance in wild birds, and the

extensive reassortment (Lu et al., 2014) i.e., swapping of individual segments into different progeny

viruses during propagation after mixed infection with two or more viruses. It would be speculative to

propose likely wild bird species donor however. Studies on avian influenza have previously shown

that the transmission between birds is directional, e.g., usually in the direction from anseriformes

such as ducks into galliformes (chicken) or charadriiformes (gulls) (Venkatesh et al., 2018). In the

case of the current infection as well, where we have found unambiguously closest related

sequences, they have tended to have been isolated from anseriformes species. The above findings,

along with the observation that the closest related viruses were isolated largely from the

Netherlands, it is likely that the source of the viral segments come from unsampled locally circulating

(in Northern Europe) avian viruses from an anseriformes host. The closest related wild bird viruses to

the 2011 H3N8 Harbour seal virus were also isolated from anseriformes species (American black

duck). In contrast with the our Grey seal virus however, phylogenetic analysis reveals a highly

overlapping set of wild bird strains that are most closely related with Harbour seal virus, an

observation that suggests a single spill-over event into a susceptible host.

H3N8 viruses currently circulate in horses and dogs but not humans or pigs. However, different H3

viruses have been found in several species including humans, pigs, horses, dogs, cats, seals,

poultry, and wild aquatic birds. They have thus been noted for a particular ability to cross species

barrier and cause productive infections. One study that examined the ability of H3N8 viruses from

canine, equine, avian, and seal origin to productively infect pigs, demonstrated that avian and seal

viruses replicated substantially and caused detectable lesions in inoculated pigs without prior

adaptation (Solórzano et al., 2015). It is possible that the ready occurrence of PB2 701N mutation in

H3N8 viruses contributes to this ability. We do not have any biological evidence for pre-20th century

human influenza, but historical analysis suggests a long association of humans with influenza

(Hirsch, 1883; Taubenberger and Morens, 2010), and has uncovered temporal–geographic

associations between equine and human influenza-like disease activity documented in Europe in the

16th-18th centuries (Morens and Taubenberger, 2010). Such analysis has also implicated an H3N8

virus in the 1889 pandemic in humans (Morens and Taubenberger, 2010), which makes mammalian-

adapted H3N8 viruses of particular interest as IAV pandemic risk candidates.

In this paper we have presented analyses of a case of seal infection with IAV in coastal England, and

compared it with previously reported seal IAV infections. This infection provides a small but unique

window to understand the ecology of avian-origin viruses that may be circulating and maintained in

mammals. Given the mammalian adaptation activity in IAV upon transmission to seals, such

infections may be of interest to pandemic surveillance and risk, and help us better understand the

determinants of mammalian adaptation in influenza and its complex drivers.

451
452

REFERENCES

Alexander, D.J. (2007). An overview of the epidemiology of avian influenza. Vaccine *25*, 5637–5644.

Anthony, S.J., Leger, J.A.S., Pugliares, K., Ip, H.S., Chan, J.M., Carpenter, Z.W., Navarrete-Macias,
I., Sanchez-Leon, M., Saliki, J.T., Pedersen, J., et al. (2012). Emergence of Fatal Avian Influenza in
New England Harbor Seals. MBio *3*, e00166-12.

Baily, J.L. (2014). The pathology and occurrence of pathogens in Scottish grey seals (Halichoerus
grypus). Thesis. University of St Andrews.

Barnett, J.E., Woodley, A.J., Hill, T.J., and Turner, L. (2000). Conditions in grey seal pups
(Halichoerus grypus) presented for rehabilitation. Vet. Rec. *147*, 98–104.

Berta, A., Churchill, M., and Boessenecker, R.W. (2018). The Origin and Evolutionary Biology of
Pinnipeds: Seals, Sea Lions, and Walruses. Annu. Rev. Earth Planet. Sci. *46*, 203–228.

Bodewes, R., García, A.R., Brasseur, S.M., Conteras, G.J.S., Bildt, M.W.G. van de, Koopmans,
M.P.G., Osterhaus, A.D.M.E., and Kuiken, T. (2015). Seroprevalence of Antibodies against Seal
Influenza A(H10N7) Virus in Harbor Seals and Gray Seals from the Netherlands. PLOS ONE *10*,
e0144899.

Bodewes, R., Zohari, S., Krog, J.S., Hall, M.D., Harder, T.C., Bestebroer, T.M., Bildt, M.W.G. van de,
Spronken, M.I., Larsen, L.E., Siebert, U., et al. (2016). Spatiotemporal Analysis of the Genetic
Diversity of Seal Influenza A(H10N7) Virus, Northwestern Europe. J. Virol. *90*, 4269.

Brookes, S.M., Núñez, A., Choudhury, B., Matrosovich, M., Essen, S.C., Clifford, D., Slomka, M.J.,
Kuntz-Simon, G., Garcon, F., Nash, B., et al. (2010). Replication, Pathogenesis and Transmission of
Pandemic (H1N1) 2009 Virus in Non-Immune Pigs. PLOS ONE *5*, e9068.

Brown, J.D., Goekjian, G., Poulson, R., Valeika, S., and Stallknecht, D.E. (2009). Avian influenza virus
in water: Infectivity is dependent on pH, salinity and temperature. Vet. Microbiol. *136*, 20–26.

Burke, D.F., and Smith, D.J. (2014). A Recommended Numbering Scheme for Influenza A HA
Subtypes. PLOS ONE *9*, e112302.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and Madden, T.L.
(2009). BLAST+: architecture and applications. BMC Bioinformatics *10*, 421.

Conenello, G.M., Zamarin, D., Perrone, L.A., Tumpey, T., and Palese, P. (2007). A Single Mutation in
the PB1-F2 of H5N1 (HK/97) and 1918 Influenza A Viruses Contributes to Increased Virulence. PLOS
Pathog. *3*, e141.

Connor, R.J., Kawaoka, Y., Webster, R.G., and Paulson, J.C. (1994). Receptor Specificity in Human,
Avian, and Equine H2 and H3 Influenza Virus Isolates. Virology *205*, 17–23.

Curran, J.M., Ellis, T.M., and Robertson, I.D. (2015). Serological Surveillance of Wild Waterfowl in
Northern Australia for Avian Influenza Virus Shows Variations in Prevalence and a Cyclical Periodicity
of Infection. Avian Dis. *59*, 492–497.

Daoust, P.-Y., Kibenge, F.S.B., Fouchier, R.A.M., van de Bildt, M.W.G., van Riel, D., and Kuiken, T.
(2011). Replication of low pathogenic avian influenza virus in naturally infected mallard ducks (Anas
platyrhynchos) causes no morphologic lesions. J. Wildl. Dis. *47*, 401–409.

Daoust, P.Y., van de Bildt, M., van Riel, D., van Amerongen, G., Bestebroer, T., Vanderstichel, R.,
Fouchier, R.A.M., and Kuiken, T. (2013). Replication of 2 Subtypes of Low-Pathogenicity Avian
Influenza Virus of Duck and Gull Origins in Experimentally Infected Mallard Ducks. Vet. Pathol. *50*,
548–559.

Dittrich, A., Scheibner, D., Salaheldin, A.H., Veits, J., Gischke, M., Mettenleiter, T.C., and Abdelwhab, E.M. (2018). Impact of Mutations in the Hemagglutinin of H10N7 Viruses Isolated from Seals on Virus Replication in Avian and Human Cells. Viruses *10*, 83.

Duignan, P.J., Saliki, J.T., St. Aubin, D.J., Early, G., Sadove, S., House, J.A., Kovacs, K., and Geraci, J.R. (1995). Epizootiology of morbillivirus infection in north american harbor seals (phoca vitulina) and gray seals (halichoerus grypus). J. Wildl. Dis. *31*, 491–501.

Duignan, P.J., Duffy, N., Rima, B.K., and Geraci, J.R. (1997). Comparative antibody response in harbour and grey seals naturally infected by a morbillivirus. Vet. Immunol. Immunopathol. *55*, 341–349.

Easterday, B.C., Trainer, D.O., Tůmová, B., and Pereira, H.G. (1968). Evidence of Infection with Influenza Viruses in Migratory Waterfowl. Nature *219*, 523.

Elbe, S., and Buckland-Merrett, G. (2017). Data, disease and diplomacy: GISAID's innovative contribution to global health. Glob. Chall. *1*, 33–46.

EU (2006). Diagnostic Manual for avian influenza.

Fereidouni, S., Munoz, O., Von Dobschuetz, S., and De Nardi, M. (2016). Influenza Virus Infection of Marine Mammals. EcoHealth *13*, 161–170.

Fereidouni, S.R., Werner, O., Starick, E., Beer, M., Harder, T.C., Aghakhan, M., Modirrousta, H., Amini, H., Moghaddam, M.K., Bozorghmehrifard, M.H., et al. (2010). Avian influenza virus monitoring in wintering waterbirds in Iran, 2003-2007. Virol. J. *7*, 43.

França, M., Stallknecht, D.E., Poulson, R., Brown, J., and Howerth, E.W. (2012). The Pathogenesis of Low Pathogenic Avian Influenza in Mallards. Avian Dis. *56*, 976–980.

Fulton, T.L., and Strobeck, C. (2006). Molecular phylogeny of the Arctoidea (Carnivora): Effect of missing data on supertree and supermatrix analyses of multiple gene data sets. Mol. Phylogenet. Evol. *41*, 165–181.

Gabriel, G., Dauber, B., Wolff, T., Planz, O., Klenk, H.-D., and Stech, J. (2005). The viral polymerase mediates adaptation of an avian influenza virus to a mammalian host. Proc. Natl. Acad. Sci. *102*, 18590–18595.

Hirsch, A. (1883). Handbook of geographical and historical pathology (New Sydenham Society).

Höfle, U., Bildt, M.W.G.V. de, Leijten, L.M., Amerongen, G.V., Verhagen, J.H., Fouchier, R.A.M., Osterhaus, A.D.M.E., and Kuiken, T. (2012). Tissue tropism and pathology of natural influenza virus infection in black-headed gulls (Chroicocephalus ridibundus). Avian Pathol. *41*, 547–553.

Ito, T., Kawaoka, Y., Nomura, A., and Otsuki, K. (1999). Receptor Specificity of Influenza A Viruses from Sea Mammals Correlates with Lung Sialyloligosaccharides in These Animals. J. Vet. Med. Sci. *61*, 955–958.

James, J., Slomka, M.J., Reid, S.M., Thomas, S.S., Mahmood, S., Byrne, A.M.P., Cooper, J., Russell, C., Mollett, B.C., Agyeman-Dua, E., et al. (2018). Proceedings Paper—Avian Diseases 10th AI Symposium Issue Development and Application of Real-Time PCR Assays for Specific Detection of Contemporary Avian Influenza Virus Subtypes N5, N6, N7, N8, and N9. Avian Dis. *63*, 209–218.

Jong, M.D. de, Simmons, C.P., Thanh, T.T., Hien, V.M., Smith, G.J.D., Chau, T.N.B., Hoang, D.M., Chau, N.V.V., Khanh, T.H., Dong, V.C., et al. (2006). Fatal outcome of human influenza A (H5N1) is associated with high viral load and hypercytokinemia. Nat. Med. *12*, 1203.

Katoh, K., and Standley, D.M. (2013). MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. Mol. Biol. Evol. *30*, 772–780.

Larsson, A. (2014). AliView: a fast and lightweight alignment viewer and editor for large datasets.
Bioinformatics *30*, 3276–3278.

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
ArXiv13033997 Q-Bio.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and
Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*, 2078–
2079.

Li, Z., Chen, H., Jiao, P., Deng, G., Tian, G., Li, Y., Hoffmann, E., Webster, R.G., Matsuoka, Y., and
Yu, K. (2005). Molecular basis of replication of duck H5N1 influenza viruses in a mammalian mouse
model. J. Virol. *79*, 12058–12064.

Liu, S., Li, R., Zhang, R., Chan, C.C.S., Xi, B., Zhu, Z., Yang, J., Poon, V.K.M., Zhou, J., Chen, M., et
al. (2011). CL-385319 inhibits H5N1 avian influenza A virus infection by blocking viral entry. Eur. J.
Pharmacol. *660*, 460–467.

Liu, S., Zhu, W., Feng, Z., Gao, R., Guo, J., Li, X., Liu, J., Wang, D., and Shu, Y. (2018). Substitution
of D701N in the PB2 protein could enhance the viral replication and pathogenicity of Eurasian avian-
like H1N1 swine influenza viruses. Emerg. Microbes Infect. *7*.

Lu, L., Lycett, S.J., and Leigh Brown, A.J. (2014). Reassortment patterns of avian influenza virus
internal segments among different subtypes. BMC Evol. Biol. *14*, 16.

Mänz, B., Schwemmle, M., and Brunotte, L. (2013). Adaptation of Avian Influenza A Virus Polymerase
in Mammals To Overcome the Host Species Barrier. J. Virol. *87*, 7200–7209.

Matrosovich, M., Tuzikov, A., Bovin, N., Gambaryan, A., Klimov, A., Castrucci, M.R., Donatelli, I., and
Kawaoka, Y. (2000). Early Alterations of the Receptor-Binding Properties of H1, H2, and H3 Avian
Influenza Virus Hemagglutinins after Their Introduction into Mammals. J. Virol. *74*, 8502–8512.

Minin, V.N., Bloomquist, E.W., and Suchard, M.A. (2008). Smooth Skyride through a Rough Skyline:
Bayesian Coalescent-Based Inference of Population Dynamics. Mol. Biol. Evol. *25*, 1459–1471.

Miotto, O., Heiny, A., Tan, T.W., August, J.T., and Brusic, V. (2008). Identification of human-to-
human transmissibility factors in PB2 proteins of influenza A by large-scale mutual information
analysis. BMC Bioinformatics *9*, S18.

Miotto, O., Heiny, A.T., Albrecht, R., García-Sastre, A., Tan, T.W., August, J.T., and Brusic, V. (2010).
Complete-Proteome Mapping of Human Influenza A Adaptive Mutations: Implications for Human
Transmissibility of Zoonotic Strains. PLOS ONE *5*, e9025.

Morens, D.M., and Taubenberger, J.K. (2010). Historical thoughts on influenza viral ecosystems, or
behold a pale horse, dead dogs, failing fowl, and sick swine. Influenza Other Respir. Viruses *4*, 327–
337.

Murphy, B.R., Harper, J., Sly, D.L., London, W.T., Miller, N.T., and Webster, R.G. (1983). Evaluation
of the A/Seal/Mass/1/80 virus in squirrel monkeys. Infect. Immun. *42*, 424–426.

Nagy, A., Vostinakova, V., Pirchanova, Z., Cernikova, L., Dirbakova, Z., Mojzis, M., Jirincova, H.,
Havlickova, M., Dan, A., Ursu, K., et al. (2010). Development and evaluation of a one-step real-time
RT-PCR assay for universal detection of influenza A viruses from avian and mammal species. Arch.
Virol. *155*, 665–673.

Nguyen, L.-T., Schmidt, H.A., von Haeseler, A., and Minh, B.Q. (2015). IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol. Biol. Evol. *32*, 268–274.

Ohishi, K., Kishida, N., Ninomiya, A., Kida, H., Takada, Y., Miyazaki, N., Boltunov, A.N., and Maruyama, T. (2004). Antibodies to human-related H3 influenza A virus in Baikal seals (Phoca sibirica) and ringed seals (Phoca hispida) in Russia. Microbiol. Immunol. *48*, 905–909.

OIE (2015). Chapter 2.3.4. Avian influenza. In Manual of Diagnostic Tests and Vaccines for Terrestrial Animals., p.

Payungporn, S., Chutinimitkul, S., Chaisingh, A., Damrongwantanapokin, S., Buranathai, C., Amonsin, A., Theamboonlers, A., and Poovorawan, Y. (2006). Single step multiplex real-time RT-PCR for H5N1 influenza A virus detection. J. Virol. Methods *131*, 143–147.

Poole, E., Elton, D., Medcalf, L., and Digard, P. (2004). Functional domains of the influenza A virus PB2 protein: identification of NP- and PB1-binding sites. Virology *321*, 120–133.

Price, M.N., Dehal, P.S., and Arkin, A.P. (2009). FastTree: Computing Large Minimum Evolution Trees with Profiles instead of a Distance Matrix. Mol. Biol. Evol. *26*, 1641–1650.

Puryear, W.B., Keogh, M., Hill, N., Moxley, J., Josephson, E., Davis, K.R., Bandoro, C., Lidgard, D., Bogomolni, A., Levin, M., et al. (2016). Prevalence of influenza A virus in live-captured North Atlantic gray seals: a possible wild reservoir. Emerg. Microbes Infect. *5*, 1–9.

Rambaut, A., Lam, T.T., Max Carvalho, L., and Pybus, O.G. (2016). Exploring the temporal structure of heterochronous sequences using TempEst (formerly Path-O-Gen). Virus Evol. *2*.

Rambaut, A., Drummond, A.J., Xie, D., Baele, G., and Suchard, M.A. (2018). Posterior Summarization in Bayesian Phylogenetics Using Tracer 1.7. Syst. Biol. *67*, 901–904.

Ramis, A.J., van Riel, D., van de Bildt, M.W.G., Osterhaus, A., and Kuiken, T. (2012). Influenza A and B virus attachment to respiratory tract in marine mammals. Emerg. Infect. Dis. *18*, 817–820.

Reperant, L.A., Rimmelzwaan, G.F., and Kuiken, T. (2009). Avian influenza viruses in mammals. Rev. Sci. Tech. Int. Off. Epizoot. *28*, 137–159.

Shu, Y., and McCauley, J. (2017). GISAID: Global initiative on sharing all influenza data – from vision to reality. Eurosurveillance *22*, 30494.

Slomka, M.J., Pavlidis, T., Banks, J., Shell, W., McNally, A., Essen, S., and Brown, I.H. (2007). Validated H5 Eurasian Real-Time Reverse Transcriptase–Polymerase Chain Reaction and Its Application in H5N1 Outbreaks in 2005–2006. Avian Dis. *51*, 373–377.

Slomka, M.J., Pavlidis, T., Coward, V.J., Voermans, J., Koch, G., Hanna, A., Banks, J., and Brown, I.H. (2009). Validated RealTime reverse transcriptase PCR methods for the diagnosis and pathotyping of Eurasian H7 avian influenza viruses. Influenza Other Respir. Viruses *3*, 151–164.

Slomka, M.J., Irvine, R.M., Pavlidis, T., Banks, J., and Brown, I.H. (2010). Role of real-time RT-PCR platform technology in the diagnosis and management of notifiable avian influenza outbreaks: experiences in Great Britain. Avian Dis. *54*, 591–596.

Solórzano, A., Foni, E., Córdoba, L., Baratelli, M., Razzuoli, E., Bilato, D., Burgo, M.Á.M. del, Perlin, D.S., Martínez, J., Martínez-Orellana, P., et al. (2015). Cross-Species Infectivity of H3N8 Influenza Virus in an Experimental Infection in Swine. J. Virol. *89*, 11190–11202.

Stallknecht, D.E., Kearney, M.T., Shane, S.M., and Zwank, P.J. (1990). Effects of pH, Temperature, and Salinity on Persistence of Avian Influenza Viruses in Water. Avian Dis. *34*, 412–418.

Steel, J., Lowen, A.C., Mubareka, S., and Palese, P. (2009). Transmission of Influenza Virus in a
Mammalian Host Is Increased by PB2 Amino Acids 627K or 627E/701N. PLOS Pathog. *5*, e1000252.

Suchard, M.A., Lemey, P., Baele, G., Ayres, D.L., Drummond, A.J., and Rambaut, A. (2018).
Bayesian phylogenetic and phylodynamic data integration using BEAST 1.10. Virus Evol. *4*.

Tamuri, A.U., Reis, M. dos, Hay, A.J., and Goldstein, R.A. (2009). Identifying Changes in Selective
Constraints: Host Shifts in Influenza. PLOS Comput. Biol. *5*, e1000564.

Taubenberger, J.K., and Morens, D.M. (2010). Influenza: The Once and Future Pandemic. Public
Health Rep. *125*, 16–26.

Van Bonn, W.G. (2015). Chapter 44 - Pinnipedia. In Fowler's Zoo and Wild Animal Medicine, Volume
8, R.E. Miller, and M.E. Fowler, eds. (St. Louis: W.B. Saunders), pp. 436–450.

Venkatesh, D., Poen, M.J., Bestebroer, T.M., Scheuer, R.D., Vuong, O., Chkhaidze, M.,
Machablishvili, A., Mamuchadze, J., Ninua, L., Fedorova, N.B., et al. (2018). Avian Influenza Viruses
in Wild Birds: Virus Evolution in a Multihost Ecosystem. J. Virol. *92*, e00433-18.

Webster, R.G., Yakhno, M., Hinshaw, V.S., Bean, W.J., and Copal Murti, K. (1978). Intestinal
influenza: Replication and characterization of influenza viruses in ducks. Virology *84*, 268–278.

Webster, R.G., Geraci, J., Petursson, G., and Skirnisson, K. (1981). Conjunctivitis in human beings
caused by influenza A virus of seals. N. Engl. J. Med. *304*.

Webster, R.G., Bean, W.J., Gorman, O.T., Chambers, T.M., and Kawaoka, Y. (1992). Evolution and
ecology of influenza A viruses. Microbiol. Mol. Biol. Rev. *56*, 152–179.

White, V.C. (2013). A review of influenza viruses in seals and the implications for public health. US
Army Med. Dep. J. 45–50.

Wilson, H.M., Hall, J.S., Flint, P.L., Franson, J.C., Ely, C.R., Schmutz, J.A., and Samuel, M.D. (2013).
High Seroprevalence of Antibodies to Avian Influenza Viruses among Wild Waterfowl in Alaska:
Implications for Surveillance. PLOS ONE *8*, e58308.

Yang, H., Nguyen, H.T., Carney, P.J., Guo, Z., Chang, J.C., Jones, J., Davis, C.T., Villanueva, J.M.,
Gubareva, L.V., and Stevens, J. (2015). Structural and Functional Analysis of Surface Proteins from
an A(H3N8) Influenza Virus Isolated from New England Harbor Seals. J. Virol. *89*, 2801–2812.

Yu, G., Smith, D.K., Zhu, H., Guan, Y., and Lam, T.T.-Y. (2017). ggtree: an r package for visualization
and annotation of phylogenetic trees with their covariates and other associated data. Methods Ecol.
Evol. *8*, 28–36.

Zerbino, D.R., and Birney, E. (2008). Velvet: Algorithms for de novo short read assembly using de
Bruijn graphs. Genome Res. *18*, 821–829.

Zohari, S., Neimanis, A., Härkönen, T., Moraeus, C., and Valarcher, J.F. (2014). Avian influenza
A(H10N7) virus involvement in mass mortality of harbour seals (Phoca vitulina) in Sweden, March
through October 2014. Eurosurveillance *19*, 20967.

**Figure 1 (A-B)**
Grey seal pup. (A) Focally extensive, chronic-active, purulent and fibrino-necrotising cellulitis in subcutaneous tissues and skeletal muscle over the thorax. (B) Immunohistochemical demonstration of Influenza A nucleoprotein antigen in the nuclei of epithelial cells within the nasal mucosa of a grey seal. (DAB chromogen and haematoxylin counterstain) (Objective: 40x).

Figure 1A

Figure 1B

## Figure 2 (A-G, H)

Maximum clade credibility summary trees for BEAST analysis of segment datasets: HA, NA, NP, NS, PA, PB1, PB2 (A-G respectively). Nodes connecting the seal virus tip with the closest related strain(s) are highlighted in grey. Black diamond (◆) shapes at nodes indicate posterior probability > 0.75. Nodes are labelled with node ages as inferred by BEAST. Figure 2H shows the maximum likelihood tree for the MP gene dataset. Support values (approximate likelihood ratio test) are labelled at nodes, and nodes connecting the seal virus tip with the closest related strain is highlighted in grey.

Figure 2A

seal – HA

Figure 2B

seal – NA

blasthit|EPI_ISL_243636|A/dunlin/Netherlands/3/2007|A_/_H3N8|2007−07−31
blasthit|EPI_ISL_243639|A/turnstone/Netherlands/2/2007|A_/_H3N8|2007−07−31
blasthit|EPI_ISL_243402|A/dunlin/Netherlands/2/2007|A_/_H3N8|2007−07−31
blasthit|EPI_ISL_30805|A/turnstone/Netherlands/1/2007|A_/_H3N8|2007−XX−XX
blasthit|EPI_ISL_243651|A/turnstone/Netherlands/4/2008|A_/_H3N8|2008−11−01
blasthit|EPI_ISL_329023|A/ruddy_turnstone/Netherlands/6/2008|A_/_H3N8|2008−11−07
blasthit|EPI_ISL_243638|A/turnstone/Netherlands/3/2008|A_/_H3N8|2008−10−31
blasthit|EPI_ISL_129672|A/duck/Germany−NW/R2185/2006|A_/_H6N8|2006−XX−XX
blasthit|EPI_ISL_73378|A/mallard/Sweden/45/2002|A_/_H11N8|2002−11−13
blasthit|EPI_ISL_189612|A/mallard/Sweden/1636/2002|A_/_H3N8|2002−12−13
blasthit|EPI_ISL_30813|A/mallard/Netherlands/33/2006|A_/_H7N8|2006−XX−XX
blasthit|EPI_ISL_30804|A/common_eider/Netherlands/1/2006|A_/_H3N8|2006−XX−XX
blasthit|EPI_ISL_84560|A/mallard/Netherlands/20/2005|A_/_H12N8|2005−XX−XX
blasthit|EPI_ISL_243633|A/mallard_duck/Netherlands/34/2006|A_/_H3N8|2006−09−11
blasthit|EPI_ISL_243390|A/mallard_duck/Netherlands/40/2006|A_/_H6N8|2006−09−16
blasthit|EPI_ISL_84554|A/common_eider/Netherlands2/2006|A_/_H4N8|2006−XX−XX
blasthit|EPI_ISL_67401|A/Mallard/65112/03|A_/_H3N8|2003−XX−XX
blasthit|EPI_ISL_267359|A/mallard_duck/Netherlands/2/2004|A_/_H6N8|2004−09−27
blasthit|EPI_ISL_243522|A/mallard_duck/Netherlands/13/2012|A_/_H3N8|2012−10−01
blasthit|EPI_ISL_243486|A/turnstone/Netherlands/1/2010|A_/_H3N8|2010−10−09
blasthit|EPI_ISL_243582|A/bewicks_swan/Netherlands/8/2009|A_/_H6N8|2009−01−04
blasthit|EPI_ISL_243640|A/mallard_duck/Netherlands/76/2008|A_/_H6N8|2008−11−27
blasthit|EPI_ISL_243386|A/mallard_duck/Netherlands/16/2007|A_/_H6N8|2007−11−15
blasthit|EPI_ISL_32005|A/mute_swan/Germany/R2927/2007|A_/_H6N8|2007−XX−XX
blasthit|EPI_ISL_243495|A/mallard_duck/Netherlands/78/2006|A_/_H6N8|2006−10−21
blasthit|EPI_ISL_74065|A/Teal/Norway/10_1575/2007|A_/_H3N8|2007−XX−XX
blasthit|EPI_ISL_129976|A/common_gull/Norway/10_1602/2009|A_/_H6N8|2009−XX−XX
query|A/grey_seal/England/027661/2017|H3N8|NA|2017−02
blasthit|EPI_ISL_267251|A/mallard_duck/Netherlands/1/2003|A_/_H3N8|2003−09−25
blasthit|EPI_ISL_267383|A/mallard_duck/Netherlands/2/2003|A_/_H3N8|2003−09−25
blasthit|EPI_ISL_267401|A/mallard_duck/Netherlands/1/2004|A_/_H3N8|2004−09−18
blasthit|EPI_ISL_243462|A/greater_white−fronted_goose/Netherlands/3/2006|A_/_H6N8|2006−01
blasthit|EPI_ISL_97364|A/goose/Germany/R1767/2007|A_/_H6N8|2007−XX−XX
blasthit|EPI_ISL_30808|A/common_teal/Netherlands/2/2005|A_/_H6N8|2005−XX−XX
blasthit|EPI_ISL_30809|A/black−headed_gull/Netherlands/1/2005|A_/_H6N8|2005−XX−XX
blasthit|EPI_ISL_3008|A/duck/Spain/543/2006|A_/_H6N8|2006−10−07
blasthit|EPI_ISL_3007|A/duck/Spain/539/2006|A_/_H6N8|2006−10−07
blasthit|EPI_ISL_243599|A/greater_white−fronted_goose/Netherlands/4/2006|A_/_H6N8|2006−01
blasthit|EPI_ISL_243606|A/gadwall_duck/Netherlands/3/2006|A_/_H3N8|2006−09−03
blasthit|EPI_ISL_267207|A/mallard_duck/Netherlands/60/2006|A_/_H3N8|2006−09−18
blasthit|EPI_ISL_243492|A/mallard_duck/Netherlands/7/2008|A_/_H3N8|2008−08−22
blasthit|EPI_ISL_31484|A/mallard/Hungary/19616/2007|A_/_H3N8|2007−XX−XX
blasthit|EPI_ISL_63529|A/mallard/Czech_Republic/14516/2007|A_/_H3N8|2007−09−17
blasthit|EPI_ISL_267292|A/white−fronted_goose/Netherlands/5/2011|A_/_H6N8|2011−12−14
blasthit|EPI_ISL_2852|A/duck/Norway/1/03|A_/_H3N8|2003−XX−XX
blasthit|EPI_ISL_243546|A/bean_goose/Netherlands/1/2008|A_/_H6N8|2008−01−22
blasthit|EPI_ISL_243540|A/greater_white−fronted_goose/Netherlands/4/2008|A_/_H6N8|2008−01
blasthit|EPI_ISL_329054|A/White−fronted_Goose/Netherlands/3/2008|A_/_H6N8|2008−01−22
blasthit|EPI_ISL_243559|A/greater_white−fronted_goose/Netherlands/2/2008|A_/_H6N8|2008−01
blasthit|EPI_ISL_15251|A/duck/Eastern_China/163/2002|A_/_H6N8|2002−XX−XX
blasthit|EPI_ISL_30793|A/mallard/Netherlands/3/2005|A_/_H3N8|2005−XX−XX

2007.46
2007.29
2006.91
2005.85
2008.28
2007.92
2001.62
1999.22
1998.2
2000.37
2001.62
2001.71
1999.61
1997.43
1999.29
2005.2
1996.97
1996.1
2007.56
1993.59
2008.11
2007.21
2001.63
2005.04
1998.35
2003.38
1999.53
1993.39
2003.59
2002.43
2005.58
2004.31
2003.64
2003.04
2006.04
1991.56
2001.75
2004.16
2005.81
3003.72
1998.82
2002.94
1993.47
1994.63
2007.93
2007.66
1992.94
2006.63
1994.13
1993.51

1991    1996    2001    2006    2011    2016

# Figure 2C

## seal – NP

Figure 2D

seal – NS

# Figure 2E

## seal – PA

# Figure 2F

## seal – PB1

# Figure 2G

## seal – PB2

2010                                    2015

# Figure 2H

seal - MP (ML tree)

0.001

Figure 3
Putative divergence times (TMRCA – time to most recent common ancestor)
of the seal sequence from the closest related wild bird sequence for each
segment. Error bars indicate 95% highest posterior density.

Table 1
Putative divergence times (TMRCA – time to most recent common ancestor) and 95% highest posterior density (HPD) of the seal sequence from the closest related wild bird sequence for each segment, and the posterior probability of the node are shown, along with information about the closest related wild bird sequence including the host, country and year of isolation and subtype of virus. For the MP gene, the closest related sequence according to the ML tree is shown.

| Segment | TMRCA | Lower 95% HPD | Upper 95% HPD | Posterior probability of node > 0.75 | Host | Country | Year | Subtype |
|---|---|---|---|---|---|---|---|---|
| HA | 2012.74 | 2011.18 | 2014.08 | y | barnacle goose | The Netherlands | 2014 | H3N6 |
| MP | n/a | n/a | n/a | n/a | mallard | The Netherlands | 2010 | H3N2 |
| NA | 1999.53 | 1994.89 | 2003.34 | n | teal/gull | Norway | 2007/2009 | H3N8/H6N8 |
| NP | 2014.98 | 2014.29 | 2015.38 | n | mallard | The Netherlands | 2015 | unknown |
| NS | 2009.56 | 2008.04 | 2010.86 | n | barnacle goose | The Netherlands | 2011 | H6N8 |
| PA | 2013.71 | 2013.1 | 2014.35 | n | mallard | The Netherlands | 2014 | H7N5 |
| PB1 | 2014.28 | 2013.43 | 2015.04 | y | duck | The Netherlands/France | 2015 | H5N1/H5N2/H5N6 |
| PB2 | 2011.45 | 2010.99 | 2011.77 | y | goose/chicken/duck | The Netherlands/France | 2011/2015/2016 | H5N2/H6N2/H7N9/H6N8 |

Table 2
Segment-wise list of substitutions found in the seal virus inferred from inspection of alignments. Only residue changes that occur in the seal virus but not in the wild bird viruses are shown. Numbers in square brackets for HA indicate the reference H3-numbering of residues in A/Aichi/2/1968 (Burke and Smith 2014) found using the tool at FluDB website: https://tinyurl.com/HAnumbering

| Segment | Amino acid substitutions in seal virus [H3 numbering] |
|---|---|
| HA | Y9C [signal peptide]<br>T64A [48]<br>I81V [65]<br>S111G [95]<br>possible A154T (but might be same RCT --> ACT (T), or GCT (A)) [138]<br>A/T176S [160]<br>S235Y [219] |
| MP | none |
| NA | N/S41D<br>T383A |
| NP | V104M |
| NS | none |
| PA | V91I<br>D/G101N<br>N/S321I |
| PB1 | I517V<br>S678N |
| PB2 | T/A105K<br>D161N<br>A/S395T<br>V667I<br>R/T676I<br>D701N<br>possible R753T (but might be same ASA --> AGA (R) or ACA (T) |