Title: Housing laboratory mice deficient for *Nod2* and *Atg16l1* in a natural environment

uncovers genetic and environmental contributions to immune variation.

Authors: Jian-Da Lin[1,*], Joseph C. Devlin[1,2,*], Frank Yeung[2,3,*], Caroline McCauley[1],

Jacqueline M. Leung[4], Ying-Han Chen[3], Alex Cronkite[1,-3], Christina Hansen[4], Charlotte

Drake-Dunn[3], Kelly V. Ruggles[5,6], Ken Cadwell[1,3,7,§,†], Andrea L. Graham[4,§ ,†] and P'ng

Loke[1,§,†]

Affiliations:

[1]Department of Microbiology, New York University School of Medicine,

[2]Sackler Institute of Graduate Biomedical Sciences,

[3]Kimmel Center for Biology and Medicine at the Skirball Institute, New York University

School of Medicine, New York, NY 10016, USA.

[4]Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ

08544, USA.

[5]Division of Translational Medicine, Department of Medicine,

[6]Applied Bioinformatics Laboratories, New York School of Medicine, New York, NY 10016,

USA

[7]Division of Gastroenterology and Hepatology, Department of Medicine, New York

University Langone Health, New York, NY 10016, USA.

*These authors contributed equally

†These authors contributed equally

§Correspondence to:

Png.Loke@nyulangone.org

Ken.Cadwell@med.nyu.edu

algraham@princeton.edu

SUMMARY

The relative contributions of genetic and environmental factors to variation in immune

responses are still poorly understood. Here, we performed a deep phenotypic analysis of

immunological parameters of laboratory mice released into an outdoor enclosure, carrying

susceptibility genes (*Nod2* and *Atg16l1*) implicated in the development of inflammatory

bowel diseases. Variations of immune cell populations were largely driven by environment,

whereas cytokine production in response to stimulation was affected more by genetic

mutations. Multi-omic models identified transcriptional signatures associated with differences

in T cell populations. Subnetworks associated with responses against *Clostridium perfringens,*

Candida albicans and *Bacteroides vulgatus* were also coupled with rewilding. Hence,

exposing laboratory mice carrying different genetic mutations to a natural environment

uncovered important contributors to immune variation.

One sentence summary

Natural environment exposure in laboratory mice primarily promotes variation in population

frequencies of immune cells, whereas cytokine responses to stimulation are affected more by

genetic susceptibility to inflammatory bowel disease.

INTRODUCTION

Variation in the magnitude of the immune response is an important determinant of susceptibility to pathogen infections, as well as a predisposition to autoimmunity and cancer (Brodin and Davis, 2017; Schirmer et al., 2018). However, immunological studies with laboratory mice typically aim to control variation in order to focus on genetic factors that alter components of both the adaptive and innate immune system. Recent advances in technology and throughput have facilitated a systems immunology approach towards deciphering the relative genetic and environmental contributors to variation of the human immune system (Bakker et al., 2018; Brodin et al., 2015; Li et al., 2016; Schirmer et al., 2016; Ter Horst et al., 2016). By altering the environment in which laboratory mice are conventionally housed, combined with deep phenotypic analysis of immune responses, we describe here a resource dataset that we used to probe the role of genetic mutations versus environmental drivers of heterogeneity in the immune system. Controlled experiments on specific groups with the same genetic mutations exposed to a new environment would be difficult to conduct in human studies.

Recently, there has been a growing effort to characterize wild mice and pet shop mice, which appear to share greater similarity to the immunological state of humans than laboratory mice (Abolins et al., 2017; Beura et al., 2016; Rosshart et al., 2019; Rosshart et al., 2017). We have developed an outdoor enclosure facility that enables us to reintroduce laboratory mice into a more natural environment, a process termed "rewilding" (Leung et al., 2018). The enclosures include soil and vegetation as well as barnlike structures but no other mammals (e.g., no wild conspecifics from whom lab-bred mice might acquire infections). Over the course of several weeks, mice dig burrows and otherwise adapt to their new environment (Supplemental Videos). Previously, we have observed C57BL/6 mice in this environment exhibit a decreased type 2 and increased type 1 response to intestinal nematode (*Trichuris muris*) infection and greater susceptibility to heavier worm burdens (Leung et al., 2018). This facility provides us with a unique opportunity to rapidly alter the housing environment of

laboratory mice that carry genetic mutations and be able to recover them for subsequent

detailed immunological profiling.

Gene-bacteria associations have been identified in both human and wild mice

indicating host genetic determinants of the gut microbial composition across mammals in

natural environments (Suzuki et al., 2019). To interrogate the impact of host genetic variants

on microbiota and inflammatory bowel disease (IBD) susceptibility in response to

environmental changes, we investigated *Atg16l1* and *Nod2* variants which are among the

highest risk factors for IBD associated with a dysregulated immune response to gut microbes

(Wlodarska et al., 2015; Wong and Cadwell, 2018). *Atg16l1* is a key component of the

autophagy pathway, and *Nod2* is involved in bacterial sensing. We and others previously

demonstrated that mice harboring mutations in *Atg16l1* or *Nod2* develop signs of

inflammation in a manner dependent on exposure to infectious entities (Biswas et al., 2010;

Cadwell et al., 2010; Lavoie et al., 2019; Matsuzawa-Ishimoto et al., 2017; Pott et al., 2018;

Ramanan et al., 2016; Ramanan et al., 2014). Mice with these particular mutations are

sensitive to the presence of murine norovirus (MNV), *Helicobacter* species, and other

commensal-like agents that are found in some animal facilities and excluded in others

(Biswas et al., 2010; Cadwell et al., 2010; Caruso et al., 2019; Kernbauer et al., 2014; Pott et

al., 2018; Ramanan et al., 2014). Since mice deficient in *Atg16l1* or *Nod2* can exhibit

dysregulated immune responses to colonization by organisms that are otherwise commensals

in wild type animals, we reasoned that they could be an interesting model system to

investigate gene-environment interactions in the outdoor enclosure.

We hypothesized that by introducing laboratory mice carrying IBD susceptibility

mutations into the outdoor enclosure, the drastic change in environment may trigger

alterations in the immune response and the microbiota, which would have greater adverse

effects on mutant mice than wildtype mice. Hence, to study the immunological consequences

of rewilding, we released *Atg16l1$^{T316A/T316A}$*, *Atg16l1$^{T316A/+}$*, *Nod2$^{-/-}$* mice in addition to

C57BL/6J wild type (WT) mice to determine whether mutations in these genes alter the

immune response to microbial exposure in a natural environment. Here we present a detailed

systems immunology profile of both rewilded and laboratory mice to further investigate

variation in immune response and the relationship with environment and genetic mutations.

RESULTS

Study design for immune profiling of rewilded and laboratory mice

From 116 mice released into the outdoor enclosure for 6-7 weeks we recovered 104 mice

(25 WT, 28 *Nod2$^{-/-}$*, 27 *Atg16l1$^{T316A/+}$*, and 24 *Atg16l1$^{T316A/T316A}$*) in time for analysis and

compared them to 80 matched controls (19 WT, 19 *Nod2$^{-/-}$*, 20 *Atg16l1$^{T316A/+}$*, 22

Atg16l1$^{T316A/T316A}$) maintained under specific pathogen free (SPF) conditions (herein referred

to as lab mice) (Figure 1A). Blood samples were collected at the time of sacrifice and

analyzed by flow cytometry with a lymphocyte panel (Table S1). Additionally, cytokine

production in the plasma was assayed (Figure 1A). Fecal samples and cecal contents were

collected for microbial profiling, and for reconstitution experiments (Companion study,

Yeung *et. al.*). Mesenteric lymph node (MLN) cells were isolated for flow cytometry analysis

with the same lymphocyte panel as the blood and an additional myeloid cell panel (STAR

METHODS). Immune activity in the MLN was measured through transcriptional profiles by

RNA-seq and cytokine production assays in response to microbial stimulations (STAR

METHODS). All of the lab mice were sacrificed in one week, whereas the rewilded mice

were sacrificed over a 2-week period based on trapping frequency. This collection of multi-

parameter datasets from nearly 200 mice provides an opportunity to examine inter-individual

variation in innate and adaptive immune cell populations by flow cytometry analysis, evaluate

cytokine responses to microbial stimulation, and integrate highly dimensional transcriptomics

and microbiota data. In order to perform multi-omic analysis, 81 mice were completely

profiled with measurements from all four data types; flow cytometry, MLN RNA-Seq, fecal

microbiota and MLN restimulation and cytokine profiles. A systems level analysis also

enabled us to quantify effects of environmental influences and IBD susceptibility gene

mutations, *Nod2* and *Atg16l1*, on the heterogeneity of immunological parameters among

individual mice.

Variation of immune cell populations in the blood is largely driven by the environment

In line with previous studies exposing lab mice to feral and pet shop mice (Beura et al., 2016), we observed major differences in CD44^hi^CD62L^lo^ or CD44^hi^CD62L^hi^ T cells (Companion study, Yeung *et. al.*). To visualize the overall landscape of flow cytometry data in an unbiased manner, phenotypic heterogeneity was analyzed on total CD45+ cells per mouse across the blood lymphocyte panel (Table S1) in lab and rewilded mice. We down sampled to 1,000 CD45+ cells from each mouse for nearly 180,000 single CD45+ cell events and single cell flow cytometry data was visualized by uniform manifold approximation and projection (UMAP) (Becht et al., 2019). UMAP analysis was superior to *t*-distributed stochastic neighborhood embedding (t-SNE) (Becht et al., 2019) (Figure S1A) for identifying major immune cell subsets (Figures 1B and 1C; Figures S1B-E). Unexpected clusters of CD44^hi^CD19+ cells were increased in the rewilded mice compared to lab mice (Figure 1B), while lab mice had more CD62L^hi^CD4+ cells (Figure 1C). This unsupervised visualization strategy was useful for identifying unexpected differences (e.g. in the CD19+ compartment and other populations not investigated by (Beura et al., 2016; Rosshart et al., 2019; Rosshart et al., 2017)) in cell clusters of the peripheral blood at a single cell level, which might have been overlooked by traditional flow cytometry gating.

In order to identify known immune cell populations, we employed a traditional gating strategy to systematically quantify the frequencies of immune cell subsets with a lymphocyte panel (Figure S2; Table S2). The blood lymphocyte panel identified 13 lymphocyte populations, a total myeloid cell population (CD11b/CD11c/DX5), and a total CD45+ cell population. From the proportions of these immune cell populations, we examined the inter-individual variation in lab and rewilded mice by principal component analysis (PCA) (Figure 1D). As shown by the separation of lab and rewilded mice along principle component (PC) 1 and 2, there is a strong effect of the environment on lymphocyte populations, and major differences between these populations can be attributed to CD44^hi^CD62L^hi^ CD4 and CD44^hi^CD62L^hi^ CD8 T cells as inferred by examining loading factors along PC1 and PC2

(Figure 1D). The reduced proportion of CD44^{lo}CD62L^{hi} T cells is consistent with some

previous reports on wild mice (Abolins et al., 2017; Beura et al., 2016). We also quantified

the extent of variability between lab and rewilded mice by measuring the Euclidean distances

between mice in our PCA. Notably, the within group distances in lab mice were significantly

less than rewilded mice (Figure 1E) suggesting that the natural environment increases

variability in immune cell populations. We then evaluated several population factors to

identify covariates with the largest effect sizes on immune cell populations (STAR

METHODS) (Figure 1F). As expected, the composition and activation state of immune cell

populations as determined by cell surface marker expression in the peripheral blood are

driven most strongly by environmental differences between the SPF facility and the outdoor

enclosure. These differences far outweighed any effects of genetic deficiency in *Nod2* or

Atg16l1 in both environments (Figure 1F). These findings are consistent with the concept that

non-heritable influences explain the majority of the variation in the proportion of immune cell

subsets as reported in twin studies (Brodin et al., 2015).

Variation of cytokine production in response to stimulations is driven more by genetic

mutations than by the environment.

Cytokine production capacity in humans has been reported to be more strongly influenced

by genetic than environmental factors (Li et al., 2016). To investigate the relative roles of

Nod2/*Atg16l1* deficiency and environment on induced responses, we performed *ex vivo*

stimulation experiments on total MLN cells from rewilded and lab mice. To examine cytokine

responses to microbial and T-cell specific antigens we stimulated with six different microbes

(*C. perfringens*, *P. aeruginosa*, *B. vulgatus*, *B. subtilis, S. aureus* and *C. albicans*),

αCD3/CD28 beads to activate T cells as well as PBS as a control (Figures 1A and 2A).

Despite no significant differences for PBS control samples between lab and rewilded mice

(data not shown), we chose to normalize cytokine production per mouse by expressing

cytokine data as a fold change levels over PBS (STAR METHODS). A broad array of 13

cytokines was measured: IFN-γ, IL-1α, IL-1ß, IL-13 IL-5, IL-10, IL-17, IL-6, CCL2, CXCL1,

CCL3, CCL4 and TNF-α. Average fold changes in cytokine production as compared to PBS

controls for both lab and rewilded mice indicate increases in IFN-γ and IL-17 after

αCD3/CD28 activation, consistent with the production of these cytokines by T cells in the

MLNs (Figure 2A). In general, bacterial antigens induced CCL3, IFN-γ, TNF-α, CCL4, and

IL-6 production, while the fungus *C. albicans* induced the regulatory cytokine IL-10 (Figure

2A). Bacterial antigens also induced CXCL1, which was not induced by *C. albicans* or

αCD3/CD28 (Figure 2A). As expected αCD3/CD28 was one of the most potent inducers of

cytokine production, especially for IFN-γ, CCL4, TNF-α and IL-5, and rewilded mice

produced higher levels of these cytokines than lab mice post stimulation (Figure 2B). IL-1α,

IL-1β, IL-13 and CCL2 were not appreciably altered by any stimulation (Figure 2A).

We next used PCA to assess the sources of variation in stimulated cytokine production for

both lab and rewilded mice (Figure 2C). Lab and rewilded mice do not clearly segregate

along PC1 or PC2 suggesting that environment is not a major source of variation. Effect size

measures based on the Euclidean distance indicated genotype as a much stronger source of

variation (Figure 2D). The differences in genotype are driven by IL-10 and IL-6 production as

indicated by the PCA loading factors (Figure 2C). More specifically, the rewilded *Nod2*$^{-/-}$

mice appear to cluster distinctly from other genotypes, suggesting key differences in cytokine

production of rewilded *Nod2*$^{-/-}$ mice (Figure 2C). Genotype also had the greatest effect size on

variation in plasma cytokine levels of individual mice (Figure S3).

Rewilded *Nod2*$^{-/-}$ mice exhibit increased cytokine responses compared to wild type

rewilded mice.

To better understand how *Nod2* and *Atg16l1* deficiency was affecting cytokine production, we

performed two-way comparisons and calculated a p-value for each comparison to determine

which cytokine producing conditions were most significantly different between WT mice and

mice with *Nod2* and *Atg16l1* deficiencies from laboratory and rewilded environments

(Figures 3A and 3B). Compared to the lab mice, the rewilded mice had many more significant

changes in stimulated cytokine production across *Nod2* and *Atg16l1* deficient mice (Figure

3A and 3B). These changes were largely driven by significant differences in rewilded *Nod2*[-/-]

mice against rewilded WT mice (Figure 3A). Specifically, the *Nod2*[-/-] mice exhibited

increases in IL-17, IL-5 and IL-10 in response to *C. perfringens* suggesting an increased

responsiveness to this bacterial stimulant that was only observed in the rewilded condition

(Figures 3A and 3C). Production of IL-10 in response to *C. albicans* was also elevated in

Nod2 and *Atg16l1* deficient mice, although *Nod2-/-* did not exhibit greater fungal colonization

than WT mice (Figure 3C) (see companion study Yeung. et.al.). Despite these differences in

cytokine responses, the rewilded *Nod2-/-* did not shown any signs of increased intestinal

inflammation by histology or changes in goblet cell numbers (data not shown), which we

have previously shown to be associated with susceptibility to *B. vulgatus* colonization

(Ramanan et al., 2016; Ramanan et al., 2014). Together these results indicate that exposure to

the outdoor environment per se that induces elevated cytokine responses is not a sufficient hit

by itself to trigger intestinal inflammation. Additional insults (i.e. a multi-hit model) may be

needed to trigger full blown pathology (Ramanan et al., 2016).

An integrated classification model identifies features predictive of environment and

genotype-specific effects.

In addition to stimulated cytokine profiles, MLN cells were assessed for immune cell

populations by flow cytometry analysis (Figure S4 and Table S2) and gene expression by

RNA-seq analysis (Figures S5A and S5B; Table S2). Consistent with the peripheral blood,

immune cell frequencies in MLNs showed a strong effect of environment (Figure S6).

Interestingly, a group of rewilded mice all sacrificed at week 7 of rewilding clustered

separately in the PCA analysis of lymphocyte populations (Figure S6B). Between week 6 and

7 of rewilding it is possible specific events in the enclosure (e.g., a severe thunderstorm, high

humidity and fungal blooms observed during that week), could have resulted in changes in T

cell activation. Similar to blood lymphocyte populations, RNA-seq analysis (Figures S5A and

S5B) and 16S rRNA profiling (Figures S5C-E; see companion study by *Yeung et al.*) showed

some environmental effects on variations. This large amount of data prompted us to use a

multi-omic classification model that could prioritize features predictive of environment and

genotype-specific effects, as well as identify relationships between these features.

We integrated multi-omic data from flow cytometry populations, gene expression,

cytokine, and microbial profiles (Table S2) in a subset of mice (n-=81) which had measures

for all four data types. A concatenated matrix (Table S3) of these features was used to build a

random forest classification model (STAR METHODS) to distinguish samples by

environment or genotype (Figure 4). The contribution of each feature to the model was also

assessed to identify features predictive of environment or genotype-specific effects (Figures

4A and 4C). When classifying the environment, a hub of transcription factors (including

Zfp36, *Atf4*, *Fos*, several *Jun* and *Klf* family members) and *Cd69* are strong predictive

features of environment (Figure 4A). Pairwise correlation analysis between these top features

also reveals associations to CD44^{hi}CD62L^{hi} populations for CD4+ and CD8+ T cells in the

blood and MLN, as well as IL-5 and TNF-α production in response to anti CD3/CD28

stimulation (Figure 4B). For the genotype model, our classification was not as accurate as the

environment, as measured by area under the receiver operator curve (AUC) of 0.78 versus 1

on the test set (STAR METHODS). Pairwise correlation analysis revealed fewer associations

between these features and overall less connected nodes (Figure 4D). Instead of cytokines

associated with T cell activation by αCD3/CD28, IL-17 responses to microbial antigen (*B.*

vulgatus and *P. aeruginosa*) stimulation were correlated (Figures 4B and 4D). Interestingly,

expressions of a number of ribosomal proteins (*Rps* and *Rpl* proteins) were also predictive of

genotype (Figures 4C and 4D). In summary, we identified a network of transcription factor

associated with T cells in the MLNs based on transcriptional profiling data that is strongly

predictive of the effects of the different environments.

Integrative network analysis identifies co-regulated modules in rewilded mice

To evaluate the interactions between data types, we next utilized an unsupervised sparse

partial least squares (sPLS) regression model to integrate the different data types into a multi-

component network (Figures 5A and S7). To condense the size of our network we collapsed

our gene expression profiles according to known Gene Ontologies (STAR METHODS) (2015;

Ashburner et al., 2000) for immune system processes in *Mus musculus*, which also focuses

our attention on the most relevant genes. In short, we compared our gene expression profiles

to all child terms of the Gene Ontology term "immune system process" (GO:0002376) and

collapsed our genes into modules based on these pre-defined ontologies. 954 genes were used

to generate 91 specific gene ontologies from our gene expression data all related to the parent

gene ontology "immune system process". These modules were used as inputs alongside the

cytokine profiles, microbial profiles and flow cytometry populations to generate an

unsupervised multi-omic network. sPLS-regression models were built pairwise between each

data type to generate a 188-node covariance network with 577 total connections between the

four different types of features (Figures 5A and S7). In the resulting model, all microbial taxa

fell below our 0.6 covariance threshold and were therefore removed (Table S4). This

indicated that bacterial composition, determined by 16S sequencing alone, was not an

important component of the network. Perhaps other strategies to assess microbial function

(e.g. transcriptomics, metagenomics or metabolomics) would have yielded more connections

to immune function.

While total CD4+ and CD8+ T cell populations in the MLN were among the most

connected nodes of the network specific gene ontologies, cytokine responses and other

immune populations were also strongly interconnected (Figure 5A). As expected from earlier

analysis (Figure 1), the blood CD44^{hi}CD62L^{hi} CD4 T cell population was strongly

interconnected and enriched in rewilded mice (Figures 5A and 5B). Additionally, by

collapsing our gene expression profiles into annotated gene ontologies, we identified a

module (GO:0061844: antimicrobial humoral immune response mediated by antimicrobial

peptides) as the most connected gene module that was enriched in rewilded mice (Figures 5A

and 5B). Interestingly, IL-5 production in response to *C. perfrigens* stimulation was the most

connected cytokine response and was found to be higher in rewilded mice (Figures 5A and 5B;

Figure S7). This node is part of a 15-node subnetwork connected to total CD4 and CD8 T cell

populations in the MLN, several other cytokine responses and the GO:0030593: neutrophil

chemotaxis module (Figure 5C). This module is of particular interest because of the increased

neutrophilia observed in the rewilded mice (Yeung et.al. companion paper), which is driven

by increased fungal colonization. Indeed, this module is tightly linked with cytokine

responses to *C. albicans* antigen stimulation (Figure 5C). While expressions of genes in this

module are increased by the rewilding environment, there does not appear to be genotype

specific differences regulating expression of the genes in this module (Figure 5D). This multi-

omics network analysis therefore substantiated the omics-by-omics inferences (Figures 1 and

2) but also provided new insight by identifying an important role for *C. albicans* responses

during rewilding, which is more fully addressed in the companion study (Yeung et.al.

companion paper).

DISCUSSION

In summary, we found that changing the environment profoundly contributes to the

inter-individual variations on immune cell frequencies, whereas cytokine responses to

pathogen stimulation were more affected by genetic deficiencies in the IBD susceptibility

genes *Nod2* and *Atg16l1*. These observations are consistent with human twin studies whereby

the majority of cell population influences (measured by flow and mass cytometry) are

determined by non-heritable factors (Brodin et al., 2015), whereas cytokine production

capacity in response to stimulation may be more strongly influenced by genetic factors (Li et

al., 2016). Another large human study indicated that features of innate immunity were more

strongly controlled by genetic variation than lymphocytes, which were driven by

environmental effects (Patin et al., 2018). Thus, although studies of lab mice, by design,

often emphasize genetic effects on immune phenotype (including cell population

distributions), here we show that even brief exposure to a natural environment renders

predictors of immune phenotype in mice a better match for predictors of human immune

phenotypes. Additionally, our dataset described here will be a useful resource for other

investigators to delve into the immunological consequences of rewilding.

An attractive hypothesis arising from these findings is that perhaps environment is a

primary driver of the composition of the immune system, but genetics is a stronger driver of

per-cell responsiveness. However, in this study we only investigated how *Nod2* and *Atg16l1*

influenced the immunological parameters we measured. To test this hypothesis further would

require us to release mice from diverse genetic backgrounds into the outdoor enclosure in a

similar experiment. This experiment only examined the effects of *Nod2* and *Atg16l1*

deficiency in the C57BL/6 background. For example, analysis of macrophage activation from

five different strains of mice that provided genetic variation on the order of the human

population yielded considerable insights into how genetic variation affects transcriptional

regulation mechanisms (Link et al., 2018). Alternatively, mice from the collaborative cross

may enable high-resolution genome mapping for such complex traits as cytokine production

in response to stimulation (Noll et al., 2019). Studies of wild animal populations may also

determine if per cell cytokine responses show a higher genetic variance than do immune cell

population distributions. Hence, the combination of new environmental challenge strategies

such as rewilding of mouse models, combined with modern multi-omic approaches in fully

wild systems, should enable us to better define determinants of immune variation at a

molecular level.

This study was initially designed to test the hypothesis that *Nod2* and *Atg16l1*

deficient mice may respond to the re-wilding environment in an adverse way. Our previous

studies had found that *B. vulgatus* colonization of *Nod2* and norovirus colonization of *Atg16l1*

deficient mice predisposed them to intestinal inflammation. Hence, we wanted to examine if

these mice with mutations in genes associated with the development of IBD would be

associated with environmental triggering of intestinal inflammation. However, we did not

observe significant differences in intestinal inflammation based on histology for either the

Nod2 or *Atg16l1* deficient mice, although we find an elevated production of cytokines in the

Nod2 deficient mice in response to the rewilding environment. While there were not

significant differences in intestinal inflammation, it is possible that additional insults (i.e. a

multi-hit model) are needed to trigger intestinal inflammation that is sufficiently severe to be

discernable by histologic examination. In our previous studies, an additional piroxicam (for

the *Nod2* deficient mice (Ramanan et al., 2016) or dextran sodium sulfate (DSS) (for the

Atg16l1 deficient mice (Matsuzawa-Ishimoto et al., 2017)) insult was required to drive

pathogenesis of the *B. vulgatus*/norovirus colonized mice.

A systems level analysis identified interconnected networks of transcriptional

signatures, immune cell populations and cytokine profiles to microbial stimulation,

highlighting a potentially important role for fungal stimulation from rewilding (Companion

study, Yeung *et. al.*). This unanticipated result may have been missed through conventional

comparisons. Increased colonization by commensal fungi is perhaps the most striking

environmental effect of the rewilding experiment that we performed. While *Nod2* deficiency

did not affect fungal colonization, *Atg16l1* deficient mice are more susceptible (Companion

study, Yeung *et. al.*). Future studies with mice from more genetically diverse backgrounds

will better define the genetic basis of host susceptibility to commensal fungi colonization

through rewilding and determine how this may affect the innate and adaptive immune

responses.

Hence, rewilding laboratory mice of different genetic backgrounds and careful

monitoring of other non-heritable influences (e.g. differential behavior via which divergent

immunological experience of individuals may accrue, by analogy with causes of divergence

in neurological phenotype (Freund et al., 2013) observed during rewilding (Cope et al., 2019)

could be a powerful new approach towards dissecting drivers of immune response

heterogeneity under homeostatic as well as during disease settings or infectious challenges.

References:

(2015). Gene Ontology Consortium: going forward. Nucleic Acids Res. 43(Database
issue), D1049-1056. Published online 2014/11/28 DOI: gku1179 [pii]
10.1093/nar/gku1179.

Abolins, S., King, E.C., Lazarou, L., Weldon, L., Hughes, L., Drescher, P., Raynes, J.G.,
Hafalla, J.C.R., Viney, M.E., and Riley, E.M. (2017). The comparative immunology of
wild and laboratory mice, Mus musculus domesticus. Nat Commun. 8, 14811. Published
online 2017/05/04 DOI: ncomms14811 [pii]
10.1038/ncomms14811.

Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P.,
Dolinski, K., Dwight, S.S., Eppig, J.T., et al. (2000). Gene ontology: tool for the
unification of biology. The Gene Ontology Consortium. Nat Genet. 25(1), 25-29.
Published online 2000/05/10 DOI: 10.1038/75556.

Bakker, O.B., Aguirre-Gamboa, R., Sanna, S., Oosting, M., Smeekens, S.P., Jaeger, M.,
Zorro, M., Vosa, U., Withoff, S., Netea-Maier, R.T., et al. (2018). Integration of multi-
omics data and deep phenotyping enables prediction of cytokine responses. Nat
Immunol. 19(7), 776-786. Published online 2018/05/23 DOI: 10.1038/s41590-018-0121-3
10.1038/s41590-018-0121-3 [pii].

Becht, E., McInnes, L., Healy, J., Dutertre, C.A., Kwok, I.W.H., Ng, L.G., Ginhoux, F.,
and Newell, E.W. (2019). Dimensionality reduction for visualizing single-cell data using
UMAP. Nat Biotechnol. Published online 2018/12/12 DOI: nbt.4314 [pii]
10.1038/nbt.4314.

Beura, L.K., Hamilton, S.E., Bi, K., Schenkel, J.M., Odumade, O.A., Casey, K.A.,
Thompson, E.A., Fraser, K.A., Rosato, P.C., Filali-Mouhim, A., et al. (2016).
Normalizing the environment recapitulates adult human immune traits in laboratory
mice. Nature. 532(7600), 512-516. Published online 2016/04/21 DOI: nature17655 [pii]
10.1038/nature17655.

Biswas, A., Liu, Y.J., Hao, L., Mizoguchi, A., Salzman, N.H., Bevins, C.L., and
Kobayashi, K.S. (2010). Induction and rescue of Nod2-dependent Th1-driven
granulomatous inflammation of the ileum. Proc Natl Acad Sci U S A. 107(33), 14739-
14744. Published online 2010/08/04 DOI: 1003363107 [pii]
10.1073/pnas.1003363107.

Brodin, P., and Davis, M.M. (2017). Human immune system variation. Nat Rev
Immunol. 17(1), 21-29. Published online 2016/12/06 DOI: nri.2016.125 [pii]
10.1038/nri.2016.125.

Brodin, P., Jojic, V., Gao, T., Bhattacharya, S., Angel, C.J., Furman, D., Shen-Orr, S.,
Dekker, C.L., Swan, G.E., Butte, A.J., et al. (2015). Variation in the human immune
system is largely driven by non-heritable influences. Cell. 160(1-2), 37-47. DOI:
10.1016/j.cell.2014.12.020.

Cadwell, K., Patel, K.K., Maloney, N.S., Liu, T.C., Ng, A.C., Storer, C.E., Head, R.D.,
Xavier, R., Stappenbeck, T.S., and Virgin, H.W. (2010). Virus-plus-susceptibility gene
interaction determines Crohn's disease gene Atg16L1 phenotypes in intestine. Cell.
141(7), 1135-1145. Published online 2010/07/07 DOI: S0092-8674(10)00545-3 [pii]

10.1016/j.cell.2010.05.009.

Caruso, R., Mathes, T., Martens, E.C., Kamada, N., Nusrat, A., Inohara, N., and Nunez,
G. (2019). A specific gene-microbe interaction drives the development of Crohn's
disease-like colitis in mice. Sci Immunol. 4(34). Published online 2019/04/21 DOI:
4/34/eaaw4341 [pii]10.1126/sciimmunol.aaw4341.

Cope, E.C., Opendak, M., LaMarca, E.A., Murthy, S., Park, C.Y., Olson, L.B., Martinez,
S., Leung, J.M., Graham, A.L., and Gould, E. (2019). The effects of living in an outdoor
enclosure on hippocampal plasticity and anxiety-like behavior in response to nematode
infection. Hippocampus. 29(4), 366-377. Published online 2018/09/27 DOI:
10.1002/hipo.23033.

Freund, J., Brandmaier, A.M., Lewejohann, L., Kirste, I., Kritzler, M., Kruger, A.,
Sachser, N., Lindenberger, U., and Kempermann, G. (2013). Emergence of individuality
in genetically identical mice. Science. 340(6133), 756-759. Published online 2013/05/11
DOI: 340/6133/756 [pii]10.1126/science.1235294.

Kernbauer, E., Ding, Y., and Cadwell, K. (2014). An enteric virus can replace the
beneficial function of commensal bacteria. Nature. 516(7529), 94-98. DOI:
10.1038/nature13960.

Lavoie, S., Conway, K.L., Lassen, K.G., Jijon, H.B., Pan, H., Chun, E., Michaud, M.,
Lang, J.K., Gallini Comeau, C.A., Dreyfuss, J.M., et al. (2019). The Crohn's disease
polymorphism, ATG16L1 T300A, alters the gut microbiota and enhances the local
Th1/Th17 response. Elife. 8. Published online 2019/01/23 DOI: 10.7554/eLife.39982
39982 [pii].

Leung, J.M., Budischak, S.A., Chung The, H., Hansen, C., Bowcutt, R., Neill, R.,
Shellman, M., Loke, P., and Graham, A.L. (2018). Rapid environmental effects on gut
nematode susceptibility in rewilded mice. PLoS Biol. 16(3), e2004108. Published online
2018/03/09 DOI: 10.1371/journal.pbio.2004108
pbio.2004108 [pii].

Li, Y., Oosting, M., Smeekens, S.P., Jaeger, M., Aguirre-Gamboa, R., Le, K.T.T., Deelen,
P., Ricano-Ponce, I., Schoffelen, T., Jansen, A.F.M., et al. (2016). A Functional Genomics
Approach to Understand Variation in Cytokine Production in Humans. Cell. 167(4),
1099-1110 e1014. Published online 2016/11/05 DOI: S0092-8674(16)31400-3 [pii]
10.1016/j.cell.2016.10.017.

Link, V.M., Duttke, S.H., Chun, H.B., Holtman, I.R., Westin, E., Hoeksema, M.A., Abe,
Y., Skola, D., Romanoski, C.E., Tao, J., et al. (2018). Analysis of Genetically Diverse
Macrophages Reveals Local and Domain-wide Mechanisms that Control Transcription
Factor Binding and Function. Cell. 173(7), 1796-1809 e1717. Published online 2018/05/22
DOI: S0092-8674(18)30511-7 [pii]10.1016/j.cell.2018.04.018.

Matsuzawa-Ishimoto, Y., Shono, Y., Gomez, L.E., Hubbard-Lucey, V.M., Cammer, M.,
Neil, J., Dewan, M.Z., Lieberman, S.R., Lazrak, A., Marinis, J.M., et al. (2017).
Autophagy protein ATG16L1 prevents necroptosis in the intestinal epithelium. J Exp
Med. 214(12), 3687-3705. Published online 2017/11/02 DOI: jem.20170558 [pii]
10.1084/jem.20170558.

Noll, K.E., Ferris, M.T., and Heise, M.T. (2019). The Collaborative Cross: A Systems
Genetics Resource for Studying Host-Pathogen Interactions. Cell Host Microbe. 25(4),
484-498. Published online 2019/04/12 DOI: S1931-3128(19)30161-1 [pii]

10.1016/j.chom.2019.03.009.

Patin, E., Hasan, M., Bergstedt, J., Rouilly, V., Libri, V., Urrutia, A., Alanio, C.,
Scepanovic, P., Hammer, C., Jonsson, F., et al. (2018). Natural variation in the
parameters of innate immune cells is preferentially driven by genetic factors. Nat
Immunol. 19(3), 302-314. Published online 2018/02/25 DOI: 10.1038/s41590-018-0049-7
10.1038/s41590-018-0049-7 [pii].

Pott, J., Kabat, A.M., and Maloy, K.J. (2018). Intestinal Epithelial Cell Autophagy Is
Required to Protect against TNF-Induced Apoptosis during Chronic Colitis in Mice.
Cell Host Microbe. 23(2), 191-202 e194. DOI: 10.1016/j.chom.2017.12.017.

Ramanan, D., Bowcutt, R., Lee, S.C., Tang, M.S., Kurtz, Z.D., Ding, Y., Honda, K.,
Gause, W.C., Blaser, M.J., Bonneau, R.A., et al. (2016). Helminth infection promotes
colonization resistance via type 2 immunity. Science. 352(6285), 608-612. Published
online 2016/04/16 DOI: science.aaf3229 [pii]10.1126/science.aaf3229.

Ramanan, D., Tang, M.S., Bowcutt, R., Loke, P., and Cadwell, K. (2014). Bacterial
sensor Nod2 prevents inflammation of the small intestine by restricting the expansion of
the   commensal   Bacteroides   vulgatus.   Immunity.   41(2),   311-324.   DOI:
10.1016/j.immuni.2014.06.015.

Rosshart, S.P., Herz, J., Vassallo, B.G., Hunter, A., Wall, M.K., Badger, J.H.,
McCulloch, J.A., Anastasakis, D.G., Sarshad, A.A., Leonardi, I., et al. (2019).
Laboratory mice born to wild mice have natural microbiota and model human immune
responses. Science. 365(6452). DOI: 10.1126/science.aaw4361.

Rosshart, S.P., Vassallo, B.G., Angeletti, D., Hutchinson, D.S., Morgan, A.P., Takeda, K.,
Hickman, H.D., McCulloch, J.A., Badger, J.H., Ajami, N.J., et al. (2017). Wild Mouse
Gut Microbiota Promotes Host Fitness and Improves Disease Resistance. Cell. 171(5),
1015-1028 e1013. DOI: 10.1016/j.cell.2017.09.016.

Schirmer, M., Kumar, V., Netea, M.G., and Xavier, R.J. (2018). The causes and
consequences of variation in human cytokine production in health. Curr Opin Immunol.
54, 50-58. Published online 2018/06/19 DOI: S0952-7915(17)30173-5 [pii]
10.1016/j.coi.2018.05.012.

Schirmer, M., Smeekens, S.P., Vlamakis, H., Jaeger, M., Oosting, M., Franzosa, E.A.,
Horst, R.T., Jansen, T., Jacobs, L., Bonder, M.J., et al. (2016). Linking the Human Gut
Microbiome   to   Inflammatory   Cytokine   Production   Capacity.   Cell.   167(7),   1897.
Published online 2016/12/17 DOI: S0092-8674(16)31658-0 [pii]
10.1016/j.cell.2016.11.046.

Suzuki, T.A., Phifer-Rixey, M., Mack, K.L., Sheehan, M.J., Lin, D., Bi, K., and
Nachman, M.W. (2019). Host genetic determinants of the gut microbiota of wild mice.
Mol Ecol. 28(13), 3197-3207. Published online 2019/05/30 DOI: 10.1111/mec.15139.

Ter Horst, R., Jaeger, M., Smeekens, S.P., Oosting, M., Swertz, M.A., Li, Y., Kumar, V.,
Diavatopoulos,   D.A.,   Jansen,   A.F.M.,   Lemmers,   H.,   et   al.   (2016).   Host   and
Environmental Factors Influencing Individual Human Cytokine Responses. Cell. 167(4),
1111-1124 e1113. Published online 2016/11/05 DOI: S0092-8674(16)31401-5 [pii]
10.1016/j.cell.2016.10.018.
540

541 **Wlodarska, M., Kostic, A.D., and Xavier, R.J. (2015). An integrative view of**
542 **microbiome-host interactions in inflammatory bowel diseases. Cell Host Microbe. 17(5),**
543 **577-591. DOI: 10.1016/j.chom.2015.04.008.**
544
545 **Wong, S.Y., and Cadwell, K. (2018). There was collusion: Microbes in inflammatory**
546 **bowel disease. PLoS Pathog. 14(9), e1007215. DOI: 10.1371/journal.ppat.1007215.**
547
548
549

Acknowledgements

We wish to thank William Craigens, Daniel Navarrete Prado, Allison Lee, and Veena Chittamuri for assistance with trapping and husbandry in the field, the PU Lab Animal Resources staff for logistical support. We wish to thank the NYU School of Medicine Flow Cytometry and Cell Sorting, Microscopy, Genome Technology, and Histology Cores for use of their instruments and technical assistance (supported in part by National Institute of Health (NIH) grant P31CA016087, S10OD01058, and and S10OD018338). **Funding:** this research was supported by US National Institute of Health (NIH) grants DK103788 (K.C. and P.L.), AI121244 (K.C.), HL123340 (K.C.), DK093668 (K.C.), AI130945 (P.L.), R01 HL125816 (K.C.), HL084312, AI133977 (P.L.), research station and research rebate awards from PU EEB (A.L.G.), pilot award from the NYU CTSA grant UL1TR001445 from the National Center for Advancing Translational Sciences (NCATS) (K.C., P.L.), pilot award from the NYU Cancer Center grant P30CA016087 (K.C.), AI100853 (Y.C.), and DK122698 (F.Y.). This work was also supported by the Department of Defense grant W81XWH-16-1-0256 (P.L.), Faculty Scholar grant from the Howard Hughes Medical Institute (K.C.), Crohn's & Colitis Foundation (K.C.), Merieux Institute (K.C.), Kenneth Rainin Foundation (K.C.), Stony-Wold Herbert Fund (K.C.), and Bernard Levine Postdoctoral Research Fellowship in Immunology (Y.C.). K.C. is a Burroughs Wellcome Fund Investigator in the Pathogenesis of Infectious Diseases. **Author contributions:** Design of experiments, data analysis, data discussion, and interpretation: J.D.L., J.C.D., F.Y., K.C., A.L.G, and P.L.; primary responsibility for execution of experiments: J.D.L., F.Y., C.M., J.M.L., Y.H.C., A.C., C.H., and C.D.D.; MLN cell RNA and 16S analysis: J.C.D., K.V.R., J.D.L., and F.Y. Supervised and Unsupervised machine learning model analysis: J.C.D., K.V.R., and J.D.L. All authors discussed data and commented on the manuscript. **Competing interests:** K.C. and P.L. receive research funding from Pfizer. K.C. has consulted for or received an honorarium from Puretech Health, Genentech, and Abbvie. P.L. consults for and has equity in Toilabs. **Data and materials availability:** Raw sequence data from 16S, ITS, and RNA sequencing experiments are deposited in the NCBI Sequence Read Archive under BioProject accession number PRJNA559026 and gene expression omnibus (GEO) accession number GSE135472.

     Figure Legends:

Figure 1. Environmental change drives inter-individual variation in immune cell

populations. (**A**) 25 C57BL/6$^{+/+}$, 28 *Nod2*$^{-/-}$, 27 *Atg16l1*$^{T316A/+}$, 24 *Atg16l1*$^{T316A/T316A}$ mice

(total=104) were housed in the wild enclosure (Rewilded) for 6-7 weeks and successfully

trapped for flow cytometry analysis of lymphoid and myeloid cell populations in the blood

and mesenteric lymph nodes (MLNs). Plasma was also collected for cytokine profiles. Age

matched 19 C57BL/6$^{+/+}$, 19 *Nod2*$^{-/-}$, 20 *Atg16l1*$^{T316A/+}$, 22 *Atg16l1*$^{T316A/T316A}$ mice (total=80)

housed under SPF conditions (Lab) were analyzed as controls. Feces were collected for 16S

rRNA sequencing. MLN cells were cultured with indicated stimulates for 48 hours and

supernatants were collected for cytokine profiling. Samples that fail quality control are not

included in downstream analyses. (**B** and **C**) UMAP projections of ~180,000 CD45+,

~110,000 CD19+, and ~36,000 CD4+ cells on flow cytometry data of the blood from lab and

rewilded mice. Events are color-coded according to CD19, CD44 (B), CD4, and CD62L (C)

fluorescence level. Box plots show the abundance of CD44^{hi} CD19+ cells (B) and CD62L^{hi}

CD4+ cells (C) in the blood of individual lab and rewilded mice. (**D**) Principal component

analysis (PCA) of gated immune cell populations in the blood and the density of each

population along the principal component (PC). Right panel indicate biplots of the gated

immune cell populations that are projected onto PC1 and PC2. (**E**) Box plot of pairwise

Euclidean distance measures based on blood immune cell population in the blood of lab

versus lab, rewilded versus rewilded and lab versus rewilded mice. (**F**) Bar plot showing the

pseudo R^2 measure of effect size on the entire distance matrix used to calculate the PCA of

immune cell populations in the blood (D). ***$P < 0.001$ by Kruskal-Wallis test between

groups (B, C) and ***$P < 0.001$ by Mann-Whitney-Wilcoxon test between groups (E).

Figure 2. The genetic contribution towards variation in cytokine production in response

to microbial stimulation is masked when mice are rewilded. (**A** to **D**) MLN cells from lab

and rewilded mice were *ex vivo* cultured with *B. subtilis*, *S. aureus*, *C. perfringens*, *B.*

vulgatus, *P. aeruginosa*, *C. albicans*, and αCD3/CD28 beads for 48 hours and the

supernatants were assayed for 13 cytokines. (**A**) Heat map of average fold changes compared

to cytokine levels of MLN cells cultured with PBS controls from all mice. (**B**) Box plot

showing the top 10 log$_2$ fold changes of cytokine responses to stimulation for MLN cells

from lab and rewilded mice. (**C**) PCA plot of of fold change cytokine stimulation profiles

with histograms of the total variance explained by each PC indicated on the axis. Right panel

indicates the loading factor of each stimulated cytokine profile that contributes to each PC. (**D**)

Bar plot showing the pseudo R^2 measure of effect size on the entire distance matrix used to

calculate the PCA of cytokine profiles (C). * P <0.05, ** P <0.01, *** $P < 0.001$ by Mann-

Whitney-Wilcoxon test between groups (B).

Figure 3. Rewilded *Nod2$^{-/-}$* mice display greater changes in cytokine production

compared to rewilded WT mice. (**A** to **C**) MLN cells from lab and rewilded mice were *ex*

vivo cultured with *B. subtilis*, *S. aureus*, *C. perfringens*, *B. vulgatus*, *P. aeruginosa*, *C.*

albicans, and αCD3/CD28 beads for 48 hours and the supernatants were assayed for 13

cytokines. (**A** and **B**) Heatmap of significant p-values from Mann-Whitney-Wilcoxon test

between group results comparing fold changes of stimulated cytokine production in rewilded

(**A**) and lab (**B**) WT mice against *Nod2$^{-/-}$*, *Atg16l1$^{T316A/+}$* and *Atg16l1$^{T316A/T316A}$* mice

respectively. Within each cytokine condition the order of stimulation remains constant as

follows; *B. subtilis*, *B. vulgatus. C. albicans*, *αCD3/CD28*, *C. perfringens*, *P. aeruginosa*, *S.*

aureus. (**C**) Representative box plots of the most significant changes between WT and *Nod2$^{-/-}$*

mice. * $P <0.05$, ** $P <0.01$, *** $P < 0.001$ by Mann-Whitney-Wilcoxon test between groups

(C).

Figure 4. Multi-omic classification models identify predictive features of environment

and genotype. (**A**) Bar plot of the top 40 features predictive of the environment and (**B**)

interaction network analysis of correlations ($R^2 > 0.6$ or $R^2 < -0.6$) between features (genes,

immune cell populations, OTUs or cytokines) that are predictive of environmental changes.

(**C**) Bar plot of the top 40 features predictive of the genotype and (**D**) network analysis of the

associations between features that are associated with genotype differences. Data types are

colored according to cytokine profiles (green), genes (red), immune populations (blue), and

OTUs (purple). The size of the nodes in B and D are scaled proportionally to the number of

connections with other nodes in each network.

Figure 5. An unsupervised multi-component model identifies a *C. albicans* and *C.*

***perfringens* response network associated with environmental changes.** (**A**) Data matrices

from flow cytometry data, transcriptional profiles, microbial profiles and stimulated cytokine

profiles were used as inputs for a sPLS-regression model to generate a covariance network. (**B**)

The most connected nodes, Blood CD4 T-cells CD44hi CD62Lhi, IL-5 in response to *C.*

perfringens, and GO:0061844 (antimicrobial humoral immune response mediated by

antimicrobial peptides) were evaluated for a difference between lab and rewilded mice

populations. (**C**) The IL-5 *C. perfringens* subnetwork was derived from the network and re-

calculated to highlight the connections most related to IL-5 production. (D) GO: 0030593

(neutrophil chemotaxis) was identified as a significant connection in the subnetwork and the

genes found in that GO term were plotted as a heatmap. Color bars indicate environment and
genotype. The size of the nodes in (**A**) and (**C**) are scaled proportionally to the number of
connections with other nodes in each network. Data types are colored according to cytokine
profiles (green), GO term (red), immune populations (blue). * P <0.05, ** P <0.01, *** *P* <
0.001 by Mann-Whitney-Wilcoxon test between groups (B).

Figure S1: Identification of immune cell populations that differ between lab and
rewilded mice with UMAP. (**A**) Projection of ~180,000 CD45+ cells visualized by tSNE vs
UMAP. (**B**) Identification of the major lymphocyte populations on UMAP based on CD19,
CD3, CD4 and CD8 expression. (**C** to **E**) Expression of cell surface molecules on CD19+
cells (C), CD4+ cells (D) and CD8+ cells (E). In (C), (D) and (E), expression of 1B11, PD1,
CD25, CD44, CD127, CXCR3, KLRG1, CD27, CD69, and CD62L are shown as high (purple)
or low (grey) from cut-off values based on overall distribution of fluorescence.

Figure S2: Flow cytometry gating strategy for the quantification of lymphoid
populations in the blood of lab and rewilded mice. (**A** to **H**) Representative flow cytometry
plots illustrating gating strategy (A) of lymphoid populations in the blood including the total
CD45+ cells (B), total myeloid cells (C), total B and T cells (D), total CD4 and CD8 T cells
(E), CD44^{lo}CD62L^{hi}, CD44^{hi}CD62L^{hi}, CD44^{hi}CD62L^{lo}, CD44^{lo}CD62L^{lo} CD8 T cells (F),
CD44^{lo}CD62L^{hi}, CD44^{hi}CD62L^{hi}, CD44^{hi}CD62L^{lo}, CD44^{lo}CD62L^{lo} CD4 T cells (G), and
CD25+CD4+ T cells (H).

Figure S3: A genetic contribution towards variation in plasma cytokine levels of lab and
rewilded mice. (**A** and **B**) Levels of RANTES, MIP-3a, KC, TNF-α, MCP-1, IP-10, MIP-1a,
MIP-1b, IL-1α, IL-6 cytokines were measured from the plasma of lab and rewilded mice at
the time of sacrifice with bead-based immunoassays (LEGENDplex). (**A**) PCA of plasma
cytokine levels in the blood of individual mice and the density of each population along the
principal components (PC). Right panel indicate biplots of the plasma cytokines are projected
onto PC1 and PC2. (**B**) Bar plot showing the pseudo R^2 measure of the effect size of different
variables on the variance of plasma cytokine levels.

Figure S4: Memory CD4+ and CD8+ T cells are increased in the mesenteric lymph
nodes (MLNs) of rewilded mice. (**A** to **F**) Representative flow cytometry plots for immune
cell populations after gating for CD45+ cells in the MLNs of lab or rewilded C57BL/6 mice;
for the total abundance of myeloid cells (A), B and T cells (B), CD4 and CD8 T cells (C),
CD44^{lo}CD62L^{hi}, CD44^{hi}CD62L^{hi}, CD44^{hi}CD62L^{lo}, CD44^{lo}CD62L^{lo} CD4 T cells (D),

CD44^{lo}CD62L^{hi}, CD44^{hi}CD62L^{hi}, CD44^{hi}CD62L^{lo}, CD44^{lo}CD62L^{lo} CD8 T cells (E), and

CD25+CD4+ T cells (F).

Figure S5: Environment has the greatest effect on variation of transcriptional profiles

and microbial composition. (**A** to **B**) Transcriptional profiles of MLNs were determined by

Celseq. (**A**) PCA based on gene profiles in the MLN and the density of each population along

the principal components (PC). Right panel indicates biplots of the genes projected onto PC1

and PC2. (**B**) Bar plot showing the pseudo R^2 measure of effect size on the entire distance

matrix used to calculate the PCA of gene expression levels. (**C**) PCoA based on Bray Curtis

distances of 16s rRNA sequencing profiles in all lab and rewilded mice across all genotypes,

(**D**) lab wild type (WT) and *Atg16l*$^{+/+}$ only, (**E**) lab wild type (WT), *Atg16l*$^{+/+}$ and *Nod2*$^{-/-}$ and

(**F**) lab wild type (WT), *Atg16l*$^{+/+}$ and *Nod2*$^{-/-}$ by respective cages. P-values were determined

by an Adonis test from Bray Curtis distances.

Figure S6: Unmeasured environmental effects drive additional variation of MLN

immune cell populations for the rewilded mice. (**A** and **B**) PCA based on lymphoid

populations in the MLN and the density of each population along the principal components

(PC). In plot (A), dot color coded by rewilded (red) and lab (purple) and in plot (B), dot color

coded by lab (green), rewilded week 6 (yellow), rewilded week 7 (blue) (**C**) Biplots of the

MLN lymphocyte populations are projected onto PC1 and PC2. (**D**) Bar plot showing the

pseudo R^2 measure of effect size on the entire distance matrix used to calculate the PCA of

immune populations.

Figure S7: An unsupervised PLS-regression multi-component network model for multi-

omic integration. A fully annotated multi-component network as shown in Figure 5A.

712
713

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

A

Serum Cytokines

PC2 20.32% expl. variation

PC1 31.43% expl. variation

● $Atg16l1^{T316A/+}$
▲ $Atg16l1^{T316A/T316A}$
■ WT
+ $Nod2^{-/-}$

● Rewilded
● Lab

TNF-α

KC

RANTES

MIP-1a   MIP-3a

IP-10

MCP-1   MIP-1b

IL-1α

B

Genotype
Environment
WeightGain
Sex
Diarrhea

Effect Size

0.00   0.01   0.02   0.03

Supplementary Figure 5

Supplementary Figure 6

Supplementary Figure 7