Alzheimer's disease neuropathology, inflammation and epigenetic age: do sex and APOE matter?

Paula Duarte-Guterman[1*], Amy M. Inkster[2,3*], Arianne Y. Albert[4], Cindy K. Barha[5], Wendy P. Robinson[2,3], Liisa A.M. Galea[1], on behalf of the Alzheimer's Disease Neuroimaging Initiative[δ]

1. Djavad Mowafaghian Centre for Brain Health and Department of Psychology, University of British Columbia, Vancouver, BC, Canada

2. Department of Medical Genetics, University of British Columbia, Vancouver, BC, Canada.

3. BC Children's Hospital Research Institute, Vancouver, BC, Canada

4. Women's Health Research Institute of British Columbia, Vancouver, BC, Canada

5. Djavad Mowafaghian Centre for Brain Health and Department of Physical Therapy, University of British Columbia, Vancouver, BC, Canada

Address all correspondence and requests for reprints to:
L. A. M. Galea, PhD
Djavad Mowafaghian Centre for Brain Health
2215 Wesbrook Mall
Vancouver, British Columbia
V6T 1Z3, Canada
E-mail: lgalea@psych.ubc.ca.

* Considered co-first authors.

[δ] Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

ABSTRACT

Alzheimer's disease (AD) disproportionately affects females and the progression from

cognitively normal to mild cognitive impairment (MCI) and to AD may be sex-specific. Using

the ADNI database, we analysed the effect of sex and APOE genotype and sex and diagnosis

(cognitively normal (CN), MCI, AD) on: AD-related endpoints (cognition, hippocampal volume,

amyloid beta, tau-pathology), immune, neuroplasticity, and stress markers, and blood epigenetic

age. More males were diagnosed with MCI but there was no sex difference in those diagnosed

with AD, suggesting the progression from CN, MCI to AD may be sex-specific. Several

biomarkers may influence sex differences in incidence and progression of MCI and AD. We

found that females had higher levels of plasma CRP and lower levels of CSF IL-8, IL-16, IgA,

and ICAM1 than males. Increasing number of APOE ε4 alleles and diagnosis also decreased

plasma CRP in both sexes. Sex, diagnosis, or APOE genotype did not affect epigenetic age

acceleration. We found, consistent with other studies, that females had higher levels of CSF tau-

pathology that was disproportionately affected by APOE genotype compared to males. Females

also had better memory (including verbal) scores than males, which may suggest a delay in the

onset of cognitive decline or diagnosis. Sex differences in immune biomarkers and hippocampal

volume (direction was dependent on method of correction) indicate that the underlying

physiology during aging may be sex-dependent.

Keywords: Sex differences; Alzheimer's disease; APOE genotype; inflammation;

neuroplasticity; epigenetic age; hippocampus; memory.

INTRODUCTION

Alzheimer's disease (AD) is a neurodegenerative disease characterized by severe cognitive decline [1]. Modifiable risk factors associated with AD include stress [2], sociocultural or lifestyle factors (e.g., education, marital status, exercise), and medical conditions [diabetes, obesity, and cardiovascular disease; ,3–5]. Non-modifiable risk factors include age, biological sex, and APOE genotype [6]. In some studies, females show greater neuropathology (brain atrophy, neurofibrillary tangles) and cognitive decline than males which depends on whether females are already cognitively impaired (mild cognitive impairment (MCI) or AD) and also on APOE genotype, a strong genetic risk factor for AD [7–14].

The hippocampus is one of the first brain areas to show atrophy with AD [15–17] and hippocampal atrophy correlates with cognitive decline [18] and AD pathology [neurofibrillary tangles; ,19]. Previous studies using the Alzheimer's Disease Neuroimaging Initiative (ADNI) indicate that females with AD have greater brain atrophy rates and cognitive decline than males with AD [9,10,13]. However, there is limited research into the role of sex and interactions with APOE ε4, a well-known genetic risk factor of AD [20], in the possible mechanisms underlying AD. In females between 65 and 75 years, one allele of APOE ε4 increases the risk of AD by 4-fold relative to males, indicating that the APOE genotype affects males and females differently [21]. Understanding why females are at a higher risk and have a higher burden of the disease is important for the development of tailored treatments based on sex and genetics.

Using the ADNI database, studies have examined the interaction of APOE genotype and biological sex on cognitive function and AD biomarkers but findings are not consistent. For example, some studies using ADNI have found a stronger association between APOEε4 and CSF tau levels in females compared to males [22] but this is not found in all ADNI studies [23,24]. Differences in type of statistical analyses, sample size, and how many groups were included in

the analyses (cognitively healthy, MCI, AD), are likely reasons for the discrepancies between

studies. In addition, some studies have focused only on one aspect of AD, for example some

studies have investigated how sex and APOE genotype affect either cognition/hippocampal

volumes [14,25–27] or AD biomarkers [23,28]. Aging and AD are multifaceted processes,

therefore one objective of our study is to employ a multivariate approach examining how sex,

APOE genotype and diagnosis affect physiological and aging markers, AD biomarkers,

hippocampal volumes and cognitive endpoints.

Chronic inflammation is a hallmark of AD, as evidenced by increased expression of

proinflammatory cytokines in the brains of AD patients without regard to sex, which can

exacerbate AD pathology [29–31]. There is however increasing evidence that there are sex

differences in immune responses with females mounting a stronger response compared to males

after an acute challenge [32,33]. In response to an endotoxin, females had higher levels of

proinflammatory plasma cytokines (TNF-α and IL-6) while males had higher levels of anti-

inflammatory plasma IL-10 [32,34]. Systemic inflammation, indicated by levels of C-reactive

protein (CRP), is higher in females than in males [35]. Higher systemic inflammation (including

plasma CRP) in midlife are also associated with higher brain amyloid levels later in life in

healthy males but not females [36], 2018) and with cognitive decline later in life in both sexes

[37]. Sex differences in the immune and inflammatory systems may differentially affect

neuroplasticity [38,39]. In addition, peripheral cortisol, the main stress hormone in humans, is

elevated in AD [40] and is associated with higher amyloid levels in the brain [41], a reduction in

hippocampal volume and negative effects on cognition in older individuals [42] that may depend

on MCI status [43]. Peripheral cortisol is also associated with elevated pro-inflammatory

cytokines [32,44]. However, it is not known how sex differences in immune (cytokines and

CRP), stress hormone (cortisol) and neuroplasticity markers may be related to sex differences in

AD or AD biomarkers. Biomarkers are highly sought after to predict disease onset and

progression and to understand the possible underlying mechanisms of AD to develop better

treatments. Therefore, we investigated potential physiological biomarkers (immune, stress,

neuroplasticity) that may explain sex differences in AD and in people at risk for AD, possessing

the APOEε4 allele, using the ADNI database.

Aging biomarkers also include epigenetic alterations, and are associated with a variety of

pathologies and adverse health conditions, including normal cognitive aging and

neurodegenerative phenotypes such as AD [45–48]. Recently, molecular biomarkers of aging

known as "epigenetic clocks" have been developed based on DNA methylation signatures

[45,46]. Epigenetic age or "DNAmAge" is a measure of the biological age of a sample (cell or

tissue), and can be calculated across a range of tissues and time points, providing an accurate

estimation of a sample's chronological age based on the presence or absence of methylation at

the 5' carbon of informative CpG dinucleotides throughout the human genome [46]. Positive

deviations of epigenetic age from chronological age (positive epigenetic age acceleration) reflect

more rapid biological aging and have been associated with numerous factors including smoking,

obesity, Parkinson's disease, Trisomy 21, and cancer [46,49–51], while negative deviations of

epigenetic age from chronological age (negative epigenetic age acceleration) have been

associated with high life-expectancy populations and memory retention [52,53]. In AD,

epigenetic age acceleration of the frontal cortex was associated with amyloid load, neuritic

plates, and cognitive decline [47]. Intra-individual DNA methylation profiles in peripheral tissue

are correlated with the epigenetic signature in the brain, likely due both to identical genetic

background affecting DNAme, and common signatures of epigenetic aging [54], thus it is

reasonable to hypothesize that epigenetic age acceleration may also be detectable in peripheral

tissues such as blood in AD participants. In healthy individuals, aging males exhibit more

positive epigenetic age acceleration than females in blood and buccal tissue, and multiple brain

regions [55]; in AD and other diseases with a sex difference, it is possible that the underlying

sex-specific pathological mechanisms may be reflected in epigenetic age acceleration measures –

for example, in AD females could potentially have more positive epigenetic age acceleration

than males.

Our aims were to first examine sex differences in cognitive ability, volume of the

hippocampus (using different correction factors), neuropathological markers of AD (amyloid

beta and tau), potential underlying physiological mechanisms (neuroplasticity, immune, stress)

and epigenetic age in peripheral tissue, and how these may be affected by APOE genotype

(number of ε4 alleles), and secondly by dementia status (cognitively healthy (CN), MCI, AD).

METHODS

ADNI database

Data used in the preparation of this article were obtained from the Alzheimer's Disease

Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). The ADNI was launched in

2003 as a public-private partnership, led by Principal Investigator Michael W. Weiner,

MD. The primary goal of ADNI has been to test whether serial magnetic resonance imaging

(MRI), positron emission tomography (PET), other biological markers, and clinical and

neuropsychological assessment can be combined to measure the progression of mild cognitive

impairment (MCI) and early Alzheimer's disease (AD). For up-to-date information, see

www.adni-info.org. Data used in this article were downloaded on or before Jan 16, 2019.

Statistical Methods: Sex and APOE genotype and sex and diagnosis

We included all participants who had the following baseline data: diagnosis in the ADNI

database, cerebrospinal fluid (CSF) levels for amyloid beta, tau and phosphorylated tau (ptau),

underwent a battery of neuropsychological tests, and had a brain MRI scan (total n = 1,460, n=

630 females, n=830 males; Table 1). Data included in our analyses were: demographics (age,

years of education, and ethnicity), baseline diagnosis (cognitively normal, CN; early MCI,

EMCI; late MCI, LMCI; or AD), number of APOE ε4 alleles (0, 1 or 2), ADNI executive

function z-scores, ADNI memory z-scores [using data from the ADNI neuropsychological

battery and validated in 56,57], hippocampal volume (mm^3), CSF amyloid beta (pg/ml), CSF tau

(pg/ml), and CSF p-tau (pg/ml). The executive function score included WAIS-R Digit Symbol

Substitution, Digit Span Backwards, Trails A and B, Category Fluency, and Clock Drawing [57].

The composite memory score included Rey Auditory Verbal Learning Test, AD Assessment

Schedule - Cognition, Mini-Mental State Examination, and Logical Memory data [56]. A small

subset of participants also had inflammatory markers measured in CSF (N = 279) and plasma (N

= 527) listed in Table 2A. Plasma and CSF samples from the ADNI study were collected at

baseline in the morning after an overnight fast. Processing, aliquoting and storage were

performed according to the ADNI Biomarker Core Laboratory Standard Operating Procedures.

Inflammatory markers were measured using a commercially available multiplex proteomic panel

(Human Discovery Multi-Analyte Profile; Luminex xMAP) developed by Rules-Based Medicine

(Austin, TX), that measures a variety of markers including cytokines, metabolic markers, and

growth factors. The protocols used to quantify plasma and CSF analytes are described in Craig-

Schapiro et al. [58] and Hu et al. [59]. We used the ADNI quality-controlled data for plasma and

CSF provided by the ADNI Consortium. Sex differences in hippocampal volume are influenced

by controlling factors such as intracranial volume [60,61]. In the present study we corrected

hippocampus volume using two different methods. The first method used the method in

Mormino et al. [62] and Jack et al. [63] using the residuals of the linear regression between

hippocampal volume and total intracranial volume. We also compared another widely used

volume correction method [27,meta analysis by 61 and references herein,64] by dividing

hippocampal volume with intracranial volume (the ratio method) to compare the different

correction methods.

We compared all available data for each study variable between the sexes using the

Wilcoxon rank sum test for continuous variables and Fisher's exact test for categorical variables.

We used general linear models to determine the relationships between (1) sex and APOE

genotype or (2) sex and dementia diagnosis as predictor variables, and cognitive ability,

corrected hippocampal volume, and biomarkers as dependent variables. Due to the sample size,

we were not able to study sex, APOE and diagnosis in one model. All models included age as a

covariate. All models initially included an interaction between sex and APOE genotype or sex

and dementia diagnosis; if this interaction was not significant, it was removed from the model to

estimate the main effects of sex and APOE genotype or diagnosis. Significance was based on the

likelihood ratio test, and all p-values for comparisons of sex and either APOE or diagnosis for all

outcomes combined were corrected for multiple testing using the Benjamini-Hochberg false

discovery rate method with the family-wise error rate set to 0.05 [65]. In total, three p-values per

dependent variable were included in each set of models (interaction term and main effects of sex

and APOE or diagnosis) resulting in 78 p-values corrected (26 dependent variables; Tables S1 to

S4) for each of the models (sex and APOE and sex and diagnosis). All regression analyses were

carried out in R v3.5.1 [66].

Statistical Methods: Epigenetic Age

We used DNAme data quantified with the Illumina Infinium HumanMethylationEPIC BeadChip array ("EPIC" array) for 1905 blood samples from 640 unique ADNI participants (n=284 females, n= 356 males) [67] with CN, MCI and AD diagnosis. DNAme IDAT files were read into R v3.5.1 [66] using the 'minfi' package, and annotated with the most recent version of the EPIC manifest, the Infinium MethylationEPIC v1.0 B4 Manifest File, (available from https://support.illumina.com/downloads.html) [68,69]. We excluded 11 low quality samples from 9 unique participants from further analyses on the basis of having a median methylated or unmethylated probe intensity <10.5 [68,69], the remaining samples were background normalized and dye-bias adjusted with normal exponential out-of-band ("noob") normalization [70]. DNAme data were converted to beta values and the biological sex of all samples was confirmed by clustering samples on all array probes mapping to the X and Y chromosomes. Beta values were calibrated to Horvath's 21,368-probe training dataset, and epigenetic age was calculated using R code modified for compatibility with the EPIC array using the 334/353 epigenetic clock probes present on the array from https://horvath.genetics.ucla.edu/ [46,71]. The missing DNAme values at these CpG sites can also be imputed based on the k-nearest neighbors method. We observed a high correlation between epigenetic age values calculated with the missing probes removed versus imputed with k=10 (R=0.99, p<2.2e-16), in agreement with previous reports [72,73]; we therefore chose to remove missing probes.

Prior to statistical analyses we removed all technical replicate samples. Epigenetic age acceleration was calculated as the residual of epigenetic age regressed on chronological age and technical batch covariates: the laboratory collection site at which blood samples were drawn, and

EPIC microarray chip and row. Intrinsic epigenetic age acceleration, a measure designed to be

independent of age-related changes in whole blood cell-type proportions, was calculated as

described in Chen et al. [74] as the residual of epigenetic age regressed on chronological age,

technical covariates of collection site, row, and chip, and the proportions of six blood cell types

(CD8T, CD4T, NK, B cells, monocytes, and granulocytes) estimated from noob-normalized

DNA methylation data with the Houseman algorithm [75]. ADNI is a longitudinal study, and

participants contributed between 1-3 blood DNAme samples within the 2-year collection period.

Because these data collected over two years fell within the median 3.6-year error of Horvath's

epigenetic clock, it therefore could not be meaningfully evaluated longitudinally. As such, for all

epigenetic age measures reported and analyzed herein, we have calculated the average epigenetic

age acceleration measure per participant from all available time points and performed all

statistical analyses on these mean values.

Statistical analyses of epigenetic age acceleration were conducted using data from the

remaining 640 participants (see Table 2B). To determine if epigenetic age acceleration or

intrinsic epigenetic age acceleration differed by sex or dementia diagnosis, we used unbalanced

two-way ANOVA designs. APOEε4 genotype and CSF biomarker (amyloid beta, tau and p-tau)

data was available from the ADNI repository for a smaller subset of participants with matched

EPIC DNAme data, (n=533, see Table 2C). Due to the small size of this cohort, we collapsed

APOE genotype into two groups: (1) samples carrying any ε4 alleles (homozygous ε4/ε4 and

heterozygous ε4/-) and (2) samples with no ε4 risk alleles (-/-). We subsequently used linear

regression to test whether presence of an APOEε4 allele, tau, dementia diagnosis, or sex were

significantly associated with epigenetic age acceleration. Amyloid beta and p-tau were not

included in this model due to collinearity between amyloid beta, tau, and p-tau CSF

concentrations (see Supplementary Table S5).

RESULTS

Demographic and biomarker information

Table 1 gives a summary of the variables for the overall data set (N=1460). Overall,

females were significantly younger and had fewer years of education than males (P<0.0001 for

both). There were more white males than white females in our sample and there were more non-

white females compared to non-white males (P<0.05). In terms of APOE genotype, there were

no sex differences in distribution of APOE genotype with 11% females and 12% of males

possessing two alleles of APOEε4. In the overall data set, the proportion of participants in each

of the diagnosis categories was significantly different for females and males (P<0.05). More

females were cognitively normal than males (26.7% compared to 20.8%, unadjusted P = 0.01)

and although there were more females with a baseline diagnosis of AD compared to males

(23.7% compared to 21.7%, unadjusted P = 0.41), this was not significant. However, there were

more males with a diagnosis of late MCI (39.5% versus 32.5%, unadjusted P=0.007) and early

MCI (18.0% versus 17.1%, unadjusted P=0.74) compared to females, although not significantly.

Because not all data were available for each subject we created a summary table for the

participants: with CSF biomarkers (Table 2A; N=279), with whole blood EPIC DNAme data

(Table 2B; N=640) and with matched EPIC data and measured CSF biomarkers (Table 2C;

N=533). Among those with measured CSF biomarkers, demographics were very similar as per

results from overall data set in Table 1 (see also legend of Table 2). For the data applicable to the

participants with available EPIC DNAme data (Table 2B) and participants with EPIC DNAme

data and CSF biomarkers (Table 2C), most of the demographics were similar to the entire data

set except the proportion of participants in each of the diagnosis categories was not significantly

different between females and males.

In the overall data set (Table 1), females had a smaller uncorrected (P<0.0001) and

corrected hippocampal volume using the regression method in [62] (P<0.0001) but larger

corrected hippocampal volume using the ratio method with intracranial volume (P<0.0001)

[27,61,64]. Females also had greater CSF amyloid beta (P=0.016), tau and p-tau (P=0.002 and

0.007 respectively), and higher memory function z-scores than males (P=0.0006; Table 1).

Biomarkers in the CSF were measured in a subset of participants (Table 2A). In this smaller

cohort, females and males had similar levels of CSF CRP, CD 40 antigen and IL-6 receptor.

However, females had lower CSF cortisol (P=0.008), interleukin-3 (P=0.001), interleukin 8

(P=0.001), interleukin-16 (P=0.004), immunoglobulin A (P<0.0001), and intercellular adhesion

molecule (P=0.0001) compared to males (Table 2A).

Sex and APOE genotype are associated with changes in memory, hippocampus volume, AD and

CSF inflammatory markers

Our first aim was to investigate whether sex and APOE genotype interact to influence

cognitive ability, volume of the hippocampus, and biomarkers of AD and inflammation. There

were significant interactions between sex and APOEε4 genotype for CSF tau, p-tau, and IL-16

(P=0.001, 0.002 and 0.03 respectively; Table 3). Tau and p-tau levels were significantly higher

in females with one or two alleles of APOEε4 compared to males (Fig 1 A and B). Although

CSF p-tau and tau levels also increase in males with APOEε4 genotype, they do not rise to the

same extent as in females. CSF IL-16 levels were significantly lower in females with no

APOEε4 alleles compared to males, whereas levels were similar between the sexes with one or

two APOEε4 alleles (Fig 1 C).

Both sex and APOE genotype were independently (main effects of sex or APOE

genotype) associated with memory z-scores and corrected hippocampal volume (Table 3).

Females had higher memory z-scores across all APOE genotypes (P<0.0001; Fig 1 D).

Curiously, females had smaller or larger hippocampi than males depending on the correction

method used: females had smaller corrected hippocampal volume based on a regression method

[62] (P<0.0001; Fig 1 E) but larger hippocampal/intracranial volume compared to males

(P<0.0001; Supplementary Table S1). These data suggest it is important to consider different

correction factors when studying sex differences in volumes of brain structures. Lower memory

z-scores were associated with increasing number of APOEε4 alleles in both sexes (P<0.0001).

Similarly, corrected hippocampus volume (independent of method of correction) was

significantly lower with increasing number of APOEε4 alleles in both sexes (Ps<0.0001).

Increasing APOEε4 alleles was also associated with lower executive function z-scores

(P<0.0001), lower CSF amyloid beta (P<0.0001), and lower CSF CRP levels (P=0.009) (Table 3;

Fig 1 F-H), however there was no additional association of these variables with sex. Finally,

results were similar for biomarkers in plasma (Supplementary Table S3) with the exception of a

significant relationship between plasma CRP and sex. Males had lower plasma levels of CRP

compared to females (adjusted P=0.003).

Sex and diagnosis are associated with changes in memory, hippocampus volume, AD and CSF

inflammatory markers

We next tested whether sex and dementia status (CN, MCI, and AD) influenced cognitive

ability, corrected hippocampal volume, and CSF biomarkers of AD and inflammation. There

were no significant interactions between sex and diagnosis for any of the tested variables

(memory, executive function, corrected hippocampal volume, CSF tau, p-tau, amyloid beta, and

CSF and plasma inflammatory markers). However, overall both sex and diagnosis were

independently associated with memory z-scores, corrected hippocampal volume and CSF tau and

p-tau (Table 4). Females had higher memory z-scores (P<0.0001), smaller corrected

hippocampus volume (P<0.0001) but larger hippocampus/intracranial volume ratio (P<0.0001;

Supplementary Table S2), and higher tau and p-tau compared to males (Ps<0.0001), irrespective

of diagnosis (Table 4). As expected, increasing severity of diagnosis was associated with lower

memory and executive function z-scores (Ps<0.0001), smaller corrected hippocampus volume

(Ps<0.0001; independent of correction method), lower CSF amyloid beta (P<0.0001), and higher

CSF tau and p-tau irrespective of sex (Ps<0.0001; Fig 2 A-F).

We found that although females had lower CSF levels of interleukin 8 (IL-8; P=0.03),

immunoglobulin A (IgA; P<0.0001), interleukin 16 (IL-16; P=0.07), and intercellular adhesion

molecule 1 (ICAM1; P=0.008), controlling for age, compared to males, there was no association

between these variables and diagnosis (Table 4; Fig 2 G-J). The results for biomarkers and

inflammatory markers were different in plasma (Supplementary Table S4) with females having

higher levels of plasma ICAM1 compared to males (P=0.05) and no sex differences in plasma

IL-8, IgA and IL-6 levels. We found a significant relationship between plasma CRP and sex

(adjusted P=0.01), immunoglobulin E (IgE) and sex (P<0.0001), IL-18 and sex (P=0.001), and

also between plasma cortisol and baseline diagnosis (adjusted P=0.005; Fig 2 K and L). Males

had lower levels of plasma CRP compared to females and late MCI and AD resulted in lower

plasma CRP levels compared to CN (adjusted P=0.04). Plasma cortisol was lower in late MCI

compared to CN but higher in AD compared to CN. In summary, although we detected

associations between sex and diagnosis and various parameters, we did not find evidence for a

clear sex and diagnosis interaction.

Epigenetic age, sex, dementia diagnosis, and AD biomarkers

We investigated the hypothesis that sex and dementia diagnosis affect epigenetic age

acceleration in blood samples of ADNI participants (see Table 5). Epigenetic age acceleration

was not associated with sex, dementia diagnosis (CN, EMCI, LMCI, and AD), or the interaction

of sex and diagnosis after multiple test correction (Fig 3). Intrinsic epigenetic age acceleration

was also not significantly associated with participant sex, diagnosis, or their interaction term.

To assess the effect of sex and more broadly defined dementia-associated cognitive

impairment on epigenetic age acceleration, we compared epigenetic age acceleration between

participants with any form of clinically ascertained cognitive impairment (AD + LMCI + EMCI,

n=423, proportion female 41%) and those without (CN, n=217, proportion female 50%). By two-

way unbalanced ANOVA neither sex, dementia status, nor their interaction were significantly

associated with epigenetic age acceleration after correction for multiple comparisons.

Matched biochemical data including APOEε4 genotype and CSF concentrations of

amyloid beta, tau, and p-tau was available for a subset of participants with EPIC DNAme data

(n=533). Based on the hypothesis that epigenetic age acceleration may be more strongly

associated with concentrations of pathologically relevant compounds than with diagnosis, we

assessed the impact of sex, APOEε4 genotype, CSF amyloid beta concentration, CSF tau and p-

tau concentration on epigenetic age acceleration and intrinsic epigenetic age acceleration with

linear regression. None of these variables were significantly associated with epigenetic age

acceleration (Table 6; results for intrinsic epigenetic age acceleration not shown).

In addition to dementia diagnosis for all participants, we also had access to two

composite scores designed by ADNI collaborators to reflect executive function and memory;

these z-scores have been demonstrated to be independently predictive of the transition from MCI

to a formal diagnosis of AD [56,57]. By a two-way unbalanced ANOVA models investigating

the effect of sex and memory z-scores on epigenetic age acceleration, neither sex (P=0.248),

memory score (P=0.486), nor their interaction (P=0.227) were associated with epigenetic age

acceleration. In a similar model, neither sex (P=0.260), executive function (P=0.105), or the

interaction term of sex and executive function (P=0.153) were associated with epigenetic age

acceleration.

DISCUSSION

In the present study, we found that tau-related pathology in the CSF was

disproportionately elevated by APOEε4 genotype in females compared to males. However,

diagnosis and APOE genotype were independently associated with reduced memory scores,

corrected hippocampal volume and reduced CSF amyloid beta which was similar in males and

females. Furthermore, there were main effects of sex as females had lower CSF and plasma

cytokines (IL-8, IL-16, IL-18) and CSF and plasma immunoglobulins (IgA, IgE, respectively)

but higher plasma CRP and tau-related pathology compared to males, regardless of diagnosis and

APOE genotype. Interestingly, depending on the correction method, females had either smaller

or larger hippocampus volume compared to males. Females had better memory scores which

may contribute to their delayed diagnosis [76]. Finally, we found no differences in epigenetic age

acceleration by dementia diagnosis or sex in this cohort of samples with available whole blood

EPIC DNAme data. In this ADNI cohort, significantly more males presented with a diagnosis of

MCI while there was no sex difference in the number of participants diagnosed with AD,

suggestive of sex differences in transitions to MCI and to AD. It is interesting to note that

population-based studies find males have more MCI whereas females have more AD [77–79],

although not all studies agree in the higher prevalence or incidence of AD in females [reviewed

in 4,80]. Previous work has demonstrated sex differences in rates of AD and symptoms of AD

[4,77,reviewed in 81], and the current study also suggests that biomarkers of AD may be

different between males and females between genotypes, and this should be considered in future

studies and researchers should be cautioned to use sex as a biological variable in all analyses.

Females show greater tau neuropathology disproportionately affected by APOE genotype

In the present study, we found that females have significantly higher baseline tau and p-

tau levels in CSF than males and these are indicative of the formation of neurofibrillary tangles

and AD pathology [82,83]. This is in agreement with a recent ADNI study [64, but see an earlier

ADNI study 9] and with animal models [84]. Intriguingly, we also found that levels of tau and p-

tau were disproportionately elevated with APOEε4 allele expression in females compared to

males and this is in agreement with a previous ADNI study that included cognitively healthy

individuals only [22]. A meta-analysis that included data from ADNI also found a stronger

association between APOEε4 and tau levels in females compared to males [85] but interestingly

this association is not present in all ADNI studies which may depend on sample size, type of

statistical analysis, and the diagnosis groups included in the analyses [23,24]. Previous studies

indicate that females with the APOEε4 allele are at a greater risk for developing AD than are

males with this allele [86], and sex differences in tau and p-tau may be one underlying

mechanism by which this occurs but further replication with additional populations is needed. In

females (65-75 years of age) one allele of ε4 increases the risk of AD by 4-fold relative to males,

indicating that genotype may affect females differently [21]. Levels of CSF tau are hypothesized

to increase after CSF amyloid beta declines and amyloid beta aggregates and deposits in the

brain [82]. However, in this study although we found sex differences in CSF tau and p-tau levels,

no significant differences were seen in CSF amyloid beta after controlling for age (see below)

indicating that the pathway may be different in females compared to males or that the timeline of

tau and amyloid beta deposition may not be consistent.

Females are more likely to be diagnosed with AD in Europe and Asia, although this sex

difference may depend in part on geographic location as the sex difference is not always

observed in studies from the United States [4,77,reviewed by 81]. Nevertheless, the

disproportionate effect of APOEε4 genotype on tau-related pathology in females supports the

idea that females have a higher burden of the disease dependent on APOE genotype. On the

other hand, males are more likely to be diagnosed with MCI compared to females [18,77,80],

while females progress faster from MCI to AD [13]. Sex differences in tau-related pathology

found in the current study may be the underlying mechanism for this accelerated transition.

Sex differences in hippocampal volume depend on correction method. Females have better

memory scores than males that may have been driven by verbal memory

In the present study, we found that increasing APOEε4 alleles and AD diagnosis was

associated with reduced corrected hippocampal volume, memory and executive function scores

consistent with past literature [15,16,18,87–92]. Surprisingly, although females have higher

levels of tau and p-tau, they presented with better memory and executive function scores than

males, regardless of diagnosis and APOE genotype. Depending on the correction method,

females had larger or smaller hippocampal volume compared to males. Females had smaller

corrected hippocampal volume using the residuals of the linear regression between hippocampal

volume and total intracranial volume [method by 62,63] but larger ratio of

hippocampal/intracranial volume compared to males [method used in 27,61,64]. These results

have important implications for future studies trying to understand sex differences in AD (and

other neurodegenerative diseases) as they lead to very different conclusions regarding

hippocampal plasticity. We suggest future research should report uncorrected as well as

corrected volumes using multiple methods. Previous studies have suggested that there are sex

differences in hippocampal volume, favoring males, but the sex differences depend on whether

hippocampal volume is corrected for by intracranial volume or total brain volume and the

method of correction [61]. In a number of studies, including the present study, males have a

larger hippocampus without correcting for intracranial volume [27,61,64,93–96]. However after

correcting for intracranial volume using the regression model, studies have found females have

larger [94], smaller (present study) or no difference [61,93] in corrected hippocampal volume

compared to males. Using the ratio method, Sohn et al. [27], Sundermann et al. [64], Murphy et

al. [95] and the current study found that females had larger hippocampal volumes compared to

males. In their meta-analysis, Tan et al. [61] conclude that sex differences in hippocampal

volume are modest and highly depend on correction method and this is supported by our

findings. Regardless of hippocampal volume, volume loss is greater in aging females [7,26,95]

and in females with one or two APOEε4 alleles [97]. Although in the present study we did not

examine longitudinal data, we found that increasing APOEε4 alleles reduced corrected

hippocampal volume similarly in males and females. In contrast, when CN, MCI and AD

individuals were analysed separately in the ADNI database, APOEε4 was associated with a

smaller corrected hippocampal volume in CN males only, controlling for age and education [64].

In addition, also using the ADNI database, Koran et al. [26] found that females with low CSF

amyloid beta had more hippocampal atrophy and faster decline in memory and executive

function than males and this sex difference was more pronounced in APOEε4 carriers. Therefore,

sex and APOE genotype can interact to affect corrected hippocampal volume reduction with age

in certain subgroups and across time (e.g., in CN or individuals with low CSF amyloid beta).

Differences in results between studies are likely due to differences in statistical analyses (e.g.,

analysing diagnosis groups separately, partitioning the data based on amyloid beta levels, and

differences in covariates included) and/or whether longitudinal data analyses are included.

We found females had better composite memory scores (but not executive function

scores) than males, regardless of diagnosis and APOE genotype. Previous studies have found

that females have better verbal memory in cognitively normal individuals [94], and in MCI and

AD ADNI cohorts compared to males [64,98]. Here we used the ADNI memory score developed

by Crane et al. [56] to detect abnormal memory including language, attention, and logical

memory so it is possible that verbal memory may be driving the sex difference favouring females

in the present study. In contrast, Buckley et al.[99] found no sex differences using a composite

cognitive score that includes memory and executive function (Preclinical Alzheimer's Cognitive

Composite score with semantic processing, PACC5) using ADNI and two other cohorts. In this

study using the current ADNI cohort, males were slightly more educated than females, and

although we did not use education as a covariate, one would expect education levels would have

positive effects on memory, suggesting that education is not a factor for the observed sex

difference in memory. Altogether, we found that in females tau pathology was increased but

memory scores, which included verbal memory, were higher compared to males suggesting

females have a reserve against brain damage that delays either the onset of cognitive decline

[100] or diagnosis [76]. However, once cognitive decline begins, females show higher rates of

declines compared to males [9,10,99] perhaps because the underlying pathology is elevated in

females.

AD affects amyloid beta similarly in both sexes

We found that AD diagnosis was associated with lower CSF amyloid beta, as expected,

and this was irrespective of sex, which indicates greater amyloid deposition with AD [83]. These

findings are consistent with data from studies in cognitively healthy individuals [94,99]. Other

studies have found using PET that males have higher amyloid beta levels or lower amyloid beta

burden compared to females dependent on APOE genotype [64] or in cognitively normal adults

in the anterior cingulate [93]. In this study, we used CSF amyloid beta data which detect

abnormal amyloid deposition earlier than amyloid beta by PET [reviewed in 82]. Thus, taken

together, sex differences in amyloid beta may be detected in specific brain regions and later in

the disease, although more research is needed investigating sex differences in AD biomarkers.

Females have higher CRP levels but lower cytokine and immunoglobulin levels compared to

males

In this study, we investigated whether sex interacted with APOE genotype or dementia

diagnosis to influence inflammatory, neurotrophic and neuroplasticity markers. We found that

plasma CRP, a widely used inflammatory and cardiovascular marker [101,102], was affected by

sex and APOE genotype. Females, regardless of diagnosis or APOE genotype, had significantly

higher plasma CRP relative to males, consistent with findings in healthy individuals [35]. Higher

levels of peripheral CRP may suggest higher systemic inflammation in females, which is

associated with an increased risk in all-cause dementia [103]. In contrast, APOEɛ4 genotype and

diagnosis decreased circulating CRP levels. The effect of APOEɛ4 genotype is consistent with

previous research in large population studies [104,105]. Recent meta-analyses, without regard to

sex, did not find differences in peripheral levels of CRP in AD compared to control patients

[106,107]. However, in patients with mild and moderate dementia only, CRP levels were lower

compared to the healthy control group [106]. In healthy individuals, higher levels of plasma CRP

in midlife are associated with a higher amyloid burden later in life in males but not females [36].

However, despite this finding, higher systemic inflammation (including CRP) was associated

with greater cognitive decline later in life in both sexes in healthy individuals [37]. It is possible

that sex differences in inflammation earlier in life have long term effects on the transition to MCI

and/or AD. To our knowledge, no other study has examined sex differences in CRP in relation to

AD.

We also found that CSF IL-16 was affected by sex and APOE genotype. CSF IL-16

levels were lower in females with no APOEɛ4 alleles compared to males, but with increasing

number of ɛ4 alleles, no sex differences were detected. IL-16 has been implicated in AD [108]

and IL-16 levels decrease with disease severity [analysis without regard to sex; ,109]. In this

ADNI cohort, IL-16 levels were not affected by diagnosis but our results suggest that APOE

genotype can modulate levels in a sex-dependent way. We also found biomarkers that were

affected by sex but not diagnosis or APOE genotype for example, females had lower CSF levels

of ICAM1 compared to males, but there was no influence of APOE genotype or diagnosis.

Consistent with our findings, ICAM1 serum levels were lower in healthy females compared to

males [110]. ICAM1 is a type of adhesion molecule associated with microvascular endothelial

activation [111] and plasma ICAM1 levels [but not CSF levels; ,112] were higher in patients

with AD [112–114]. However, it is intriguing that females have lower CSF levels of cytokines

(IL-8, IL-16, IL-18), and immunoglobulins (IgE and IgA) but higher tau pathology compared to

males. Neuroinflammation is associated with AD but it can have both beneficial and detrimental

roles [115]. Increased expression of pro-inflammatory cytokines contributes to neuronal loss,

while anti-inflammatory effects contribute to amyloid beta clearance [116]. In AD mouse

models, some pro-inflammatory mechanisms reduced plaque pathology, while anti-inflammatory

cytokines increased amyloid beta deposition [117–122]. It has been suggested that there are

beneficial pro-inflammatory mechanisms and detrimental anti-inflammatory mechanisms in AD

[116]. It is possible that males and females have varying levels of beneficial vs detrimental

immune responses which can affect how the disease progresses in each of the sexes but it is also

important to remember that CSF levels may not match levels in different regions of the brain.

Sex, AD and biochemical markers do not affect blood epigenetic age acceleration

We did not observe an association between either sex or diagnosis and epigenetic age

acceleration or intrinsic epigenetic age acceleration. To our knowledge, no other study has

similarly probed epigenetic age acceleration in peripheral tissue in the presence of AD, or

whether epigenetic age acceleration in AD is associated with sex.

This study was partially undertaken to investigate whether epigenetic age acceleration

that has been associated with the AD brain is reflected in peripheral tissues. Levine et al. [47]

have previously demonstrated increased epigenetic age acceleration in AD, however Levine's

study was conducted on post-mortem prefrontal cortex tissue, and did not explicitly investigate

the role of sex in epigenetic age acceleration [47]. While brain-blood methylation profiles are

reasonably correlated (r=0.86) [54], DNA methylation profiles of peripheral tissues are imperfect

representatives of the brain, and do not recapitulate all epigenetic alterations with high fidelity.

Thus, our findings do not contradict the finding of increased epigenetic age acceleration in the

presence of AD in the prefrontal cortex, but suggest that accelerated epigenetic aging in AD is

not a pan-tissue phenomenon. Our finding of a lack of significant association between AD,

biological sex, and epigenetic age acceleration in whole blood DNA methylation profiles could

suggest a tissue-specific dysregulation of an epigenetic maintenance system, in which the brain

epigenome is most strongly affected by AD [47]. The phenotype of patients affected by AD and

global gene expression patterns of the APOE protein, with high expression in brain, and low

expression in whole blood [123] further support this hypothesis.

Intriguingly, epigenetic age was observed to be lower on average than chronological age.

Horvath's epigenetic clock was trained on DNAme data from older versions of the Illumina

DNAme arrays with more limited genomic coverage; 19 of the CpG probes required to calculate

epigenetic age via this method do not exist on the EPIC array. Two previous studies investigated

the application of Horvath's epigenetic clock to EPIC data with conflicting results [73,124], the

largest issue being chronic underestimation of epigenetic age due to the positive linear regression

coefficients associated with the missing probes [124]. Both imputing and removing the missing

probes from the array resulted in a chronic underprediction of epigenetic age with Horvath's

clock, suggesting that this is likely an artefact of the array platform and probe-set rather than the

method chosen to deal with missing values, although it is possible that an adjustment factor

could be devised to more accurately apply Horvath's clock to EPIC data. In future explorations

of epigenetic age with EPIC DNAme array data this should be considered, as there are other

epigenetic age predictors available that have been trained on EPIC data such as the PhenoAge

and GrimAge clocks, although these tools have limitations as well; for example, both PhenoAge

and GrimAge were trained only on blood DNAme data, as compared to the original pan-tissue

epigenetic clock, and therefore may have limited applicability and relevance in other tissues

[125,126].

Limitations

The ADNI cohort is not ethnically or socioeconomically diverse, being mostly composed

of white (only 12 individuals were not-white) and highly educated individuals (average 15.69

years of education). As incidence, prevalence, and age of onset of AD varies by ethnicity [127–

129] and education [130], our conclusions may not apply to more ethnically and socially diverse

populations. In addition to sex, it is possible the underlying mechanisms of AD are different

depending on ethnicity. APOEε4 genotype is a well-known AD risk factor and in this cohort the

proportion of participants in the different APOEε4 allele groups is correlated with diagnosis.

Future research with a larger cohort is required to test how sex, APOE genotype and diagnosis

interact together in one model. Small sample size is also a limitation of the epigenetic analyses

presented. Even in the larger 640-participant cohort, only 37 participants (5.78%) had an AD

diagnosis, so statistical analyses were underpowered to detect subtle differences by diagnosis

group. Additionally other pathologies in these participants, such as cancer, cardiovascular

disease, smoking status, or obesity may have influenced AD neuropathology, biomarkers and

epigenomes and limited our interpretations.

CONCLUSION

585   As expected, more females presented with a diagnosis of AD whereas more males

presented with MCI diagnosis compared to the opposite sex. AD biomarkers (CSF tau and p-tau

but not amyloid beta) were disproportionately affected by APOE genotype in females compared

to males supporting the idea that females share a higher burden of the disease. Interestingly,

although females in this cohort had elevated AD biomarkers, they also had higher memory

function scores compared to males, regardless of APOE genotype and dementia diagnosis.

Therefore, it is possible that females may have a reserve that protects the brain from damage to

delay cognitive decline or delay diagnosis. Finally, we found that females had lower cytokine

and immunoglobulin levels but higher CRP levels compared to males. Together our work

suggests that that the underlying physiology of aging and AD may be sex-specific.

ACKNOWLEDGMENTS

597   Data collection and sharing for this project was funded by the Alzheimer's Disease

Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD

ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the

National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering,

and through generous contributions from the following: AbbVie, Alzheimer's Association;

Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-

Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli

Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company

Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy

Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development

LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx

Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal

Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian

Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private

sector contributions are facilitated by the Foundation for the National Institutes of Health

(www.fnih.org). The grantee organization is the Northern California Institute for Research and

Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the

University of Southern California. ADNI data are disseminated by the Laboratory for Neuro

Imaging at the University of Southern California. Funding for this study was provided by a

Canadian Institutes of Health Research (CHIR) grant to LAMG (PJT-148662) and a CIHR sex

catalyst grant to WPR (Reference No. 158613). WPR receives salary support through an

investigatorship award from the BC Children's Hospital Research Institute. PDG is funded by

the Alzheimer's Association of the USA and Brain Canada with the financial support of Health

Canada through the Brain Canada Research Fund (AARF-17-529705). The views expressed

herein do not necessarily represent the views of the Minister of Health or the Government of

Canada. AMI is funded by a University of British Columbia Four Year Doctoral Fellowship and

the CIHR Frederick Banting and Charles Best Masters Research Award. We thank Elizabeth

Perez for help with data organization.

CONFLICT OF INTEREST

The authors have no conflict of interest to report.

FIGURE CAPTIONS

Figure 1. A. CSF tau (pg/ml), B. CSF p-tau (pg/ml), C. CSF IL-16 (pg/ml), D. ADNI memory z-

scores, E. corrected hippocampal volume (hippocampal volume/intracranial volume), F. ADNI

executive function z-scores, G. CSF amyloid beta (pg/ml), and H. CSF C-reactive protein (CRP;

µg/ml) in ADNI participants by sex and number of APOEε4 alleles (0, 1, 2 alelles).

Figure 2. A. ADNI memory z-scores, B. corrected hippocampal volume (hippocampal volume/

intracranial volume), C. CSF tau (pg/ml), D. CSF p-tau (pg/ml), E. CSF IL-16 (pg/ml), F. CSF

IL-8 (pg/ml), G. CSF IgA (mg/ml), H. CSF Intercellular adhesion molecule (ICAM1; ng/ml), I.

ADNI executive function z-scores, J. CSF amyloid beta (pg/ml), K. plasma C-reactive protein

(CRP; µg/ml), and L. plasma cortisol (ng/ml) in ADNI participants by sex and diagnosis (CN,

EMCI, LMCI, AD). CN, cognitively normal; EMCI, early mild cognitive impairment; LMCI,

late mild cognitive impairment; AD, Alzheimer's disease.

Figure 3. Universal epigenetic age acceleration does not differ statistically significantly by

participant sex or diagnosis (CN, EMCI, LMCI, AD) in this ADNI cohort. CN, cognitively

normal; EMCI, early mild cognitive impairment; LMCI, late mild cognitive impairment; AD,

Alzheimer's disease.

REFERENCES

[1] Alzheimer's Association (2017) 2017 Alzheimer's disease facts and figures. *Alzheimer's & Dementia* **13**, 325–373.

[2] Caruso A, Nicoletti F, Mango D, Saidi A, Orlando R, Scaccianoce S (2018) Stress as risk factor for Alzheimer's disease. *Pharmacological Research* **132**, 130–134.

[3] Baumgart M, Snyder HM, Carrillo MC, Fazio S, Kim H, Johns H (2015) Summary of the evidence on modifiable risk factors for cognitive decline and dementia: A population-based perspective. *Alzheimers Dement* **11**, 718–726.

[4] Nebel RA, Aggarwal NT, Barnes LL, Gallagher A, Goldstein JM, Kantarci K, Mallampalli MP, Mormino EC, Scott L, Yu WH, Maki PM, Mielke MM (2018) Understanding the impact of sex and gender in Alzheimer's disease: A call to action. *Alzheimer's & Dementia* **14**, 1171–1183.

[5] Xu W, Tan L, Wang H-F, Jiang T, Tan M-S, Tan L, Zhao Q-F, Li J-Q, Wang J, Yu J-T (2015) Meta-analysis of modifiable risk factors for Alzheimer's disease. *J Neurol Neurosurg Psychiatry* **86**, 1299–1306.

[6] Riedel BC, Thompson PM, Brinton RD (2016) Age, APOE and sex: Triad of risk of Alzheimer's disease. *The Journal of Steroid Biochemistry and Molecular Biology* **160**, 134–147.

[7] Ardekani BA, Convit A, Bachman AH (2016) Analysis of the MIRIAD Data Shows Sex Differences in Hippocampal Atrophy Progression. *J. Alzheimers Dis.* **50**, 847–857.

[8] Barnes LL, Wilson RS, Bienias JL, Schneider JA, Evans DA, Bennett DA (2005) Sex differences in the clinical manifestations of Alzheimer disease pathology. *Arch. Gen. Psychiatry* **62**, 685–691.

[9] Holland D, Desikan RS, Dale AM, McEvoy LK, Alzheimer's Disease Neuroimaging Initiative (2013) Higher rates of decline for women and apolipoprotein E epsilon4 carriers. *AJNR Am J Neuroradiol* **34**, 2287–2293.

[10] Hua X, Hibar DP, Lee S, Toga AW, Jack CR, Weiner MW, Thompson PM (2010) Sex and age differences in atrophic rates: an ADNI study with n=1368 MRI scans. *Neurobiology of Aging* **31**, 1463–1480.

[11] Irvine K, Laws KR, Gale TM, Kondel TK (2012) Greater cognitive deterioration in women than men with Alzheimer's disease: a meta analysis. *J Clin Exp Neuropsychol* **34**, 989–998.

[12] Koran MEI, Hohman TJ (2017) Sex differences in the association between AD biomarkers and cognitive decline. *Brain Imaging and Behavior* **11**, 205–213.

[13] Lin KA, Choudhury KR, Rathakrishnan BG, Marks DM, Petrella JR, Doraiswamy PM (2015) Marked gender differences in progression of mild cognitive impairment over 8 years. *Alzheimer's & Dementia: Translational Research & Clinical Interventions* **1**, 103–110.

[14] Wang X, Zhou W, Ye T, Lin X, Zhang J, Initiative for ADN (2019) Sex Difference in the Association of APOE4 with Memory Decline in Mild Cognitive Impairment. *Journal of Alzheimer's Disease* **69**, 1161–1169.

[15] Apostolova LG, Dutton RA, Dinov ID, Hayashi KM, Toga AW, Cummings JL, Thompson PM (2006) Conversion of mild cognitive impairment to Alzheimer disease predicted by hippocampal atrophy maps. *Arch. Neurol.* **63**, 693–699.

692	[16]	Jack CR, Petersen RC, Xu Y, O'Brien PC, Smith GE, Ivnik RJ, Boeve BF, Tangalos EG,
693		Kokmen E (2000) Rates of hippocampal atrophy correlate with change in clinical status
694		in aging and AD. *Neurology* **55**, 484–489.
695	[17]	Kidron D, Black SE, Stanchev P, Buck B, Szalai JP, Parker J, Szekely C, Bronskill MJ
696		(1997) Quantitative MR volumetry in Alzheimer's disease: Topographic markers and the
697		effects of sex and education. *Neurology* **49**, 1504–1512.
698	[18]	Petersen RC, Jack CR, Xu YC, Waring SC, O'Brien PC, Smith GE, Ivnik RJ, Tangalos
699		EG, Boeve BF, Kokmen E (2000) Memory and MRI-based hippocampal volumes in
700		aging and AD. *Neurology* **54**, 581–587.
701	[19]	Jack CR, Dickson DW, Parisi JE, Xu YC, Cha RH, O'Brien PC, Edland SD, Smith GE,
702		Boeve BF, Tangalos EG, Kokmen E, Petersen RC (2002) Antemortem MRI Findings
703		Correlate with Hippocampal Neuropathology in Typical Aging and Dementia. *Neurology*
704		**58**, 750–757.
705	[20]	Ossenkoppele R, Jansen WJ, Rabinovici GD, Knol DL, van der Flier WM, van Berckel
706		BNM, Scheltens P, Visser PJ, Verfaillie SCJ, Zwan MD, Adriaanse SM, Lammertsma
707		AA, Barkhof F, Jagust WJ, Miller BL, Rosen HJ, Landau SM, Villemagne VL, Rowe
708		CC, Lee DY, Na DL, Seo SW, Sarazin M, Roe CM, Sabri O, Barthel H, Koglin N,
709		Hodges J, Leyton CE, Vandenberghe R, van Laere K, Drzezga A, Forster S, Grimmer T,
710		Sánchez-Juan P, Carril JM, Mok V, Camus V, Klunk WE, Cohen AD, Meyer PT,
711		Hellwig S, Newberg A, Frederiksen KS, Fleisher AS, Mintun MA, Wolk DA, Nordberg
712		A, Rinne JO, Chételat G, Lleo A, Blesa R, Fortea J, Madsen K, Rodrigue KM, Brooks DJ
713		(2015) Prevalence of Amyloid PET Positivity in Dementia Syndromes: A Meta-analysis.
714		*JAMA* **313**, 1939.
715	[21]	Neu SC, Pa J, Kukull W, Beekly D, Kuzma A, Gangadharan P, Wang L-S, Romero K,
716		Arneric SP, Redolfi A, Orlandi D, Frisoni GB, Au R, Devine S, Auerbach S, Espinosa A,
717		Boada M, Ruiz A, Johnson SC, Koscik R, Wang J-J, Hsu W-C, Chen Y-L, Toga AW
718		(2017) Apolipoprotein E Genotype and Sex Risk Factors for Alzheimer Disease: A Meta-
719		analysis. *JAMA Neurology* **74**, 1178.
720	[22]	Damoiseaux JS, Seeley WW, Zhou J, Shirer WR, Coppola G, Karydas A, Rosen HJ,
721		Miller BL, Kramer JH, Greicius MD, Alzheimer's Disease Neuroimaging Initiative
722		(2012) Gender modulates the APOE ε4 effect in healthy older adults: convergent
723		evidence from functional brain connectivity and spinal fluid tau levels. *J. Neurosci.* **32**,
724		8254–8262.
725	[23]	Buckley RF, Mormino EC, Chhatwal J, Schultz AP, Rabin JS, Rentz DM, Acar D,
726		Properzi MJ, Dumurgier J, Jacobs H, Gomez-Isla T, Johnson KA, Sperling RA,
727		Hanseeuw BJ (2019) Associations between baseline amyloid, sex, and APOE on
728		subsequent tau accumulation in cerebrospinal fluid. *Neurobiology of Aging* **78**, 178–185.
729	[24]	Sampedro F, Vilaplana E, de Leon MJ, Alcolea D, Pegueroles J, Montal V, Carmona-
730		Iragui M, Sala I, Sánchez-Saudinos M-B, Antón-Aguirre S, Morenas-Rodríguez E,
731		Camacho V, Falcón C, Pavía J, Ros D, Clarimón J, Blesa R, Lleó A, Fortea J,
732		Alzheimer's Disease Neuroimaging Initiative (2015) APOE-by-sex interactions on brain
733		structure and metabolism in healthy elderly controls. *Oncotarget* **6**, 26663–26674.
734	[25]	Caldwell JZK, Berg J-L, Shan G, Cummings JL, Banks SJ (2018) Sex Moderates the
735		Impact of Diagnosis and Amyloid PET Positivity on Hippocampal Subfield Volume. *J*
736		*Alzheimers Dis* **64**, 79–89.

737   [26]   Koran MEI, Wagener M, Hohman TJ, Alzheimer's Neuroimaging Initiative (2017) Sex
738   differences in the association between AD biomarkers and cognitive decline. *Brain*
739   *Imaging Behav* **11**, 205–213.
740   [27]   Sohn D, Shpanskaya K, Lucas JE, Petrella JR, Saykin AJ, Tanzi RE, Samatova NF,
741   Doraiswamy PM (2018) Sex Differences in Cognitive Decline in Subjects with High
742   Likelihood of Mild Cognitive Impairment due to Alzheimer's disease. *Scientific Reports*
743   **8**, 7490.
744   [28]   Liu M, Paranjpe MD, Zhou X, Duy PQ, Goyal MS, Benzinger TLS, Lu J, Wang R, Zhou
745   Y (2019) Sex modulates the ApoE ε4 effect on brain tau deposition measured by 18F-
746   AV-1451 PET in individuals with mild cognitive impairment. *Theranostics* **9**, 4959–
747   4970.
748   [29]   Heppner FL, Ransohoff RM, Becher B (2015) Immune attack: the role of inflammation
749   in Alzheimer disease. *Nature Reviews Neuroscience* **16**, 358–372.
750   [30]   Kinney JW, Bemiller SM, Murtishaw AS, Leisgang AM, Salazar AM, Lamb BT (2018)
751   Inflammation as a central mechanism in Alzheimer's disease. *Alzheimer's & Dementia:*
752   *Translational Research & Clinical Interventions* **4**, 575–590.
753   [31]   Swardfager W, Lanctôt K, Rothenburg L, Wong A, Cappell J, Herrmann N (2010) A
754   Meta-Analysis of Cytokines in Alzheimer's Disease. *Biological Psychiatry* **68**, 930–941.
755   [32]   Engler H, Benson S, Wegner A, Spreitzer I, Schedlowski M, Elsenbruch S (2016) Men
756   and women differ in inflammatory and neuroendocrine responses to endotoxin but not in
757   the severity of sickness symptoms. *Brain, Behavior, and Immunity* **52**, 18–26.
758   [33]   Klein SL, Flanagan KL (2016) Sex differences in immune responses. *Nature Reviews*
759   *Immunology* **16**, 626–638.
760   [34]   Wegner A, Benson S, Rebernik L, Spreitzer I, Jäger M, Schedlowski M, Elsenbruch S,
761   Engler H (2017) Sex differences in the pro-inflammatory cytokine response to endotoxin
762   unfold in vivo but not ex vivo in healthy humans. *Innate Immun* **23**, 432–439.
763   [35]   Khera A, McGuire DK, Murphy SA, Stanek HG, Das SR, Vongpatanasin W, Wians FH,
764   Grundy SM, de Lemos JA (2005) Race and gender differences in C-reactive protein
765   levels. *J. Am. Coll. Cardiol.* **46**, 464–469.
766   [36]   Walker KA, Windham BG, Brown IV CH, Knopman DS, Jack Jr CR, Mosley Jr TH,
767   Selvin E, Wong DF, Hughes TM, Zhou Y, Gross AL, Gottesman RF (2018) The
768   Association of Mid- and Late-Life Systemic Inflammation with Brain Amyloid
769   Deposition: The ARIC-PET Study. *Journal of Alzheimer's Disease* **66**, 1041–1052.
770   [37]   Walker KA, Gottesman RF, Wu A, Knopman DS, Gross AL, Mosley TH, Selvin E,
771   Windham BG (2019) Systemic inflammation during midlife and cognitive change over
772   20 years. *Neurology* **92**, e1256–e1267.
773   [38]   Dantzer R (2018) Neuroimmune Interactions: From the Brain to the Immune System and
774   Vice Versa. *Physiol. Rev.* **98**, 477–504.
775   [39]   de Miranda AS, Zhang C-J, Katsumoto A, Teixeira AL (2017) Hippocampal adult
776   neurogenesis: Does the immune system matter? *Journal of the Neurological Sciences*
777   **372**, 482–495.
778   [40]   Doecke JD, Laws SM, Faux NG, Wilson W, Burnham SC, Lam C-P, Mondal A, Bedo J,
779   Bush AI, Brown B, Ruyck KD, Ellis KA, Fowler C, Gupta VB, Head R, Macaulay SL,
780   Pertile K, Rowe CC, Rembach A, Rodrigues M, Rumble R, Szoeke C, Taddei K, Taddei
781   T, Trounson B, Ames D, Masters CL, Martins RN, Group for the ADNI and AIB and

LR (2012) Blood-Based Protein Biomarkers for Diagnosis of Alzheimer Disease. *Arch Neurol* **69**, 1318–1325.

[41] Toledo JB, Toledo E, Weiner MW, Jack CR, Jagust W, Lee VM-Y, Shaw LM, Trojanowski JQ (2012) Cardiovascular risk factors, cortisol, and amyloid-β deposition in Alzheimer's Disease Neuroimaging Initiative. *Alzheimers Dement* **8**, 483–489.

[42] Lupien SJ, de Leon M, de Santi S, Convit A, Tarshish C, Nair NP, Thakur M, McEwen BS, Hauger RL, Meaney MJ (1998) Cortisol levels during human aging predict hippocampal atrophy and memory deficits. *Nat. Neurosci.* **1**, 69–73.

[43] Murphy KJ, Sheppard PAS, Troyer AK, Hampson E, Galea LAM (2019) Sex differences in cortisol and memory following acute social stress in amnestic mild cognitive impairment. *bioRxiv* 757484.

[44] McEwen BS, Biron CA, Brunson KW, Bulloch K, Chambers WH, Dhabhar FS, Goldfarb RH, Kitson RP, Miller AH, Spencer RL, Weiss JM (1997) The role of adrenocorticoids as modulators of immune function in health and disease: neural, endocrine and immune interactions. *Brain Research Reviews* **23**, 79–133.

[45] Hannum G, Guinney J, Zhao L, Zhang L, Hughes G, Sadda S, Klotzle B, Bibikova M, Fan J-B, Gao Y, Deconde R, Chen M, Rajapakse I, Friend S, Ideker T, Zhang K (2013) Genome-wide Methylation Profiles Reveal Quantitative Views of Human Aging Rates. *Molecular Cell* **49**, 359–367.

[46] Horvath S (2013) DNA methylation age of human tissues and cell types. *Genome Biol* **14**, R115.

[47] Levine ME, Lu AT, Bennett DA, Horvath S (2015) Epigenetic age of the pre-frontal cortex is associated with neuritic plaques, amyloid load, and Alzheimer's disease related cognitive functioning. *Aging (Albany NY)* **7**, 1198–1211.

[48] Yokoyama AS, Rutledge JC, Medici V (2017) DNA methylation alterations in Alzheimer's disease. *Environ Epigenet* **3**,.

[49] Gale CR, Marioni RE, Harris SE, Starr JM, Deary IJ (2018) DNA methylation and the epigenetic clock in relation to physical frailty in older people: the Lothian Birth Cohort 1936. *Clin Epigenetics* **10**,.

[50] Horvath S, Garagnani P, Bacalini MG, Pirazzini C, Salvioli S, Gentilini D, Di Blasio AM, Giuliani C, Tung S, Vinters HV, Franceschi C (2015) Accelerated epigenetic aging in Down syndrome. *Aging Cell* **14**, 491–495.

[51] Horvath S, Ritz BR (2015) Increased epigenetic age and granulocyte counts in the blood of Parkinson's disease patients. *Aging (Albany NY)* **7**, 1130–1142.

[52] Degerman S, Josefsson M, Nordin Adolfsson A, Wennstedt S, Landfors M, Haider Z, Pudas S, Hultdin M, Nyberg L, Adolfsson R (2017) Maintained memory in aging is associated with young epigenetic age. *Neurobiology of Aging* **55**, 167–171.

[53] McEwen LM, Morin AM, Edgar RD, MacIsaac JL, Jones MJ, Dow WH, Rosero-Bixby L, Kobor MS, Rehkopf DH (2017) Differential DNA methylation and lymphocyte proportions in a Costa Rican high longevity region. *Epigenetics Chromatin* **10**,.

[54] Braun PR, Han S, Hing B, Nagahama Y, Gaul LN, Heinzman JT, Grossbach AJ, Close L, Dlouhy BJ, Howard MA, Kawasaki H, Potash JB, Shinozaki G (2019) Genome-wide DNA methylation comparison between live human brain and peripheral tissues within individuals. *Translational Psychiatry* **9**, 47.

[55] Horvath S, Gurven M, Levine ME, Trumble BC, Kaplan H, Allayee H, Ritz BR, Chen B, Lu AT, Rickabaugh TM, Jamieson BD, Sun D, Li S, Chen W, Quintana-Murci L, Fagny

M, Kobor MS, Tsao PS, Reiner AP, Edlefsen KL, Absher D, Assimes TL (2016) An
epigenetic clock analysis of race/ethnicity, sex, and coronary heart disease. *Genome Biol*
17,.

[56] Crane PK, Carle A, Gibbons LE, Insel P, Mackin RS, Gross A, Jones RN, Mukherjee S,
Curtis SM, Harvey D, Weiner M, Mungas D, Alzheimer's Disease Neuroimaging
Initiative (2012) Development and assessment of a composite score for memory in the
Alzheimer's Disease Neuroimaging Initiative (ADNI). *Brain Imaging Behav* **6**, 502–516.

[57] Gibbons LE, Carle AC, Mackin RS, Harvey D, Mukherjee S, Insel P, Curtis SM, Mungas
D, Crane PK, Alzheimer's Disease Neuroimaging Initiative (2012) A composite score for
executive functioning, validated in Alzheimer's Disease Neuroimaging Initiative (ADNI)
participants with baseline mild cognitive impairment. *Brain Imaging Behav* **6**, 517–527.

[58] Craig-Schapiro R, Kuhn M, Xiong C, Pickering EH, Liu J, Misko TP, Perrin RJ, Bales
KR, Soares H, Fagan AM, Holtzman DM (2011) Multiplexed Immunoassay Panel
Identifies Novel CSF Biomarkers for Alzheimer's Disease Diagnosis and Prognosis.
PLOS ONE **6**, e18850.

[59] Hu WT, Holtzman DM, Fagan AM, Shaw LM, Perrin R, Arnold SE, Grossman M,
Xiong C, Craig-Schapiro R, Clark CM, Pickering E, Kuhn M, Chen Y, Van Deerlin VM,
McCluskey L, Elman L, Karlawish J, Chen-Plotkin A, Hurtig HI, Siderowf A, Swenson
F, Lee VM-Y, Morris JC, Trojanowski JQ, Soares H, Alzheimer's Disease Neuroimaging
Initiative (2012) Plasma multianalyte profiling in mild cognitive impairment and
Alzheimer disease. *Neurology* **79**, 897–905.

[60] Lotze M, Domin M, Gerlach FH, Gaser C, Lueders E, Schmidt CO, Neumann N (2019)
Novel findings from 2,838 Adult Brains on Sex Differences in Gray Matter Brain
Volume. *Scientific Reports* **9**, 1671.

[61] Tan A, Ma W, Vira A, Marwha D, Eliot L (2016) The human hippocampus is not
sexually-dimorphic: Meta-analysis of structural MRI volumes. *NeuroImage* **124**, 350–
366.

[62] Mormino EC, Betensky RA, Hedden T, Schultz AP, Amariglio RE, Rentz DM, Johnson
KA, Sperling RA (2014) Synergistic Effect of β-Amyloid and Neurodegeneration on
Cognitive Decline in Clinically Normal Individuals. *JAMA Neurol* **71**, 1379–1385.

[63] Jack CR, Knopman DS, Weigand SD, Wiste HJ, Vemuri P, Lowe V, Kantarci K, Gunter
JL, Senjem ML, Ivnik RJ, Roberts RO, Rocca WA, Boeve BF, Petersen RC (2012) An
operational approach to National Institute on Aging–Alzheimer's Association criteria for
preclinical Alzheimer disease. *Annals of Neurology* **71**, 765–775.

[64] Sundermann EE, Tran M, Maki PM, Bondi MW (2018) Sex differences in the
association between apolipoprotein E ε4 allele and Alzheimer's disease markers.
Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring **10**, 438–447.

[65] Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B*
(Methodological) **57**, 289–300.

[66] R Core Team (2018) *R: A language and environment for statistical   computing. R*
Foundation for Statistical Computing, Vienna, Austria., R Foundation for Statistical
Computing, Vienna, Austria.

[67] Vasanthakumar A, Davis JW, Idler K, Kim S, Nho K, Nudelman KNH, Faber K, Bai Y,
Foroud TM, Estrada K, Li Q, Saykin AJ, Waring J (2017) DNA METHYLATION

DYNAMICS IN ALZHEIMER'S DISEASE DIAGNOSIS AND PROGRESSION.
Alzheimer's & Dementia **13**, P297.
[68] Aryee MJ, Jaffe AE, Corrada-Bravo H, Ladd-Acosta C, Feinberg AP, Hansen KD,
Irizarry RA (2014) Minfi: a flexible and comprehensive Bioconductor package for the
analysis of Infinium DNA methylation microarrays. *Bioinformatics* **30**, 1363–1369.
[69] Fortin J-P, Triche TJ, Hansen KD (2017) Preprocessing, normalization and integration of
the Illumina HumanMethylationEPIC array with minfi. *Bioinformatics* **33**, 558–560.
[70] Triche TJ, Weisenberger DJ, Van Den Berg D, Laird PW, Siegmund KD (2013) Low-
level processing of Illumina Infinium DNA Methylation BeadArrays. *Nucleic Acids Res*
41, e90.
[71] Teschendorff AE, Marabita F, Lechner M, Bartlett T, Tegner J, Gomez-Cabrero D, Beck
S (2013) A beta-mixture quantile normalization method for correcting probe design bias
in Illumina Infinium 450 k DNA methylation data. *Bioinformatics* **29**, 189–196.
[72] Fiorito G, Polidoro S, Dugué P-A, Kivimaki M, Ponzi E, Matullo G, Guarrera S,
Assumma MB, Georgiadis P, Kyrtopoulos SA, Krogh V, Palli D, Panico S, Sacerdote C,
Tumino R, Chadeau-Hyam M, Stringhini S, Severi G, Hodge AM, Giles GG, Marioni R,
Linnér RK, O'Halloran AM, Kenny RA, Layte R, Baglietto L, Robinson O, McCrory C,
Milne RL, Vineis P (2017) Social adversity and epigenetic aging: a multi-cohort study on
socioeconomic differences in peripheral blood DNA methylation. *Scientific Reports* **7**,
16266.
[73] McEwen LM, Jones MJ, Lin DTS, Edgar RD, Husquin LT, MacIsaac JL, Ramadori KE,
Morin AM, Rider CF, Carlsten C, Quintana-Murci L, Horvath S, Kobor MS (2018)
Systematic evaluation of DNA methylation age estimation with common preprocessing
methods and the Infinium MethylationEPIC BeadChip array. *Clinical Epigenetics* **10**,
123.
[74] Chen BH, Marioni RE, Colicino E, Peters MJ, Ward-Caviness CK, Tsai P-C, Roetker
NS, Just AC, Demerath EW, Guan W, Bressler J, Fornage M, Studenski S, Vandiver AR,
Moore AZ, Tanaka T, Kiel DP, Liang L, Vokonas P, Schwartz J, Lunetta KL, Murabito
JM, Bandinelli S, Hernandez DG, Melzer D, Nalls M, Pilling LC, Price TR, Singleton
AB, Gieger C, Holle R, Kretschmer A, Kronenberg F, Kunze S, Linseisen J, Meisinger
C, Rathmann W, Waldenberger M, Visscher PM, Shah S, Wray NR, McRae AF, Franco
OH, Hofman A, Uitterlinden AG, Absher D, Assimes T, Levine ME, Lu AT, Tsao PS,
Hou L, Manson JE, Carty CL, LaCroix AZ, Reiner AP, Spector TD, Feinberg AP, Levy
D, Baccarelli A, van Meurs J, Bell JT, Peters A, Deary IJ, Pankow JS, Ferrucci L,
Horvath S (2016) DNA methylation-based measures of biological age: meta-analysis
predicting time to death. *Aging (Albany NY)* **8**, 1844–1859.
[75] Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson HH,
Wiencke JK, Kelsey KT (2012) DNA methylation arrays as surrogate measures of cell
mixture distribution. *BMC Bioinformatics* **13**, 86.
[76] Sundermann EE, Biegon A, Rubin LH, Lipton RB, Landau S, Maki PM, Alzheimer's
Disease Neuroimaging Initiative (2017) Does the Female Advantage in Verbal Memory
Contribute to Underestimating Alzheimer's Disease Pathology in Women versus Men? *J.*
Alzheimers Dis. **56**, 947–957.
[77] Mielke MM, Vemuri P, Rocca WA (2014) Clinical epidemiology of Alzheimer's
disease: assessing sex and gender differences. *Clin Epidemiol* **6**, 37–48.

[78] Petersen RC, Roberts RO, Knopman DS, Geda YE, Cha RH, Pankratz VS, Boeve BF, Tangalos EG, Ivnik RJ, Rocca WA (2010) Prevalence of mild cognitive impairment is higher in men. The Mayo Clinic Study of Aging. *Neurology* **75**, 889–897.

[79] Winblad B, Amouyel P, Andrieu S, Ballard C, Brayne C, Brodaty H, Cedazo-Minguez A, Dubois B, Edvardsson D, Feldman H, Fratiglioni L, Frisoni GB, Gauthier S, Georges J, Graff C, Iqbal K, Jessen F, Johansson G, Jönsson L, Kivipelto M, Knapp M, Mangialasche F, Melis R, Nordberg A, Rikkert MO, Qiu C, Sakmar TP, Scheltens P, Schneider LS, Sperling R, Tjernberg LO, Waldemar G, Wimo A, Zetterberg H (2016) Defeating Alzheimer's disease and other dementias: a priority for European science and society. *The Lancet Neurology* **15**, 455–532.

[80] Jack CR, Therneau TM, Weigand SD, Wiste HJ, Knopman DS, Vemuri P, Lowe VJ, Mielke MM, Roberts RO, Machulda MM, Graff-Radford J, Jones DT, Schwarz CG, Gunter JL, Senjem ML, Rocca WA, Petersen RC (2019) Prevalence of Biologically vs Clinically Defined Alzheimer Spectrum Entities Using the National Institute on Aging–Alzheimer's Association Research Framework. *JAMA Neurol.*

[81] Ferretti MT, Iulita MF, Cavedo E, Chiesa PA, Schumacher Dimech A, Santuccione Chadha A, Baracchi F, Girouard H, Misoch S, Giacobini E, Depypere H, Hampel H (2018) Sex differences in Alzheimer disease — the gateway to precision medicine. *Nature Reviews Neurology* **14**, 457–469.

[82] Blennow K, Dubois B, Fagan AM, Lewczuk P, de Leon MJ, Hampel H (2015) Clinical utility of cerebrospinal fluid biomarkers in the diagnosis of early Alzheimer's disease. *Alzheimer's & Dementia* **11**, 58–69.

[83] Henriques AD, Benedet AL, Camargos EF, Rosa-Neto P, Nóbrega OT (2018) Fluid and imaging biomarkers for Alzheimer's disease: Where we stand and where to head to. *Experimental Gerontology* **107**, 169–177.

[84] Lewis J, Dickson DW, Lin W-L, Chisholm L, Corral A, Jones G, Yen S-H, Sahara N, Skipper L, Yager D, Eckman C, Hardy J, Hutton M, McGowan E (2001) Enhanced Neurofibrillary Degeneration in Transgenic Mice Expressing Mutant Tau and APP. *Science, New Series* **293**, 1487–1491.

[85] Hohman TJ, Dumitrescu L, Barnes LL, Thambisetty M, Beecham G, Kunkle B, Gifford KA, Bush WS, Chibnik LB, Mukherjee S, Jager PLD, Kukull W, Crane PK, Resnick SM, Keene CD, Montine TJ, Schellenberg GD, Haines JL, Zetterberg H, Blennow K, Larson EB, Johnson SC, Albert M, Bennett DA, Schneider JA, Jefferson AL (2018) Sex-Specific Association of Apolipoprotein E With Cerebrospinal Fluid Levels of Tau. *JAMA Neurol* **75**, 989–998.

[86] Altmann A, Tian L, Henderson VW, Greicius MD (2014) Sex modifies the APOE-related risk of developing Alzheimer disease. *Annals of Neurology* **75**, 563–573.

[87] Buckner RL (2004) Memory and Executive Function in Aging and AD: Multiple Factors that Cause Decline and Reserve Factors that Compensate. *Neuron* **44**, 195–208.

[88] Ewers M, Walsh C, Trojanowski JQ, Shaw LM, Petersen RC, Jack CR, Feldman HH, Bokde ALW, Alexander GE, Scheltens P, Vellas B, Dubois B, Weiner M, Hampel H (2012) Prediction of conversion from mild cognitive impairment to Alzheimer's disease dementia based upon biomarkers and neuropsychological test performance. *Neurobiology of Aging* **33**, 1203-1214.e2.

[89]   Li B, Shi J, Gutman BA, Baxter LC, Thompson PM, Caselli RJ, Wang Y (2016) Influence of APOE Genotype on Hippocampal Atrophy over Time - An N=1925 Surface-Based ADNI Study. *PLoS One* **11**,.

[90]   Mungas D, Beckett L, Harvey D, Farias ST, Reed B, Carmichael O, Olichney J, Miller J, DeCarli C (2010) Heterogeneity of Cognitive Trajectories in Diverse Older Person. *Psychol Aging* **25**, 606–619.

[91]   Pievani M, Galluzzi S, Thompson PM, Rasser PE, Bonetti M, Frisoni GB (2011) APOE4 is associated with greater atrophy of the hippocampal formation in Alzheimer's disease. *Neuroimage* **55**, 909–919.

[92]   Shi J, Leporé N, Gutman BA, Thompson PM, Baxter LC, Caselli RJ, Wang Y, Alzheimer's Disease Neuroimaging Initiative (2014) Genetic influence of apolipoprotein E4 genotype on hippocampal morphometry: An N = 725 surface-based Alzheimer's disease neuroimaging initiative study. *Hum Brain Mapp* **35**, 3903–3918.

[93]   Cavedo E, Chiesa PA, Houot M, Ferretti MT, Grothe MJ, Teipel SJ, Lista S, Habert M-O, Potier M-C, Dubois B, Hampel H, Bakardjian H, Benali H, Bertin H, Bonheur J, Boukadida L, Boukerrou N, Cavedo E, Chiesa P, Colliot O, Dubois B, Dubois M, Epelbaum S, Gagliardi G, Genthon R, Habert M-O, Hampel H, Houot M, Kas A, Lamari F, Levy M, Lista S, Metzinger C, Mochel F, Nyasse F, Poisson C, Potier M-C, Revillon M, Santos A, Andrade KS, Sole M, Surtee M, de Schotten MT, Vergallo A, Younsi N, Aguilar LF, Babiloni C, Baldacci F, Benda N, Black KL, Bokde ALW, Bonuccelli U, Broich K, Bun RS, Cacciola F, Castrillo J, Cavedo E, Ceravolo R, Chiesa PA, Colliot O, Coman C-M, Corvol J-C, Cuello AC, Cummings JL, Depypere H, Dubois B, Duggento A, Durrleman S, Escott-Price V, Federoff H, Ferretti MT, Fiandaca M, Frank RA, Garaci F, Genthon R, George N, Giorgi FS, Graziani M, Haberkamp M, Habert M-O, Hampel H, Herholz K, Karran E, Kim SH, Koronyo Y, Koronyo-Hamaoui M, Lamari F, Langevin T, Lehéricy S, Lista S, Lorenceau J, Mapstone M, Neri C, Nisticò R, Nyasse-Messene F, O'bryant SE, Perry G, Ritchie C, Rojkova K, Rossi S, Saidi A, Santarnecchi E, Schneider LS, Sporns O, Toschi N, Verdooner SR, Vergallo A, Villain N, Welikovitch LA, Woodcock J, Younesi E (2018) Sex differences in functional and molecular neuroimaging biomarkers of Alzheimer's disease in cognitively normal older adults with subjective memory complaints. *Alzheimer's & Dementia* **14**, 1204–1215.

[94]   Jack CR, Wiste HJ, Weigand SD, Knopman DS, Vemuri P, Mielke MM, Lowe V, Senjem ML, Gunter JL, Machulda MM, Gregg BE, Pankratz VS, Rocca WA, Petersen RC (2015) Age, Sex, and APOE ε4 Effects on Memory, Brain Structure, and β-Amyloid Across the Adult Life Span. *JAMA Neurol* **72**, 511–519.

[95]   Murphy DGM, DeCarli C, Mclntosh AR, Daly E, Mentis MJ, Pietrini P, Szczepanik J, Schapiro MB, Grady CL, Horwitz B, Rapoport SI (1996) Sex Differences in Human Brain Morphometry and Metabolism: An In Vivo Quantitative Magnetic Resonance Imaging and Positron Emission Tomography Study on the Effect of Aging. *Arch Gen Psychiatry* **53**, 585–594.

[96]   Ritchie SJ, Cox SR, Shen X, Lombardo MV, Reus LM, Alloza C, Harris MA, Alderson HL, Hunter S, Neilson E, Liewald DCM, Auyeung B, Whalley HC, Lawrie SM, Gale CR, Bastin ME, McIntosh AM, Deary IJ (2018) Sex Differences in the Adult Human Brain: Evidence from 5216 UK Biobank Participants. *Cereb Cortex* **28**, 2959–2975.

[97]   Fleisher A, Grundman M, Jack CR, Petersen RC, Taylor C, Kim HT, Schiller DHB, Bagwell V, Sencakova D, Weiner MF, DeCarli C, DeKosky ST, Dyck CH van, Thal LJ

(2005) Sex, Apolipoprotein E ε4 Status, and Hippocampal Volume in Mild Cognitive
Impairment. *Arch Neurol* **62**, 953–957.
[98] Sundermann EE, Biegon A, Rubin LH, Lipton RB, Mowrey W, Landau S, Maki PM
(2016) Better verbal memory in women than men in MCI despite similar levels of
hippocampal atrophy. *Neurology* **86**, 1368–1376.
[99] Buckley RF, Mormino EC, Amariglio RE, Properzi MJ, Rabin JS, Lim YY, Papp KV,
Jacobs HIL, Burnham S, Hanseeuw BJ, Doré V, Dobson A, Masters CL, Waller M,
Rowe CC, Maruff P, Donohue MC, Rentz DM, Kirn D, Hedden T, Chhatwal J, Schultz
AP, Johnson KA, Villemagne VL, Sperling RA (2018) Sex, amyloid, and APOE ε4 and
risk of cognitive decline in preclinical Alzheimer's disease: Findings from three well-
characterized cohorts. *Alzheimer's & Dementia* **14**, 1193–1203.
[100] Stern Y (2002) What is cognitive reserve? Theory and research application of the reserve
concept. *J Int Neuropsychol Soc* **8**, 448–460.
[101] Koenig W, Sund M, Fröhlich M, Fischer HG, Löwel H, Döring A, Hutchinson WL,
Pepys MB (1999) C-Reactive protein, a sensitive marker of inflammation, predicts future
risk of coronary heart disease in initially healthy middle-aged men: results from the
MONICA (Monitoring Trends and Determinants in Cardiovascular Disease) Augsburg
Cohort Study, 1984 to 1992. *Circulation* **99**, 237–242.
[102] Ridker PM, Buring JE, Shih J, Matias M, Hennekens CH (1998) Prospective study of C-
reactive protein and the risk of future cardiovascular events among apparently healthy
women. *Circulation* **98**, 731–733.
[103] Koyama A, O'Brien J, Weuve J, Blacker D, Metti AL, Yaffe K (2013) The role of
peripheral inflammatory markers in dementia and Alzheimer's disease: a meta-analysis.
J. Gerontol. A Biol. Sci. Med. Sci. **68**, 433–440.
[104] Hubacek JA, Peasey A, Pikhart H, Stavek P, Kubinova R, Marmot M, Bobak M (2010)
APOE polymorphism and its effect on plasma C-reactive protein levels in a large general
population sample. *Hum Immunol* **71**, 304–308.
[105] Yun Y-W, Kweon S-S, Choi J-S, Rhee J-A, Lee Y-H, Nam H-S, Jeong S-K, Park K-S,
Ryu S-Y, Choi S-W, Kim HN, Cauley JA, Shin M-H (2015) APOE Polymorphism Is
Associated with C-reactive Protein Levels but Not with White Blood Cell Count: Dong-
gu Study and Namwon Study. *J Korean Med Sci* **30**, 860–865.
[106] Gong C, Wei D, Wang Y, Ma J, Yuan C, Zhang W, Yu G, Zhao Y (2016) A Meta-
Analysis of C-Reactive Protein in Patients With Alzheimer's Disease. *Am J Alzheimers*
Dis Other Demen **31**, 194–200.
[107] Ng A, Tam WW, Zhang MW, Ho CS, Husain SF, McIntyre RS, Ho RC (2018) IL-1β,
IL-6, TNF- α and CRP in Elderly Patients with Depression or Alzheimer's disease:
Systematic Review and Meta-Analysis. *Scientific Reports* **8**, 12050.
[108] Rosa MD, Dell'Ombra N, Zambito AM, Malaguarnera M, Nicoletti F, Malaguarnera L
(2006) Chitotriosidase and inflammatory mediator levels in Alzheimer's disease and
cerebrovascular dementia. *European Journal of Neuroscience* **23**, 2648–2656.
[109] Motta M, Imbesi R, Di Rosa M, Stivala F, Malaguarnera L (2007) Altered plasma
cytokine levels in Alzheimer's disease: Correlation with the disease progression.
Immunology Letters **114**, 46–51.
[110] Ponthieux A, Herbeth B, Droesch S, Lambert D, Visvikis S (2003) Age- and Sex-related
Reference Values for Serum Adhesion Molecule Concentrations in Healthy Individuals:

Intercellular Adhesion Molecule-1 and E-, P-, and L-Selectin. *Clinical Chemistry* **49**, 1544–1546.

[111] Zenaro E, Piacentino G, Constantin G (2017) The blood-brain barrier in Alzheimer's disease. *Neurobiology of Disease* **107**, 41–56.

[112] Nielsen HM, Londos E, Minthon L, Janciauskiene SM (2007) Soluble adhesion molecules and angiotensin-converting enzyme in dementia. *Neurobiol. Dis.* **26**, 27–35.

[113] Huang C-W, Tsai M-H, Chen N-C, Chen W-H, Lu Y-T, Lui C-C, Chang Y-T, Chang W-N, Chang AYW, Chang C-C (2015) Clinical significance of circulating vascular cell adhesion molecule-1 to white matter disintegrity in Alzheimer's dementia. *Thrombosis and Haemostasis* **114**, 1230–1240.

[114] Rentzos M, Michalopoulou M, Nikolaou C, Cambouri C, Rombos A, Dimitrakopoulos A, Kapaki E, Vassilopoulos D (2004) Serum Levels of Soluble Intercellular Adhesion Molecule-1 and Soluble Endothelial Leukocyte Adhesion Molecule-1 in Alzheimer's Disease. *J Geriatr Psychiatry Neurol* **17**, 225–231.

[115] Walters A, Phillips E, Zheng R, Biju M, Kuruvilla T (2016) Evidence for neuroinflammation in Alzheimer's disease: Neuroinflammation in Alzheimer's. *Progress in Neurology and Psychiatry* **20**, 25–31.

[116] Heneka MT, Carson MJ, Khoury JE, Landreth GE, Brosseron F, Feinstein DL, Jacobs AH, Wyss-Coray T, Vitorica J, Ransohoff RM, Herrup K, Frautschy SA, Finsen B, Brown GC, Verkhratsky A, Yamanaka K, Koistinaho J, Latz E, Halle A, Petzold GC, Town T, Morgan D, Shinohara ML, Perry VH, Holmes C, Bazan NG, Brooks DJ, Hunot S, Joseph B, Deigendesch N, Garaschuk O, Boddeke E, Dinarello CA, Breitner JC, Cole GM, Golenbock DT, Kummer MP (2015) Neuroinflammation in Alzheimer's disease. *The Lancet Neurology* **14**, 388–405.

[117] Chakrabarty P, Tianbai L, Herring A, Ceballos-Diaz C, Das P, Golde TE (2012) Hippocampal expression of murine IL-4 results in exacerbation of amyloid deposition. *Mol Neurodegener* **7**, 36.

[118] Chakrabarty P, Herring A, Ceballos-Diaz C, Das P, Golde TE (2011) Hippocampal expression of murine TNFα results in attenuation of amyloid deposition in vivo. *Molecular Neurodegeneration* **6**, 16.

[119] Chakrabarty P, Ceballos-Diaz C, Beccard A, Janus C, Dickson D, Golde TE, Das P (2010) IFN-gamma promotes complement expression and attenuates amyloid plaque deposition in amyloid beta precursor protein transgenic mice. *J. Immunol.* **184**, 5333–5343.

[120] Chakrabarty P, Jansen-West K, Beccard A, Ceballos-Diaz C, Levites Y, Verbeeck C, Zubair AC, Dickson D, Golde TE, Das P (2010) Massive gliosis induced by interleukin-6 suppresses Aβ deposition in vivo: evidence against inflammation as a driving force for amyloid deposition. *FASEB J* **24**, 548–559.

[121] Ghosh S, Wu MD, Shaftel SS, Kyrkanides S, LaFerla FM, Olschowka JA, O'Banion MK (2013) Sustained interleukin-1β overexpression exacerbates tau pathology despite reduced amyloid burden in an Alzheimer's mouse model. *J. Neurosci.* **33**, 5053–5064.

[122] Shaftel SS, Carlson TJ, Olschowka JA, Kyrkanides S, Matousek SB, O'Banion MK (2007) Chronic interleukin-1beta expression in mouse brain leads to leukocyte infiltration and neutrophil-independent blood brain barrier permeability without overt neurodegeneration. *J. Neurosci.* **27**, 9301–9309.

[123]   GTEx Project (2018) *The Genotype-Tissue Expression (GTEx) Project was supported by the Common Fund of the Office of the Director of the National Institutes of Health, and by NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS. The data used for the analyses described in this manuscript were obtained from the GTEx Portal on 11/04/18.*

[124]   Dhingra R, Kwee LC, Diaz-Sanchez D, Devlin RB, Cascio W, Hauser ER, Gregory S, Shah S, Kraus WE, Olden K, Ward-Caviness CK (2019) Evaluating DNA methylation age on the Illumina MethylationEPIC Bead Chip. *PLOS ONE* **14**, e0207834.

[125]   Levine ME, Lu AT, Quach A, Chen BH, Assimes TL, Bandinelli S, Hou L, Baccarelli AA, Stewart JD, Li Y, Whitsel EA, Wilson JG, Reiner AP, Aviv A, Lohman K, Liu Y, Ferrucci L, Horvath S (2018) An epigenetic biomarker of aging for lifespan and healthspan. *Aging (Albany NY)* **10**, 573–591.

[126]   Lu AT, Quach A, Wilson JG, Reiner AP, Aviv A, Raj K, Hou L, Baccarelli AA, Li Y, Stewart JD, Whitsel EA, Assimes TL, Ferrucci L, Horvath S (2019) DNA methylation GrimAge strongly predicts lifespan and healthspan. *Aging (Albany NY)* **11**, 303–327.

[127]   Fitten LJ, Ortiz F, Fairbanks L, Bartzokis G, Lu P, Klein E, Coppola G, Ringman J (2014) Younger age of dementia diagnosis in a Hispanic population in southern California. *Int J Geriatr Psychiatry* **29**, 586–593.

[128]   Mayeda ER, Glymour MM, Quesenberry CP, Whitmer RA (2016) Inequalities in dementia incidence between six racial and ethnic groups over 14 years. *Alzheimer's & Dementia* **12**, 216–224.

[129]   Steenland K, Goldstein FC, Levey A, Wharton W (2016) A Meta-Analysis of Alzheimer's Disease Incidence and Prevalence Comparing African-Americans and Caucasians. *J. Alzheimers Dis.* **50**, 71–76.

[130]   Sharp ES, Gatz M (2011) The Relationship between Education and Dementia An Updated Systematic Review. *Alzheimer Dis Assoc Disord* **25**, 289–304.

Figure 1

Figure 2

Figure 3

Table 1. Demographic and clinical information for all participants and subdivided by sex. Biomarkers for AD are from cerebrospinal fluid. P-values adjusted for age are presented for easier comparison and are taken from the linear model of sex and APOE genotype (Table 3). Volume of the hippocampus was corrected using a regression method or using a ratio with intracranial volume (see methods for more details).

| | Total | Sex | | P-value | P-value (adjusted for age) |
	No. 1,460	Female No. 630	Male No. 830		
Age					
Mean (SD)	74.13 (±7.25)	73.15 (±7.28)	74.87 (±7.14)	< 0.0001	
Education (years)					
Mean (SD)	15.83 (±2.88)	15.15 (±2.79)	16.34 (±2.85)	< 0.0001	
Ethnicity					
White	1,352 (92.60%)	573 (90.95%)	779 (93.86%)	0.043	
Not white	108 (7.40%)	57 (9.05%)	51 (6.14%)		
Baseline diagnosis					
CN	341 (23.4%)	168 (26.7%)	173 (20.8%)	0.013	
EMCI	257 (17.6%)	108 (17.1%)	149 (18.0%)		
LMCI	533 (36.5%)	205 (32.5%)	328 (39.5%)		
AD	329 (22.5%)	149 (23.7%)	180 (21.7%)		
APOEε4 allele number					
0	702 (48.08%)	300 (47.62%)	402 (48.43%)	0.8	
1	574 (39.32%)	252 (40.00%)	322 (38.80%)		
2	170 (11.64%)	70 (11.11%)	100 (12.05%)		
Missing	14 (0.96%)	8 (1.27%)	6 (0.72%)		
Volume of hippocampus (uncorrected, mm^3)					
Mean (SD)	6659.47 (±1176.42)	6446.71 (±1169.97)	6822.86 (±1155.87)	< 0.0001	
Missing	226 (15.48%)	94 (14.92%)	132 (15.90%)		
Volume of hippocampus (regression corrected)					
Mean (SD)	6659.47 (±1174.34)	6447.09 (±1167.90)	6822.57 (±1153.83)	**< 0.0001**	**<0.0001**
Missing	226 (15.48%)	94 (14.92%)	132 (15.90%)		
Volume of hippocampus/intracranial volume					
Mean (SD)	0.00436 (±0.00080)	0.00454 (±0.00082)	0.00423 (±0.00076)	**< 0.0001**	**<0.0001**
Missing	226 (15.48%)	94 (14.92%)	132 (15.90%)		
Amyloid Beta					
Mean (SD)	830.97 (±358.04)	856.41 (±346.87)	812.44 (±365.16)	**0.016**	0.38
Missing	513 (35.14%)	231 (36.67%)	282 (33.98%)		
Tau					
Mean (SD)	294.38 (±137.27)	314.56 (±152.70)	279.70 (±122.91)	**0.002**	**<0.0001**
Missing	513 (35.14%)	231 (36.67%)	282 (33.98%)		
PTau					
Mean (SD)	28.89 (±15.31)	30.87 (±16.95)	27.44 (±13.83)	**0.007**	**<0.0001**
Missing	513 (35.14%)	231 (36.67%)	282 (33.98%)		
Executive Function (ADNI_EF)					
Mean (SD)	0.02 (±0.96)	0.06 (±0.97)	-0.00 (±0.95)	0.20	**<0.0001**
Missing	311 (21.30%)	145 (23.02%)	166 (20.00%)		
Memory (ADNI_MEM)					
Mean (SD)	0.10 (±0.87)	0.21 (±0.94)	0.02 (±0.80)	**0.0006**	**<0.0001**
Missing	310 (21.23%)	145 (23.02%)	165 (19.88%)		

P-values are from Wilcoxon rank sum tests for continuous variables and Fisher's exact tests for categorical variables. Missing refers to number of individuals and the percent of the total cohort that had missing data for that variable

Table 2. Demographic and clinical information for subset of ADNI data subdivided by sex. A. Participants with measured biomarkers in cerebrospinal fluid (CSF), B. Participants with available whole blood Illumina HumanMethylationEPIC DNA methylation data, C. Participants with matched Illumina HumanMethylationEPIC DNA methylation array data and measured CSF biomarkers. In all three subdata sets, females were significantly younger and had fewer years of education than males. In data set A (but not B and C), more females (24.0 % compared to 21.8%) were diagnosed with AD, more females were cognitively normal (26.5% compared to 22.9%) and fewer females were diagnosed with late MCI compared to males (49.5% compared to 55.3%). In data set A, females had lower CSF cortisol, interleukin-3, interleukin 8, interleukin-16, immunoglobulin A, and intercellular adhesion molecule compared to males. Empty cells indicate data not available.

	A				B				C			
		Sex				Sex				Sex		
	Total No. 279	Female No. 109	Male No. 170	P-value	Total No. 640	Female No. 284	Male No. 356	P-value	Total No. 533	Female No. 243	Male No. 290	P-value
Age												
Mean (SD)	75.15 (±6.86)	73.75 (±6.69)	76.04 (±6.83)	0.007	75.63 (±7.68)	74.78 (±8.03)	76.31 (±7.32)	<0.0001	75.01 (±7.61)	74.31 (±8.10)	75.61 (±7.11)	0.0019
Education (years)												
Mean (SD)	15.69 (±2.95)	14.68 (±2.74)	16.34 (±2.90)	< 0.0001	16.21 (±2.70)	15.53 (±2.59)	16.75 (±2.68)	<0.0001	16.24 (±2.64)	15.57 (±2.49)	16.83 (±2.64)	< 0.0001
Ethnicity												
White	267 (95.70%)	103 (94.50%)	164 (96.47%)	0.55	627 (97.97%)	279 (98.23 %)	348 (97.75%)	0.78	521 (97.75 %)	238 (97.94%)	283 (97.94 %)	0.99
Not White[1]	12 (4.30%)	6 (5.50%)	6 (3.53%)		13 (2.03%)	5 (1.76%)	8 (2.25%)		12 (2.25 %)	5 (2.06 %)	7 (2.41 %)	
Baseline diagnosis												
CN	74 (26.5%)	35 (32.1%)	39 (22.9%)	0.051	217 (33.9%)	109 (38.38%)	108 (30.34%)	0.11	171 (32.08%)	88 (36.21%)	83 (28.62 %)	0.19
EMCI					186 (29.06%)	83 (29.23%)	103 (28.93%)		173 (32.46 %)	79 (32.51 %)	94 (32.41%)	
LMCI	138 (49.5%)	44 (40.4%)	94 (55.3%)		200 (31.25%)	78 (27.46%)	122 (34.27%)		155 (29.08 %)	94 (38.68 %)	92 (31.72%)	
AD	67 (24.0%)	30 (27.5%)	37 (21.8%)		37 (5.78%)	14 (4.23 %)	23 (6.46%)		34 (6.38 %)	13 (5.35 %)	21 (7.24 %)	
APOEε4 allele number												
0	134 (48.03%)	51 (46.79%)	83 (48.82%)	0.78	369 (57.66 %)	169 (59.51%)	200 (56.18%)	0.37	313 (58.72 %)	146 (60.08 %)	167 (57.59 %)	0.45
1	109 (39.07%)	42 (38.53%)	67 (39.41%)		220 (34.38%)	97 (34.15%)	123 (34.55%)		173 (32.46%)	80 (32.92 %)	93 (32.07 %)	
2	36 (12.90%)	16 (14.68%)	20 (11.76%)		51 (7.97%)	18 (6.34%)	33 (9.27%)		47 (8.82%)	17 (7.00 %)	30 (10.34 %)	
Cortisol (ng/mL)												
Mean (SD)	16.05 (±6.04)	14.92 (±6.01)	16.78 (±5.96)	0.008								
C reactive protein (ug/mL)												
Mean (SD)	-2.83 (±0.56)	-2.77 (±0.64)	-2.87 (±0.51)	0.23								
CD40 antigen (ng/mL)												
Mean (SD)	-0.65 (±0.12)	-0.66 (±0.10)	-0.64 (±0.14)	0.12								
Interleukin 16 (pg/mL)												
Mean (SD)	0.91 (±0.18)	0.87 (±0.17)	0.94 (±0.19)	0.004								
Interleukin 3 (ng/mL)												
Mean (SD)	-2.22 (±0.32)	-2.28 (±0.29)	-2.17 (±0.34)	0.001								
Interleukin 6 receptor (ng/mL)												
Mean (SD)	-0.01 (±0.15)	-0.02 (±0.14)	-0.00 (±0.15)	0.30								
Interleukin 8 (pg/mL)												
Mean (SD)	1.68 (±0.15)	1.64 (±0.11)	1.70 (±0.16)	0.001								
Intercellular adhesion molecule (ng/mL)												
Mean (SD)	0.96 (±0.44)	0.83 (±0.33)	1.04 (±0.48)	0.0001								
Immunoglobulin A (mg/mL)												
Mean (SD)	-2.54 (±0.31)	-2.68 (±0.26)	-2.45 (±0.31)	< 0.0001								
Executive Function Score												
Mean (SD)					0.36 (±0.98)	0.38 (±1.01)	0.34 (±0.95)	0.17				
Memory Score												
Mean (SD)					0.40 (±0.92)	0.57 (±1.01)	0.26 (±0.82)	<0.0001				
Amyloid Beta												
Mean (SD)									1040.98 (±454.72)	1055.50 (±449.23)	1028.35 (±459.36)	0.18
Tau												
Mean (SD)									289.80 (±124.68)	300.90 (±139.07)	280.13 (±109.82)	0.072
PTau												
Mean (SD)									27.47 (±13.65)	28.25 (±15.08)	26.78 (±12.24)	0.36

P-values are from Wilcoxon rank sum tests for continuous variables and Fisher's exact tests for categorical variables. ‡Includes self-reported Black, Asian, American Indian/Alaskan, and >1 ethnicity.

Table 3. Linear regression results for models with sex and APOE status. Only shown are the models with significant associations. P-values are for overall tests and are FDR-adjusted. All model summaries are available in Supplementary Table S1.

Predictors	ADNI MEM Estimates (CI)	adjusted p	ADNI EF Estimates (CI)	adjusted p	ABETA Estimates (CI)	adjusted p	Hippocampus volume (corrected) Estimates (CI)	adjusted p	TAU Estimates (CI)	adjusted p	PTAU Estimates (CI)	adjusted p
(Intercept)	1.63 (1.12 – 2.14)		2.05 (1.47 – 2.62)		1458.67 (1247.77 – 1669.57)		11550.67 (10935.47 – 12165.88)		56.15 (-30.03 – 142.34)		4.75 (-4.89 – 14.38)	
AGE (years)	-0.02 (-0.02 – -0.01)		-0.02 (-0.03 – -0.02)		-6.19 (-9.01 – -3.36)		-65.72 (-73.95 – -57.49)		2.68 (1.54 – 3.81)		0.26 (0.13 – 0.39)	
Male (ref = Female)	-0.17 (-0.26 – -0.07)	**0.003**	-0.03 (-0.14 – 0.08)	0.73	-29.77 (-71.55 – 12.01)	0.33	482.18 (365.03 – 599.32)	**<0.0001**	-7.37 (-31.43 – 16.70)		-0.34 (-3.03 – 2.35)	
APOE status (ref = 0 alleles)		**<0.0001**		**<0.0001**		**<0.0001**		**<0.0001**				
1 allele	-0.45 (-0.55 – -0.34)		-0.3 (-0.42 – -0.19)		-240.23 (-284.27 – -196.20)		-518.73 (-642.15 – -395.31)		104.14 (77.21 – 131.06)		11.73 (8.72 – 14.74)	
2 alleles	-0.69 (-0.85 – -0.53)		-0.46 (-0.64 – -0.28)		-455.95 (-521.02 – -390.88)		-946.01 (-1136.77 – -755.25)		178.88 (137.45 – 220.31)		19.81 (15.18 – 24.44)	
Interaction term										**0.001**		**0.002**
Male:1 allele									-49.76 (-85.30 – -14.22)		-5.58 (-9.55 – -1.60)	
Male:2 alleles									-101.56 (-153.97 – -49.16)		-10.78 (-16.64 – -4.92)	
Observations	1145		1144		947		1224		947		947	
R^2 / adjusted R^2	0.106 / 0.103		0.058 / 0.055		0.203 / 0.199		0.233/0.231		0.140 / 0.134		0.136 / 0.130	

Table 3. Continued

Predictors	C Reactive Protein ug/ml Estimates (CI)	adjusted p	Interleukin 16 pg/ml Estimates (CI)	adjusted p	Interleukin 8 pg/m L Estimates (CI)	adjusted p	Immunoglobulin A mg/ml Estimates (CI)	adjusted p	Intercellular Adhesion Molecule 1 ng/ml Estimates (CI)	adjusted p
(Intercept)	-3.05 (-3.79 – -2.32)		0.35 (0.10 – 0.59)		1.38 (1.18 – 1.57)		-2.82 (-3.22 – -2.43)		-0.19 (-0.75 – 0.36)	
AGE (years)	0.01 (-0.00 – 0.02)		0.01 (0.00 – 0.01)		0 (0.00 – 0.01)		0 (-0.00 – 0.01)		0.01 (0.01 – 0.02)	
Male (ref = Female)	-0.12 (-0.26 – 0.01)	0.19	0.12 (0.06 – 0.18)		0.1 (0.05 – 0.15)	**0.02**	0.21 (0.14 – 0.29)	**<0.0001**	0.18 (0.07 – 0.28)	**0.003**
APOE status (ref = 0 alleles)		**0.009**				0.39		0.33		0.38
1 allele	-0.19 (-0.33 – -0.05)		0.08 (0.01 – 0.16)		0.04 (-0.02 – 0.10)		0.02 (-0.05 – 0.10)		0.09 (-0.01 – 0.20)	
2 alleles	-0.31 (-0.52 – -0.10)		0.06 (-0.04 – 0.16)		0.01 (-0.07 – 0.09)		-0.09 (-0.20 – 0.02)		0.02 (-0.13 – 0.18)	
Interaction term				**0.03**						
Male:1 allele			-0.13 (-0.22 – -0.03)							
Male:2 alleles			-0.15 (-0.28 – -0.02)							
Observations	279		279		279		279		279	
R^2 / adjusted R^2	0.058 / 0.045		0.117 / 0.098		0.092 / 0.072		0.135 / 0.122		0.107 / 0.094	

Table 4. Linear regression results for models with sex and baseline diagnosis. Only shown are the models with significant associations. P-values are for overall tests and are FDR-adjusted. All model summaries are available in Supplementary Table S2.

Predictors	ADNI MEM Estimates (CI)	adjusted p	ADNI EF Estimates (CI)	adjusted p	ABETA Estimates (CI)	adjusted p	Hippocampus volume (corrected) Estimates (CI)	adjusted p	TAU Estimates (CI)	adjusted p	PTAU Estimates (CI)	adjusted p
(Intercept)	1.79 (1.45 – 2.13)		2.26 (1.80 – 2.73)		1161.57 (944.04 – 1379.10)		11424.98 (10886.11 – 11963.86)		154.16 (70.57 – 237.75)		15.72 (6.36 – 25.09)	
AGE (years)	-0.01 (-0.01 – -0.00)		-0.02 (-0.03 – -0.01)		-1.64 (-4.52 – 1.24)		-57.95 (-65.07 – -50.83)		1.25 (0.15 – 2.36)		0.1 (-0.02 – 0.23)	
Male (ref = Female)	-0.16 (-0.23 – -0.09)	**<0.0001**	-0.04 (-0.13 – 0.05)	0.56	-26.62 (-69.46 – 16.22)	0.42	516.27 (414.05 – 618.49)	**<0.0001**	-42.59 (-59.05 – -26.13)	**<0.0001**	-4.22 (-6.06 – -2.38)	**<0.0001**
Diagnosis (ref = CN)		**<0.0001**		**<0.0001**		**<0.0001**		**<0.0001**		**<0.0001**		**<0.0001**
EMCI	-0.5 (-0.61 – -0.39)		-0.42 (-0.56 – -0.27)		-85.2 (-148.82 – -21.59)		-340.14 (-495.23 – -185.04)		37.85 (13.41 – 62.30)		4.22 (1.48 – 6.96)	
LMCI	-1.08 (-1.16 – -1.00)		-0.79 (-0.90 – -0.67)		-256.85 (-315.81 – -197.89)		-1049.98 (-1183.33 – -916.63)		93.34 (70.69 – 116.00)		10.58 (8.05 – 13.12)	
AD	-1.84 (-1.94 – -1.75)		-1.63 (-1.76 – -1.50)		-390.48 (-453.59 – -327.37)		-1603.56 (-1751.83 – -1455.29)		143.6 (119.35 – 167.86)		15.81 (13.10 – 18.53)	
Observations	1150		1149		947		1234		947		947	
R² / adjusted R²	0.589 / 0.588		0.380 / 0.377		0.168 / 0.164		0.416/ 0.414		0.164 / 0.160		0.156 / 0.152	

Table 4. Continued

Predictors	Interleukin 16 pg/ml Estimates (CI)	adjusted p	Interleukin 8 pg/ml Estimates (CI)	adjusted p	Immunoglobulin A mg/ml Estimates (CI)	adjusted p	Intercellular Adhesion Molecule 1 ng/ml Estimates (CI)	adjusted p
(Intercept)	0.42 (0.18 – 0.65)		1.4 (1.20 – 1.59)		-2.9 (-3.29 – -2.50)		-0.22 (-0.77 – 0.33)	
AGE (years)	0.01 (0.00 – 0.01)		0 (0.00 – 0.01)		0 (-0.00 – 0.01)		0.01 (0.01 – 0.02)	
Male (ref = Female)	0.05 (0.01 – 0.10)	0.07	0.05 (0.01 – 0.09)	**0.03**	0.21 (0.14 – 0.29)	**<0.0001**	0.17 (0.06 – 0.27)	**0.008**
Diagnosis (ref = CN)		0.66		0.94		0.98		0.71
EMCI								
LMCI	0.01 (-0.04 – 0.06)		0.01 (-0.03 – 0.05)		0 (-0.08 – 0.09)		0.06 (-0.06 – 0.18)	
AD	-0.03 (-0.08 – 0.03)		0.01 (-0.04 – 0.06)		-0.01 (-0.11 – 0.09)		0.02 (-0.12 – 0.16)	
Observations	279		279		279		279	
R² / adjusted R²	0.089 / 0.075		0.059 / 0.045		0.123 / 0.111		0.101 / 0.088	

Table 5. Results of epigenetic age and epigenetic age acceleration calculation for all DNAme analyses, for both the larger DNAme cohort and the subset of samples with matched CSF biomarker data.

| | DNAme Cohort | | | | DNAme & CSF Biomarker Data Cohort | | | |
| | | Sex | | | | | Sex | |
	Total No. 640	Female No. 284	Male No. 356	P-value	Total No. 533	Female No. 243	Male No. 290	P-value
Age								
Mean (SD)	75.63 (±7.68)	74.78 (±8.03)	76.31 (±7.32)	<0.0001	75.01 (±7.61)	74.31 (±8.10)	75.61 (±7.11)	0.0019
Epigenetic Ag e (years)								
Mean (SD)	69.92 (±8.06)	67.45 (±8.15)	70.11 (±7.79)	<0.0001	68.47 (±8.17)	67.05 (±8.33)	69.72 (±7.82)	<0.0001
Epigenetic Age Acceleration (years)								
Mean (SD)	0.025 (±4.22)	-0.14 (±4.16)	0.16 (±4.26)	0.1	0.027 (±4.30)	-0.18 (±4.23)	0.20 (4.35)	0.057
Intrinsic Age Acceleration (years)								
Mean (SD)	0.026 (±4.11)	-0.19 (±4.06)	0.20 (±4.15)	0.019	0.020 (±4.18)	-0.25 (±4.14)	0.26 (±4.21)	0.021

P-values are from Wilcoxon rank sum tests for continuous variables and Fisher's exact tests for categorical variables

Table 6. Linear model for assessment of relationship of biochemical concentrations and **presence of APOE e4 risk allele** on universal epigenetic age acceleration. Intrinsic epigenetic age acceleration linear model not shown.

	Age Acceleration & CSF Biomarkers		
Predictors	*Estimates*	*CI*	*adjusted p*
(Intercept)	-0.61	-1.67 – 0.46	0.61
Male (ref = Female)	0.6	-0.11 – 1.31	0.34
Diagnosis (ref = CN)			
EMCI	0.71	-0.17 – 1.60	0.39
LMCI	0.4	-0.55 – 1.36	0.64
AD	0.53	-1.07 – 2.14	0.64
APOE risk allele			
1 or 2 alleles (ref = 0 alleles)	-0.24	-1.01 – 1.31	0.813
CSF Tau	0.0001	-0.0077 – 0.023	0.95
Observations			533
R^2 / adjusted R^2	0.0108/-0.000438		

Supplemental File

Alzheimer's disease neuropathology, inflammation and epigenetic age: do sex and APOE

matter?

Paula Duarte-Guterman[1*], Amy M. Inkster[2,3*], Arianne Y. Albert[4], Cindy K. Barha[5], Wendy P.

Robinson[2,3], Liisa A.M. Galea[1], on behalf of the Alzheimer's Disease Neuroimaging Initiative[δ]

1. Djavad Mowafaghian Centre for Brain Health and Department of Psychology, University of
   British Columbia, Vancouver, BC, Canada
2. Department of Medical Genetics, University of British Columbia, Vancouver, BC, Canada.
3. BC Children's Hospital Research Institute, Vancouver, BC, Canada
4. Women's Health Research Institute of British Columbia, Vancouver, BC, Canada
5. Djavad Mowafaghian Centre for Brain Health and Department of Physical Therapy,
   University of British Columbia, Vancouver, BC, Canada

Address all correspondence and requests for reprints to:
L. A. M. Galea, PhD
Djavad Mowafaghian Centre for Brain Health
2215 Wesbrook Mall
Vancouver, British Columbia
V6T 1Z3, Canada
E-mail: lgalea@psych.ubc.ca.

* Considered co-first authors.

[δ] Data used in preparation of this article were obtained from the Alzheimer's Disease
Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within
the ADNI contributed to the design and implementation of ADNI and/or provided data but did
not participate in analysis or writing of this report. A complete listing of ADNI investigators can
be found at: http://adni.loni.usc.edu/wp-
content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Table S1. Linear regression results for all variables investigated by sex and APOE status. Markers in CSF

Predictors	ADNI MEM Estimates (CI)	adjusted p	ADNI EF Estimates (CI)	adjusted p	ABETA Estimates (CI)	adjusted p	Hippocampus volume (corrected) Estimates (CI)	adjusted p	Hippocampus/Intracranial volume Estimates (CI)	adjusted p	TAU Estimates (CI)	adjusted p	PTAU Estimates (CI)	adjusted p	Cortisol Cortisol ng ml Estimates (CI)	adjusted p
(Intercept)	1.63 (1.12 – 2.14)		2.05 (1.47 – 2.62)		1458.67 (1247.77 – 1669.57)		(10935.47 – 12165.88)		0.00752 (0.00709 – 0.00795)		56.15 (-30.03 – 142.34)		4.75 (-4.89 – 14.38)		-1.38 (-9.15 – 6.39)	
AGE (years)	-0.02 (-0.02 – -0.01)		-0.02 (-0.03 – -0.02)		-6.19 (-9.01 – -3.36)		-65.72 (-73.95 – -57.49)		-0.00004 (-0.0004 – -0.0003)		2.68 (1.54 – 3.81)		0.26 (0.13 – 0.39)		0.21 (0.11 – 0.32)	
Male (ref = Female)	-0.17 (-0.26 – -0.07)	**0.003**	-0.03 (-0.14 – 0.08)	0.73	-29.77 (-71.55 – 12.01)	0.33	482.18 (365.03 – 599.32)	**<0.0001**	-0.00024 (-0.00033 – -0.00016)	**<0.0001**	-7.37 (-31.43 – 16.70)		-0.34 (-3.03 – 2.35)		1.37 (-0.05 – 2.78)	0.16
APOE status (ref = 0 alleles)		**<0.0001**		**<0.0001**		**<0.0001**		**<0.0001**		**<0.0001**						0.61
1 allele	-0.45 (-0.55 – -0.34)		-0.3 (-0.42 – -0.19)		-240.23 (-284.27 – -196.20)		-518.73 (-642.15 – -395.31)		-0.00031 (--0.0004 – -0.00022)		104.14 (77.21 – 131.06)		11.73 (8.72 – 14.74)		1.03 (-0.44 – 2.50)	
2 alleles	-0.69 (-0.85 – -0.53)		-0.46 (-0.64 – -0.28)		-455.95 (-521.02 – -390.88)		-946.01 (-1136.77 – -755.25)		-0.00057 (-0.00071 – --0.00044)		178.88 (137.45 – 220.31)		19.81 (15.18 – 24.44)		0.5 (-1.68 – 2.68)	
Interaction term												**0.001**		**0.002**		
Male:1 allele											-49.76 (-85.30 – -14.22)		-5.58 (-9.55 – -1.60)			
Male:2 alleles											-101.56 (-153.97 – -49.16)		-10.78 (-16.64 – -4.92)			
Observations		1145		1144		947		1224		1224		947		947		279
R² / adjusted R²	0.106 / 0.103		0.058 / 0.055		0.203 / 0.199		0.233/0.231		0.191/0.189		0.140 / 0.134		0.136 / 0.130		0.086 / 0.073	

Table S1 (continued). Linear regression results for all variables investigated by sex and APOE status. Markers in CSF

Predictors	C Reactive Protein ug/ml Estimates (CI)	adjusted p	CD 40 antigen ng/ml Estimates (CI)	adjusted p	Interleukin 16 pg/ml Estimates (CI)	adjusted p	Interleukin 3 ng/ml Estimates (CI)	adjusted p	Interleukin 6 receptor ng/ml Estimates (CI)	adjusted p	Interleukin 8 pg/ml Estimates (CI)	adjusted p	Immunoglobulin A mg/ml Estimates (CI)	adjusted p	Intercellular Adhesion Molecule ng/ml Estimates (CI)	adjusted p
(Intercept)	-3.05 (-3.79 – -2.32)		-1.13 (-1.29 – -0.98)		0.35 (0.10 – 0.59)		-2.96 (-3.38 – -2.54)		-0.29 (-0.48 – -0.10)		1.38 (1.18 – 1.57)		-2.82 (-3.22 – -2.43)		-0.19 (-0.75 – 0.36)	
AGE (years)	0.01 (-0.00 – 0.02)		0.01 (0.00 – 0.01)		0.01 (0.00 – 0.01)		0.01 (0.00 – 0.02)		0 (0.00 – 0.01)		0 (0.00 – 0.01)		0 (-0.00 – 0.01)		0.01 (0.01 – 0.02)	
Male (ref = Female)	-0.12 (-0.26 – 0.01)	0.19	0.01 (-0.02 – 0.04)	0.69	0.12 (0.06 – 0.18)		0.08 (0.00 – 0.16)	0.11	0.01 (-0.02 – 0.05)	0.64	0.1 (0.05 – 0.15)	**0.02**	0.21 (0.14 – 0.29)	**<0.0001**	0.18 (0.07 – 0.28)	**0.003**
APOE status (ref = 0 alleles)		**0.009**		0.8				0.41		0.17		0.39		0.33		0.38
1 allele	-0.19 (-0.33 – -0.05)		0.01 (-0.02 – 0.03)		0.08 (0.01 – 0.16)		-0.03 (-0.10 – 0.05)		0.04 (0.00 – 0.08)		0.04 (-0.02 – 0.10)		0.02 (-0.05 – 0.10)		0.09 (-0.01 – 0.20)	
2 alleles	-0.31 (-0.52 – -0.10)		0.02 (-0.03 – 0.06)		0.06 (-0.04 – 0.16)		-0.1 (-0.22 – 0.02)		0.04 (-0.02 – 0.09)		0.01 (-0.07 – 0.09)		-0.09 (-0.20 – 0.02)		0.02 (-0.13 – 0.18)	
Interaction term						**0.03**										
Male:1 allele					-0.13 (-0.22 – -0.03)											
Male:2 alleles					-0.15 (-0.28 – -0.02)											
Observations		279		279		279		279		279		279		279		279
R² / adjusted R²	0.058 / 0.045		0.124 / 0.111		0.117 / 0.098		0.081 / 0.068		0.045 / 0.031		0.092 / 0.072		0.135 / 0.122		0.107 / 0.094	

Table S2. Linear regression results for all variables investigated by sex and baseline diagnosis. Markers in CSF

Predictors	ADNI MEM Estimates (CI)	adjusted p	ADNI EF Estimates (CI)	adjusted p	ABETA Estimates (CI)	adjusted p	Hippocampus volume (corrected) Estimates (CI)	adjusted p	Hippocampus/Intracranial volume Estimates (CI)	adjusted p	TAU Estimates (CI)	adjusted p	PTAU Estimates (CI)	adjusted p	Cortisol Cortisol ng ml Estimates (CI)	adjusted p
(Intercept)	1.79 (1.45 – 2.13)		2.26 (1.80 – 2.73)		1161.57 (944.04 – 1379.10)		11424.98 (10886.11 – 11963.86)		0.00747 (0.00710 – 0.00785)		154.16 (70.57 – 237.75)		15.72 (6.36 – 25.09)		-1.87 (-9.56 – 5.81)	
AGE (years)	-0.01 (-0.01 – -0.00)		-0.02 (-0.03 – -0.01)		-1.64 (-4.52 – 1.24)		-57.95 (-65.07 – -50.83)		-0.00003 (-0.00004 – --0.00003)		1.25 (0.15 – 2.36)		0.1 (-0.02 – 0.23)		0.22 (0.12 – 0.32)	
Male (ref = Female)	-0.16 (-0.23 – -0.09)	**<0.0001**	-0.04 (-0.13 – 0.05)	0.56	-26.62 (-69.46 – 16.22)	0.42	516.27 (414.05 – 618.49)	**<0.0001**	-0.00022 (-0.00029 – -0.00015)	**<0.0001**	-42.59 (-59.05 – -26.13)	**<0.0001**	-4.22 (-6.06 – -2.38)	**<0.0001**	1.18 (-0.25 – 2.61)	0.25
Diagnosis (ref = CN)		**<0.0001**		**<0.0001**		**<0.0001**		**<0.0001**		**<0.0001**		**<0.0001**		**<0.0001**		0.5
EMCI	-0.5 (-0.61 – -0.39)		-0.42 (-0.56 – -0.27)		-85.2 (-148.82 – -21.59)		-340.14 (-495.23 – -185.04)		-0.00016 (-0.00027 – -0.00005)		37.85 (13.41 – 62.30)		4.22 (1.48 – 6.96)			
LMCI	-1.08 (-1.16 – -1.00)		-0.79 (-0.90 – -0.67)		-256.85 (-315.81 – -197.89)		-1049.98 (-1183.33 – -916.63)		-0.00073 (-0.00083 – -0.00064)		93.34 (70.69 – 116.00)		10.58 (8.05 – 13.12)		1.24 (-0.42 – 2.90)	
AD	-1.84 (-1.94 – -1.75)		-1.63 (-1.76 – -1.50)		-390.48 (-453.59 – -327.37)		-1603.56 (-1751.83 – -1455.29)		-0.00106 (-0.00116 – -0.00096)		143.6 (119.35 – 167.86)		15.81 (13.10 – 18.53)		0.24 (-1.68 – 2.16)	
Observations	1150		1149		947		1234		1234		947		947		279	
R^2 / adjusted R^2	0.589 / 0.588		0.380 / 0.377		0.168 / 0.164		0.416/ 0.414		0.398 / 0.396		0.164 / 0.160		0.156 / 0.152		0.088 / 0.075	

Table S2 (continued). Linear regression results for all variables investigated by sex and baseline diagnosis. Markers in CSF

Predictors	C Reactive Protein ug/ml Estimates (CI)	adjusted p	CD 40 antigen ng/ml Estimates (CI)	adjusted p	Interleukin 16 pg/ml Estimates (CI)	adjusted p	Interleukin 3 ng/ml Estimates (CI)	adjusted p	Interleukin 6.receptor ng/ml Estimates (CI)	adjusted p	Interleukin 8 pg/ml Estimates (CI)	adjusted p	Immunoglobulin A mg/ml Estimates (CI)	adjusted p	Intercellular Adhesion Molecule ng/ml Estimates (CI)	adjusted p
(Intercept)	-3.19 (-3.93 – -2.45)		-1.11 (-1.27 – -0.96)		0.42 (0.18 – 0.65)		-3 (-3.42 – -2.59)		-0.26 (-0.45 – -0.07)		1.4 (1.20 – 1.59)		-2.9 (-3.29 – -2.50)		-0.22 (-0.77 – 0.33)	
AGE (years)	0.01 (-0.00 – 0.02)		0.01 (0.00 – 0.01)		0.01 (0.00 – 0.01)		0.01 (0.00 – 0.02)		0 (0.00 – 0.01)		0 (0.00 – 0.01)		0 (-0.00 – 0.01)		0.01 (0.01 – 0.02)	
Male (ref = Female)	-0.1 (-0.24 – 0.03)	0.31	0.01 (-0.02 – 0.03)	0.81	0.05 (0.01 – 0.10)	0.07	0.08 (0.01 – 0.16)	0.11	0.01 (-0.02 – 0.05)	0.71	0.05 (0.01 – 0.09)	**0.03**	0.21 (0.14 – 0.29)	**<0.0001**	0.17 (0.06 – 0.27)	**0.008**
Diagnosis (ref = CN)		0.32		0.2		0.66		0.66		0.67		0.94		0.98		0.71
EMCI																
LMCI	-0.15 (-0.31 – 0.01)		0.01 (-0.03 – 0.04)		0.01 (-0.04 – 0.06)		-0.04 (-0.13 – 0.05)		0.01 (-0.03 – 0.05)		0.01 (-0.03 – 0.05)		0 (-0.08 – 0.09)		0.06 (-0.06 – 0.18)	
AD	-0.15 (-0.33 – 0.04)		-0.03 (-0.07 – 0.01)		-0.03 (-0.08 – 0.03)		-0.06 (-0.17 – 0.04)		-0.02 (-0.07 – 0.03)		0.01 (-0.04 – 0.06)		-0.01 (-0.11 – 0.09)		0.02 (-0.12 – 0.16)	
Observations	279		279		279		279		279		279		279		279	
R^2 / adjusted R^2	0.030 / 0.016		0.138 / 0.125		0.089 / 0.075		0.077 / 0.063		0.031 / 0.017		0.059 / 0.045		0.123 / 0.111		0.101 / 0.088	

Table S3. Linear regression results for all plasma variables investigated by sex and APOE genotype

Predictors	Interleukin 18 pg/ml Estimates (CI)	adjusted p	Cortisol Cortisol ng/ml Estimates (CI)	adjusted p	C Reactive Protein ug/ml Estimates (CI)	adjusted p	Intercellular Adhesion Molecule ng/ml Estimates (CI)	adjusted p	Immunoglobulin E ng/ml Estimates (CI)	adjusted p	Interleukin 8 pg/ml Estimates (CI)	adjusted p
(Intercept)	2.41 (2.25 – 2.57)		2.04 (1.92 – 2.16)		0.06 (-0.39 – 0.52)		1.91 (1.78 – 2.04)		1.66 (1.18 – 2.14)		0.79 (0.62 – 0.96)	
AGE (years)	0 (-0.00 – 0.00)		0 (-0.00 – 0.00)		0 (-0.00 – 0.01)		0 (-0.00 – 0.00)		0 (-0.01 – 0.01)		0 (0.00 – 0.01)	
Male (ref = Female)	0.06 (0.03 – 0.09)	**0.0007**	0.01 (-0.01 – 0.04)	0.45	-0.15 (-0.25 – -0.06)	**0.003**	-0.04 (-0.06 – -0.01)	**0.03**	0.25 (0.15 – 0.34)	**<0.0001**	-0.01 (-0.04 – 0.03)	0.76
APOE status (ref = 0 alleles)		0.17		0.33		**<0.0001**		0.83		0.36		0.29
1 allele	-0.03 (-0.07 – -0.00)		0.01 (-0.01 – 0.04)		-0.28 (-0.37 – -0.18)		0.01 (-0.02 – 0.04)		0.03 (-0.06 – 0.13)		-0.03 (-0.07 – 0.00)	
2 alleles	-0.04 (-0.09 – 0.01)		0.03 (-0.00 – 0.07)		-0.36 (-0.50 – -0.23)		0 (-0.04 – 0.04)		-0.1 (-0.25 – 0.04)		-0.03 (-0.09 – 0.02)	
Observations	527		527		526		527		527		527	
R² / adjusted R²	0.038 / 0.031		0.014 / 0.007		0.105 / 0.098		0.019 / 0.012		0.056 / 0.048		0.027 / 0.019	

Table S3 (continued). Linear regression results for all plasma variables investigated by sex and APOE genotype

Predictors	CD 40 antigen ng/ml Estimates (CI)	adjusted p	Interleukin 16 pg/ml Estimates (CI)	adjusted p	Interleukin 3 ng/ml Estimates (CI)	adjusted p	Interleukin 6 receptor ng/ml Estimates (CI)	adjusted p	Immunoglobulin A mg/ml Estimates (CI)	adjusted p
(Intercept)	-0.58 (-0.68 – -0.47)		2.32 (2.18 – 2.47)		-1.71 (-1.98 – -1.45)		1.42 (1.30 – 1.55)		0.48 (0.28 – 0.69)	
AGE (years)	0.01 (0.00 – 0.01)		0 (0.00 – 0.00)		0 (-0.00 – 0.00)		0 (-0.00 – 0.00)		0 (-0.00 – 0.00)	
Male (ref = Female)	-0.01 (-0.03 – 0.01)	0.58	0.01 (-0.02 – 0.04)	0.65	0 (-0.06 – 0.05)	0.89	-0.03 (-0.05 – -0.00)	0.1	0.02 (-0.02 – 0.06)	0.61
APOE status (ref = 0 alleles)		0.61		0.69		0.7		0.4		0.65
1 allele	-0.01 (-0.03 – 0.01)		-0.02 (-0.05 – 0.01)		0.01 (-0.05 – 0.06)		0.01 (-0.01 – 0.04)		-0.01 (-0.05 – 0.03)	
2 alleles	0.01 (-0.02 – 0.04)		-0.01 (-0.06 – 0.03)		0.04 (-0.03 – 0.12)		-0.02 (-0.06 – 0.02)		-0.04 (-0.10 – 0.02)	
Observations	526		527		527		527		527	
R² / adjusted R²	0.133 / 0.126		0.023 / 0.015		0.002 / -0.005		0.016 / 0.008		0.009 / 0.002	

Table S4. Linear regression results for all plasma variables investigated by sex and baseline diagnosis.

Predictors	Interleukin 18 pg/ml Estimates (CI)	adjusted p	Cortisol Cortisol ng/ml Estimates (CI)	adjusted p	C Reactive Protein ug/ml Estimates (CI)	adjusted p	Intercellular Adhesion Molecule ng/ml Estimates (CI)	adjusted p	Immunoglobulin E ng/ml Estimates (CI)	adjusted p	Interleukin 8 pg/ml Estimates (CI)	adjusted p
(Intercept)	2.33 (2.17 – 2.49)		2.08 (1.96 – 2.20)		-0.07 (-0.56 – 0.41)		1.92 (1.78 – 2.05)		1.6 (1.11 – 2.09)		0.78 (0.61 – 0.96)	
AGE (years)	0 (-0.00 – 0.00)		0 (-0.00 – 0.00)		0.01 (0.00 – 0.01)		0 (-0.00 – 0.00)		0 (-0.01 – 0.01)		0 (0.00 – 0.01)	
Male (ref = Female)	0.06 (0.03 – 0.09)	**0.001**	0.02 (-0.01 – 0.04)	0.35	-0.14 (-0.24 – -0.05)	**0.01**	-0.03 (-0.06 – -0.01)	0.05	0.25 (0.15 – 0.34)	**<0.0001**	0 (-0.04 – 0.03)	0.89
Diagnosis (ref = CN)		0.56		**0.005**		**0.04**		0.18		0.91		0.28
LMCI	0.04 (-0.01 – 0.10)		-0.02 (-0.07 – 0.02)		-0.26 (-0.44 – -0.09)		-0.01 (-0.06 – 0.03)		-0.01 (-0.18 – 0.17)		-0.04 (-0.10 – 0.02)	
AD	0.03 (-0.03 – 0.10)		0.03 (-0.02 – 0.08)		-0.22 (-0.41 – -0.03)		0.02 (-0.03 – 0.08)		-0.04 (-0.23 – 0.16)		0 (-0.07 – 0.06)	
Observations	527		527		526		527		527		527	
R^2 / adjusted R^2	0.032 / 0.025		0.034 / 0.026		0.043 / 0.036		0.029 / 0.021		0.050 / 0.043		0.027 / 0.020	

Table S4 (continued). Linear regression results for all plasma variables investigated by sex and baseline diagnosis.

Predictors	CD 40 antigen ng/ml Estimates (CI)	adjusted p	Interleukin 16 pg/ml Estimates (CI)	adjusted p	Interleukin 3 ng/ml Estimates (CI)	adjusted p	Interleukin 6 receptor ng/ml Estimates (CI)	adjusted p	Immunoglobulin A mg/ml Estimates (CI)	adjusted p
(Intercept)	-0.56 (-0.67 – -0.46)		2.38 (2.23 – 2.53)		-1.62 (-1.89 – -1.35)		1.44 (1.32 – 1.56)		0.42 (0.22 – 0.63)	
AGE (years)	0.01 (0.00 – 0.01)		0 (0.00 – 0.00)		0 (-0.00 – 0.00)		0 (-0.00 – 0.00)		0 (-0.00 – 0.00)	
Male (ref = Female)	-0.01 (-0.03 – 0.01)	0.62	0.01 (-0.02 – 0.04)	0.58	-0.01 (-0.06 – 0.05)	0.91	-0.02 (-0.05 – -0.00)	0.14	0.02 (-0.02 – 0.06)	0.61
Diagnosis (ref = CN)		0.06		0.06		0.1		0.56		0.84
LMCI	-0.02 (-0.05 – 0.02)		-0.08 (-0.13 – -0.02)		-0.04 (-0.14 – 0.06)		-0.03 (-0.08 – 0.01)		0.03 (-0.05 – 0.10)	
AD	0.02 (-0.02 – 0.06)		-0.08 (-0.14 – -0.02)		-0.12 (-0.22 – -0.01)		-0.03 (-0.08 – 0.02)		0.02 (-0.07 – 0.10)	
Observations	526		527		527		527		527	
R^2 / adjusted R^2	0.143 / 0.137		0.036 / 0.029		0.013 / 0.006		0.014 / 0.006		0.008 / -0.000	

Supplementary Table S5. Correlation matrix of biochemical variables for those ADNI samples with available DNAme and biochemical data (n=533). Correlation coefficients are Spearman's rho.

	Amyloid Beta	p value	Tau	p value	pTau	p value
Amyloid Beta	1.00	<0.0001	-0.18	<0.0001	-0.27	<0.0001
Tau	-0.18	<0.0001	1.00	<0.0001	0.98	<0.0001
pTau	-0.27	<0.0001	0.98	<0.0001	1.00	<0.0001