Inflammation and epigenetic age in Alzheimer's disease: do sex and APOE matter?

Paula Duarte-Guterman[1*], Amy M. Inkster[2,3*], Arianne Y. Albert[4], Cindy K. Barha[5], Wendy P. Robinson[2,3], Liisa A.M. Galea[1], on behalf of the Alzheimer's Disease Neuroimaging Initiative[δ]

1. Djavad Mowafaghian Centre for Brain Health and Department of Psychology, University of British Columbia, Vancouver, BC, Canada

2. Department of Medical Genetics, University of British Columbia, Vancouver, BC, Canada.

3. BC Children's Hospital Research Institute, Vancouver, BC, Canada

4. Women's Health Research Institute of British Columbia, Vancouver, BC, Canada

5. Djavad Mowafaghian Centre for Brain Health and Department of Physical Therapy, University of British Columbia, Vancouver, BC, Canada

Address all correspondence and requests for reprints to:
L. A. M. Galea, PhD
Djavad Mowafaghian Centre for Brain Health
2215 Wesbrook Mall
Vancouver, British Columbia
V6T 1Z3, Canada
E-mail: lgalea@psych.ubc.ca.

* Considered co-first authors.

[δ] Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

ABSTRACT

Alzheimer's disease (AD) disproportionately affects females that carry the APOEε4

allele, and the progression from cognitively normal to mild cognitive impairment (MCI) and to

AD may be sex-specific. The objective of this study was to investigate underlying physiological

inflammatory and epigenetic mechanisms that may help explain why there are sex differences in

AD and APOEε4 carriers. Inflammation is observed in AD and is related to aging, stress and

neuroplasticity, and aging biomarkers also include changes in DNA methylation patterns which

have been used to determine epigenetic age. Although studies are scarce, sex differences are

noted in inflammation and in epigenetic aging. We investigated using the ADNI database the

effect of sex and APOE genotype (non-carriers or carriers of 1 or 2 APOEε4 alleles) and sex and

diagnosis (cognitively normal (CN), MCI, AD) on CSF and plasma markers of stress and

inflammation, and blood epigenetic age. We found sex and APOE genotype interactions on CSF

levels of IL-8 and IL-16 cytokines suggesting that the presence of APOEε4 alleles differentially

affects male and female physiology. Furthermore, females had higher levels of plasma CRP and

ICAM1 but lower levels of CSF ICAM1, IL-8, IL-16, and IgA than males, showing in part that

results may differ by sample source (plasma and CSF). Carrying 1 or 2 of APOEε4 alleles and

diagnosis (MCI and AD) decreased plasma CRP in both sexes. Neither sex, diagnosis, nor

presence of APOEε4 alleles affected epigenetic age acceleration. Sex differences in

inflammatory biomarkers support that the underlying physiological changes during aging differ

by sex and tissue origin.

Keywords: Sex differences; Alzheimer's disease; APOE genotype; inflammation; cytokines;

epigenetic age.

INTRODUCTION

Alzheimer's disease (AD) is a neurodegenerative disease characterized by severe cognitive decline (Alzheimer's Association, 2017). Modifiable risk factors associated with AD include stress (Caruso et al., 2018), sociocultural or lifestyle factors (e.g., education, marital status, exercise), and medical conditions (diabetes, obesity, and cardiovascular disease) (Baumgart et al., 2015; Nebel et al., 2018; Xu et al., 2015). Non-modifiable risk factors include age, biological sex, and APOE genotype (Riedel et al., 2016). In some studies, females show greater neuropathology (brain atrophy, neurofibrillary tangles) and cognitive decline than males, although this depends on whether females are already cognitively impaired (mild cognitive impairment (MCI) or AD) and on the presence of an APOEε4 allele, a well-known genetic risk factor of AD (Ardekani et al., 2016; Barnes et al., 2005; Holland et al., 2013; Hua et al., 2010; Irvine et al., 2012; Koran and Hohman, 2017; Lin et al., 2015; Wang et al., 2019). Previous studies using data from the Alzheimer's Disease Neuroimaging Initiative (ADNI) indicate that females with AD have greater brain atrophy rates and cognitive decline than males with AD (Holland et al., 2013; Hua et al., 2010; Lin et al., 2015). However, there is limited research into the role of sex and its interaction with APOE genotype in the possible mechanisms underlying AD. However, females with one APOEε4 allele present have a greater risk to develop AD at an earlier age (Neu et al., 2017), show greater cognitive decline and neuropathology associated with AD relative to males (Beydoun et al., 2012; Hohman et al., 2018; Koran et al., 2017), indicating that the APOE genotype interacts with sex on various factors related to AD. Understanding why females are at a higher risk and have a higher burden of the disease is important for the development of tailored treatments. Biomarkers are highly sought after to predict disease onset and progression and to understand the underlying mechanisms of diseases in order to develop or improve treatments.

Chronic inflammation is a hallmark of AD, as evidenced by increased expression of

proinflammatory cytokines in the brains of AD patients (not analyzed by sex), which can

exacerbate AD pathology (Heppner et al., 2015; Kinney et al., 2018; Swardfager et al., 2010).

There is, however, increasing evidence that there are sex differences in immune responses in

healthy adults with females mounting a stronger response compared to males after an acute

challenge (Engler et al., 2016; Klein and Flanagan, 2016). In response to an endotoxin, females

have higher levels of proinflammatory plasma cytokines (TNF-α and IL-6) while males have

higher plasma levels of anti-inflammatory IL-10 (Engler et al., 2016; Wegner et al., 2017). In

addition, sex differences in inflammation are associated with AD pathology (Walker et al., 2018)

but not cognitive decline in normal aging (Walker et al., 2019). Specifically, higher systemic

inflammation beginning in mid-life is associated with higher brain amyloid levels later in life in

healthy males, but not females (Walker et al., 2018). To our knowledge, very few studies have

stratified by sex and APOE genotype or sex and diagnosis of cognitive status on potential

biomarkers of AD.

Sex differences in inflammatory biomarker systems may also differentially affect

neuroplasticity (Dantzer, 2018; de Miranda et al., 2017), which is reduced in AD and correlates

with cognitive decline (Apostolova et al., 2006; Petersen et al., 2000). In addition, peripheral

cortisol, the main stress hormone in humans, is elevated in AD (Doecke et al., 2012) and is

associated with higher amyloid levels in the brain (Toledo et al., 2012), a reduction in

hippocampal volume, and cognitive impairment in older individuals (Lupien et al., 1998) that

may depend on MCI status (Murphy et al., 2019). Peripheral cortisol is also associated with

elevated pro-inflammatory cytokines (Engler et al., 2016; McEwen et al., 1997). However, it is

not known how sex differences in markers of inflammation (e.g., cytokines, immunoglobulins,

CRP, intercellular adhesion molecule, ICAM1), and stress hormones (cortisol) may be related to

sex differences in AD.

Aging biomarkers include epigenetic alterations that are associated with a variety of

pathologies and adverse health conditions, including normal cognitive aging and

neurodegenerative phenotypes such as AD (Hannum et al., 2013; Horvath, 2013; Levine et al.,

2015; Yokoyama et al., 2017). Recently, molecular biomarkers of aging known as "epigenetic

clocks" have been developed based on DNA methylation (DNAme) signatures (Hannum et al.,

2013; Horvath, 2013). Epigenetic age or "DNAmAge" is a measure of the biological age of a

sample (cell or tissue), and can be calculated across a range of tissues and time points, providing

an accurate estimation of a sample's chronological age based on the presence or absence of

methylation at the 5' carbon of cytosine in informative cytosine-guanine (CpG) dinucleotides

throughout the human genome (Horvath, 2013). Positive deviations of epigenetic age from

chronological age (positive epigenetic age acceleration) reflect more rapid biological aging and

have been associated with numerous factors including smoking, obesity, Parkinson's disease,

Trisomy 21, and cancer (Gale et al., 2018; Horvath, 2013; Horvath et al., 2015; Horvath and

Ritz, 2015), while negative deviations of epigenetic age from chronological age (negative

epigenetic age acceleration) have been associated with high life-expectancy populations and

memory retention (Degerman et al., 2017; McEwen et al., 2017). In AD, epigenetic age

acceleration of the frontal cortex was associated with amyloid load, neuritic plates, and cognitive

decline (Levine et al., 2015). Within an individual, the DNA methylation profile in peripheral

tissues is correlated with the epigenetic signature in the brain, likely due to identical genetic

background affecting DNAme and common signatures of epigenetic aging (Braun et al., 2019),

thus it is reasonable to hypothesize that epigenetic age acceleration may also be detectable in

peripheral tissues such as blood in AD participants. In healthy individuals, aging males exhibit

more positive epigenetic age acceleration than females in blood, buccal tissue, and in multiple

brain regions (Horvath et al., 2016). In AD, it is possible that the underlying sex-specific

pathological mechanisms may be reflected in epigenetic age acceleration measures, and given

the complexity of the sex-specific phenotypes of AD, this may be observed as either sex showing

accelerated epigenetic age – for example, AD females could potentially have more positive

epigenetic age acceleration than males.

Using the ADNI database, we examined sex differences in plasma and CSF physiological

biomarkers (inflammation, stress, related) and blood epigenetic age, and how these may be

affected by APOE genotype (non-carriers or carriers of 1 or 2 APOEε4 alleles), and dementia

status (cognitively healthy (CN), MCI, AD). We tested the hypotheses that (1) females have

higher levels of inflammation and stress hormones compared to males and these levels are

disproportionately affected by the presence of APOEε4 alleles and AD diagnosis; and (2)

females have more positive epigenetic age acceleration and this is exacerbated by APOEε4

carriership and AD diagnosis.

<u>METHODS</u>

ADNI database

Data used in the preparation of this article were obtained from the Alzheimer's Disease

Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). The ADNI was launched in

2003 as a public-private partnership, led by Principal Investigator Michael W. Weiner,

MD. The primary goal of ADNI has been to test whether serial magnetic resonance imaging

(MRI), positron emission tomography (PET), other biological markers, and clinical and

neuropsychological assessment can be combined to measure the progression of mild cognitive

impairment (MCI) and early Alzheimer's disease (AD). For up-to-date information, see

www.adni-info.org. Data used in this article were downloaded on or before Jan 16, 2019.

Inclusion and exclusion criteria of participants (Petersen et al., 2010; Aisen et al., 2010) and

general procedures are detailed online (http://adni.loni.usc.edu/methods/documents/). Briefly,

cognitively normal (CN) participants had normal memory function based on education-adjusted

scores on the Wechsler Memory Scale Logical Memory II and a Clinical Dementia Rating

(CDR) of 0. Amnestic late MCI (LMCI) participants had objective memory loss (measured by

education-adjusted scores from Wechsler Memory Scale Logical Memory II), a CDR of 0.5,

preserved daily activities, and absence of dementia. Early MCI (EMCI) participants had a milder

episodic memory impairment compared to LMCI and were recruited using the Wechsler

Memory Scale Logical Memory II scores (between ~0.5 and 1.5 SD below the mean of CN

participants). All AD participants met NINCDS/ADRDA Alzheimer's Criteria and a CDR of 0.5

or 1.0.

Statistical Methods: Inflammatory markers

We included all ADNI participants that had inflammatory markers measured in CSF (N =

279) and plasma (N = 527) listed in Table 1. Data included in our analyses were: demographics

(age, years of education, and ethnicity), baseline diagnosis (cognitively normal, CN; late MCI,

LMCI; or AD), and number of APOEε4 alleles. We collapsed APOE genotype into two groups:

(1) participants carrying any ε4 alleles (homozygous ε4/ε4 and heterozygous ε4/-) and (2)

participants with no ε4 risk alleles (-/-). Plasma and CSF samples from the ADNI study were

collected in CN, LMCI and AD participants at baseline in the morning after an overnight fast.

Processing, aliquoting and storage were performed according to the ADNI Biomarker Core

Laboratory Standard Operating Procedures. Inflammatory markers were measured using a

commercially available multiplex proteomic panel (Human Discovery Multi-Analyte Profile;

Luminex xMAP) developed by Rules-Based Medicine (Austin, TX), that measures a variety of

markers including cytokines, metabolic markers, and growth factors. The protocols used to

quantify plasma and CSF analytes are described in Craig-Schapiro et al. (2011) and Hu et al.

(2012). We used the ADNI quality-controlled data for plasma and CSF provided by the ADNI

Consortium. For plasma IL-16, we removed one outlier that was more than two times lower than

the 25^{th} percentile in the plasma data. Sensitivity analysis with the outlier present suggested that

it was disproportionately influencing the results.

We compared all available data for each study variable between the sexes using the

Wilcoxon rank sum test for continuous variables and Fisher's exact test for categorical variables.

We used general linear models to determine the relationships between (1) sex and APOE

genotype (non-carriers or carriers of 1 or 2 APOEε4 alleles) or (2) sex and baseline diagnosis as

predictor variables, and biomarkers as dependent variables. Due to the limited sample size, we

were not able to study sex, APOE genotype and baseline diagnosis in one model. All models

included age and education as covariates. All models initially included an interaction between

sex and presence of APOEε4 alleles or sex and baseline diagnosis; if this interaction was not

significant, it was removed from the model to estimate the main effects of sex and APOE

genotype or diagnosis. Significance was based on the likelihood ratio test, and all P-values for

comparisons of sex and either APOE genotype or diagnosis for all outcomes combined were

corrected for multiple testing using the Benjamini-Hochberg false discovery rate method with the

family-wise error rate set to 0.05 (Benjamini and Hochberg, 1995). In total, three P-values per

dependent variable were included in each set of models (interaction term and main effects of sex

and APOE or diagnosis) resulting in 27 P-values corrected in CSF (9 dependent variables) and

33 P-values corrected in plasma (11 dependent variables; Supplementary Tables S1 to S4) for

each of the models (sex and APOE and sex and diagnosis). Significant interaction terms were

followed up using pairwise simple-effects tests with Benjamini-Hochberg p-value correction. We

report significance differences (adjusted $P<0.05$) and trends (adjusted $P<0.08$). All regression

analyses were carried out in R v3.5.1 (R Core Team, 2018).

Statistical Methods: Epigenetic Age

We used DNAme data quantified with the Illumina Infinium HumanMethylationEPIC

BeadChip array ("EPIC" array) for 1905 longitudinal blood samples from 640 unique ADNI

participants with CN, EMCI, LMCI, and AD diagnosis (n=284 females, n= 356 males)

(Vasanthakumar et al., 2017). DNAme IDAT files were read into R v3.5.1 (R Core Team, 2018)

using the 'minfi' package, and annotated with the most recent version of the EPIC manifest, the

Infinium MethylationEPIC v1.0 B4 Manifest File, (available from

https://support.illumina.com/downloads.html) (Aryee et al., 2014; Fortin et al., 2017). We

excluded 11 low quality samples from 9 unique participants from further analyses on the basis of

having median methylated and unmethylated probe intensity <10.5 (Aryee et al., 2014; Fortin et

al., 2017), the remaining samples were background normalized and dye-bias adjusted with

normal exponential out-of-band ("noob") normalization (Triche et al., 2013). DNAme data were

converted to beta values and the biological sex of all samples was confirmed by clustering

samples on all array probes mapping to the X and Y chromosomes. Beta values were calibrated

to Horvath's 21,368-probe training dataset using the calibrated beta-mixture quantile (BMIQ)

normalization procedure suggested by Horvath, and epigenetic age was calculated using R code

modified for compatibility with the EPIC array using the 334/353 epigenetic clock probes

present on the array from https://horvath.genetics.ucla.edu/ (Horvath, 2013; Teschendorff et al.,

2013). Epigenetic age could have been calculated by imputing missing DNAme values at the 19

clock CpG sites not covered by the EPIC array based on the k-nearest neighbors method, but as

we observed a high correlation between epigenetic age values calculated with the missing probes

removed versus imputed with k=10 (R=0.99, p<2.2e-16), in agreement with previous reports

(Fiorito et al., 2017; McEwen et al., 2018), we therefore chose to remove missing probes.

Prior to statistical analyses we removed all technical replicate samples. Epigenetic age

acceleration was calculated as the residual of epigenetic age regressed on chronological age and

technical batch covariates: the laboratory collection site at which blood samples were drawn and

EPIC microarray chip and row. Intrinsic epigenetic age acceleration, a measure designed to be

independent of age-related changes in whole blood cell-type proportions, was calculated as

described in Chen et al. (2016) as the residual of epigenetic age regressed on chronological age,

technical covariates of collection site, row, and chip, and the proportions of six blood cell types

(CD8T, CD4T, NK, B cells, monocytes, and granulocytes) estimated from noob-normalized

DNA methylation data with the Houseman algorithm (Houseman et al., 2012). ADNI is a

longitudinal study, and participants contributed between 1-3 blood DNAme samples within the

2-year collection period. Because these data collected over two years fell within the median 3.6-

year error of Horvath's epigenetic clock, longitudinal evaluation could not be meaningfully

conducted. As such, for all epigenetic age measures reported and analyzed herein, we have

calculated the average epigenetic age acceleration measure per participant from all available time

points, and performed all statistical analyses on these mean values.

To determine if epigenetic age acceleration or intrinsic epigenetic age acceleration

differed by sex or dementia diagnosis, we used unbalanced two-way ANOVA designs. APOEε4

genotype and CSF biomarker (amyloid beta, tau and p-tau) data were available from the ADNI

repository for a smaller subset of participants with matched EPIC DNAme data, (n=533, see

Table 1C). As above, due to the small size of this cohort, we collapsed APOE genotype into two

groups: (1) non-carriers of ε4 risk alleles and (2) carriers of ε4 alleles. We subsequently used

linear regression to test whether presence of an APOEε4 allele, tau, dementia diagnosis, or sex

were significantly associated with epigenetic age acceleration. Amyloid beta and p-tau were not

included in this model due to collinearity between amyloid beta, tau, and p-tau CSF

concentrations (see Supplementary Table S5).

RESULTS

Demographic information

Table 1 gives a summary of the variables for the participants with: CSF biomarkers

(Table 1A; N=279), whole blood EPIC DNAme data (Table 1B; N=640) and matched EPIC data

and measured CSF biomarkers (Table 1C; N=533). Given the differences in sample sizes, we

performed demographic analyses on the three datasets. In the biomarker dataset, females were

significantly younger and had fewer years of education than males ($P<0.01$ and $P<0.0001$,

respectively). Although there were no sex differences in distribution of APOEε4 alleles, the

proportion of participants in each of the diagnosis categories was significantly different for

females and males ($P=0.051$; Table 1A). For the data applicable to the participants with available

EPIC DNAme data (Table 1B) and participants with matched EPIC DNAme data and CSF

biomarkers (Table 1C), most of the demographics were similar to the biomarker dataset with the

exception that the proportion of participants in each of the diagnosis categories was not

significantly different between females and males.

Sex and presence of APOEε4 alleles are associated with changes in inflammatory markers

Our first aim was to investigate whether sex and APOE genotype interact to influence

inflammation using biomarkers, which we analysed separately in CSF and plasma

(Supplementary Table S1 and S3, respectively). Caution should be noted as inflammatory

signalling can differ depending on tissue examined (Bettcher et al., 2018; Setiawan et al., 2015).

For inflammatory markers measured in CSF, only IL-16 and IL-8 elicited a significant

interaction between sex and APOE genotype for CSF IL-16 and IL-8 (P= 0.016 and P=0.035,

respectively; Table 2). CSF IL-16 and IL-8 levels were significantly lower in females non-

carriers of APOEε4 alleles compared to males (both P's<0.001), whereas levels were similar

between the sexes in carriers of APOEε4 alleles (P's>0.9; Fig 1 A and B). Furthermore, in

females with APOEε4 alleles, IL-16 was significantly higher than in non-APOEε4 female

carriers (P=0.050), while a trend was observed in males (P=0.062). Whereas for IL-8, males with

APOEε4 alleles had lower levels of IL-8 compared to males with no APOEε4 alleles (P=0.014)

but there was no difference in females (P>0.3). Regardless of sex, CSF CRP levels were lower in

individuals with 1 or 2 APOEε4 alleles compared to non-carriers (main effect of genotype:

P=0.009; Table 2; Fig 1 C). There was a trend for an increase in IL-6 receptor levels in APOEε4

carriers regardless of sex compared to non APOEε4 carriers (main effect of genotype: P=0.071;

Table 2). Lastly females had significantly lower CSF levels of IgA and ICAM1 and a trend for

lower CSF cortisol levels compared to males (main effect of sex: P<0.001; P=0.009, and

P=0.070, respectively; Table 2). There were no other significant main or interaction effects on

any other CSF biomarkers.

For biomarkers measured in plasma, there were no significant interactions between sex

and APOE genotype (Supplementary Table S3). However, females had higher plasma CRP

levels (main effect of sex: adjusted P=0.048; Fig 1 D) and ICAM1 (main effect of sex: P=0.051)

compared to males and significantly lower levels of IL-18 (main effect of sex: P=0.001; Fig 1 E)

and immunoglobulin E (IgE: main effect of sex: P<0.001; Fig 1 F) compared to males.

Furthermore, plasma CRP decreased in carriers of APOEε4 alleles compared to non-carriers

(main effect of genotype: P<0.001; Fig 1 D).

Sex and baseline diagnosis are associated with changes in inflammatory markers

We next tested whether sex and baseline dementia status (CN, MCI, and AD) influenced

CSF and plasma biomarkers of inflammation (Supplementary Table S2 and S4, respectively).

There were no significant interactions between sex and diagnosis for any of the tested variables

in CSF (Table 3 and Supplementary Table S2) or plasma (Supplementary Table S4). For CSF

levels, females had significantly lower levels of IgA (main effect of sex: P<0.001) and ICAM1

(main effect of sex: P=0.026) and a trend for lower IL-16 levels (main effect of sex: P=0.055)

compared to males (Table 3; Fig 2 A-C), but we did not observe any significant main effects of

diagnosis for any CSF variable.

In plasma, we found that females had lower levels of IgE (main effect of sex: P<0.001)

and IL-18 compared to males (main effect of sex: P=0.004) and trends for females to have higher

levels of ICAM1 (main effect of sex: P=0.056) and CRP (main effect of sex: P=0.056; Fig. 2 D)

compared to males. In addition, we found diagnosis significantly influenced plasma cortisol

(main effect of baseline diagnosis: P=0.01) with lower levels in LMCI compared to AD

(P<0.001; Fig 2 E). We found trends for diagnosis to influence plasma IL-16, CRP and CD 40

levels (main effect of diagnosis: P=0.054, P=0.056; P=0.067). Plasma IL-16 (P's=0.006) and

CRP (P=0.006 and P=0.02) levels were lower in LMCI and AD compared to CN (Fig 2 D and

F). For plasma CD 40, levels were lower in LMCI compared to AD (P=0.01; Supplementary

Table S4). In summary, although we detected associations between sex and diagnosis and

various biomarkers, we did not find evidence of a sex and diagnosis interaction on any variables

examined. Furthermore, the results for inflammatory markers in plasma did not always match

results in CSF (Supplementary Tables S2 and S4).

Epigenetic age, sex, dementia diagnosis, and AD biomarkers

We investigated the hypothesis that sex and dementia diagnosis affect epigenetic age

acceleration in blood samples of ADNI participants (Table 4). Epigenetic age acceleration was

not associated with sex, dementia diagnosis (CN, EMCI, LMCI, and AD), nor the interaction of

sex and diagnosis after multiple test correction (Fig 3). Intrinsic epigenetic age acceleration was

also not significantly associated with participant sex, diagnosis, or their interaction term.

To assess the effect of sex and more broadly defined dementia-associated cognitive

impairment on epigenetic age acceleration, we compared epigenetic age acceleration between

participants with any form of clinically ascertained cognitive impairment (AD + LMCI + EMCI,

N=423, proportion female 41%) and those without (CN, N=217, proportion female 50%). By

two-way unbalanced ANOVA neither sex, dementia status, nor their interaction were

significantly associated with epigenetic age acceleration after correction for multiple

comparisons.

Matched biochemical data including APOEε4 genotype and CSF concentrations of

amyloid beta, tau, and p-tau was available for a subset of participants with EPIC DNAme data

(N=533). Based on the hypothesis that epigenetic age acceleration may be more strongly

associated with concentrations of pathologically relevant compounds than with diagnosis, we

assessed the impact of sex, APOEε4 genotype, and CSF tau concentration on epigenetic age

acceleration and intrinsic epigenetic age acceleration with linear regression. None of these

variables were significantly associated with epigenetic age acceleration (Table 5; results for

intrinsic epigenetic age acceleration not shown).

In addition to dementia diagnosis for all participants, we also had access to two

composite scores designed by ADNI collaborators to reflect (i) executive function and (ii)

memory; these z-scores have been demonstrated to be independently predictive of the transition

from MCI to a formal diagnosis of AD (Crane et al., 2012; Gibbons et al., 2012). By two-way

unbalanced ANOVA models investigating the effect of sex and memory z-score on epigenetic

age acceleration, neither sex (P=0.248), memory score (P=0.486), nor their interaction (P=0.227)

were associated with epigenetic age acceleration. In a similar model, neither sex (P=0.260),

executive function (P=0.105), or the interaction term of sex and executive function (P=0.153)

were associated with epigenetic age acceleration.

DISCUSSION

In the present study using ADNI data from CN, MCI and AD participants we found

interactions between sex and APOE genotype (but not between sex and diagnosis) on CSF and

plasma levels of IL-8 and IL-16. CSF levels of IL-8 and IL-16 were lower in female APOEε4

non-carriers compared to males but similar levels were found between the sexes with 1 or 2

APOEε4 alleles. Regardless of sex, possessing 1 or 2 APOEε4 alleles was associated with

decreased levels of CSF and plasma CRP. Sex differences were seen in inflammatory markers,

regardless of diagnosis or genotype, as females had lower CSF cytokines (IL-16, IL-18), CSF

ICAM1, CSF and plasma immunoglobulins (IgA, IgE) and plasma IL-18. However, tissue (CSF,

plasma) mattered for results from certain inflammatory markers (ICAM1 and to a lesser extent

CRP) as females had higher plasma CRP and ICAM1 compared to males, opposite to what was

found in CSF, supporting previous studies suggesting differences in inflammatory signals

between CSF and blood (Bettcher et al., 2018). Finally, we found no significant differences in

epigenetic age acceleration by dementia diagnosis or sex in this cohort of samples. Sex and

genotype differences in CSF and plasma inflammatory markers suggest differences in underlying

physiology that may affect aging and the progression of AD and this should be considered in

future studies. Researchers should be cautioned to use sex as a biological variable in all analyses.

Sex interacts with presence of APOEε4 alleles to affect levels of IL-8 and IL-16

In this study, we found that sex interacted with genotype to influence CSF IL-16 and IL-

8. CSF IL-16 and IL-8 levels were lower in females with no APOEε4 alleles compared to males,

but no sex differences in these cytokine levels were detected in participants with 1 or 2 ε4 alleles.

Our results suggest that presence of APOEε4 alleles can modulate CSF (and potentially plasma)

cytokine levels in a sex-dependent way. The APOE protein can regulate transcription in vitro

(Theendakara et al., 2018) and APOE4, but not APOE3, increases levels of IL-6 and IL-8 in vitro

(Theendakara et al., 2016). In the current study, we found that the sex differences in IL-16 and

IL-8 levels disappeared with the presence of 1 or 2 APOEε4 alleles. One possibility is that the

APOE4 protein regulates cytokine levels differently in males and females. IL-16 has been

implicated in AD (Rosa et al., 2006) and plasma IL-16 levels decreased with diagnosis (in males

and females; current study) and AD severity (analysis without regard to sex; Motta et al., 2007).

On the other hand, levels of IL-8 were not affected by diagnosis, consistent with a meta-analysis

of cytokines in AD (Swardfager et al., 2010). It is unclear what the impact of regulation of CSF

cytokine levels by sex and APOEε4 has on AD symptoms or pathology, however given that

females with APOEε4 alleles are disproportionally affected by AD (Hohman et al., 2018; Neu et

al., 2017), IL-16 and IL-8 levels are unlikely to be a mechanism for this effect as differences in

sex by genotype were noticed in the absence not presence of APOEε4 alleles.

Females have higher CRP levels compared to males and CRP levels are lower in APOEε4

carriers

We found that plasma and CSF levels of CRP, a widely used inflammatory and

cardiovascular marker (Koenig et al., 1999; Ridker et al., 1998), were independently affected by

sex and APOE genotyoe. Females, regardless of diagnosis or APOEε4 alleles, had significantly

higher plasma CRP relative to males, consistent with findings in healthy individuals (Khera et

al., 2005). Higher levels of peripheral CRP may suggest higher systemic inflammation in

females, which is associated with an increased risk in all-cause dementia (Koyama et al., 2013).

Higher levels of serum CRP are also associated with higher levels of serum estradiol in

postmenopausal healthy females (Eldridge et al., 2020) which suggests that sex differences in

CRP levels may be partly due to sex differences in estradiol levels. In addition, we found that the

presence of APOEε4 alleles decreased plasma and CSF CRP levels consistent with previous

research in large population studies (Hubacek et al., 2010; Yun et al., 2015). In our study, we

also found a trend for lower levels of plasma CRP with LMCI and AD compared to CN. Recent

meta-analyses did not find differences in peripheral levels of CRP in AD compared to control patients (Gong et al., 2016; Ng et al., 2018). However, in patients with mild and moderate dementia only, serum CRP levels were lower compared to the cognitively normal group (Gong et al., 2016). In healthy individuals, higher levels of plasma CRP in mid-life are associated with a higher amyloid burden later in life in males but not females (Walker et al., 2018). However, despite this finding, higher systemic inflammation (including CRP) was associated with greater cognitive decline later in life in both sexes in healthy individuals (Walker et al., 2019). It is possible that sex differences in inflammation earlier in life have long-term effects on the transition to MCI and/or AD.

Females have lower cytokine and immunoglobulin levels compared to males

We found some biomarkers that were affected by sex, but not diagnosis or presence of APOEε4 alleles. For example, females had lower CSF levels of ICAM1 compared to males, regardless of APOE genotype or diagnosis, but, although a trend, the opposite effect was seen in plasma. In contrast, in healthy adults (18-55 years old), serum levels of ICAM1 are lower in females compared to males (Ponthieux et al., 2003). ICAM1 is a type of adhesion molecule associated with microvascular endothelial activation (Zenaro et al., 2017) and plasma ICAM1 levels (but not CSF levels, Nielsen et al., 2007) were higher in patients with AD (Huang et al., 2015; Nielsen et al., 2007; Rentzos et al., 2004). Although in the present study we did not observe a significant effect of plasma ICAM1 with diagnosis, the unadjusted P-value was 0.063 with higher levels in LMCI and AD groups. It is intriguing that females have lower CSF levels of cytokines (IL-16, IL-18), ICAM1, and immunoglobulins (IgE and IgA) but higher plasma CRP and ICAM1 levels. Neuroinflammation is associated with AD but it can have both beneficial and

detrimental roles (Walters et al., 2016) as increased expression of pro-inflammatory cytokines

contributes to neuronal loss, while anti-inflammatory effects contribute to amyloid beta clearance

(Heneka et al., 2015). It is possible that males and females have varying levels of beneficial vs

detrimental immune responses which can affect how the disease progresses in each of the sexes

but it is also important to remember that CSF levels may not match levels in different tissues

(blood) or regions of the brain.

Sex, AD and biochemical markers do not affect blood epigenetic age acceleration

432   We did not observe an association between either sex or diagnosis and epigenetic age

433   acceleration or intrinsic epigenetic age acceleration. To our knowledge, no other study has

434   similarly probed epigenetic age acceleration in a peripheral tissue in the presence of AD, or

435   whether epigenetic age acceleration in AD is sex-specific.

436   This study was partially undertaken to investigate whether epigenetic age acceleration

that has been associated with the AD brain is reflected in peripheral tissues. Levine et al. (Levine

et al., 2015) have previously demonstrated increased epigenetic age acceleration in AD, however

Levine's study was conducted on post-mortem prefrontal cortex tissue, and did not explicitly

investigate the role of sex in epigenetic age acceleration (Levine et al., 2015). While brain-blood

global DNA methylation profiles are reasonably correlated (r=0.86) (Braun et al., 2019), DNA

methylation profiles of peripheral tissues are imperfect representatives of the brain, and do not

recapitulate all epigenetic alterations with high fidelity. Thus, our findings do not contradict the

finding of increased epigenetic age acceleration in the presence of AD in the prefrontal cortex,

but suggest that accelerated epigenetic aging in AD is not a pan-tissue phenomenon. Our finding

of a lack of significant association between AD, biological sex, and epigenetic age acceleration

in whole blood DNA methylation profiles could suggest a tissue-specific dysregulation of an

epigenetic maintenance system, in which the brain epigenome is most strongly affected by AD

(Levine et al., 2015). The phenotype of patients affected by AD and global gene expression

patterns of the APOE protein, with high expression in brain, and low expression in whole blood

(GTEx Project, 2018) further support this hypothesis.

Intriguingly, epigenetic age was observed to be lower on average than chronological age.

Horvath's epigenetic clock was trained on DNAme data from older versions of the Illumina

DNAme arrays with more limited genomic coverage; 19 of the 353 CpG sites required to

calculate epigenetic age via Horvath's algorithm do not exist on the newer EPIC array. Two

previous studies investigated the application of Horvath's epigenetic clock to EPIC data with

conflicting results (Dhingra et al., 2019; McEwen et al., 2018), the largest issue being chronic

underestimation of epigenetic age due to the positive linear regression coefficients associated

with the missing probes (Dhingra et al., 2019). Both imputing and removing the missing probes

from the array resulted in a chronic underprediction of epigenetic age with Horvath's clock

suggesting that this is likely an artefact of the array platform and truncated probe-set rather than

the method chosen to deal with missing values, although it is possible that an adjustment factor

could be devised to more accurately apply Horvath's clock to EPIC data. In future explorations

of epigenetic age with EPIC DNAme array data this should be considered, as there are other

epigenetic age predictors available that have been trained on EPIC data such as the PhenoAge

and GrimAge clocks, although these tools have limitations as well; for example, both PhenoAge

and GrimAge were trained only on blood DNAme data, as compared to the original pan-tissue

epigenetic clocks, and therefore may have limited applicability and relevance in other tissues

(Levine et al., 2018; Lu et al., 2019).

Limitations

The ADNI cohort is not ethnically or socioeconomically diverse, being mostly composed

of self-reported white (only 12 individuals were not-white) and highly educated individuals

(average 15.69 years of education). As AD incidence, prevalence, and age of onset varies by

ethnicity (Fitten et al., 2014; Mayeda et al., 2016; Steenland et al., 2016) and education (Sharp

and Gatz, 2011), our conclusions may not apply to more ethnically and socially diverse

populations. In addition to sex, it is possible the underlying mechanisms of AD are different

depending on ethnicity. APOEε4 genotype is a well-known AD risk factor and in this cohort the

proportion of participants in the different APOEε4 allele groups is correlated with diagnosis

(Table S6). Future research with a larger cohort is required to test how sex, APOE genotype and

diagnosis interact together in one model. Even in the larger 640-participant cohort, only 37

participants (5.78%) had an AD diagnosis, so statistical analyses were underpowered to detect

subtle differences by diagnosis group, which may account for our lack of interaction between sex

and diagnosis. Additionally, other pathologies in these participants, such as cancer,

cardiovascular disease, smoking status, or obesity may have influenced inflammatory markers

and epigenomes and limited our interpretations.

CONCLUSION

The current study provides evidence that sex and presence of APOEε4 alleles are

associated with CSF levels of the inflammatory markers IL-16 and IL-8. We found sex

differences indicating that females had lower cytokine and immunoglobulin levels but higher

plasma CRP and ICAM1 levels compared to males, although the direction of the ICAM1 finding

was tissue-dependent. There were no significant associations between sex, APOEε4 alleles or

diagnosis and epigenetic age. Together, our work suggests that that presence of APOEε4 alleles

can affect cytokine levels differently in males and females and the underlying pathophysiology

of aging and AD may be tissue- and sex-specific.

ACKNOWLEDGMENTS

Data collection and sharing for this project was funded by the Alzheimer's Disease

Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD

ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the

National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering,

and through generous contributions from the following: AbbVie, Alzheimer's Association;

Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-

Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli

Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company

Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy

Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development

LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx

Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal

Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian

Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private

sector contributions are facilitated by the Foundation for the National Institutes of Health

(www.fnih.org). The grantee organization is the Northern California Institute for Research and

Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the

University of Southern California. ADNI data are disseminated by the Laboratory for Neuro

Imaging at the University of Southern California. Funding for this study was provided by a

Canadian Institutes of Health Research (CHIR) grant to LAMG (PJT-148662) and a CIHR sex

catalyst grant to WPR (Reference No. 158613). WPR receives salary support through an

investigatorship award from the BC Children's Hospital Research Institute. PDG is funded by

the Alzheimer's Association of the USA and Brain Canada with the financial support of Health

Canada through the Brain Canada Research Fund (AARF-17-529705). The views expressed

herein do not necessarily represent the views of the Minister of Health or the Government of

Canada. AMI is funded by a University of British Columbia Four Year Doctoral Fellowship and

the CIHR Frederick Banting and Charles Best Masters Research Award. We thank Elizabeth

Perez for help with data organization.

CONFLICT OF INTEREST

529   The authors have no conflict of interest to report.

REFERENCES

Alzheimer's Association, 2017. 2017 Alzheimer's disease facts and figures. Alzheimer's & Dementia 13, 325–373. https://doi.org/10.1016/j.jalz.2017.02.001

Apostolova, L.G., Dutton, R.A., Dinov, I.D., Hayashi, K.M., Toga, A.W., Cummings, J.L., Thompson, P.M., 2006. Conversion of mild cognitive impairment to Alzheimer disease predicted by hippocampal atrophy maps. Arch. Neurol. 63, 693–699. https://doi.org/10.1001/archneur.63.5.693

Ardekani, B.A., Convit, A., Bachman, A.H., 2016. Analysis of the MIRIAD Data Shows Sex Differences in Hippocampal Atrophy Progression. J. Alzheimers Dis. 50, 847–857. https://doi.org/10.3233/JAD-150780

Aryee, M.J., Jaffe, A.E., Corrada-Bravo, H., Ladd-Acosta, C., Feinberg, A.P., Hansen, K.D., Irizarry, R.A., 2014. Minfi: a flexible and comprehensive Bioconductor package for the analysis of Infinium DNA methylation microarrays. Bioinformatics 30, 1363–1369. https://doi.org/10.1093/bioinformatics/btu049

Barnes, L.L., Wilson, R.S., Bienias, J.L., Schneider, J.A., Evans, D.A., Bennett, D.A., 2005. Sex
differences in the clinical manifestations of Alzheimer disease pathology. Arch. Gen.
Psychiatry 62, 685–691. https://doi.org/10.1001/archpsyc.62.6.685
Baumgart, M., Snyder, H.M., Carrillo, M.C., Fazio, S., Kim, H., Johns, H., 2015. Summary of
the evidence on modifiable risk factors for cognitive decline and dementia: A population-
based perspective. Alzheimers Dement 11, 718–726.
https://doi.org/10.1016/j.jalz.2015.05.016
Benjamini, Y., Hochberg, Y., 1995. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society. Series B
(Methodological) 57, 289–300.
Bettcher, B.M., Johnson, S.C., Fitch, R., Casaletto, K.B., Heffernan, K.S., Asthana, S.,
Zetterberg, H., Blennow, K., Carlsson, C.M., Neuhaus, J., Bendlin, B.B., Kramer, J.H.,
2018. CSF and Plasma Levels of Inflammation Differentially Relate to CNS Markers of
Alzheimer's Disease Pathology and Neuronal Damage. J Alzheimers Dis 62, 385–397.
https://doi.org/10.3233/JAD-170602
Beydoun, M.A., Boueiz, A., Abougergi, M.S., Kitner-Triolo, M.H., Beydoun, H.A., Resnick,
S.M., O'Brien, R., Zonderman, A.B., 2012. Sex differences in the association of the
apolipoprotein E epsilon 4 allele with incidence of dementia, cognitive impairment, and
decline. Neurobiol. Aging 33, 720-731.e4.
https://doi.org/10.1016/j.neurobiolaging.2010.05.017
Braun, P.R., Han, S., Hing, B., Nagahama, Y., Gaul, L.N., Heinzman, J.T., Grossbach, A.J.,
Close, L., Dlouhy, B.J., Howard, M.A., Kawasaki, H., Potash, J.B., Shinozaki, G., 2019.
Genome-wide DNA methylation comparison between live human brain and peripheral
tissues within individuals. Translational Psychiatry 9, 47. https://doi.org/10.1038/s41398-
019-0376-y
Caruso, A., Nicoletti, F., Mango, D., Saidi, A., Orlando, R., Scaccianoce, S., 2018. Stress as risk
factor for Alzheimer's disease. Pharmacological Research 132, 130–134.
https://doi.org/10.1016/j.phrs.2018.04.017
Chen, B.H., Marioni, R.E., Colicino, E., Peters, M.J., Ward-Caviness, C.K., Tsai, P.-C., Roetker,
N.S., Just, A.C., Demerath, E.W., Guan, W., Bressler, J., Fornage, M., Studenski, S.,
Vandiver, A.R., Moore, A.Z., Tanaka, T., Kiel, D.P., Liang, L., Vokonas, P., Schwartz,
J., Lunetta, K.L., Murabito, J.M., Bandinelli, S., Hernandez, D.G., Melzer, D., Nalls, M.,
Pilling, L.C., Price, T.R., Singleton, A.B., Gieger, C., Holle, R., Kretschmer, A.,
Kronenberg, F., Kunze, S., Linseisen, J., Meisinger, C., Rathmann, W., Waldenberger,
M., Visscher, P.M., Shah, S., Wray, N.R., McRae, A.F., Franco, O.H., Hofman, A.,
Uitterlinden, A.G., Absher, D., Assimes, T., Levine, M.E., Lu, A.T., Tsao, P.S., Hou, L.,
Manson, J.E., Carty, C.L., LaCroix, A.Z., Reiner, A.P., Spector, T.D., Feinberg, A.P.,
Levy, D., Baccarelli, A., van Meurs, J., Bell, J.T., Peters, A., Deary, I.J., Pankow, J.S.,
Ferrucci, L., Horvath, S., 2016. DNA methylation-based measures of biological age:
meta-analysis predicting time to death. Aging (Albany NY) 8, 1844–1859.
https://doi.org/10.18632/aging.101020
Craig-Schapiro, R., Kuhn, M., Xiong, C., Pickering, E.H., Liu, J., Misko, T.P., Perrin, R.J.,
Bales, K.R., Soares, H., Fagan, A.M., Holtzman, D.M., 2011. Multiplexed Immunoassay
Panel Identifies Novel CSF Biomarkers for Alzheimer's Disease Diagnosis and
Prognosis. PLOS ONE 6, e18850. https://doi.org/10.1371/journal.pone.0018850

Crane, P.K., Carle, A., Gibbons, L.E., Insel, P., Mackin, R.S., Gross, A., Jones, R.N., Mukherjee, S., Curtis, S.M., Harvey, D., Weiner, M., Mungas, D., Alzheimer's Disease Neuroimaging Initiative, 2012. Development and assessment of a composite score for memory in the Alzheimer's Disease Neuroimaging Initiative (ADNI). Brain Imaging Behav 6, 502–516. https://doi.org/10.1007/s11682-012-9186-z

Dantzer, R., 2018. Neuroimmune Interactions: From the Brain to the Immune System and Vice Versa. Physiol. Rev. 98, 477–504. https://doi.org/10.1152/physrev.00039.2016

de Miranda, A.S., Zhang, C.-J., Katsumoto, A., Teixeira, A.L., 2017. Hippocampal adult neurogenesis: Does the immune system matter? Journal of the Neurological Sciences 372, 482–495. https://doi.org/10.1016/j.jns.2016.10.052

Degerman, S., Josefsson, M., Nordin Adolfsson, A., Wennstedt, S., Landfors, M., Haider, Z., Pudas, S., Hultdin, M., Nyberg, L., Adolfsson, R., 2017. Maintained memory in aging is associated with young epigenetic age. Neurobiology of Aging 55, 167–171. https://doi.org/10.1016/j.neurobiolaging.2017.02.009

Dhingra, R., Kwee, L.C., Diaz-Sanchez, D., Devlin, R.B., Cascio, W., Hauser, E.R., Gregory, S., Shah, S., Kraus, W.E., Olden, K., Ward-Caviness, C.K., 2019. Evaluating DNA methylation age on the Illumina MethylationEPIC Bead Chip. PLOS ONE 14, e0207834. https://doi.org/10.1371/journal.pone.0207834

Doecke, J.D., Laws, S.M., Faux, N.G., Wilson, W., Burnham, S.C., Lam, C.-P., Mondal, A., Bedo, J., Bush, A.I., Brown, B., Ruyck, K.D., Ellis, K.A., Fowler, C., Gupta, V.B., Head, R., Macaulay, S.L., Pertile, K., Rowe, C.C., Rembach, A., Rodrigues, M., Rumble, R., Szoeke, C., Taddei, K., Taddei, T., Trounson, B., Ames, D., Masters, C.L., Martins, R.N., Group, for the A.D.N.I. and A.I.B. and L.R., 2012. Blood-Based Protein Biomarkers for Diagnosis of Alzheimer Disease. Arch Neurol 69, 1318–1325. https://doi.org/10.1001/archneurol.2012.1282

Eldridge, R.C., Wentzensen, N., Pfeiffer, R.M., Brinton, L.A., Hartge, P., Guillemette, C., Kemp, T.J., Pinto, L.A., Trabert, B., 2020. Endogenous estradiol and inflammation biomarkers: potential interacting mechanisms of obesity-related disease. Cancer Causes Control. https://doi.org/10.1007/s10552-020-01280-6

Engler, H., Benson, S., Wegner, A., Spreitzer, I., Schedlowski, M., Elsenbruch, S., 2016. Men and women differ in inflammatory and neuroendocrine responses to endotoxin but not in the severity of sickness symptoms. Brain, Behavior, and Immunity 52, 18–26. https://doi.org/10.1016/j.bbi.2015.08.013

Fiorito, G., Polidoro, S., Dugué, P.-A., Kivimaki, M., Ponzi, E., Matullo, G., Guarrera, S., Assumma, M.B., Georgiadis, P., Kyrtopoulos, S.A., Krogh, V., Palli, D., Panico, S., Sacerdote, C., Tumino, R., Chadeau-Hyam, M., Stringhini, S., Severi, G., Hodge, A.M., Giles, G.G., Marioni, R., Linnér, R.K., O'Halloran, A.M., Kenny, R.A., Layte, R., Baglietto, L., Robinson, O., McCrory, C., Milne, R.L., Vineis, P., 2017. Social adversity and epigenetic aging: a multi-cohort study on socioeconomic differences in peripheral blood DNA methylation. Scientific Reports 7, 16266. https://doi.org/10.1038/s41598-017-16391-5

Fitten, L.J., Ortiz, F., Fairbanks, L., Bartzokis, G., Lu, P., Klein, E., Coppola, G., Ringman, J., 2014. Younger age of dementia diagnosis in a Hispanic population in southern California. Int J Geriatr Psychiatry 29, 586–593. https://doi.org/10.1002/gps.4040

Fortin, J.-P., Triche, T.J., Hansen, K.D., 2017. Preprocessing, normalization and integration of
the Illumina HumanMethylationEPIC array with minfi. Bioinformatics 33, 558–560.
https://doi.org/10.1093/bioinformatics/btw691
Gale, C.R., Marioni, R.E., Harris, S.E., Starr, J.M., Deary, I.J., 2018. DNA methylation and the
epigenetic clock in relation to physical frailty in older people: the Lothian Birth Cohort
1936. Clin Epigenetics 10. https://doi.org/10.1186/s13148-018-0538-4
Gibbons, L.E., Carle, A.C., Mackin, R.S., Harvey, D., Mukherjee, S., Insel, P., Curtis, S.M.,
Mungas, D., Crane, P.K., Alzheimer's Disease Neuroimaging Initiative, 2012. A
composite score for executive functioning, validated in Alzheimer's Disease
Neuroimaging Initiative (ADNI) participants with baseline mild cognitive impairment.
Brain Imaging Behav 6, 517–527. https://doi.org/10.1007/s11682-012-9176-1
Gong, C., Wei, D., Wang, Y., Ma, J., Yuan, C., Zhang, W., Yu, G., Zhao, Y., 2016. A Meta-
Analysis of C-Reactive Protein in Patients With Alzheimer's Disease. Am J Alzheimers
Dis Other Demen 31, 194–200. https://doi.org/10.1177/1533317515602087
GTEx Project, 2018. The Genotype-Tissue Expression (GTEx) Project was supported by the
Common Fund of the Office of the Director of the National Institutes of Health, and by
NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS. The data used for the analyses
described in this manuscript were obtained from the GTEx Portal on 11/04/18.
Hannum, G., Guinney, J., Zhao, L., Zhang, L., Hughes, G., Sadda, S., Klotzle, B., Bibikova, M.,
Fan, J.-B., Gao, Y., Deconde, R., Chen, M., Rajapakse, I., Friend, S., Ideker, T., Zhang,
K., 2013. Genome-wide Methylation Profiles Reveal Quantitative Views of Human
Aging Rates. Molecular Cell 49, 359–367. https://doi.org/10.1016/j.molcel.2012.10.016
Heneka, M.T., Carson, M.J., Khoury, J.E., Landreth, G.E., Brosseron, F., Feinstein, D.L., Jacobs,
A.H., Wyss-Coray, T., Vitorica, J., Ransohoff, R.M., Herrup, K., Frautschy, S.A., Finsen,
B., Brown, G.C., Verkhratsky, A., Yamanaka, K., Koistinaho, J., Latz, E., Halle, A.,
Petzold, G.C., Town, T., Morgan, D., Shinohara, M.L., Perry, V.H., Holmes, C., Bazan,
N.G., Brooks, D.J., Hunot, S., Joseph, B., Deigendesch, N., Garaschuk, O., Boddeke, E.,
Dinarello, C.A., Breitner, J.C., Cole, G.M., Golenbock, D.T., Kummer, M.P., 2015.
Neuroinflammation in Alzheimer's disease. The Lancet Neurology 14, 388–405.
https://doi.org/10.1016/S1474-4422(15)70016-5
Heppner, F.L., Ransohoff, R.M., Becher, B., 2015. Immune attack: the role of inflammation in
Alzheimer disease. Nature Reviews Neuroscience 16, 358–372.
https://doi.org/10.1038/nrn3880
Hohman, T.J., Dumitrescu, L., Barnes, L.L., Thambisetty, M., Beecham, G., Kunkle, B., Gifford,
K.A., Bush, W.S., Chibnik, L.B., Mukherjee, S., Jager, P.L.D., Kukull, W., Crane, P.K.,
Resnick, S.M., Keene, C.D., Montine, T.J., Schellenberg, G.D., Haines, J.L., Zetterberg,
H., Blennow, K., Larson, E.B., Johnson, S.C., Albert, M., Bennett, D.A., Schneider, J.A.,
Jefferson, A.L., 2018. Sex-Specific Association of Apolipoprotein E With Cerebrospinal
Fluid Levels of Tau. JAMA Neurol 75, 989–998.
https://doi.org/10.1001/jamaneurol.2018.0821
Holland, D., Desikan, R.S., Dale, A.M., McEvoy, L.K., Alzheimer's Disease Neuroimaging
Initiative, 2013. Higher rates of decline for women and apolipoprotein E epsilon4
carriers. AJNR Am J Neuroradiol 34, 2287–2293. https://doi.org/10.3174/ajnr.A3601
Horvath, S., 2013. DNA methylation age of human tissues and cell types. Genome Biol 14,
R115. https://doi.org/10.1186/gb-2013-14-10-r115

Horvath, S., Garagnani, P., Bacalini, M.G., Pirazzini, C., Salvioli, S., Gentilini, D., Di Blasio, A.M., Giuliani, C., Tung, S., Vinters, H.V., Franceschi, C., 2015. Accelerated epigenetic aging in Down syndrome. Aging Cell 14, 491–495. https://doi.org/10.1111/acel.12325

Horvath, S., Gurven, M., Levine, M.E., Trumble, B.C., Kaplan, H., Allayee, H., Ritz, B.R., Chen, B., Lu, A.T., Rickabaugh, T.M., Jamieson, B.D., Sun, D., Li, S., Chen, W., Quintana-Murci, L., Fagny, M., Kobor, M.S., Tsao, P.S., Reiner, A.P., Edlefsen, K.L., Absher, D., Assimes, T.L., 2016. An epigenetic clock analysis of race/ethnicity, sex, and coronary heart disease. Genome Biol 17. https://doi.org/10.1186/s13059-016-1030-0

Horvath, S., Ritz, B.R., 2015. Increased epigenetic age and granulocyte counts in the blood of Parkinson's disease patients. Aging (Albany NY) 7, 1130–1142.

Houseman, E.A., Accomando, W.P., Koestler, D.C., Christensen, B.C., Marsit, C.J., Nelson, H.H., Wiencke, J.K., Kelsey, K.T., 2012. DNA methylation arrays as surrogate measures of cell mixture distribution. BMC Bioinformatics 13, 86. https://doi.org/10.1186/1471-2105-13-86

Hu, W.T., Holtzman, D.M., Fagan, A.M., Shaw, L.M., Perrin, R., Arnold, S.E., Grossman, M., Xiong, C., Craig-Schapiro, R., Clark, C.M., Pickering, E., Kuhn, M., Chen, Y., Van Deerlin, V.M., McCluskey, L., Elman, L., Karlawish, J., Chen-Plotkin, A., Hurtig, H.I., Siderowf, A., Swenson, F., Lee, V.M.-Y., Morris, J.C., Trojanowski, J.Q., Soares, H., Alzheimer's Disease Neuroimaging Initiative, 2012. Plasma multianalyte profiling in mild cognitive impairment and Alzheimer disease. Neurology 79, 897–905. https://doi.org/10.1212/WNL.0b013e318266fa70

Hua, X., Hibar, D.P., Lee, S., Toga, A.W., Jack, C.R., Weiner, M.W., Thompson, P.M., 2010. Sex and age differences in atrophic rates: an ADNI study with n=1368 MRI scans. Neurobiology of Aging 31, 1463–1480. https://doi.org/10.1016/j.neurobiolaging.2010.04.033

Huang, C.-W., Tsai, M.-H., Chen, N.-C., Chen, W.-H., Lu, Y.-T., Lui, C.-C., Chang, Y.-T., Chang, W.-N., Chang, A.Y.W., Chang, C.-C., 2015. Clinical significance of circulating vascular cell adhesion molecule-1 to white matter disintegrity in Alzheimer's dementia. Thrombosis and Haemostasis 114, 1230–1240. https://doi.org/10.1160/TH14-11-0938

Hubacek, J.A., Peasey, A., Pikhart, H., Stavek, P., Kubinova, R., Marmot, M., Bobak, M., 2010. APOE polymorphism and its effect on plasma C-reactive protein levels in a large general population sample. Hum Immunol 71, 304–308. https://doi.org/10.1016/j.humimm.2010.01.008

Irvine, K., Laws, K.R., Gale, T.M., Kondel, T.K., 2012. Greater cognitive deterioration in women than men with Alzheimer's disease: a meta analysis. J Clin Exp Neuropsychol 34, 989–998. https://doi.org/10.1080/13803395.2012.712676

Khera, A., McGuire, D.K., Murphy, S.A., Stanek, H.G., Das, S.R., Vongpatanasin, W., Wians, F.H., Grundy, S.M., de Lemos, J.A., 2005. Race and gender differences in C-reactive protein levels. J. Am. Coll. Cardiol. 46, 464–469. https://doi.org/10.1016/j.jacc.2005.04.051

Kinney, J.W., Bemiller, S.M., Murtishaw, A.S., Leisgang, A.M., Salazar, A.M., Lamb, B.T., 2018. Inflammation as a central mechanism in Alzheimer's disease. Alzheimer's & Dementia: Translational Research & Clinical Interventions 4, 575–590. https://doi.org/10.1016/j.trci.2018.06.014

Klein, S.L., Flanagan, K.L., 2016. Sex differences in immune responses. Nature Reviews Immunology 16, 626–638. https://doi.org/10.1038/nri.2016.90

Koenig, W., Sund, M., Fröhlich, M., Fischer, H.G., Löwel, H., Döring, A., Hutchinson, W.L.,
 Pepys, M.B., 1999. C-Reactive protein, a sensitive marker of inflammation, predicts
 future risk of coronary heart disease in initially healthy middle-aged men: results from
 the MONICA (Monitoring Trends and Determinants in Cardiovascular Disease)
 Augsburg Cohort Study, 1984 to 1992. Circulation 99, 237–242.
 https://doi.org/10.1161/01.cir.99.2.237
Koran, M.E.I., Hohman, T.J., 2017. Sex differences in the association between AD biomarkers
 and cognitive decline. Brain Imaging and Behavior 11, 205–213.
 https://doi.org/10.1007/s11682-016-9523-8
Koran, M.E.I., Wagener, M., Hohman, T.J., Alzheimer's Neuroimaging Initiative, 2017. Sex
 differences in the association between AD biomarkers and cognitive decline. Brain
 Imaging Behav 11, 205–213. https://doi.org/10.1007/s11682-016-9523-8
Koyama, A., O'Brien, J., Weuve, J., Blacker, D., Metti, A.L., Yaffe, K., 2013. The role of
 peripheral inflammatory markers in dementia and Alzheimer's disease: a meta-analysis.
 J. Gerontol. A Biol. Sci. Med. Sci. 68, 433–440. https://doi.org/10.1093/gerona/gls187
Levine, M.E., Lu, A.T., Bennett, D.A., Horvath, S., 2015. Epigenetic age of the pre-frontal
 cortex is associated with neuritic plaques, amyloid load, and Alzheimer's disease related
 cognitive functioning. Aging (Albany NY) 7, 1198–1211.
Levine, M.E., Lu, A.T., Quach, A., Chen, B.H., Assimes, T.L., Bandinelli, S., Hou, L.,
 Baccarelli, A.A., Stewart, J.D., Li, Y., Whitsel, E.A., Wilson, J.G., Reiner, A.P., Aviv,
 A., Lohman, K., Liu, Y., Ferrucci, L., Horvath, S., 2018. An epigenetic biomarker of
 aging for lifespan and healthspan. Aging (Albany NY) 10, 573–591.
 https://doi.org/10.18632/aging.101414
Lin, K.A., Choudhury, K.R., Rathakrishnan, B.G., Marks, D.M., Petrella, J.R., Doraiswamy,
 P.M., 2015. Marked gender differences in progression of mild cognitive impairment over
 8 years. Alzheimer's & Dementia: Translational Research & Clinical Interventions 1,
 103–110. https://doi.org/10.1016/j.trci.2015.07.001
Lu, A.T., Quach, A., Wilson, J.G., Reiner, A.P., Aviv, A., Raj, K., Hou, L., Baccarelli, A.A., Li,
 Y., Stewart, J.D., Whitsel, E.A., Assimes, T.L., Ferrucci, L., Horvath, S., 2019. DNA
 methylation GrimAge strongly predicts lifespan and healthspan. Aging (Albany NY) 11,
 303–327. https://doi.org/10.18632/aging.101684
Lupien, S.J., de Leon, M., de Santi, S., Convit, A., Tarshish, C., Nair, N.P., Thakur, M.,
 McEwen, B.S., Hauger, R.L., Meaney, M.J., 1998. Cortisol levels during human aging
 predict hippocampal atrophy and memory deficits. Nat. Neurosci. 1, 69–73.
 https://doi.org/10.1038/271
Mayeda, E.R., Glymour, M.M., Quesenberry, C.P., Whitmer, R.A., 2016. Inequalities in
 dementia incidence between six racial and ethnic groups over 14 years. Alzheimer's &
 Dementia 12, 216–224. https://doi.org/10.1016/j.jalz.2015.12.007
McEwen, B.S., Biron, C.A., Brunson, K.W., Bulloch, K., Chambers, W.H., Dhabhar, F.S.,
 Goldfarb, R.H., Kitson, R.P., Miller, A.H., Spencer, R.L., Weiss, J.M., 1997. The role of
 adrenocorticoids as modulators of immune function in health and disease: neural,
 endocrine and immune interactions. Brain Research Reviews 23, 79–133.
 https://doi.org/10.1016/S0165-0173(96)00012-4
McEwen, L.M., Jones, M.J., Lin, D.T.S., Edgar, R.D., Husquin, L.T., MacIsaac, J.L., Ramadori,
 K.E., Morin, A.M., Rider, C.F., Carlsten, C., Quintana-Murci, L., Horvath, S., Kobor,
 M.S., 2018. Systematic evaluation of DNA methylation age estimation with common

preprocessing methods and the Infinium MethylationEPIC BeadChip array. Clinical Epigenetics 10, 123. https://doi.org/10.1186/s13148-018-0556-2

McEwen, L.M., Morin, A.M., Edgar, R.D., MacIsaac, J.L., Jones, M.J., Dow, W.H., Rosero-Bixby, L., Kobor, M.S., Rehkopf, D.H., 2017. Differential DNA methylation and lymphocyte proportions in a Costa Rican high longevity region. Epigenetics Chromatin 10. https://doi.org/10.1186/s13072-017-0128-2

Motta, M., Imbesi, R., Di Rosa, M., Stivala, F., Malaguarnera, L., 2007. Altered plasma cytokine levels in Alzheimer's disease: Correlation with the disease progression. Immunology Letters 114, 46–51. https://doi.org/10.1016/j.imlet.2007.09.002

Murphy, K.J., Sheppard, P.A.S., Troyer, A.K., Hampson, E., Galea, L.A.M., 2019. Sex differences in cortisol and memory following acute social stress in amnestic mild cognitive impairment. bioRxiv 757484. https://doi.org/10.1101/757484

Nebel, R.A., Aggarwal, N.T., Barnes, L.L., Gallagher, A., Goldstein, J.M., Kantarci, K., Mallampalli, M.P., Mormino, E.C., Scott, L., Yu, W.H., Maki, P.M., Mielke, M.M., 2018. Understanding the impact of sex and gender in Alzheimer's disease: A call to action. Alzheimer's & Dementia 14, 1171–1183. https://doi.org/10.1016/j.jalz.2018.04.008

Neu, S.C., Pa, J., Kukull, W., Beekly, D., Kuzma, A., Gangadharan, P., Wang, L.-S., Romero, K., Arneric, S.P., Redolfi, A., Orlandi, D., Frisoni, G.B., Au, R., Devine, S., Auerbach, S., Espinosa, A., Boada, M., Ruiz, A., Johnson, S.C., Koscik, R., Wang, J.-J., Hsu, W.-C., Chen, Y.-L., Toga, A.W., 2017. Apolipoprotein E Genotype and Sex Risk Factors for Alzheimer Disease: A Meta-analysis. JAMA Neurology 74, 1178. https://doi.org/10.1001/jamaneurol.2017.2188

Ng, A., Tam, W.W., Zhang, M.W., Ho, C.S., Husain, S.F., McIntyre, R.S., Ho, R.C., 2018. IL-1β, IL-6, TNF- α and CRP in Elderly Patients with Depression or Alzheimer's disease: Systematic Review and Meta-Analysis. Scientific Reports 8, 12050. https://doi.org/10.1038/s41598-018-30487-6

Nielsen, H.M., Londos, E., Minthon, L., Janciauskiene, S.M., 2007. Soluble adhesion molecules and angiotensin-converting enzyme in dementia. Neurobiol. Dis. 26, 27–35. https://doi.org/10.1016/j.nbd.2006.11.011

Petersen, R.C., Jack, C.R., Xu, Y.C., Waring, S.C., O'Brien, P.C., Smith, G.E., Ivnik, R.J., Tangalos, E.G., Boeve, B.F., Kokmen, E., 2000. Memory and MRI-based hippocampal volumes in aging and AD. Neurology 54, 581–587. https://doi.org/10.1212/wnl.54.3.581

Ponthieux, A., Herbeth, B., Droesch, S., Lambert, D., Visvikis, S., 2003. Age- and Sex-related Reference Values for Serum Adhesion Molecule Concentrations in Healthy Individuals: Intercellular Adhesion Molecule-1 and E-, P-, and L-Selectin. Clinical Chemistry 49, 1544–1546. https://doi.org/10.1373/49.9.1544

R Core Team, 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. R Foundation for Statistical Computing, Vienna, Austria.

Rentzos, M., Michalopoulou, M., Nikolaou, C., Cambouri, C., Rombos, A., Dimitrakopoulos, A., Kapaki, E., Vassilopoulos, D., 2004. Serum Levels of Soluble Intercellular Adhesion Molecule-1 and Soluble Endothelial Leukocyte Adhesion Molecule-1 in Alzheimer's Disease. J Geriatr Psychiatry Neurol 17, 225–231. https://doi.org/10.1177/0891988704269822

Ridker, P.M., Buring, J.E., Shih, J., Matias, M., Hennekens, C.H., 1998. Prospective study of C-
reactive protein and the risk of future cardiovascular events among apparently healthy
women. Circulation 98, 731–733. https://doi.org/10.1161/01.cir.98.8.731
Riedel, B.C., Thompson, P.M., Brinton, R.D., 2016. Age, APOE and sex: Triad of risk of
Alzheimer's disease. The Journal of Steroid Biochemistry and Molecular Biology 160,
134–147. https://doi.org/10.1016/j.jsbmb.2016.03.012
Rosa, M.D., Dell'Ombra, N., Zambito, A.M., Malaguarnera, M., Nicoletti, F., Malaguarnera, L.,
2006. Chitotriosidase and inflammatory mediator levels in Alzheimer's disease and
cerebrovascular dementia. European Journal of Neuroscience 23, 2648–2656.
https://doi.org/10.1111/j.1460-9568.2006.04780.x
Setiawan, E., Wilson, A.A., Mizrahi, R., Rusjan, P.M., Miler, L., Rajkowska, G., Suridjan, I.,
Kennedy, J.L., Rekkas, P.V., Houle, S., Meyer, J.H., 2015. Role of translocator protein
density, a marker of neuroinflammation, in the brain during major depressive episodes.
JAMA Psychiatry 72, 268–275. https://doi.org/10.1001/jamapsychiatry.2014.2427
Sharp, E.S., Gatz, M., 2011. The Relationship between Education and Dementia An Updated
Systematic Review. Alzheimer Dis Assoc Disord 25, 289–304.
https://doi.org/10.1097/WAD.0b013e318211c83c
Steenland, K., Goldstein, F.C., Levey, A., Wharton, W., 2016. A Meta-Analysis of Alzheimer's
Disease Incidence and Prevalence Comparing African-Americans and Caucasians. J.
Alzheimers Dis. 50, 71–76. https://doi.org/10.3233/JAD-150778
Swardfager, W., Lanctôt, K., Rothenburg, L., Wong, A., Cappell, J., Herrmann, N., 2010. A
Meta-Analysis of Cytokines in Alzheimer's Disease. Biological Psychiatry 68, 930–941.
https://doi.org/10.1016/j.biopsych.2010.06.012
Teschendorff, A.E., Marabita, F., Lechner, M., Bartlett, T., Tegner, J., Gomez-Cabrero, D.,
Beck, S., 2013. A beta-mixture quantile normalization method for correcting probe
design bias in Illumina Infinium 450 k DNA methylation data. Bioinformatics 29, 189–
196. https://doi.org/10.1093/bioinformatics/bts680
Theendakara, V., Peters-Libeu, C.A., Bredesen, D.E., Rao, R.V., 2018. Transcriptional Effects of
ApoE4: Relevance to Alzheimer's Disease. Molecular Neurobiology 55, 5243–5254.
https://doi.org/10.1007/s12035-017-0757-2
Theendakara, V., Peters-Libeu, C.A., Spilman, P., Poksay, K.S., Bredesen, D.E., Rao, R.V.,
2016. Direct Transcriptional Effects of Apolipoprotein E. The Journal of Neuroscience
36, 685–700. https://doi.org/10.1523/JNEUROSCI.3562-15.2016
Toledo, J.B., Toledo, E., Weiner, M.W., Jack, C.R., Jagust, W., Lee, V.M.-Y., Shaw, L.M.,
Trojanowski, J.Q., 2012. Cardiovascular risk factors, cortisol, and amyloid-β deposition
in Alzheimer's Disease Neuroimaging Initiative. Alzheimers Dement 8, 483–489.
https://doi.org/10.1016/j.jalz.2011.08.008
Triche, T.J., Weisenberger, D.J., Van Den Berg, D., Laird, P.W., Siegmund, K.D., 2013. Low-
level processing of Illumina Infinium DNA Methylation BeadArrays. Nucleic Acids Res
41, e90. https://doi.org/10.1093/nar/gkt090
Vasanthakumar, A., Davis, J.W., Idler, K., Kim, S., Nho, K., Nudelman, K.N.H., Faber, K., Bai,
Y., Foroud, T.M., Estrada, K., Li, Q., Saykin, A.J., Waring, J., 2017. DNA
METHYLATION DYNAMICS IN ALZHEIMER'S DISEASE DIAGNOSIS AND
PROGRESSION. Alzheimer's & Dementia 13, P297.
https://doi.org/10.1016/j.jalz.2017.06.209

Walker, K.A., Gottesman, R.F., Wu, A., Knopman, D.S., Gross, A.L., Mosley, T.H., Selvin, E.,
Windham, B.G., 2019. Systemic inflammation during midlife and cognitive change over
20 years. Neurology 92, e1256–e1267.
Walker, K.A., Windham, B.G., Brown IV, C.H., Knopman, D.S., Jack Jr, C.R., Mosley Jr, T.H.,
Selvin, E., Wong, D.F., Hughes, T.M., Zhou, Y., Gross, A.L., Gottesman, R.F., 2018.
The Association of Mid- and Late-Life Systemic Inflammation with Brain Amyloid
Deposition: The ARIC-PET Study. Journal of Alzheimer's Disease 66, 1041–1052.
https://doi.org/10.3233/JAD-180469
Walters, A., Phillips, E., Zheng, R., Biju, M., Kuruvilla, T., 2016. Evidence for
neuroinflammation in Alzheimer's disease: Neuroinflammation in Alzheimer's. Progress
in Neurology and Psychiatry 20, 25–31. https://doi.org/10.1002/pnp.444
Wang, X., Zhou, W., Ye, T., Lin, X., Zhang, J., Initiative, for A.D.N., 2019. Sex Difference in
the Association of APOE4 with Memory Decline in Mild Cognitive Impairment. Journal
of Alzheimer's Disease 69, 1161–1169. https://doi.org/10.3233/JAD-181234
Wegner, A., Benson, S., Rebernik, L., Spreitzer, I., Jäger, M., Schedlowski, M., Elsenbruch, S.,
Engler, H., 2017. Sex differences in the pro-inflammatory cytokine response to endotoxin
unfold in vivo but not ex vivo in healthy humans. Innate Immun 23, 432–439.
https://doi.org/10.1177/1753425917707026
Xu, W., Tan, Lan, Wang, H.-F., Jiang, T., Tan, M.-S., Tan, Lin, Zhao, Q.-F., Li, J.-Q., Wang, J.,
Yu, J.-T., 2015. Meta-analysis of modifiable risk factors for Alzheimer's disease. J
Neurol Neurosurg Psychiatry 86, 1299–1306. https://doi.org/10.1136/jnnp-2015-310548
Yokoyama, A.S., Rutledge, J.C., Medici, V., 2017. DNA methylation alterations in Alzheimer's
disease. Environ Epigenet 3. https://doi.org/10.1093/eep/dvx008
Yun, Y.-W., Kweon, S.-S., Choi, J.-S., Rhee, J.-A., Lee, Y.-H., Nam, H.-S., Jeong, S.-K., Park,
K.-S., Ryu, S.-Y., Choi, S.-W., Kim, H.N., Cauley, J.A., Shin, M.-H., 2015. APOE
Polymorphism Is Associated with C-reactive Protein Levels but Not with White Blood
Cell Count: Dong-gu Study and Namwon Study. J Korean Med Sci 30, 860–865.
https://doi.org/10.3346/jkms.2015.30.7.860
Zenaro, E., Piacentino, G., Constantin, G., 2017. The blood-brain barrier in Alzheimer's disease.
Neurobiology of Disease, Brain Barriers in Health and Disease 107, 41–56.
https://doi.org/10.1016/j.nbd.2016.07.007
892

FIGURE CAPTIONS

Figure 1. Marginal mean (±95% confidence interval) of **CSF levels** of A. IL-16 (pg/ml), B. IL-8

(pg/ml), C. C-reactive protein (CRP; μg/ml), and **plasma levels** of D. CRP (CRP; μg/ml), E. IL-

18 (pg/ml), and F. IgE (ng/ml) in ADNI participants by sex and presence or absence of APOEε4

alleles (none, or 1 and 2 alleles).

Figure 2. Marginal mean (± 95% confidence interval) of **CSF levels** of A. IL-16 (pg/ml), B.

IgA (mg/ml), C. Intercellular adhesion molecule (ICAM1; ng/ml), and **plasma levels** of D. C-

reactive protein (CRP; μg/ml), E. cortisol (ng/ml), and F. IL-16 in ADNI participants by sex and

diagnosis (CN, cognitively normal; LMCI, late mild cognitive impairment; and AD, Alzheimer's

disease).

Figure 3. Universal epigenetic age acceleration does not differ statistically significantly by

participant sex or diagnosis (CN, EMCI, LMCI, AD) in this ADNI cohort. CN, cognitively

normal; EMCI, early mild cognitive impairment; LMCI, late mild cognitive impairment; AD,

Alzheimer's disease.

Figure 1

CSF

Plasma

Figure 2

CSF

Plasma

Figure 3

Table 1. Demographic and clinical information for all ADNI participants subdivided by sex. A. Participants with measured biomarkers in cerebrospinal fluid (CSF), B. Participants with available whole blood Illumina HumanMethylationEPIC DNA methylation data, C. Participants with matched Illumina HumanMethylationEPIC DNA methylation array data and measured CSF biomarkers. In all three subdata sets, females were significantly younger and had fewer years of education than males. In data set A (but not B and C), more females (24.0 % compared to 21.8%) were diagnosed with AD, more females were cognitively normal (26.5% compared to 22.9%) and fewer females were diagnosed with late MCI compared to males (49.5% compared to 55.3%). In data set A, females had lower CSF cortisol, interleukin-3, interleukin 8, interleukin-16, immunoglobulin A, and intercellular adhesion molecule 1 compared to males. Empty cells indicate data not available or not analysed.

	A				**B**				**C**			
		Sex				**Sex**				**Sex**		
	Total	**Female**	**Male**	**P-value**	**Total**	**Female**	**Male**	**P-value**	**Total**	**Female**	**Male**	**P-value**
	No. 279	**No. 109**	**No. 170**		**No. 640**	**No. 284**	**No. 356**		**No. 533**	**No. 243**	**No. 290**	
Age												
Mean (SD)	75.15 (±6.86)	73.75 (±6.69)	76.04 (±6.83)	0.007	75.63 (±7.68)	74.78 (±8.03)	76.31 (±7.32)	<0.0001	75.01 (±7.61)	74.31 (±8.10)	75.61 (±7.11)	0.0019
Education (years)												
Mean (SD)	15.69 (±2.95)	14.68 (±2.74)	16.34 (±2.90)	< 0.0001	16.21 (±2.70)	15.53 (±2.59)	16.75 (±2.68)	<0.0001	16.24 (±2.64)	15.57 (±2.49)	16.83 (±2.64)	< 0.0001
Ethnicity												
White	267 (95.70%)	103 (94.50%)	164 (96.47%)	0.55	627 (97.97%)	279 (98.23 %)	348 (97.75%)	0.78	521 (97.75 %)	238 (97.94 %)	283 (97.94 %)	0.99
Not White[1]	12 (4.30%)	6 (5.50%)	6 (3.53%)		13 (2.03%)	5 (1.76%)	8 (2.25%)		12 (2.25 %)	5 (2.06%)	7 (2.41 %)	
Baseline diagnosis												
CN	74 (26.5%)	35 (32.1%)	39 (22.9%)	0.051	217 (33.9%)	109 (38.38%)	108 (30.34%)	0.11	171 (32.08 %)	88 (36.21 %)	83 (28.62 %)	0.19
EMCI					186 (29.06%)	83 (29.23%)	103 (28.93%)		173 (32.46%)	79 (32.51%)	94 (32.41 %)	
LMCI	138 (49.5%)	44 (40.4%)	94 (55.3%)		200 (31.25%)	78 (27.46%)	122 (34.27%)		155 (29.08%)	94 (38.68%)	92 (31.72 %)	
AD	67 (24.0%)	30 (27.5%)	37 (21.8%)		37 (5.78%)	14 (4.23%)	23 (6.46%)		34 (6.38%)	13 (5.35 %)	21 (7.24 %)	
APOEε4 allele number												
0	134 (48.03%)	51 (46.79%)	83 (48.82%)	0.78	369 (57.66 %)	169 (59.51%)	200 (56.18%)	0.37	313 (58.72 %)	146 (60.08 %)	167 (57.59 %)	0.45
1	109 (39.07%)	42 (38.53%)	67 (39.41%)		220 (34.38%)	97 (34.15%)	123 (34.55%)		173 (32.46%)	80 (32.92 %)	93 (32.07 %)	
2	36 (12.90%)	16 (14.68%)	20 (11.76%)		51 (7.97%)	18 (6.34%)	33 (9.27%)		47 (8.82%)	17 (7.00%)	30 (10.34 %)	
Cortisol (ng/mL)												
Mean (SD)	16.05 (±6.04)	14.92 (±6.01)	16.78 (±5.96)	0.008								
C reactive protein (ug/mL)												
Mean (SD)	-2.83 (±0.56)	-2.77 (±0.64)	-2.87 (±0.51)	0.23								
CD40 antigen (ng/mL)												
Mean (SD)	-0.65 (±0.12)	-0.66 (±0.10)	-0.64 (±0.14)	0.12								
Interleukin 16 (pg/mL)												
Mean (SD)	0.91 (±0.18)	0.87 (±0.17)	0.94 (±0.19)	0.004								
Interleukin 3 (ng/mL)												
Mean (SD)	-2.22 (±0.32)	-2.28 (±0.29)	-2.17 (±0.34)	0.001								
Interleukin 6 receptor (ng/mL)												
Mean (SD)	-0.01 (±0.15)	-0.02 (±0.14)	-0.00 (±0.15)	0.30								
Interleukin 8 (pg/mL)												
Mean (SD)	1.68 (±0.15)	1.64 (±0.11)	1.70 (±0.16)	0.001								
Intercellular adhesion molecule 1 (ng/mL)												
Mean (SD)	0.96 (±0.44)	0.83 (±0.33)	1.04 (±0.48)	0.0001								
Immunoglobulin A (mg/mL)												
Mean (SD)	-2.54 (±0.31)	-2.68 (±0.26)	-2.45 (±0.31)	< 0.0001								
Amyloid Beta												
Mean (SD)									1040.98 (±454.72)	1055.50 (±449.23)	1028.35 (±459.36)	0.18
Tau												
Mean (SD)									289.80 (±124.68)	300.90 (±139.07)	280.13 (±109.82)	0.072
PTau												
Mean (SD)									27.47 (±13.65)	28.25 (±15.08)	26.78 (±12.24)	0.36

P-values are from Wilcoxon rank sum tests for continuous variables and Fisher's exact tests for categorical variables. ⅼIncludes self-reported Black, Asian, American Indian/Alaskan, and >1 ethnicity.

Table 2. Linear regression results for models with sex and APOE status. P-values are for overall tests and are FDR-adjusted. Only shown are the models with significant associations (adjusted P<0.05) and trends (adjusted P<0.08). All model summaries are available in Supplementary Table S1.

Predictors	Cortisol Cortisol ng ml Estimates (CI)	p	adjusted p	C Reactive Protein ug/ml Estimates (CI)	p	adjusted p	Interleukin 16 pg/ml Estimates (CI)	p	adjusted p	Interleukin 6 receptor ng/ml Estimates (CI)	p	adjusted p	Interleukin 8 pg/ml Estimates (CI)	p	adjusted p	Immunoglobulin A mg/ml Estimates (CI)	p	adjusted p	Intercellular Adhesion Molecule ng/ml Estimates (CI)	p	adjusted p
(Intercept)	1.43 (-7.00 – 9.87)			-3.03 (-3.84 – -2.23)			0.42 (0.15 – 0.68)			-0.27 (-0.48 – -0.06)			1.33 (1.11 – 1.54)			-2.91 (-3.35 – -2.48)			-0.23 (-0.83 – 0.38)		
AGE (years)	0.22 (0.12 – 0.32)			0.01 (-0.00 – 0.02)			0.01 (0.00 – 0.01)			0.00 (0.00 – 0.01)			0.00 (0.00 – 0.01)			0.00 (-0.00 – 0.01)			0.01 (0.01 – 0.02)		
EDUCATION (years)	-0.21 (-0.45 – 0.03)			-0.01 (-0.03 – 0.02)			-0.01 (-0.01 – 0.00)			-0.00 (-0.01 – 0.00)			0.00 (-0.00 – 0.01)			0.00 (-0.01 – 0.01)			-0.00 (-0.02 – 0.02)		
Male (ref = Female)	1.72 (0.25 – 3.19)	0.022	**0.07**	-0.11 (-0.25 – 0.03)	0.126	0.22	0.13 (0.07 – 0.19)			0.02 (-0.02 – 0.05)	0.372	0.5	0.09 (0.04 – 0.14)	<0.001		0.21 (0.14 – 0.29)	<0.001	**<0.001**	0.18 (0.07 – 0.28)	0.001	**0.009**
APOE4 1 or 2 alleles (ref = 0 alleles)	0.82 (-0.55 – 2.19)	0.241	0.35	-0.22 (-0.35 – -0.09)	0.001	**0.009**	0.08 (0.01 – 0.14)			0.04 (0.00 – 0.07)	0.025	**0.071**	0.03 (-0.02 – 0.09)	0.261		-0.00 (-0.07 – 0.07)	0.898	0.97	0.08 (-0.02 – 0.18)	0.128	0.22
Interaction: Male by 1 or 2 alleles							-0.13 (-0.22 – -0.05)	0.003	**0.016**				-0.09 (-0.16 – -0.02)	**0.008**	**0.035**						
Observations	279			279			279			279			279			279			279		
R² / adjusted R²	0.095 / 0.082			0.055 / 0.042			0.122 / 0.106			0.046 / 0.032			0.092 / 0.076			0.123 / 0.111			0.105 / 0.092		

Table 3. Linear regression results for models with sex and baseline diagnosis. Only shown are the models with significant associations. P-values are for overall tests and are FDR-adjusted. All model summaries are available in Supplementary Table S2.

Predictors	Interleukin 16 pg/ml			Immunoglobulin A mg/ml			Intercellular Adhesion Molecule ng/ml		
	Estimates (CI)	*p*	*adjusted p*	*Estimates (CI)*	*p*	*adjusted p*	*Estimates (CI)*	*p*	*adjusted p*
(Intercept)	0.51 (0.26 – 0.77)			-2.92 (-3.35 – -2.48)			-0.19 (-0.80 – 0.42)		
AGE (years)	0.01 (0.00 – 0.01)			0.00 (-0.00 – 0.01)			0.01 (0.01 – 0.02)		
EDUCATION (years)	-0.01 (-0.01 – 0.00)			0.00 (-0.01 – 0.01)			-0.00 (-0.02 – 0.02)		
Male (ref = Female)	0.06 (0.02 – 0.11)	0.007	**0.055**	0.21 (0.14 – 0.29)	<0.001	**<0.001**	0.17 (0.06 – 0.28)	0.002	**0.026**
Diagnosis (ref = CN)		0.4	0.61		0.99	0.99		0.54	0.67
LMCI	0.01 (-0.05 – 0.06)			0.00 (-0.08 – 0.09)			0.06 (-0.06 – 0.18)		
AD	-0.03 (-0.09 – 0.03)			-0.01 (-0.11 – 0.09)			0.02 (-0.12 – 0.16)		
Observations	279			279			279		
R^2 / adjusted R^2	0.099 / 0.082			0.124 / 0.107			0.101 / 0.085		

Table 4. Results of epigenetic age and epigenetic age acceleration calculation for all DNAme analyses, for both the larger DNAme cohort and the subset of samples with matched CSF biomarker data.

| | DNAme Cohort | | | | DNAme & CSF Biomarker Data Cohort | | | |
| | | Sex | | | | | Sex | |
	Total No. 640	Female No. 284	Male No. 356	P-value	Total No. 533	Female No. 243	Male No. 290	P-value
Age								
Mean (SD)	75.63 (±7.68)	74.78 (±8.03)	76.31 (±7.32)	<0.0001	75.01 (±7.61)	74.31 (±8.10)	75.61 (±7.11)	0.0019
Epigenetic Age (years)								
Mean (SD)	69.92 (±8.06)	67.45 (±8.15)	70.11 (±7.79)	<0.0001	68.47 (±8.17)	67.05 (±8.33)	69.72 (±7.82)	<0.0001
Epigenetic Age Acceleration (years)								
Mean (SD)	0.025 (±4.22)	-0.14 (±4.16)	0.16 (±4.26)	0.1	0.027 (±4.30)	-0.18 (±4.23)	0.20 (4.35)	0.057
Intrinsic Age Acceleration (years)								
Mean (SD)	0.026 (±4.11)	-0.19 (±4.06)	0.20 (±4.15)	0.019	0.020 (±4.18)	-0.25 (±4.14)	0.26 (±4.21)	0.021

P-values are from Wilcoxon rank sum tests for continuous variables and Fisher's exact tests for categorical variables

Table 5. Linear model for assessment of relationship of biochemical concentrations and presence of APOE ε4 risk allele on universal epigenetic age acceleration. Intrinsic epigenetic age acceleration linear model not shown.

Predictors	Estimates	CI	adjusted p
Age Acceleration & CSF Biomarkers			
(Intercept)	-0.61	-1.67 – 0.46	0.61
Male (ref = Female)	0.6	-0.11 – 1.31	0.34
Diagnosis (ref = CN)			
EMCI	0.71	-0.17 – 1.60	0.39
LMCI	0.4	-0.55 – 1.36	0.64
AD	0.53	-1.07 – 2.14	0.64
APOE risk allele			
1 or 2 alleles (ref = 0 alleles)	-0.24	-1.01 – 1.31	0.813
CSF Tau	0.0001	-0.0077 – 0.023	0.95
Observations	533		
R^2 / adjusted R^2	0.0108/-0.000438		

Supplemental File

Inflammation and epigenetic age in Alzheimer's disease: do sex and APOE matter?

Paula Duarte-Guterman[1*], Amy M. Inkster[2,3*], Arianne Y. Albert[4], Cindy K. Barha[5], Wendy P.

Robinson[2,3], Liisa A.M. Galea[1], on behalf of the Alzheimer's Disease Neuroimaging Initiative[δ]

1. Djavad Mowafaghian Centre for Brain Health and Department of Psychology, University of British Columbia, Vancouver, BC, Canada
2. Department of Medical Genetics, University of British Columbia, Vancouver, BC, Canada.
3. BC Children's Hospital Research Institute, Vancouver, BC, Canada
4. Women's Health Research Institute of British Columbia, Vancouver, BC, Canada
5. Djavad Mowafaghian Centre for Brain Health and Department of Physical Therapy, University of British Columbia, Vancouver, BC, Canada

Address all correspondence and requests for reprints to:
L. A. M. Galea, PhD
Djavad Mowafaghian Centre for Brain Health
2215 Wesbrook Mall
Vancouver, British Columbia
V6T 1Z3, Canada
E-mail: lgalea@psych.ubc.ca.

* Considered co-first authors.

[δ] Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Table S1. Linear regression results for all variables investigated by sex and APOE status. Markers in CSF

Predictors	Cortisol Cortisol ng ml Estimates (CI)	p	adjusted p	C Reactive Protein ug/ml Estimates (CI)	p	adjusted p	CD 40 antigen ng/ml Estimates (CI)	p	adjusted p	Interleukin 16 pg/ml Estimates (CI)	p	adjusted p	Interleukin 3 ng/ml Estimates (CI)	p	adjusted p
(Intercept)	1.43 (-7.00 – 9.87)			-3.03 (-3.84 – -2.23)			-1.10 (-1.27 – -0.93)			0.42 (0.15 – 0.68)			-3.03 (-3.49 – -2.57)		
AGE (years)	0.22 (0.12 – 0.32)			0.01 (-0.00 – 0.02)			0.01 (0.00 – 0.01)			0.01 (0.00 – 0.01)			0.01 (0.00 – 0.02)		
EDUCATION (years)	-0.21 (-0.45 – 0.03)			-0.01 (-0.03 – 0.02)			-0.00 (-0.01 – 0.00)			-0.01 (-0.01 – 0.00)			0.00 (-0.01 – 0.01)		
Male (ref = Female)	1.72 (0.25 – 3.19)	0.022	**0.07**	-0.11 (-0.25 – 0.03)	0.126	0.22	0.01 (-0.02 – 0.04)	0.432	0.55	0.13 (0.07 – 0.19)			0.08 (-0.00 – 0.16)	0.057	0.15
APOE4 1 or 2 alleles (ref = 0 alleles)	0.82 (-0.55 – 2.19)	0.241	0.35	-0.22 (-0.35 – -0.09)	0.001	**0.009**	0.01 (-0.02 – 0.04)	0.59	0.72	0.08 (0.01 – 0.14)			-0.04 (-0.12 – 0.03)	0.262	0.37
Interaction: Male by 1 or 2 alleles										-0.13 (-0.22 – -0.05)	0.003	**0.016**			
Observations	279			279			279			279			279		
R^2 / adjusted R^2	0.095 / 0.082			0.055 / 0.042			0.125 / 0.112			0.122 / 0.106			0.076 / 0.063		

Table S1. Continued

Predictors	Interleukin 6 receptor ng/ml Estimates (CI)	p	adjusted p	Interleukin 8 pg/ml Estimates (CI)	p	adjusted p	Immunoglobulin A mg/ml Estimates (CI)	p	adjusted p	Intercellular Adhesion Molecule ng/ml Estimates (CI)	p	adjusted p
(Intercept)	-0.27 (-0.48 – -0.06)			1.33 (1.11 – 1.54)			-2.91 (-3.35 – -2.48)			-0.23 (-0.83 – 0.38)		
AGE (years)	0.00 (0.00 – 0.01)			0.00 (0.00 – 0.01)			0.00 (-0.00 – 0.01)			0.01 (0.01 – 0.02)		
EDUCATION (years)	-0.00 (-0.01 – 0.00)			0.00 (-0.00 – 0.01)			0.00 (-0.01 – 0.01)			-0.00 (-0.02 – 0.02)		
Male (ref = Female)	0.02 (-0.02 – 0.05)	0.372	0.5	0.09 (0.04 – 0.14)	<0.001		0.21 (0.14 – 0.29)	<0.001	**<0.001**	0.18 (0.07 – 0.28)	0.001	**0.009**
APOE4 1 or 2 alleles (ref = 0 alleles)	0.04 (0.00 – 0.07)	0.025	**0.071**	0.03 (-0.02 – 0.09)	0.261		-0.00 (-0.07 – 0.07)	0.898	0.97	0.08 (-0.02 – 0.18)	0.128	0.22
Interaction: Male by 1 or 2 alleles				-0.09 (-0.16 – -0.02)	0.008	**0.035**						
Observations	279			279			279			279		
R^2 / adjusted R^2	0.046 / 0.032			0.092 / 0.076			0.123 / 0.111			0.105 / 0.092		

Table S2. Linear regression results for all variables investigated by sex and baseline diagnosis. Markers in CSF

Predictors	Cortisol Cortisol ng ml Estimates (CI)	p	adjusted p	C Reactive Protein ug/ml Estimates (CI)	p	adjusted p	CD 40 antigen ng/ml Estimates (CI)	p	adjusted p	Interleukin 16 pg/ml Estimates (CI)	p	adjusted p	Interleukin 3 ng/ml Estimates (CI)	p	adjusted p
(Intercept)	1.49 (-6.96 – 9.95)			-3.12 (-3.94 – -2.30)			-1.08 (-1.25 – -0.91)			0.51 (0.26 – 0.77)			-3.03 (-3.49 – -2.57)		
AGE (years)	0.22 (0.12 – 0.32)			0.01 (-0.00 – 0.02)			0.01 (0.00 – 0.01)			0.01 (0.00 – 0.01)			0.01 (0.00 – 0.02)		
EDUCATION (years)	-0.23 (-0.47 – 0.01)			-0.00 (-0.03 – 0.02)			-0.00 (-0.01 – 0.00)			-0.01 (-0.01 – 0.00)			0.00 (-0.01 – 0.01)		
Male (ref = Female)	1.55 (0.07 – 3.04)	0.041	0.18	-0.10 (-0.24 – 0.05)	0.19	0.42	0.01 (-0.02 – 0.04)	0.51	0.65	0.06 (0.02 – 0.11)	0.007	**0.055**	0.08 (-0.00 – 0.16)	0.052	0.18
Diagnosis (ref = CN)		0.24	0.49		0.15	0.36		0.067	0.2		0.4	0.61		0.5	0.65
LMCI	1.23 (-0.43 – 2.88)			-0.15 (-0.31 – 0.01)			0.01 (-0.03 – 0.04)			0.01 (-0.05 – 0.06)			-0.04 (-0.13 – 0.05)		
AD	0.10 (-1.83 – 2.02)			-0.15 (-0.34 – 0.03)			-0.03 (-0.07 – 0.00)			-0.03 (-0.09 – 0.03)			-0.06 (-0.17 – 0.04)		
Observations	279			279			279			279			279		
R^2 / adjusted R^2	0.100 / 0.084			0.030 / 0.012			0.141 / 0.125			0.099 / 0.082			0.077 / 0.060		

Table S2. Continued

Predictors	Interleukin 6 receptor ng/ml Estimates (CI)	p	adjusted p	Interleukin 8 pg/ml Estimates (CI)	p	adjusted p	Immunoglobulin A mg/ml Estimates (CI)	p	adjusted p	Intercellular Adhesion Molecule ng/ml Estimates (CI)	p	adjusted p
(Intercept)	-0.22 (-0.43 – -0.01)			1.35 (1.14 – 1.56)			-2.92 (-3.35 – -2.48)			-0.19 (-0.80 – 0.42)		
AGE (years)	0.00 (0.00 – 0.01)			0.00 (0.00 – 0.01)			0.00 (-0.00 – 0.01)			0.01 (0.01 – 0.02)		
EDUCATION (years)	-0.00 (-0.01 – 0.00)			0.00 (-0.00 – 0.01)			0.00 (-0.01 – 0.01)			-0.00 (-0.02 – 0.02)		
Male (ref = Female)	0.02 (-0.02 – 0.05)	0.42	0.61	0.04 (0.01 – 0.08)	0.019	0.12	0.21 (0.14 – 0.29)	<0.001	**<0.001**	0.17 (0.06 – 0.28)	0.002	**0.026**
Diagnosis (ref = CN)		0.46	0.65		0.85	0.96		0.99	0.99		0.54	0.67
LMCI	0.01 (-0.04 – 0.05)			0.01 (-0.03 – 0.05)			0.00 (-0.08 – 0.09)			0.06 (-0.06 – 0.18)		
AD	-0.02 (-0.07 – 0.03)			0.01 (-0.03 – 0.06)			-0.01 (-0.11 – 0.09)			0.02 (-0.12 – 0.16)		
Observations	279			279			279			279		
R^2 / adjusted R^2	0.034 / 0.016			0.062 / 0.045			0.124 / 0.107			0.101 / 0.085		

Table S3. Linear regression results for all plasma variables investigated by sex and APOE genotype

Predictors	Cortisol Cortisol ng ml			C Reactive Protein ug/ml			CD 40 antigen ng/ml			Interleukin 16 pg/ml			Interleukin 3 ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	2.06 (1.92 – 2.19)			0.46 (-0.05 – 0.97)			-0.52 (-0.65 – -0.39)			2.42 (2.27 – 2.58)			-1.86 (-2.16 – -1.57)		
AGE (years)	0.00 (-0.00 – 0.00)			0.00 (-0.00 – 0.01)			0.01 (0.00 – 0.01)			0.00 (0.00 – 0.00)			0.00 (-0.00 – 0.00)		
EDUCATION (years)	-0.00 (-0.00 – 0.00)			-0.03 (-0.04 – -0.01)			-0.00 (-0.01 – 0.00)			-0.01 (-0.01 – -0.00)			0.01 (0.00 – 0.02)		
Male (ref = Female)	0.01 (-0.01 – 0.04)	0.26	0.57	-0.13 (-0.22 – -0.04)	0.006	**0.048**	-0.04 (-0.07 – -0.01)			0.01 (-0.02 – 0.04)	0.50	0.79	-0.02 (-0.07 – 0.04)	0.57	0.8
APOE4 1 or 2 alleles (ref = 0 alleles)	0.02 (-0.00 – 0.04)	0.11	0.29	-0.31 (-0.39 – -0.22)	<0.001	**<0.001**	-0.03 (-0.07 – 0.00)			-0.01 (-0.04 – 0.01)	0.31	0.66	0.02 (-0.03 – 0.07)	0.43	0.67
Interaction: Male by 1 or 2 alleles							0.05 (0.01 – 0.10)	**0.022**	0.11						
Observations	527			526			527			527			527		
R² / adjusted R²	0.013 / 0.005			0.123 / 0.117			0.126 / 0.118			0.042 / 0.033			0.011 / 0.004		

Table S3. Continued

Predictors	Interleukin 6 receptor ng/ml			Interleukin 8 pg/ml			Immunoglobulin A mg/ml			Intercellular Adhesion Molecule ng/ml			Interleukin 18 pg/ml			Immunoglobulin E ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	1.43 (1.29 – 1.57)			0.79 (0.60 – 0.99)			0.48 (0.25 – 0.71)			1.89 (1.75 – 2.04)			2.42 (2.24 – 2.59)			1.42 (0.88 – 1.97)		
AGE (years)	0.00 (-0.00 – 0.00)			0.00 (0.00 – 0.01)			0.00 (-0.00 – 0.00)			0.00 (0.00 – 0.00)			-0.00 (-0.00 – 0.00)			0.00 (-0.00 – 0.01)		
EDUCATION (years)	-0.00 (-0.01 – 0.00)			0.00 (-0.00 – 0.01)			-0.00 (-0.01 – 0.01)			0.00 (-0.00 – 0.00)			-0.00 (-0.01 – 0.00)			0.01 (-0.01 – 0.02)		
Male (ref = Female)	-0.02 (-0.05 – -0.00)	0.049	0.14	-0.05 (-0.10 – 0.00)			0.02 (-0.02 – 0.06)	0.371	0.67	-0.04 (-0.06 – -0.01)	0.008	**0.051**	0.06 (0.03 – 0.09)	<0.001	**0.001**	0.24 (0.14 – 0.33)	<0.001	**<0.001**
APOE4 1 or 2 alleles (ref = 0 alleles)	0.00 (-0.02 – 0.03)	0.70	0.84	-0.08 (-0.13 – -0.02)			-0.02 (-0.06 – 0.02)	0.387	0.67	0.01 (-0.02 – 0.03)	0.586	0.80	-0.04 (-0.07 – -0.01)	0.019	0.10	0.01 (-0.09 – 0.10)	0.91	0.94
Interaction: Male by 1 or 2 alleles				0.07 (0.00 – 0.14)	0.046	0.14												
Observations	527			527			527			527			527			527		
R² / adjusted R²	0.011 / 0.004			0.036 / 0.026			0.019 / 0.012			0.008 / 0.000			0.038 / 0.031			0.052 / 0.045		

Table S4. Linear regression results for all plasma variables investigated by sex and baseline diagnosis.

Predictors	Cortisol Cortisol ng ml			C Reactive Protein ug/ml			CD 40 antigen ng/ml			Interleukin 16 pg/ml			Interleukin 3 ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	2.09 (1.95 – 2.22)			0.33 (-0.21 – 0.88)			-0.55 (-0.68 – -0.42)			2.48 (2.31 – 2.65)			-1.78 (-2.08 – -1.47)		
AGE (years)	0.00 (-0.00 – 0.00)			0.01 (0.00 – 0.01)			0.01 (0.00 – 0.01)			0.00 (0.00 – 0.00)			0.00 (-0.00 – 0.00)		
EDUCATION (years)	-0.00 (-0.00 – 0.00)			-0.02 (-0.04 – -0.01)			-0.00 (-0.00 – 0.00)			-0.01 (-0.01 – -0.00)			0.01 (0.00 – 0.02)		
Male (ref = Female)	0.02 (-0.01 – 0.04)	0.18	0.44	-0.12 (-0.21 – -0.03)	0.012	**0.056**	-0.01 (-0.03 – 0.01)	0.38	0.59	0.01 (-0.02 – 0.04)	0.39	0.57	-0.02 (-0.07 – 0.04)	0.56	0.75
Diagnosis (ref = CN)		0.0009	**0.010**		0.007	**0.056**		0.016	**0.067**		0.009	**0.054**		0.044	0.17
LMCI	-0.02 (-0.07 – 0.02)			-0.27 (-0.44 – -0.10)			-0.02 (-0.06 – 0.02)			-0.08 (-0.13 – -0.03)			-0.04 (-0.13 – 0.06)		
AD	0.03 (-0.02 – 0.08)			-0.24 (-0.43 – -0.05)			0.02 (-0.02 – 0.07)			-0.08 (-0.14 – -0.02)			-0.11 (-0.21 – -0.00)		
Observations	527			526			527			526			527		
R^2 / adjusted R^2	0.034 / 0.024			0.061 / 0.052			0.131 / 0.123			0.048 / 0.039			0.022 / 0.013		

Table S4. Continued

Predictors	Interleukin 6 receptor ng/ml			Interleukin 8 pg/ml			Immunoglobulin A mg/ml			Intercellular Adhesion Molecule ng/ml			Interleukin 18 pg/ml			Immunoglobulin E ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	1.47 (1.33 – 1.61)			0.73 (0.54 – 0.93)			0.43 (0.20 – 0.67)			1.90 (1.75 – 2.05)			2.33 (2.15 – 2.51)			1.44 (0.89 – 2.00)		
AGE (years)	0.00 (-0.00 – 0.00)			0.00 (0.00 – 0.01)			0.00 (-0.00 – 0.00)			0.00 (-0.00 – 0.00)			0.00 (-0.00 – 0.00)			0.00 (-0.00 – 0.01)		
EDUCATION (years)	-0.00 (-0.01 – 0.00)			0.00 (-0.00 – 0.01)			-0.00 (-0.01 – 0.01)			0.00 (-0.00 – 0.01)			-0.00 (-0.01 – 0.00)			0.01 (-0.01 – 0.02)		
Male (ref = Female)	-0.02 (-0.05 – 0.00)	0.063	0.19	-0.01 (-0.04 – 0.03)	0.65	0.79	0.02 (-0.02 – 0.06)	0.42	0.59	-0.03 (-0.06 – -0.01)	0.011	**0.056**	0.06 (0.03 – 0.09)	<0.001	**0.004**	0.24 (0.14 – 0.33)	<0.001	**<0.001**
Diagnosis (ref = CN)		0.34	0.56		0.11	0.30		0.72	0.82		0.063	0.19		0.32	0.56		0.88	0.94
LMCI	-0.03 (-0.08 – 0.01)			-0.04 (-0.10 – 0.02)			0.03 (-0.05 – 0.10)			-0.01 (-0.06 – 0.03)			0.04 (-0.01 – 0.10)			-0.00 (-0.18 – 0.17)		
AD	-0.03 (-0.08 – 0.02)			-0.00 (-0.07 – 0.07)			0.02 (-0.07 – 0.10)			0.02 (-0.03 – 0.08)			0.03 (-0.03 – 0.10)			-0.03 (-0.23 – 0.16)		
Observations	527			527			527			527			527			527		
R^2 / adjusted R^2	0.015 / 0.006			0.029 / 0.020			0.008 / -0.002			0.029 / 0.019			0.032 / 0.023			0.053 / 0.043		

Table S5. Correlation matrix of biochemical variables for those ADNI samples with available DNAme and biochemical data (n=533). Correlation coefficients are Spearman's rho.

	Amyloid Beta	p value	Tau	p value	pTau	p value
Amyloid Beta	1.00	<0.0001	-0.18	<0.0001	-0.27	<0.0001
Tau	-0.18	<0.0001	1.00	<0.0001	0.98	<0.0001
pTau	-0.27	<0.0001	0.98	<0.0001	1.00	<0.0001

Table S6. Contingency table for the distribution of the APOEε4 alleles (0 or 1 and 2 ε4 alleles) in each of the diagnosis groups (CN, late MCI and AD).

	CN	LMCI	AD
0	55 (74.32%)	60 (43.48%)	19(28.36%)
1 and 2	19 (25.68%)	78 (56.52%)	48 (71.64%)