Inflammation and epigenetic age in Alzheimer's disease: do sex and APOE matter?

Paula Duarte-Guterman[1*], Amy M. Inkster[2,3*], Arianne Y. Albert[4], Cindy K. Barha[5], Wendy P.

Robinson[2,3], Liisa A.M. Galea[1], on behalf of the Alzheimer's Disease Neuroimaging Initiative[δ]

1. Djavad Mowafaghian Centre for Brain Health and Department of Psychology, University of

British Columbia, Vancouver, BC, Canada

2. Department of Medical Genetics, University of British Columbia, Vancouver, BC, Canada.

3. BC Children's Hospital Research Institute, Vancouver, BC, Canada

4. Women's Health Research Institute of British Columbia, Vancouver, BC, Canada

5. Djavad Mowafaghian Centre for Brain Health and Department of Physical Therapy,

University of British Columbia, Vancouver, BC, Canada

13
Address all correspondence and requests for reprints to:
L. A. M. Galea, PhD
Djavad Mowafaghian Centre for Brain Health
2215 Wesbrook Mall
Vancouver, British Columbia
V6T 1Z3, Canada
E-mail: lgalea@psych.ubc.ca.
21
22
23 * Considered co-first authors.
24
25
[δ] Data used in preparation of this article were obtained from the Alzheimer's Disease
Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within
the ADNI contributed to the design and implementation of ADNI and/or provided data but did
not participate in analysis or writing of this report. A complete listing of ADNI investigators can
be found at: http://adni.loni.usc.edu/wp-
content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf
32
33

34 **Running title:** Inflammation and epigenetic age in AD

ABSTRACT

Alzheimer's disease (AD) disproportionately affects females with steeper cognitive decline and more neuropathology compared to males, which is exacerbated in females carrying the APOEε4 allele. The risk of developing AD is also higher in female APOEε4 carriers in earlier age groups (aged 65-75), and the progression from cognitively normal to mild cognitive impairment (MCI) and to AD may be sex-specific. The objective of this study was to investigate underlying physiological inflammatory and epigenetic mechanisms that may help explain why there are sex differences in AD and APOEε4 carriers. Inflammation is observed in AD and is related to aging, stress and neuroplasticity, and aging biomarkers also include changes in DNA methylation patterns which have been used to determine epigenetic age. Although studies are scarce, sex differences are noted in inflammation and in epigenetic aging. We investigated, using the ADNI database, the effect of sex and APOE genotype (non-carriers or carriers of 1 or 2 APOEε4 alleles) and sex and diagnosis (cognitively normal (CN), MCI, AD) on CSF (N= 279) and plasma (N= 527) markers of stress and inflammation, and blood epigenetic age (N= 640). We found CSF IL-16 and IL-8 levels were significantly lower in females non-carriers of APOEε4 alleles compared to males, whereas levels were similar between the sexes among carriers of APOEε4 alleles. Furthermore, females had higher levels of plasma CRP and ICAM1 but lower levels of CSF ICAM1, IL-8, IL-16, and IgA than males. Carrying 1 or 2 APOEε4 alleles and diagnosis (MCI and AD) decreased plasma CRP in both sexes. Neither sex, diagnosis, nor presence of APOEε4 alleles affected epigenetic age acceleration. Sex differences in inflammatory biomarkers support that the underlying physiological changes during aging differ by sex and tissue origin.

Keywords: Sex differences; Alzheimer's disease; APOE genotype; inflammation; cytokines; epigenetic age.

INTRODUCTION

Alzheimer's disease (AD) is a neurodegenerative disease characterized by severe cognitive decline [1]. Risk factors for AD include modifiable risk factors such as sociocultural or lifestyle factors (e.g., education, marital status, exercise), chronic stress exposure [2], and medical conditions (diabetes, obesity, and cardiovascular disease) [3–5]. Non-modifiable lifetime risk factors for AD include age, female sex, and APOE genotype [6]. However, research on the effects of biological sex on risk for AD is equivocal and may depend on geographic location (reviewed in [4,7,8]). Nevertheless, females with AD show greater cognitive decline [9–11] and neuropathology compared to males (faster brain atrophy rates, neurofibrillary tangles; [10,12–15]). Intriguingly, the presence of APOEε4 alleles increases the risk to develop AD in females compared to males at an earlier age (aged 65-75; [16]), and accelerates neuropathology and cognitive decline more so in females than males [10,11,14,17–19], indicating that the APOE genotype interacts with sex on various factors related to AD. However, there is limited research into the role of sex and its interaction with APOE genotype in the possible mechanisms underlying AD. Understanding why females in general and female APOEε4 carriers have a higher burden of the disease is important for the development of tailored treatments. Biomarkers are highly sought after to predict disease onset and progression and to understand the underlying mechanisms of diseases in order to develop or improve treatments.

Chronic low grade inflammation is a hallmark of AD, as evidenced by increased expression of proinflammatory cytokines in the brains of AD patients (not analyzed by sex), which can exacerbate AD pathology [20–22]. There is, however, increasing evidence that there

are sex differences in immune responses in healthy adults with females mounting a stronger

response compared to males after an acute challenge [23,24]. In response to an endotoxin,

females have higher levels of pro-inflammatory plasma cytokines (TNF-α and IL-6) while males

have higher plasma levels of anti-inflammatory IL-10 [23,25]. Marquez et al. [26] found that

aging affects the immune system differently in males and females, with females having higher

genomic activity for adaptive cells and males having activity for monocytes and inflammation.

Although limited, there is evidence that sex differences in systemic inflammation are associated

with greater AD pathology [27] but not cognitive decline in normal aging [28]. Specifically,

higher C-reactive protein (CRP) levels in blood beginning in midlife are associated with higher

brain amyloid levels later in life in healthy males, but not in healthy females [27]. To our

knowledge, very few studies have stratified by sex and APOE genotype or sex and diagnosis of

cognitive status on potential biomarkers of AD, including inflammation.

Sex differences in inflammatory biomarker systems may also differentially affect

neuroplasticity [29,30], which is reduced in AD and correlates with cognitive decline [31,32]. In

addition, peripheral cortisol, the main stress hormone in humans, is elevated in AD [33] and is

associated with higher amyloid levels in the brain [34], a reduction in hippocampal volume, and

cognitive impairment in older individuals [35] that may depend on MCI status [36]. Peripheral

cortisol is also associated with elevated pro-inflammatory cytokines [23,37]. However, it is not

known how sex differences in markers of inflammation (e.g., cytokines, immunoglobulins, CRP,

intercellular adhesion molecule, ICAM1), and stress hormones (cortisol) may be related to sex

differences in AD.

Aging biomarkers include epigenetic alterations that are associated with a variety of

pathologies and adverse health conditions, including normal cognitive aging and

neurodegenerative phenotypes such as AD [38–41]. Recently, molecular biomarkers of aging

known as "epigenetic clocks" have been developed based on DNA methylation (DNAme)

signatures [38,39]. Positive deviations of epigenetic age from chronological age (positive

epigenetic age acceleration) reflect more rapid biological aging and have been associated with

phenotypes including smoking, obesity, Parkinson's disease, Trisomy 21, and cancer [39,42–44],

while negative epigenetic age acceleration has been associated with high life-expectancy and

memory retention [45,46]. In healthy individuals, aging males exhibit more positive epigenetic

age acceleration than females in blood, buccal tissue, and in multiple brain regions [47]. In AD,

epigenetic age acceleration of the frontal cortex has been associated with amyloid load, neuritic

plates, and cognitive decline (not analyzed by sex; [40]). Within an individual, the DNA

methylation profile in peripheral tissues is correlated with the epigenetic signature in the brain

[48], thus it is reasonable to hypothesize that epigenetic age acceleration associated with AD

may also be detectable in peripheral tissues such as blood in AD participants. In AD, it is

possible that sex-specific pathology may be further reflected in sex differences in epigenetic age

acceleration.

Using the ADNI database, we examined sex differences in CSF and plasma physiological

biomarkers (inflammation and stress related) and blood epigenetic age, and how these may be

affected by APOE genotype (non-carriers or carriers of 1 or 2 APOEε4 alleles), and dementia

status (cognitively healthy (CN), MCI, AD). We tested the hypotheses that (1) females have

higher levels of inflammation and stress hormones compared to males and these levels are

disproportionately affected by the presence of APOEε4 alleles and AD diagnosis; and (2)

females have more positive epigenetic age acceleration and this is exacerbated by APOEε4

carriership and AD diagnosis.

METHODS

ADNI database

Data used in the preparation of this article were obtained from the Alzheimer's Disease

Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). The ADNI was launched in

2003 as a public-private partnership, led by Principal Investigator Michael W. Weiner,

MD. The primary goal of ADNI has been to test whether serial magnetic resonance imaging

(MRI), positron emission tomography (PET), other biological markers, and clinical and

neuropsychological assessment can be combined to measure the progression of mild cognitive

impairment (MCI) and early Alzheimer's disease (AD). For up-to-date information, see

www.adni-info.org. Data used in this article were downloaded on or before Jan 16, 2019.

Inclusion and exclusion criteria of participants [49,50] were the same for all the datasets

analysed in the current study (biomarkers in CSF and plasma, and epigenetic data), and general

procedures are detailed online (http://adni.loni.usc.edu/methods/documents/). Briefly,

cognitively normal (CN) participants had normal memory function based on education-adjusted

scores on the Wechsler Memory Scale Logical Memory II and a Clinical Dementia Rating

(CDR) of 0. Amnestic late MCI (LMCI) participants had objective memory loss (measured by

education-adjusted scores from Wechsler Memory Scale Logical Memory II), a CDR of 0.5,

preserved daily activities, and absence of dementia. Early MCI (EMCI) participants had a milder

episodic memory impairment compared to LMCI and were recruited using the Wechsler

Memory Scale Logical Memory II scores (between ~0.5 and 1.5 SD below the mean of CN

participants). All AD participants met NINCDS/ADRDA Alzheimer's Criteria and a CDR of 0.5

or 1.0.

Statistical Methods: Inflammatory markers

We included all ADNI participants that had inflammatory markers measured in CSF (N =

279) and plasma (N = 527) listed in Tables 1 and 2. Data included in our analyses were:

demographics (age, years of education, and ethnicity), baseline diagnosis (cognitively normal,

CN; late MCI, LMCI; or AD), and number of APOEε4 alleles. We collapsed APOE genotype

into two groups: (1) participants carrying any ε4 alleles (homozygous ε4/ε4 and heterozygous

ε4/-) and (2) participants with no ε4 risk alleles (-/-). Plasma and CSF samples from the ADNI

study were collected in CN, LMCI and AD participants at baseline in the morning after an

overnight fast. Processing, aliquoting and storage were performed according to the ADNI

Biomarker Core Laboratory Standard Operating Procedures. Inflammatory markers were

measured using a commercially available multiplex proteomic panel (Human Discovery Multi-

Analyte Profile; Luminex xMAP) developed by Rules-Based Medicine (Austin, TX), that

measures a variety of markers including cytokines, metabolic markers, and growth factors. We

initially chose biomarkers available in plasma involved in inflammation and immune responses

(cytokines, immunoglobulins, CRP, and ICAM1) and stress (cortisol; Table 2). We analysed the

same biomarkers in CSF (however IgE and IL-18 are not available in CSF). The protocols used

to quantify plasma and CSF analytes are described in Craig-Schapiro et al. [51] and Hu et al.

[52]. We used the ADNI quality-controlled data for plasma and CSF provided by the ADNI

Consortium. For plasma IL-16, we removed one outlier that was more than two times lower than

the 25[th] percentile in the plasma data. Sensitivity analysis with the outlier present suggested that

it was disproportionately influencing the results.

We compared all available data for each study variable between the sexes using the

Wilcoxon rank sum test for continuous variables and Fisher's exact test for categorical variables.

Nonparametric tests are standard for comparing variables where the distribution is unknown or

expected to be non-normal. We used general linear models to determine the relationships

between (1) sex and APOE genotype (non-carriers or carriers of 1 or 2 APOEε4 alleles) or (2)

sex and baseline diagnosis as predictor variables, and biomarkers as dependent variables. Due to

the limited sample size, we were not able to study sex, APOE genotype and baseline diagnosis in

one model. All models included age and education as covariates. All models initially included an

interaction between sex and presence of APOEε4 alleles or sex and baseline diagnosis; if this

interaction was not significant, it was removed from the model to estimate the main effects of

sex and APOE genotype or diagnosis. Significance was based on the likelihood ratio test, and all

P-values for comparisons of sex and either APOE genotype or diagnosis for all outcomes

combined were corrected for multiple testing using the Benjamini-Hochberg false discovery rate

method with the family-wise error rate set to 0.05 [53]. In total, three P-values per dependent

variable were included in each set of models (interaction term and main effects of sex and APOE

or diagnosis) resulting in 27 P-values corrected in CSF (9 dependent variables) and 33 P-values

corrected in plasma (11 dependent variables; Supplementary Tables S1 to S4) for each of the

models (sex and APOE and sex and diagnosis). Significant interaction terms were followed up

using pairwise simple-effects tests with Benjamini-Hochberg P-value correction. A subset of

participants with CSF measurements had corresponding plasma measurements (N= 273 total, N=

88 females and N=149 males). For each biomarker, we calculated Pearson's correlation

coefficients between CSF and plasma levels in males and females separately. We then compared

these correlations using Fisher'r to Z transformation and a Z-test using the method by Zou [54].

We report significance differences (adjusted P≤0.05) and trends (adjusted P≤0.08). All

regression analyses were carried out in R v3.5.1 [55].

Statistical Methods: Epigenetic Age

We used DNAme data quantified with the Illumina Infinium HumanMethylationEPIC

BeadChip array ("EPIC" array) for 1905 longitudinal blood samples from 640 unique ADNI

participants with CN, EMCI, LMCI, and AD diagnosis (n=284 females, n= 356 males) [56].

DNAme IDAT files were read into R v3.5.1 [55] using the 'minfi' package, and annotated with

the most recent version of the EPIC manifest, the Infinium MethylationEPIC v1.0 B4 Manifest

File, (available from https://support.illumina.com/downloads.html) [57,58]. We excluded 11 low

quality samples from 9 unique participants from further analyses on the basis of having median

methylated and unmethylated probe intensity <10.5 [57,58], the remaining samples were

background normalized and dye-bias adjusted with normal exponential out-of-band ("noob")

normalization [59]. DNAme data were converted to beta values and the biological sex of all

samples was confirmed by clustering samples on all array probes mapping to the X and Y

chromosomes. Beta values were calibrated to Horvath's 21,368-probe training dataset using the

calibrated beta-mixture quantile (BMIQ) normalization procedure suggested by Horvath, and

epigenetic age was calculated using R code modified for compatibility with the EPIC array using

the 334/353 epigenetic clock probes present on the array from https://horvath.genetics.ucla.edu/

[39,60]. Epigenetic age could have been calculated by imputing missing DNAme values at the 19

clock CpG sites not covered by the EPIC array based on the k-nearest neighbors method, but as

we observed a high correlation between epigenetic age values calculated with the missing probes

removed versus imputed with k=10 (R=0.99, P<2.2e-16), in agreement with previous reports

[61,62], we therefore chose to remove missing probes.

Prior to statistical analyses we removed all technical replicate samples. Epigenetic age

acceleration was calculated as the residual of epigenetic age regressed on chronological age and

technical batch covariates: the laboratory collection site at which blood samples were drawn and

EPIC microarray chip and row. Intrinsic epigenetic age acceleration, a measure designed to be

independent of age-related changes in whole blood cell-type proportions, was calculated as

described in Chen et al. [63] as the residual of epigenetic age regressed on chronological age,

technical covariates of collection site, row, and chip, and the proportions of six blood cell types

(CD8T, CD4T, NK, B cells, monocytes, and granulocytes) estimated from noob-normalized

DNA methylation data with the Houseman algorithm [64]. ADNI is a longitudinal study, and

participants contributed between 1-3 blood DNAme samples within the 2-year collection period.

Because these data collected over two years fell within the median 3.6-year error of Horvath's

epigenetic clock, longitudinal evaluation could not be meaningfully conducted. As such, for all

epigenetic age measures reported and analyzed herein, we calculated the average epigenetic age

acceleration measure per participant from all available time points, and performed all statistical

analyses on these mean values.

To determine if epigenetic age acceleration or intrinsic epigenetic age acceleration

differed by sex or dementia diagnosis, we used unbalanced two-way ANOVA designs. APOE

genotype and CSF biomarker (amyloid beta, tau and p-tau) data were available from the ADNI

repository for a smaller subset of participants with matched EPIC DNAme data, (n=533, see

Table 1C). As above, due to the small size of this cohort, we collapsed APOE genotype into two

groups: (1) non-carriers of ε4 risk alleles and (2) carriers of ε4 alleles. We subsequently used

linear regression to test whether presence of APOEε4 alleles, tau, dementia diagnosis, or sex

were significantly associated with epigenetic age acceleration. Amyloid beta and p-tau were not

included in this model due to collinearity between amyloid beta, tau, and p-tau CSF

concentrations (see Supplementary Table S5).

RESULTS

Demographic information

Table 1 gives a summary of the variables for the participants with: CSF biomarkers (Table 1A; N=279), whole blood EPIC DNAme data (Table 1B; N=640) and matched EPIC data and measured CSF biomarkers (Table 1C; N=533). Given the differences in sample sizes, we performed demographic analyses on the three datasets. In the biomarker dataset, females were significantly younger and had fewer years of education than males ($P<0.01$ and $P<0.0001$, respectively). Although there were no sex differences in distribution of APOEε4 alleles, there was a strong trend for the proportion of participants in each of the diagnosis categories to be different for females and males ($P=0.051$; Table 1A). For the data applicable to the participants with available EPIC DNAme data (Table 1B) and participants with matched EPIC DNAme data and CSF biomarkers (Table 1C), most of the demographics were similar to the biomarker dataset with the exception that the proportion of participants in each of the diagnosis categories was not significantly different between females and males.

Sex and presence of APOEε4 alleles were associated with changes in inflammatory markers

Our first aim was to investigate whether sex and APOE genotype interact to influence inflammation using biomarkers, which we analysed separately in CSF and plasma (Supplementary Table S1 and S3, respectively). Caution should be noted as inflammatory signalling can differ depending on tissue examined [65,66].

For inflammatory markers measured in CSF, only IL-16 and IL-8 elicited a significant interaction between sex and APOE genotype ($P= 0.016$ and $P=0.035$, respectively; Table 3). CSF IL-16 and IL-8 levels were significantly lower in females non-carriers of APOEε4 alleles

compared to males (both P's<0.001), whereas levels were similar between the sexes in carriers of

APOEε4 alleles (P's>0.9; Fig 1 A and B). Furthermore, in females with APOEε4 alleles, IL-16

was significantly higher than in non-APOEε4 female carriers (P=0.050), while a trend was

observed in males (P=0.062). Whereas for IL-8, males with APOEε4 alleles had lower levels of

IL-8 compared to males with no APOEε4 alleles (P=0.014) but there was no difference in

females (P>0.3). Regardless of sex, CSF CRP levels were lower in individuals with 1 or 2

APOEε4 alleles compared to non-carriers (main effect of genotype: P=0.009; Table 3; Fig 1 C).

There was a trend for an increase in IL-6 receptor levels in APOEε4 carriers regardless of sex

compared to non APOEε4 carriers (main effect of genotype: P=0.071; Table 3). Lastly females

had significantly lower CSF levels of IgA and ICAM1 and a trend for lower CSF cortisol levels

compared to males (main effect of sex: P<0.001; P=0.009, and P=0.070, respectively; Table 3).

There were no other significant main or interaction effects on any other CSF biomarkers.

For biomarkers measured in plasma, there were no significant interactions between sex

and APOE genotype (Supplementary Table S3). However, females had higher plasma CRP

levels (main effect of sex: P=0.048; Fig 1 D) and ICAM1 (trend for a main effect of sex:

P=0.051) compared to males and significantly lower levels of IL-18 (main effect of sex:

P=0.001; Fig 1 E) and immunoglobulin E (IgE: main effect of sex: P<0.001; Fig 1 F) compared

to males. Furthermore, plasma CRP decreased in carriers of APOEε4 alleles compared to non-

carriers (main effect of genotype: P<0.001; Fig 1 D).

Sex and baseline diagnosis were associated with changes in inflammatory markers

We next tested whether sex and baseline dementia status (CN, MCI, and AD) influenced

CSF and plasma biomarkers of inflammation (Supplementary Table S2 and S4, respectively).

There were no significant interactions between sex and diagnosis for any of the tested variables

in CSF (Table 4 and Supplementary Table S2) or plasma (Supplementary Table S4). For CSF

levels, females had significantly lower levels of IgA (main effect of sex: P<0.001) and ICAM1

(main effect of sex: P=0.026) and a trend for lower IL-16 levels (main effect of sex: P=0.055)

compared to males (Table 4; Fig 2 A-C), but we did not observe any significant main effects of

diagnosis for any CSF variable.

In plasma, we found that females had lower levels of IgE (main effect of sex: P<0.001)

and IL-18 compared to males (main effect of sex: P=0.004) and trends for females to have higher

levels of ICAM1 (main effect of sex: P=0.056) and CRP (main effect of sex: P=0.056; Fig. 2 D)

compared to males. In addition, we found diagnosis significantly influenced plasma cortisol

(main effect of baseline diagnosis: P=0.01) with lower levels in LMCI compared to AD

(P<0.001; Fig 2 E). We found trends for diagnosis to influence plasma IL-16, CRP and CD 40

levels (main effect of diagnosis: P=0.054, P=0.056; P=0.067). Plasma IL-16 (P's=0.006) and

CRP (P=0.006 and P=0.02) levels were lower in LMCI and AD compared to CN (Fig 2 D and

F). For plasma CD 40, levels were lower in LMCI compared to AD (P=0.01; Supplementary

Table S4). In summary, although we detected associations between sex and diagnosis and

various biomarkers, we did not find evidence of a sex and diagnosis interaction on any variables

examined.

Correlations between cerebrospinal and plasma levels of biomarkers were mostly positive

The results for inflammatory markers in plasma did not always match results in CSF

(Supplementary Tables S2 and S4). We therefore investigated the relationship between plasma

and CSF biomarkers in males and females (Table 5, Fig 3). Perhaps surprisingly, we found the

majority of biomarkers were significantly positively correlated between plasma and CSF levels

in both males and females. These significant positive correlations included CRP (males, r=0.793;

females r=0.860; P's<0.0001), IL-6 receptor (males, r=0.459; females r=0.493, P's<0.0001), IgA

(males, r=0.705; females r=0.529; P's<0.0001), and cortisol in both sexes (males, r=0.176;

females, r=0.327; P=0.032 and 0.002, respectively). IL-16 was significantly correlated in females

(r=0.290, P=0.006) but only a trend in males (r=0.156, P=0.058). Plasma and CSF levels of

ICAM1 and CD 40 were positively correlated in males only (r=0.231, P=0.005 and r=0.374,

P<0.0001, respectively) whereas plasma and CSF IL-3 levels were negatively correlated in

females only (r=-0.246, P=0.021; Fig 3). There were significant sex differences, favoring males,

in the strength of correlation between the sexes for CD 40 (P=0.01) and IgA (P=0.03), with

trends for sex differences, favouring males in ICAM1 (P=0.06) and favouring females in IL-3

(P=0.06).

Epigenetic age, sex, dementia diagnosis, and AD biomarkers

We investigated the hypothesis that sex and dementia diagnosis affect epigenetic age

acceleration in blood samples of ADNI participants (Table 6). Epigenetic age acceleration was

not associated with sex, dementia diagnosis (CN, EMCI, LMCI, and AD), nor the interaction of

sex and diagnosis after multiple test correction (Fig 4). Intrinsic epigenetic age acceleration was

also not significantly associated with participant sex, diagnosis, or their interaction term.

To assess the effect of sex and more broadly defined dementia-associated cognitive

impairment on epigenetic age acceleration, we compared epigenetic age acceleration between

participants with any form of clinically ascertained cognitive impairment (AD + LMCI + EMCI,

N=423, proportion female 41%) and those without (CN, N=217, proportion female 50%). By

two-way unbalanced ANOVA neither sex, dementia status, nor their interaction were

significantly associated with epigenetic age acceleration after correction for multiple

comparisons.

Matched biochemical data including APOEε4 genotype and CSF concentrations of

amyloid beta, tau, and p-tau was available for a subset of participants with EPIC DNAme data

(N=533). Based on the hypothesis that epigenetic age acceleration may be more strongly

associated with concentrations of pathologically relevant compounds than with diagnosis, we

assessed the impact of sex, APOEε4 genotype, and CSF tau concentration on epigenetic age

acceleration and intrinsic epigenetic age acceleration with linear regression. None of these

variables were significantly associated with epigenetic age acceleration (Table 7; results for

intrinsic epigenetic age acceleration not shown).

In addition to dementia diagnosis for all participants, we also had access to two

composite scores designed by ADNI collaborators to reflect (i) executive function and (ii)

memory; these z-scores have been demonstrated to be independently predictive of the transition

from MCI to a formal diagnosis of AD [67,68]. By two-way unbalanced ANOVA models

investigating the effect of sex and memory z-score on epigenetic age acceleration, neither sex

(P=0.248), memory score (P=0.486), nor their interaction (P=0.227) were associated with

epigenetic age acceleration. In a similar model, neither sex (P=0.260), executive function

(P=0.105), or the interaction term of sex and executive function (P=0.153) were associated with

epigenetic age acceleration.

DISCUSSION

In the present study using ADNI data from CN, MCI and AD participants we found

interactions between sex and APOE genotype (but not between sex and diagnosis) on CSF and

plasma levels of IL-8 and IL-16 (see Table 2 for summary of the results). CSF levels of IL-8 and

IL-16 were lower in female APOEε4 non-carriers compared to males but similar levels were

found between the sexes with 1 or 2 APOEε4 alleles. Regardless of sex, possessing 1 or 2

APOEε4 alleles was associated with decreased levels of CSF and plasma CRP. Sex differences

were seen in inflammatory markers, regardless of diagnosis or genotype, as females had lower

CSF cytokines (IL-16, IL-18), CSF ICAM1, CSF and plasma immunoglobulins (IgA, IgE) and

plasma IL-18. However, tissue (CSF, plasma) mattered for results for certain inflammatory

markers (ICAM1 and to a lesser extent CRP) as females had higher plasma CRP and ICAM1

compared to males, opposite to what was found in CSF. Despite these differences in outcomes

between plasma and CSF biomarker analyses, plasma and CSF levels were positively correlated

for cortisol, CRP, IL-6 receptor, IgA in both sexes, whereas IL-16, and IL-8 were correlated in

females and CD 40 and ICAM1 were correlated in males, indicating good consistency between

CSF and plasma levels of these biomarkers. Intriguingly, IL-3 stood out from all these

biomarkers with a negative correlation between CSF and plasma levels in females only. Males

exhibited significantly stronger correlations between plasma and CSF levels for CD 40 and IgA

compared to females. Finally, we found no significant differences in epigenetic age acceleration

by dementia diagnosis or sex in this cohort of samples. Sex and genotype differences in CSF and

plasma inflammatory markers suggest differences in underlying physiology that may affect aging

and the progression of AD and this should be considered in future studies. Researchers should be

cautioned to use sex as a biological variable in all analyses.

Sex interacted with presence of APOEε4 alleles to affect levels of IL-8 and IL-16

In this study, we found that sex interacted with APOE genotype to influence CSF IL-16 and IL-8. CSF IL-16 and IL-8 levels were lower in females with no APOEε4 alleles compared to males, but no sex differences in these cytokine levels were detected in participants with 1 or 2 ε4 alleles. Our results suggest that presence of APOEε4 alleles can modulate CSF (and potentially plasma) cytokine levels in a sex-dependent way. The APOE protein can regulate transcription in vitro [69] and APOE4, but not APOE3, increases levels of IL-6 and IL-8 in vitro [70]. In the current study, we found that the sex differences in IL-16 and IL-8 levels disappeared with the presence of 1 or 2 APOEε4 alleles. One possibility is that the APOE4 protein regulates cytokine levels differently in males and females. IL-16 has been implicated in AD [71] and plasma IL-16 levels decreased with diagnosis (in males and females; current study) and AD severity (analysis without regard to sex; [72]). On the other hand, levels of IL-8 were not affected by diagnosis, consistent with a meta-analysis of cytokines in AD [22]. It is unclear what the impact of regulation of CSF cytokine levels by sex and APOEε4 has on AD symptoms or pathology, however given that females with APOEε4 alleles are disproportionally affected by AD during certain ages [16,18], IL-16 and IL-8 levels are unlikely to be a mechanism for this effect as differences in sex by genotype were noticed in the absence not presence of APOEε4 alleles.

Females had higher CRP levels compared to males and CRP levels were lower in APOEε4 carriers

We found that plasma and CSF levels of CRP, a widely used inflammatory and cardiovascular marker [73,74], were independently affected by sex and APOE genotype. Females, regardless of diagnosis or APOEε4 alleles, had significantly higher plasma CRP

relative to males, consistent with findings in healthy individuals [75]. Higher levels of peripheral

CRP may suggest higher systemic inflammation in females, which is associated with an

increased risk in all-cause dementia [76]. Higher levels of serum CRP are also associated with

higher levels of serum estradiol in postmenopausal healthy females [77] which suggests that sex

differences in CRP levels may be partly due to sex differences in estradiol levels. In addition, we

found that the presence of APOEε4 alleles decreased plasma and CSF CRP levels consistent with

previous research in large population studies [78,79]. In our study, we also found a trend for

lower levels of plasma CRP with LMCI and AD compared to CN. Recent meta-analyses did not

find differences in peripheral levels of CRP in AD compared to healthy controls [80,81].

However, in participants with mild and moderate dementia only, serum CRP levels were lower

compared to the cognitively healthy group [80]. In healthy individuals, higher levels of plasma

CRP in midlife are associated with a higher amyloid burden later in life in males but not females

[27]. However, despite this finding, higher systemic inflammation in midlife (including CRP) is

associated with greater cognitive decline later in life in both sexes in healthy individuals [28]. It

is important to acknowledge evidence that midlife obesity, but not later life obesity, is associated

with an increased risk to develop dementia [82,83], which may be related to altered inflammation

(e.g., cytokines and CRP) due to the accumulation of adipose tissue [84,85]. It is possible that

sex differences in inflammation and/or obesity earlier in life have long-term effects on the

transition to MCI and/or AD.

Females had lower cytokine and immunoglobulin levels compared to males

We found some biomarkers that were affected by sex, but not diagnosis or presence of

APOEε4 alleles. For example, females had lower CSF levels of ICAM1 compared to males,

regardless of APOE genotype or diagnosis, but, although a trend, the opposite effect was seen in

plasma. In contrast, in healthy adults (18-55 years old), serum levels of ICAM1 are lower in

females compared to males [86]. ICAM1 is a type of adhesion molecule associated with

microvascular endothelial activation [87] and plasma ICAM1 levels (but not CSF levels; [88])

were higher in patients with AD [88–90]. Although in the present study we did not observe a

significant effect of plasma ICAM1 with diagnosis, the unadjusted P-value was 0.063 with

higher levels in LMCI and AD groups. It is intriguing that females have lower CSF levels of

cytokines (IL-16, IL-18), ICAM1, and immunoglobulins (IgE and IgA) but higher plasma CRP

and ICAM1 levels. Although neuroinflammation is associated with AD, it may be both a product

and a driver of neurodegeneration, and it may have both beneficial and detrimental roles in AD

[91,92]. In AD mouse models, inflammatory cytokines (e.g., IL1β, IL-4, IL-6, IL-10, IFNγ,

TNFα) can both increase amyloid beta deposition and reduce amyloid plaque pathology [93–

101]. In transgenic mice, amyloid deposition is associated with low T-cell activation suggesting

that the immune system is hypo-responsive to amyloid beta [102]. Thus, increases of

inflammatory markers may not always be indicative of worse neuropathology or outcomes, but

may be contributing to reductions in AD neuropathology. It is also possible that males and

females have varying levels of beneficial vs detrimental immune responses which can

differentially affect how the disease progresses between the sexes. Indeed, we found sex

differences in the correlation between CSF and plasma biomarkers (CD 40, IgA, ICAM1 and IL-

3), which suggests that plasma and CSF levels may be regulated differently in males and

females.

Sex, AD and biochemical markers did not affect blood epigenetic age acceleration

We did not observe an association between either sex or diagnosis and epigenetic age

acceleration or intrinsic epigenetic age acceleration. To our knowledge, no other study has

similarly probed epigenetic age acceleration in a peripheral tissue in the presence of AD, or

whether epigenetic age acceleration in AD is sex-specific.

This study was partially undertaken to investigate whether epigenetic age acceleration

that has been associated with the AD brain is reflected in peripheral tissues. Levine et al. [40]

have previously demonstrated increased epigenetic age acceleration in AD, however Levine's

study was conducted on post-mortem prefrontal cortex tissue, and did not explicitly investigate

the role of sex in epigenetic age acceleration [40]. While brain-blood global DNA methylation

profiles are reasonably correlated (r=0.86) [48], DNA methylation profiles of peripheral tissues

are imperfect representatives of the brain, and do not recapitulate all epigenetic alterations with

high fidelity. Thus, our findings do not contradict the finding of increased epigenetic age

acceleration in the presence of AD in the prefrontal cortex, but suggest that accelerated

epigenetic aging in AD is not a pan-tissue phenomenon. Our finding of a lack of significant

association between AD, biological sex, and epigenetic age acceleration in whole blood DNA

methylation profiles could suggest a tissue-specific dysregulation of an epigenetic maintenance

system, in which the brain epigenome is most strongly affected by AD [40]. The phenotype of

patients affected by AD and global gene expression patterns of the APOE protein, with high

expression in brain, and low expression in whole blood [103] further support this hypothesis.

Additionally, previous work by Hogervorst et al. [104] indicates that men with AD, specifically

those carrying the e4 allele, demonstrate accelerated endocrinological aging (hypogonadism). It

is possible that the accelerated epigenetic aging observed in the AD brain is sex-specific, and

although not reflected in blood, may be reflected in other organs and tissues, such as the brain, or

potentially target organs of endocrine glands.

Intriguingly, epigenetic age in the ADNI samples was observed to be lower on average

than chronological age. Horvath's epigenetic clock was trained on DNAme data from older

versions of the Illumina DNAme arrays with more limited genomic coverage; 19 of the 353 CpG

sites required to calculate epigenetic age via Horvath's algorithm do not exist on the newer EPIC

array. Two previous studies investigated the application of Horvath's epigenetic clock to EPIC

data with conflicting results [62,105], the largest issue being chronic underestimation of

epigenetic age due to the positive linear regression coefficients associated with the missing

probes [105]. Both imputing and removing the missing probes from the array resulted in a

chronic underprediction of epigenetic age with Horvath's clock suggesting that this is likely an

artefact of the array platform and truncated probe-set rather than the method chosen to deal with

missing values, although it is possible that an adjustment factor could be devised to more

accurately apply Horvath's clock to EPIC data. In future explorations of epigenetic age with

EPIC DNAme array data this should be considered, as there are other epigenetic age predictors

available that have been trained on EPIC data such as the PhenoAge and GrimAge clocks,

although these tools have limitations as well; for example, both PhenoAge and GrimAge were

trained only on blood DNAme data, as compared to the original pan-tissue epigenetic clocks, and

therefore may have limited applicability and relevance in other tissues [106,107].

Limitations

The ADNI cohort is not ethnically or socioeconomically diverse, being mostly composed

of self-reported white (only 12 individuals were not-white) and highly educated individuals

(average 15.69 years of education). As AD incidence, prevalence, and age of onset varies by

ethnicity [108–110] and education [111], our conclusions may not apply to more ethnically and

socially diverse populations. In addition to sex, it is possible the underlying mechanisms of AD

are different depending on ethnicity. APOEε4 genotype is a well-known AD risk factor and in

this cohort the proportion of participants in the different APOEε4 allele groups is correlated with

diagnosis (Table S6). Future research with a larger cohort is required to test how sex, APOE

genotype and diagnosis interact together in one model. Even in the larger 640-participant cohort,

only 37 participants (5.78%) had an AD diagnosis, so statistical analyses were underpowered to

detect subtle differences by diagnosis group, which may account for our lack of interaction

between sex and diagnosis. Additionally, other pathologies in these participants, such as cancer,

cardiovascular disease, smoking status, or obesity may have influenced inflammatory markers

and epigenomes and limited our interpretations.

CONCLUSION

The current study provides evidence that sex and presence of APOEε4 alleles are

associated with CSF levels of the inflammatory markers IL-16 and IL-8. We found sex

differences indicating that females had lower cytokine and immunoglobulin levels but higher

plasma CRP and ICAM1 levels compared to males, although the direction of the ICAM1 finding

was tissue-dependent. There were no significant associations between sex, APOEε4 alleles or

diagnosis and epigenetic age. Together, our work suggests that that presence of APOEε4 alleles

can affect cytokine levels differently in males and females and the underlying pathophysiology

of aging and AD may be tissue- and sex-specific.

ACKNOWLEDGMENTS

Data collection and sharing for this project was funded by the Alzheimer's Disease

Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD

ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the

National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering,

and through generous contributions from the following: AbbVie, Alzheimer's Association;

Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-

Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli

Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company

Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy

Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development

LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx

Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal

Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian

Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private

sector contributions are facilitated by the Foundation for the National Institutes of Health

(www.fnih.org). The grantee organization is the Northern California Institute for Research and

Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the

University of Southern California. ADNI data are disseminated by the Laboratory for Neuro

Imaging at the University of Southern California. Funding for this study was provided by a

Canadian Institutes of Health Research (CHIR) grant to LAMG (PJT-148662) and a CIHR sex

catalyst grant to WPR (Reference No. 158613). WPR receives salary support through an

investigatorship award from the BC Children's Hospital Research Institute. PDG is funded by

the Alzheimer's Association of the USA and Brain Canada with the financial support of Health

Canada through the Brain Canada Research Fund (AARF-17-529705). The views expressed

herein do not necessarily represent the views of the Minister of Health or the Government of

Canada. AMI is funded by a University of British Columbia Four Year Doctoral Fellowship and

the CIHR Frederick Banting and Charles Best Masters Research Award. We thank Elizabeth

Perez for help with data organization.

CONFLICT OF INTEREST

The authors have no conflict of interest to report.

REFERENCES

[1]   Alzheimer's Association (2017) 2017 Alzheimer's disease facts and figures. Alzheimer's
& Dementia **13**, 325–373.
[2]   Caruso A, Nicoletti F, Mango D, Saidi A, Orlando R, Scaccianoce S (2018) Stress as risk
factor for Alzheimer's disease. Pharmacological Research **132**, 130–134.
[3]   Baumgart M, Snyder HM, Carrillo MC, Fazio S, Kim H, Johns H (2015) Summary of the
evidence on modifiable risk factors for cognitive decline and dementia: A population-
based perspective. Alzheimers Dement **11**, 718–726.
[4]   Nebel RA, Aggarwal NT, Barnes LL, Gallagher A, Goldstein JM, Kantarci K,
Mallampalli MP, Mormino EC, Scott L, Yu WH, Maki PM, Mielke MM (2018)
Understanding the impact of sex and gender in Alzheimer's disease: A call to action.
Alzheimer's & Dementia **14**, 1171–1183.
[5]   Xu W, Tan L, Wang H-F, Jiang T, Tan M-S, Tan L, Zhao Q-F, Li J-Q, Wang J, Yu J-T
(2015) Meta-analysis of modifiable risk factors for Alzheimer's disease. J Neurol
Neurosurg Psychiatry **86**, 1299–1306.
[6]   Riedel BC, Thompson PM, Brinton RD (2016) Age, APOE and sex: Triad of risk of
Alzheimer's disease. The Journal of Steroid Biochemistry and Molecular Biology **160**,
134–147.
[7]   Laws KR, Irvine K, Gale TM (2018) Sex differences in Alzheimer's disease. Current
Opinion in Psychiatry **31**, 133–139.
[8]   Mielke MM (2018) Sex and Gender Differences in Alzheimer's Disease Dementia.
Psychiatr Times **35**, 14–17.
[9]   Irvine K, Laws KR, Gale TM, Kondel TK (2012) Greater cognitive deterioration in
women than men with Alzheimer's disease: a meta analysis. J Clin Exp Neuropsychol
34, 989–998.

[10] Lin KA, Choudhury KR, Rathakrishnan BG, Marks DM, Petrella JR, Doraiswamy PM
(2015) Marked gender differences in progression of mild cognitive impairment over 8
years. Alzheimer's & Dementia: Translational Research & Clinical Interventions **1**, 103–
110.
[11] Wang X, Zhou W, Ye T, Lin X, Zhang J, Initiative for ADN (2019) Sex Difference in
the Association of APOE4 with Memory Decline in Mild Cognitive Impairment. Journal
of Alzheimer's Disease **69**, 1161–1169.
[12] Ardekani BA, Convit A, Bachman AH (2016) Analysis of the MIRIAD Data Shows Sex
Differences in Hippocampal Atrophy Progression. Journal of Alzheimer's Disease **50**,
847–857.
[13] Barnes LL, Wilson RS, Bienias JL, Schneider JA, Evans DA, Bennett DA (2005) Sex
differences in the clinical manifestations of Alzheimer disease pathology. Arch. Gen.
Psychiatry **62**, 685–691.
[14] Holland D, Desikan RS, Dale AM, McEvoy LK, Alzheimer's Disease Neuroimaging
Initiative (2013) Higher rates of decline for women and apolipoprotein E epsilon4
carriers. AJNR Am J Neuroradiol **34**, 2287–2293.
[15] Hua X, Hibar DP, Lee S, Toga AW, Jack CR, Weiner MW, Thompson PM (2010) Sex
and age differences in atrophic rates: an ADNI study with n=1368 MRI scans.
Neurobiology of Aging **31**, 1463–1480.
[16] Neu SC, Pa J, Kukull W, Beekly D, Kuzma A, Gangadharan P, Wang L-S, Romero K,
Arneric SP, Redolfi A, Orlandi D, Frisoni GB, Au R, Devine S, Auerbach S, Espinosa A,
Boada M, Ruiz A, Johnson SC, Koscik R, Wang J-J, Hsu W-C, Chen Y-L, Toga AW
(2017) Apolipoprotein E Genotype and Sex Risk Factors for Alzheimer Disease: A Meta-
analysis. JAMA Neurology **74**, 1178.
[17] Beydoun MA, Boueiz A, Abougergi MS, Kitner-Triolo MH, Beydoun HA, Resnick SM,
O'Brien R, Zonderman AB (2012) Sex differences in the association of the
apolipoprotein E epsilon 4 allele with incidence of dementia, cognitive impairment, and
decline. Neurobiol. Aging **33**, 720-731.e4.
[18] Hohman TJ, Dumitrescu L, Barnes LL, Thambisetty M, Beecham G, Kunkle B, Gifford
KA, Bush WS, Chibnik LB, Mukherjee S, Jager PLD, Kukull W, Crane PK, Resnick
SM, Keene CD, Montine TJ, Schellenberg GD, Haines JL, Zetterberg H, Blennow K,
Larson EB, Johnson SC, Albert M, Bennett DA, Schneider JA, Jefferson AL (2018) Sex-
Specific Association of Apolipoprotein E With Cerebrospinal Fluid Levels of Tau.
JAMA Neurol **75**, 989–998.
[19] Koran MEI, Wagener M, Hohman TJ, Alzheimer's Neuroimaging Initiative (2017) Sex
differences in the association between AD biomarkers and cognitive decline. Brain
Imaging Behav **11**, 205–213.
[20] Heppner FL, Ransohoff RM, Becher B (2015) Immune attack: the role of inflammation
in Alzheimer disease. Nature Reviews Neuroscience **16**, 358–372.
[21] Kinney JW, Bemiller SM, Murtishaw AS, Leisgang AM, Salazar AM, Lamb BT (2018)
Inflammation as a central mechanism in Alzheimer's disease. Alzheimer's & Dementia:
Translational Research & Clinical Interventions **4**, 575–590.
[22] Swardfager W, Lanctôt K, Rothenburg L, Wong A, Cappell J, Herrmann N (2010) A
Meta-Analysis of Cytokines in Alzheimer's Disease. Biological Psychiatry **68**, 930–941.

[23] Engler H, Benson S, Wegner A, Spreitzer I, Schedlowski M, Elsenbruch S (2016) Men
and women differ in inflammatory and neuroendocrine responses to endotoxin but not in
the severity of sickness symptoms. Brain, Behavior, and Immunity **52**, 18–26.
[24] Klein SL, Flanagan KL (2016) Sex differences in immune responses. Nature Reviews
Immunology **16**, 626–638.
[25] Wegner A, Benson S, Rebernik L, Spreitzer I, Jäger M, Schedlowski M, Elsenbruch S,
Engler H (2017) Sex differences in the pro-inflammatory cytokine response to endotoxin
unfold in vivo but not ex vivo in healthy humans. Innate Immun **23**, 432–439.
[26] Márquez EJ, Chung C, Marches R, Rossi RJ, Nehar-Belaid D, Eroglu A, Mellert DJ,
Kuchel GA, Banchereau J, Ucar D (2020) Sexual-dimorphism in human immune system
aging. Nature Communications **11**, 751.
[27] Walker KA, Windham BG, Brown IV CH, Knopman DS, Jack Jr CR, Mosley Jr TH,
Selvin E, Wong DF, Hughes TM, Zhou Y, Gross AL, Gottesman RF (2018) The
Association of Mid- and Late-Life Systemic Inflammation with Brain Amyloid
Deposition: The ARIC-PET Study. Journal of Alzheimer's Disease **66**, 1041–1052.
[28] Walker KA, Gottesman RF, Wu A, Knopman DS, Gross AL, Mosley TH, Selvin E,
Windham BG (2019) Systemic inflammation during midlife and cognitive change over
20 years. Neurology **92**, e1256–e1267.
[29] Dantzer R (2018) Neuroimmune Interactions: From the Brain to the Immune System and
Vice Versa. Physiol. Rev. **98**, 477–504.
[30] de Miranda AS, Zhang C-J, Katsumoto A, Teixeira AL (2017) Hippocampal adult
neurogenesis: Does the immune system matter? Journal of the Neurological Sciences
372, 482–495.
[31] Apostolova LG, Dutton RA, Dinov ID, Hayashi KM, Toga AW, Cummings JL,
Thompson PM (2006) Conversion of mild cognitive impairment to Alzheimer disease
predicted by hippocampal atrophy maps. Arch. Neurol. **63**, 693–699.
[32] Petersen RC, Jack CR, Xu YC, Waring SC, O'Brien PC, Smith GE, Ivnik RJ, Tangalos
EG, Boeve BF, Kokmen E (2000) Memory and MRI-based hippocampal volumes in
aging and AD. Neurology **54**, 581–587.
[33] Doecke JD, Laws SM, Faux NG, Wilson W, Burnham SC, Lam C-P, Mondal A, Bedo J,
Bush AI, Brown B, Ruyck KD, Ellis KA, Fowler C, Gupta VB, Head R, Macaulay SL,
Pertile K, Rowe CC, Rembach A, Rodrigues M, Rumble R, Szoeke C, Taddei K, Taddei
T, Trounson B, Ames D, Masters CL, Martins RN, Group for the ADNI and AIB and
LR (2012) Blood-Based Protein Biomarkers for Diagnosis of Alzheimer Disease. Arch
Neurol **69**, 1318–1325.
[34] Toledo JB, Toledo E, Weiner MW, Jack CR, Jagust W, Lee VM-Y, Shaw LM,
Trojanowski JQ (2012) Cardiovascular risk factors, cortisol, and amyloid-β deposition in
Alzheimer's Disease Neuroimaging Initiative. Alzheimers Dement **8**, 483–489.
[35] Lupien SJ, de Leon M, de Santi S, Convit A, Tarshish C, Nair NP, Thakur M, McEwen
BS, Hauger RL, Meaney MJ (1998) Cortisol levels during human aging predict
hippocampal atrophy and memory deficits. Nat. Neurosci. **1**, 69–73.
[36] Murphy KJ, Sheppard PAS, Troyer AK, Hampson E, Galea LAM (2019) Sex differences
in cortisol and memory following acute social stress in amnestic mild cognitive
impairment. bioRxiv 757484.
[37] McEwen BS, Biron CA, Brunson KW, Bulloch K, Chambers WH, Dhabhar FS, Goldfarb
RH, Kitson RP, Miller AH, Spencer RL, Weiss JM (1997) The role of adrenocorticoids

as modulators of immune function in health and disease: neural, endocrine and immune interactions. Brain Research Reviews **23**, 79–133.

[38] Hannum G, Guinney J, Zhao L, Zhang L, Hughes G, Sadda S, Klotzle B, Bibikova M, Fan J-B, Gao Y, Deconde R, Chen M, Rajapakse I, Friend S, Ideker T, Zhang K (2013) Genome-wide Methylation Profiles Reveal Quantitative Views of Human Aging Rates. Molecular Cell **49**, 359–367.

[39] Horvath S (2013) DNA methylation age of human tissues and cell types. Genome Biol **14**, R115.

[40] Levine ME, Lu AT, Bennett DA, Horvath S (2015) Epigenetic age of the pre-frontal cortex is associated with neuritic plaques, amyloid load, and Alzheimer's disease related cognitive functioning. Aging (Albany NY) **7**, 1198–1211.

[41] Yokoyama AS, Rutledge JC, Medici V (2017) DNA methylation alterations in Alzheimer's disease. Environ Epigenet **3**,.

[42] Gale CR, Marioni RE, Harris SE, Starr JM, Deary IJ (2018) DNA methylation and the epigenetic clock in relation to physical frailty in older people: the Lothian Birth Cohort 1936. Clin Epigenetics **10**,.

[43] Horvath S, Garagnani P, Bacalini MG, Pirazzini C, Salvioli S, Gentilini D, Di Blasio AM, Giuliani C, Tung S, Vinters HV, Franceschi C (2015) Accelerated epigenetic aging in Down syndrome. Aging Cell **14**, 491–495.

[44] Horvath S, Ritz BR (2015) Increased epigenetic age and granulocyte counts in the blood of Parkinson's disease patients. Aging (Albany NY) **7**, 1130–1142.

[45] Degerman S, Josefsson M, Nordin Adolfsson A, Wennstedt S, Landfors M, Haider Z, Pudas S, Hultdin M, Nyberg L, Adolfsson R (2017) Maintained memory in aging is associated with young epigenetic age. Neurobiology of Aging **55**, 167–171.

[46] McEwen LM, Morin AM, Edgar RD, MacIsaac JL, Jones MJ, Dow WH, Rosero-Bixby L, Kobor MS, Rehkopf DH (2017) Differential DNA methylation and lymphocyte proportions in a Costa Rican high longevity region. Epigenetics Chromatin **10**,.

[47] Horvath S, Gurven M, Levine ME, Trumble BC, Kaplan H, Allayee H, Ritz BR, Chen B, Lu AT, Rickabaugh TM, Jamieson BD, Sun D, Li S, Chen W, Quintana-Murci L, Fagny M, Kobor MS, Tsao PS, Reiner AP, Edlefsen KL, Absher D, Assimes TL (2016) An epigenetic clock analysis of race/ethnicity, sex, and coronary heart disease. Genome Biol **17**,.

[48] Braun PR, Han S, Hing B, Nagahama Y, Gaul LN, Heinzman JT, Grossbach AJ, Close L, Dlouhy BJ, Howard MA, Kawasaki H, Potash JB, Shinozaki G (2019) Genome-wide DNA methylation comparison between live human brain and peripheral tissues within individuals. Translational Psychiatry **9**, 47.

[49] Aisen P, Petersen R, Donohue M, Gamst A, Raman R, Thomas R, Walter S, Trojanowski J, Shaw L, Beckett L, Jack C, Jagust W, Toga A, Saykin A, Morris J, Weiner M (2010) Clinical core of the Alzheimer's Disease Neuroimaging Initiative: progress and plans. Alzheimers Dement **6**, 239–246.

[50] Petersen RC, Aisen PS, Beckett LA, Donohue MC, Gamst AC, Harvey DJ, Jack CR, Jagust WJ, Shaw LM, Toga AW, Trojanowski JQ, Weiner MW (2010) Alzheimer's Disease Neuroimaging Initiative (ADNI). Neurology **74**, 201–209.

[51] Craig-Schapiro R, Kuhn M, Xiong C, Pickering EH, Liu J, Misko TP, Perrin RJ, Bales KR, Soares H, Fagan AM, Holtzman DM (2011) Multiplexed Immunoassay Panel

Identifies Novel CSF Biomarkers for Alzheimer's Disease Diagnosis and Prognosis.
PLOS ONE **6**, e18850.

[52]   Hu WT, Holtzman DM, Fagan AM, Shaw LM, Perrin R, Arnold SE, Grossman M,
Xiong C, Craig-Schapiro R, Clark CM, Pickering E, Kuhn M, Chen Y, Van Deerlin VM,
McCluskey L, Elman L, Karlawish J, Chen-Plotkin A, Hurtig HI, Siderowf A, Swenson
F, Lee VM-Y, Morris JC, Trojanowski JQ, Soares H, Alzheimer's Disease Neuroimaging
Initiative (2012) Plasma multianalyte profiling in mild cognitive impairment and
Alzheimer disease. Neurology **79**, 897–905.

[53]   Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society. Series B
(Methodological) **57**, 289–300.

[54]   Zou GY (2007) Toward using confidence intervals to compare correlations.
Psychological Methods **12**, 399–413.

[55]   R Core Team (2018) R: A language and environment for statistical   computing. R
Foundation for Statistical Computing, Vienna, Austria., R Foundation for Statistical
Computing, Vienna, Austria.

[56]   Vasanthakumar A, Davis JW, Idler K, Kim S, Nho K, Nudelman KNH, Faber K, Bai Y,
Foroud TM, Estrada K, Li Q, Saykin AJ, Waring J (2017) DNA METHYLATION
DYNAMICS IN ALZHEIMER'S DISEASE DIAGNOSIS AND PROGRESSION.
Alzheimer's & Dementia **13**, P297.

[57]   Aryee MJ, Jaffe AE, Corrada-Bravo H, Ladd-Acosta C, Feinberg AP, Hansen KD,
Irizarry RA (2014) Minfi: a flexible and comprehensive Bioconductor package for the
analysis of Infinium DNA methylation microarrays. Bioinformatics **30**, 1363–1369.

[58]   Fortin J-P, Triche TJ, Hansen KD (2017) Preprocessing, normalization and integration of
the Illumina HumanMethylationEPIC array with minfi. Bioinformatics **33**, 558–560.

[59]   Triche TJ, Weisenberger DJ, Van Den Berg D, Laird PW, Siegmund KD (2013) Low-
level processing of Illumina Infinium DNA Methylation BeadArrays. Nucleic Acids Res
41, e90.

[60]   Teschendorff AE, Marabita F, Lechner M, Bartlett T, Tegner J, Gomez-Cabrero D, Beck
S (2013) A beta-mixture quantile normalization method for correcting probe design bias
in Illumina Infinium 450 k DNA methylation data. Bioinformatics **29**, 189–196.

[61]   Fiorito G, Polidoro S, Dugué P-A, Kivimaki M, Ponzi E, Matullo G, Guarrera S,
Assumma MB, Georgiadis P, Kyrtopoulos SA, Krogh V, Palli D, Panico S, Sacerdote C,
Tumino R, Chadeau-Hyam M, Stringhini S, Severi G, Hodge AM, Giles GG, Marioni R,
Linnér RK, O'Halloran AM, Kenny RA, Layte R, Baglietto L, Robinson O, McCrory C,
Milne RL, Vineis P (2017) Social adversity and epigenetic aging: a multi-cohort study on
socioeconomic differences in peripheral blood DNA methylation. Scientific Reports **7**,
16266.

[62]   McEwen LM, Jones MJ, Lin DTS, Edgar RD, Husquin LT, MacIsaac JL, Ramadori KE,
Morin AM, Rider CF, Carlsten C, Quintana-Murci L, Horvath S, Kobor MS (2018)
Systematic evaluation of DNA methylation age estimation with common preprocessing
methods and the Infinium MethylationEPIC BeadChip array. Clinical Epigenetics **10**,
123.

[63]   Chen BH, Marioni RE, Colicino E, Peters MJ, Ward-Caviness CK, Tsai P-C, Roetker
NS, Just AC, Demerath EW, Guan W, Bressler J, Fornage M, Studenski S, Vandiver AR,
Moore AZ, Tanaka T, Kiel DP, Liang L, Vokonas P, Schwartz J, Lunetta KL, Murabito

JM, Bandinelli S, Hernandez DG, Melzer D, Nalls M, Pilling LC, Price TR, Singleton AB, Gieger C, Holle R, Kretschmer A, Kronenberg F, Kunze S, Linseisen J, Meisinger C, Rathmann W, Waldenberger M, Visscher PM, Shah S, Wray NR, McRae AF, Franco OH, Hofman A, Uitterlinden AG, Absher D, Assimes T, Levine ME, Lu AT, Tsao PS, Hou L, Manson JE, Carty CL, LaCroix AZ, Reiner AP, Spector TD, Feinberg AP, Levy D, Baccarelli A, van Meurs J, Bell JT, Peters A, Deary IJ, Pankow JS, Ferrucci L, Horvath S (2016) DNA methylation-based measures of biological age: meta-analysis predicting time to death. Aging (Albany NY) **8**, 1844–1859.

[64] Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson HH, Wiencke JK, Kelsey KT (2012) DNA methylation arrays as surrogate measures of cell mixture distribution. BMC Bioinformatics **13**, 86.

[65] Bettcher BM, Johnson SC, Fitch R, Casaletto KB, Heffernan KS, Asthana S, Zetterberg H, Blennow K, Carlsson CM, Neuhaus J, Bendlin BB, Kramer JH (2018) CSF and Plasma Levels of Inflammation Differentially Relate to CNS Markers of Alzheimer's Disease Pathology and Neuronal Damage. J Alzheimers Dis **62**, 385–397.

[66] Setiawan E, Wilson AA, Mizrahi R, Rusjan PM, Miler L, Rajkowska G, Suridjan I, Kennedy JL, Rekkas PV, Houle S, Meyer JH (2015) Role of translocator protein density, a marker of neuroinflammation, in the brain during major depressive episodes. JAMA Psychiatry **72**, 268–275.

[67] Crane PK, Carle A, Gibbons LE, Insel P, Mackin RS, Gross A, Jones RN, Mukherjee S, Curtis SM, Harvey D, Weiner M, Mungas D, Alzheimer's Disease Neuroimaging Initiative (2012) Development and assessment of a composite score for memory in the Alzheimer's Disease Neuroimaging Initiative (ADNI). Brain Imaging Behav **6**, 502–516.

[68] Gibbons LE, Carle AC, Mackin RS, Harvey D, Mukherjee S, Insel P, Curtis SM, Mungas D, Crane PK, Alzheimer's Disease Neuroimaging Initiative (2012) A composite score for executive functioning, validated in Alzheimer's Disease Neuroimaging Initiative (ADNI) participants with baseline mild cognitive impairment. Brain Imaging Behav **6**, 517–527.

[69] Theendakara V, Peters-Libeu CA, Bredesen DE, Rao RV (2018) Transcriptional Effects of ApoE4: Relevance to Alzheimer's Disease. Molecular Neurobiology **55**, 5243–5254.

[70] Theendakara V, Peters-Libeu CA, Spilman P, Poksay KS, Bredesen DE, Rao RV (2016) Direct Transcriptional Effects of Apolipoprotein E. The Journal of Neuroscience **36**, 685–700.

[71] Rosa MD, Dell'Ombra N, Zambito AM, Malaguarnera M, Nicoletti F, Malaguarnera L (2006) Chitotriosidase and inflammatory mediator levels in Alzheimer's disease and cerebrovascular dementia. European Journal of Neuroscience **23**, 2648–2656.

[72] Motta M, Imbesi R, Di Rosa M, Stivala F, Malaguarnera L (2007) Altered plasma cytokine levels in Alzheimer's disease: Correlation with the disease progression. Immunology Letters **114**, 46–51.

[73] Koenig W, Sund M, Fröhlich M, Fischer HG, Löwel H, Döring A, Hutchinson WL, Pepys MB (1999) C-Reactive protein, a sensitive marker of inflammation, predicts future risk of coronary heart disease in initially healthy middle-aged men: results from the MONICA (Monitoring Trends and Determinants in Cardiovascular Disease) Augsburg Cohort Study, 1984 to 1992. Circulation **99**, 237–242.

[74] Ridker PM, Buring JE, Shih J, Matias M, Hennekens CH (1998) Prospective study of C-reactive protein and the risk of future cardiovascular events among apparently healthy women. Circulation **98**, 731–733.

[75] Khera A, McGuire DK, Murphy SA, Stanek HG, Das SR, Vongpatanasin W, Wians FH, Grundy SM, de Lemos JA (2005) Race and gender differences in C-reactive protein levels. J. Am. Coll. Cardiol. **46**, 464–469.

[76] Koyama A, O'Brien J, Weuve J, Blacker D, Metti AL, Yaffe K (2013) The role of peripheral inflammatory markers in dementia and Alzheimer's disease: a meta-analysis. J. Gerontol. A Biol. Sci. Med. Sci. **68**, 433–440.

[77] Eldridge RC, Wentzensen N, Pfeiffer RM, Brinton LA, Hartge P, Guillemette C, Kemp TJ, Pinto LA, Trabert B (2020) Endogenous estradiol and inflammation biomarkers: potential interacting mechanisms of obesity-related disease. Cancer Causes Control.

[78] Hubacek JA, Peasey A, Pikhart H, Stavek P, Kubinova R, Marmot M, Bobak M (2010) APOE polymorphism and its effect on plasma C-reactive protein levels in a large general population sample. Hum Immunol **71**, 304–308.

[79] Yun Y-W, Kweon S-S, Choi J-S, Rhee J-A, Lee Y-H, Nam H-S, Jeong S-K, Park K-S, Ryu S-Y, Choi S-W, Kim HN, Cauley JA, Shin M-H (2015) APOE Polymorphism Is Associated with C-reactive Protein Levels but Not with White Blood Cell Count: Dong-gu Study and Namwon Study. J Korean Med Sci **30**, 860–865.

[80] Gong C, Wei D, Wang Y, Ma J, Yuan C, Zhang W, Yu G, Zhao Y (2016) A Meta-Analysis of C-Reactive Protein in Patients With Alzheimer's Disease. Am J Alzheimers Dis Other Demen **31**, 194–200.

[81] Ng A, Tam WW, Zhang MW, Ho CS, Husain SF, McIntyre RS, Ho RC (2018) IL-1β, IL-6, TNF- α and CRP in Elderly Patients with Depression or Alzheimer's disease: Systematic Review and Meta-Analysis. Scientific Reports **8**, 12050.

[82] Albanese E, Launer LJ, Egger M, Prince MJ, Giannakopoulos P, Wolters FJ, Egan K (2017) Body mass index in midlife and dementia: Systematic review and meta-regression analysis of 589,649 men and women followed in longitudinal studies. Alzheimers Dement (Amst) **8**, 165–178.

[83] Fitzpatrick AL, Kuller LH, Lopez OL, Diehr P, O'Meara ES, Longstreth WT, Luchsinger JA (2009) Midlife and Late-Life Obesity and the Risk of Dementia: Cardiovascular Health Study. Arch Neurol **66**, 336–342.

[84] Nguyen JCD, Killcross AS, Jenkins TA (2014) Obesity and cognitive decline: role of inflammation and vascular changes. Front. Neurosci. **8**,.

[85] Parimisetty A, Dorsemans A-C, Awada R, Ravanan P, Diotel N, Lefebvre d'Hellencourt C (2016) Secret talk between adipose tissue and central nervous system via secreted factors—an emerging frontier in the neurodegenerative research. Journal of Neuroinflammation **13**, 67.

[86] Ponthieux A, Herbeth B, Droesch S, Lambert D, Visvikis S (2003) Age- and Sex-related Reference Values for Serum Adhesion Molecule Concentrations in Healthy Individuals: Intercellular Adhesion Molecule-1 and E-, P-, and L-Selectin. Clinical Chemistry **49**, 1544–1546.

[87] Zenaro E, Piacentino G, Constantin G (2017) The blood-brain barrier in Alzheimer's disease. Neurobiology of Disease **107**, 41–56.

[88] Nielsen HM, Londos E, Minthon L, Janciauskiene SM (2007) Soluble adhesion molecules and angiotensin-converting enzyme in dementia. Neurobiol. Dis. **26**, 27–35.

[89] Huang C-W, Tsai M-H, Chen N-C, Chen W-H, Lu Y-T, Lui C-C, Chang Y-T, Chang W-N, Chang AYW, Chang C-C (2015) Clinical significance of circulating vascular cell

adhesion molecule-1 to white matter disintegrity in Alzheimer's dementia. Thrombosis
and Haemostasis **114**, 1230–1240.

[90] Rentzos M, Michalopoulou M, Nikolaou C, Cambouri C, Rombos A, Dimitrakopoulos
A, Kapaki E, Vassilopoulos D (2004) Serum Levels of Soluble Intercellular Adhesion
Molecule-1 and Soluble Endothelial Leukocyte Adhesion Molecule-1 in Alzheimer's
Disease. J Geriatr Psychiatry Neurol **17**, 225–231.

[91] Heneka MT, Carson MJ, Khoury JE, Landreth GE, Brosseron F, Feinstein DL, Jacobs
AH, Wyss-Coray T, Vitorica J, Ransohoff RM, Herrup K, Frautschy SA, Finsen B,
Brown GC, Verkhratsky A, Yamanaka K, Koistinaho J, Latz E, Halle A, Petzold GC,
Town T, Morgan D, Shinohara ML, Perry VH, Holmes C, Bazan NG, Brooks DJ, Hunot
S, Joseph B, Deigendesch N, Garaschuk O, Boddeke E, Dinarello CA, Breitner JC, Cole
GM, Golenbock DT, Kummer MP (2015) Neuroinflammation in Alzheimer's disease.
The Lancet Neurology **14**, 388–405.

[92] Walters A, Phillips E, Zheng R, Biju M, Kuruvilla T (2016) Evidence for
neuroinflammation in Alzheimer's disease: Neuroinflammation in Alzheimer's. Progress
in Neurology and Psychiatry **20**, 25–31.

[93] Chakrabarty P, Li A, Ceballos-Diaz C, Eddy JA, Funk CC, Moore B, DiNunno N,
Rosario AM, Cruz PE, Verbeeck C, Sacino A, Nix S, Janus C, Price ND, Das P, Golde
TE (2015) IL-10 Alters Immunoproteostasis in APP Mice, Increasing Plaque Burden and
Worsening Cognitive Behavior. Neuron **85**, 519–533.

[94] Chakrabarty P, Tianbai L, Herring A, Ceballos-Diaz C, Das P, Golde TE (2012)
Hippocampal expression of murine IL-4 results in exacerbation of amyloid deposition.
Mol Neurodegener **7**, 36.

[95] Chakrabarty P, Herring A, Ceballos-Diaz C, Das P, Golde TE (2011) Hippocampal
expression of murine TNFα results in attenuation of amyloid deposition in vivo.
Molecular Neurodegeneration **6**, 16.

[96] Chakrabarty P, Ceballos-Diaz C, Beccard A, Janus C, Dickson D, Golde TE, Das P
(2010) IFN-gamma promotes complement expression and attenuates amyloid plaque
deposition in amyloid beta precursor protein transgenic mice. J. Immunol. **184**, 5333–
5343.

[97] Chakrabarty P, Jansen-West K, Beccard A, Ceballos-Diaz C, Levites Y, Verbeeck C,
Zubair AC, Dickson D, Golde TE, Das P (2010) Massive gliosis induced by interleukin-6
suppresses Abeta deposition in vivo: evidence against inflammation as a driving force for
amyloid deposition. FASEB J. **24**, 548–559.

[98] Ghosh S, Wu MD, Shaftel SS, Kyrkanides S, LaFerla FM, Olschowka JA, O'Banion MK
(2013) Sustained interleukin-1β overexpression exacerbates tau pathology despite
reduced amyloid burden in an Alzheimer's mouse model. J. Neurosci. **33**, 5053–5064.

[99] Guillot-Sestier M-V, Doty KR, Gate D, Rodriguez J, Leung BP, Rezai-Zadeh K, Town T
(2015) Il10 Deficiency Rebalances Innate Immunity to Mitigate Alzheimer-Like
Pathology. Neuron **85**, 534–548.

[100] Shaftel SS, Carlson TJ, Olschowka JA, Kyrkanides S, Matousek SB, O'Banion MK
(2007) Chronic interleukin-1beta expression in mouse brain leads to leukocyte
infiltration and neutrophil-independent blood brain barrier permeability without overt
neurodegeneration. J. Neurosci. **27**, 9301–9309.

[101] Town T, Laouar Y, Pittenger C, Mori T, Szekely CA, Tan J, Duman RS, Flavell RA
(2008) Blocking TGF-β–Smad2/3 innate immune signaling mitigates Alzheimer-like
pathology. Nature Medicine **14**, 681−687.
[102] Ferretti MT, Merlini M, Späni C, Gericke C, Schweizer N, Enzmann G, Engelhardt B,
Kulic L, Suter T, Nitsch RM (2016) T-cell brain infiltration and immature antigen-
presenting cells in transgenic models of Alzheimer's disease-like cerebral amyloidosis.
Brain, Behavior, and Immunity **54**, 211−225.
[103] GTEx Project (2018) The Genotype-Tissue Expression (GTEx) Project was supported by
the Common Fund of the Office of the Director of the National Institutes of Health, and
by NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS. The data used for the analyses
described in this manuscript were obtained from the GTEx Portal on 11/04/18.
[104] Hogervorst E, Combrinck M, Smith AD (2003) Testosterone and gonadotropin levels in
men with dementia. Neuro Endocrinol. Lett. **24**, 203−208.
[105] Dhingra R, Kwee LC, Diaz-Sanchez D, Devlin RB, Cascio W, Hauser ER, Gregory S,
Shah S, Kraus WE, Olden K, Ward-Caviness CK (2019) Evaluating DNA methylation
age on the Illumina MethylationEPIC Bead Chip. PLOS ONE **14**, e0207834.
[106] Levine ME, Lu AT, Quach A, Chen BH, Assimes TL, Bandinelli S, Hou L, Baccarelli
AA, Stewart JD, Li Y, Whitsel EA, Wilson JG, Reiner AP, Aviv A, Lohman K, Liu Y,
Ferrucci L, Horvath S (2018) An epigenetic biomarker of aging for lifespan and
healthspan. Aging (Albany NY) **10**, 573−591.
[107] Lu AT, Quach A, Wilson JG, Reiner AP, Aviv A, Raj K, Hou L, Baccarelli AA, Li Y,
Stewart JD, Whitsel EA, Assimes TL, Ferrucci L, Horvath S (2019) DNA methylation
GrimAge strongly predicts lifespan and healthspan. Aging (Albany NY) **11**, 303−327.
[108] Fitten LJ, Ortiz F, Fairbanks L, Bartzokis G, Lu P, Klein E, Coppola G, Ringman J
(2014) Younger age of dementia diagnosis in a Hispanic population in southern
California. Int J Geriatr Psychiatry **29**, 586−593.
[109] Mayeda ER, Glymour MM, Quesenberry CP, Whitmer RA (2016) Inequalities in
dementia incidence between six racial and ethnic groups over 14 years. Alzheimer's &
Dementia **12**, 216−224.
[110] Steenland K, Goldstein FC, Levey A, Wharton W (2016) A Meta-Analysis of
Alzheimer's Disease Incidence and Prevalence Comparing African-Americans and
Caucasians. J. Alzheimers Dis. **50**, 71−76.
[111] Sharp ES, Gatz M (2011) The Relationship between Education and Dementia An
Updated Systematic Review. Alzheimer Dis Assoc Disord **25**, 289−304.
924

Figure 1. Marginal mean (±95% confidence interval) of **CSF levels** of A. IL-16 (pg/ml), B. IL-8

(pg/ml), C. C-reactive protein (CRP; µg/ml), and **plasma levels** of D. CRP (CRP; µg/ml), E. IL-

18 (pg/ml), and F. IgE (ng/ml) in ADNI participants by sex and presence or absence of APOEε4

alleles (none or 1 and 2 alleles).

Figure 2. Marginal mean (± 95% confidence interval) of **CSF levels** of A. IL-16 (pg/ml), B.

IgA (mg/ml), C. Intercellular adhesion molecule (ICAM1; ng/ml), and **plasma levels** of D. C-

reactive protein (CRP; µg/ml), E. cortisol (ng/ml), and F. IL-16 in ADNI participants by sex and

diagnosis (CN, cognitively normal; LMCI, late mild cognitive impairment; and AD, Alzheimer's

disease).

Figure 3. Correlations between plasma and CSF levels of A. CD 40, B. ICAM1, C. IL-3, and D.

IgA in males and females separately. CD 40 and ICAM1 were positively correlated in males

while IL-3 was negatively correlated in females. IgA was more strongly correlated in males

compared to females (see Table 5 for details).

Figure 4. Universal epigenetic age acceleration does not differ statistically significantly by

participant sex or diagnosis (CN, EMCI, LMCI, AD) in this ADNI cohort. CN, cognitively

normal; EMCI, early mild cognitive impairment; LMCI, late mild cognitive impairment; AD,

Alzheimer's disease.

Figure 1

CSF

Plasma

Figure 2

CSF

Plasma

Figure 3

Figure 4

Table 1. Demographic and clinical information for all ADNI participants subdivided by sex. A. Participants with measured biomarkers in cerebrospinal fluid (CSF), B. Participants with available whole blood Illumina HumanMethylationEPIC DNA methylation data, C. Participants with matched Illumina HumanMethylationEPIC DNA methylation array data and measured CSF biomarkers. In all three subdata sets, females were significantly younger and had fewer years of education than males. In data set A (but not B and C), more females (24.0 % compared to 21.8%) were diagnosed with AD, more females were cognitively normal (26.5% compared to 22.9%) and fewer females were diagnosed with late MCI compared to males (49.5% compared to 55.3%). In data set A, females had lower CSF cortisol, interleukin-3, interleukin 8, interleukin-16, immunoglobulin A, and intercellular adhesion molecule 1 compared to males. Empty cells indicate data not available or not analysed. CN, cognitively normal; EMCI, early mild cognitive impairment; LMCI, late mild cognitive impairment; AD, Alzheimer's disease.

	A				**B**				**C**			
		Sex				**Sex**				**Sex**		
	Total	**Female**	**Male**	**P-value**	**Total**	**Female**	**Male**	**P-value**	**Total**	**Female**	**Male**	**P-value**
	No. 279	**No. 109**	**No. 170**		**No. 640**	**No. 284**	**No. 356**		**No. 533**	**No. 243**	**No. 290**	
Age												
Mean (SD)	75.15 (±6.86)	73.75 (±6.69)	76.04 (±6.83)	0.007	75.63 (±7.68)	74.78 (±8.03)	76.31 (±7.32)	<0.0001	75.01 (±7.61)	74.31 (±8.10)	75.61 (±7.11)	0.0019
Education (years)												
Mean (SD)	15.69 (±2.95)	14.68 (±2.74)	16.34 (±2.90)	< 0.0001	16.21 (±2.70)	15.53 (±2.59)	16.75 (±2.68)	<0.0001	16.24 (±2.64)	15.57 (±2.49)	16.83 (±2.64)	< 0.0001
Ethnicity												
White	267 (95.70%)	103 (94.50%)	164 (96.47%)	0.55	627 (97.97%)	279 (98.23 %)	348 (97.75%)	0.78	521 (97.75 %)	238 (97.94 %)	283 (97.94 %)	0.99
Not White[I]	12 (4.30%)	6 (5.50%)	6 (3.53%)		13 (2.03%)	5 (1.76%)	8 (2.25%)		12 (2.25 %)	5 (2.06 %)	7 (2.41 %)	
Baseline diagnosis												
CN	74 (26.5%)	35 (32.1%)	39 (22.9%)	0.051	217 (33.9%)	109 (38.38%)	108 (30.34%)	0.11	171 (32.08 %)	88 (36.21 %)	83 (28.62 %)	0.19
EMCI					186 (29.06%)	83 (29.23%)	103 (28.93%)		173 (32.46 %)	79 (32.51 %)	94 (32.41 %)	
LMCI	138 (49.5%)	44 (40.4%)	94 (55.3%)		200 (31.25%)	78 (27.46%)	122 (34.27%)		155 (29.08 %)	94 (38.68 %)	92 (31.72 %)	
AD	67 (24.0%)	30 (27.5%)	37 (21.8%)		37 (5.78%)	14 (4.23 %)	23 (6.46%)		34 (6.38 %)	13 (5.35 %)	21 (7.24 %)	
APOEε4 allele number												
0	134 (48.03%)	51 (46.79%)	83 (48.82%)	0.78	369 (57.66 %)	169 (59.51%)	200 (56.18%)	0.37	313 (58.72 %)	146 (60.08 %)	167 (57.59 %)	0.45
1	109 (39.07%)	42 (38.53%)	67 (39.41%)		220 (34.38%)	97 (34.15%)	123 (34.55%)		173 (32.46 %)	80 (32.92 %)	93 (32.07 %)	
2	36 (12.90%)	16 (14.68%)	20 (11.76%)		51 (7.97%)	18 (6.34%)	33 (9.27%)		47 (8.82%)	17 (7.00%)	30 (10.34 %)	
Cortisol (ng/mL)												
Mean (SD)	16.05 (±6.04)	14.92 (±6.01)	16.78 (±5.96)	0.008								
C reactive protein (ug/mL)												
Mean (SD)	-2.83 (±0.56)	-2.77 (±0.64)	-2.87 (±0.51)	0.23								
CD40 antigen (ng/mL)												
Mean (SD)	-0.65 (±0.12)	-0.66 (±0.10)	-0.64 (±0.14)	0.12								
Interleukin 16 (pg/mL)												
Mean (SD)	0.91 (±0.18)	0.87 (±0.17)	0.94 (±0.19)	0.004								
Interleukin 3 (ng/mL)												
Mean (SD)	-2.22 (±0.32)	-2.28 (±0.29)	-2.17 (±0.34)	0.001								
Interleukin 6 receptor (ng/mL)												
Mean (SD)	-0.01 (±0.15)	-0.02 (±0.14)	-0.00 (±0.15)	0.30								
Interleukin 8 (pg/mL)												
Mean (SD)	1.68 (±0.15)	1.64 (±0.11)	1.70 (±0.16)	0.001								
Intercellular adhesion molecule 1 (ng/mL)												
Mean (SD)	0.96 (±0.44)	0.83 (±0.33)	1.04 (±0.48)	0.0001								
Immunoglobulin A (mg/mL)												
Mean (SD)	-2.54 (±0.31)	-2.68 (±0.26)	-2.45 (±0.31)	< 0.0001								
Amyloid Beta												
Mean (SD)									1040.98 (±454.72)	1055.50 (±449.23)	1028.35 (±459.36)	0.18
Tau												
Mean (SD)									289.80 (±124.68)	300.90 (±139.07)	280.13 (±109.82)	0.072
PTau												
Mean (SD)									27.47 (±13.65)	28.25 (±15.08)	26.78 (±12.24)	0.36

P-values are from Wilcoxon rank sum tests for continuous variables and Fisher's exact tests for categorical variables. IIncludes self-reported Black, Asian, American Indian/Alaskan, and >1 ethnicity.

Table 2. List of biomarkers analysed in the current study with their main biological function and main finding in the CSF and plasma. Main effects of sex (sex difference), APOEε4 genotype (non-carriers or carriers), and diagnosis (CN, cognitively normal; LMCI, late mild cognitive impairment; AD, Alzheimer's disease) and interaction between sex and APOEε4 genotype (sex * APOEε4 genotype) are shown. Significant effects are adjusted P≤0.05 and trends are adjusted P≤0.08. See results for details. n/a - not available

Biomarker	Biological function	Results in CSF	Results in Plasma
Cortisol	Stress hormone and inflammation	Sex difference (trend): ♀ < ♂	Diagnosis: LMCI < AD
Intercellular adhesion molecule 1	Immune response, immunoglobulin family	Sex difference: ♀ < ♂	Sex difference: ♀ > ♂
C-reactive protein	Inflammation	APOEε4 genotype: non-carriers > carriers	Sex difference: ♀ > ♂ APOEε4 genotype: non-carriers > carriers Diagnosis (trend): CN > LMCI and CN > AD
CD40 antigen	Immune and inflammatory responses		Diagnosis (trend): LMCI < AD
Interleukin 3	Immune and inflammatory responses		
Interleukin 6 receptor	Immune and inflammatory responses	APOEε4 genotype (trend): non-carriers < carriers	
Interleukin 8	Immune and inflammatory responses	Sex * APOEε4 genotype: non-carriers ♀ < ♂ carriers ♀ = ♂	
Interleukin 16	Immune and inflammatory responses	Sex * APOEε4 genotype: non-carriers ♀ < ♂ carriers ♀ = ♂	Diagnosis (trend): CN > LMCI CN > AD
Immunoglobulin A	Immune and inflammatory responses	Sex difference: ♀ < ♂	
Interleukin 18	Immune and inflammatory responses	n/a	Sex difference: ♀ < ♂
Immunoglobulin E	Immune and inflammatory responses	n/a	Sex difference: ♀ < ♂

Table 3. Linear regression results for models with sex and APOE genotype (non-carriers or carriers of 1 or 2 APOEε4 alleles). P-values are for overall tests and are FDR-adjusted. Only shown are the models with significant associations (adjusted P ≤0.05) and trends (adjusted P ≤0.08). All model summaries are available in Supplementary Table S1.

Predictors	Cortisol Cortisol ng ml			C Reactive Protein ug/ml			Interleukin 16 pg/ml			Interleukin 6 receptor ng/ml			Interleukin 8 pg/ml			Immunoglobulin A mg/ml			Intercellular Adhesion Molecule ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	1.43 (-7.00 – 9.87)			-3.03 (-3.84 – -2.23)			0.42 (0.15 – 0.68)			-0.27 (-0.48 – -0.06)			1.33 (1.11 – 1.54)			-2.91 (-3.35 – -2.48)			-0.23 (-0.83 – 0.38)		
AGE (years)	0.22 (0.12 – 0.32)			0.01 (-0.00 – 0.02)			0.01 (0.00 – 0.01)			0.00 (0.00 – 0.01)			0.00 (0.00 – 0.01)			0.00 (-0.00 – 0.01)			0.01 (0.01 – 0.02)		
EDUCATION (years)	-0.21 (-0.45 – 0.03)			-0.01 (-0.03 – 0.02)			-0.01 (-0.01 – 0.00)			-0.00 (-0.01 – 0.00)			0.00 (-0.00 – 0.01)			0.00 (-0.01 – 0.01)			-0.00 (-0.02 – 0.02)		
Male (ref = Female)	1.72 (0.25 – 3.19)	0.022	**0.07**	-0.11 (-0.25 – 0.03)	0.126	0.22	0.13 (0.07 – 0.19)			0.02 (-0.02 – 0.05)	0.372	0.5	0.09 (0.04 – 0.14)	<0.001		0.21 (0.14 – 0.29)	<0.001	**<0.001**	0.18 (0.07 – 0.28)	0.001	**0.009**
APOE4 1 or 2 alleles (ref = 0 alleles)	0.82 (-0.55 – 2.19)	0.241	0.35	-0.22 (-0.35 – -0.09)	0.001	**0.009**	0.08 (0.01 – 0.14)			0.04 (0.00 – 0.07)	0.025	**0.071**	0.03 (-0.02 – 0.09)	0.261		-0.00 (-0.07 – 0.07)	0.898	0.97	0.08 (-0.02 – 0.18)	0.128	0.22
Interaction: Male by 1 or 2 alleles							-0.13 (-0.22 – -0.05)	0.003	**0.016**				-0.09 (-0.16 – -0.02)	**0.008**	**0.035**						
Observations	279			279			279			279			279			279			279		
R² / adjusted R²	0.095 / 0.082			0.055 / 0.042			0.122 / 0.106			0.046 / 0.032			0.092 / 0.076			0.123 / 0.111			0.105 / 0.092		

Table 4. Linear regression results for models with sex and baseline diagnosis (CN, cognitively normal; LMCI, late mild cognitive impairment; AD, Alzheimer's disease). Only shown are the models with significant associations (adjusted P≤0.05) and trends (adjusted P≤0.08). P-values are for overall tests and are FDR-adjusted. All model summaries are available in Supplementary Table S2. There were no significant interactions between diagnosis and sex.

Predictors	Interleukin 16 pg/ml			Immunoglobulin A mg/ml			Intercellular Adhesion Molecule ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	0.51 (0.26 – 0.77)			-2.92 (-3.35 – -2.48)			-0.19 (-0.80 – 0.42)		
AGE (years)	0.01 (0.00 – 0.01)			0.00 (-0.00 – 0.01)			0.01 (0.01 – 0.02)		
EDUCATION (years)	-0.01 (-0.01 – 0.00)			0.00 (-0.01 – 0.01)			-0.00 (-0.02 – 0.02)		
Male (ref = Female)	0.06 (0.02 – 0.11)	0.007	**0.055**	0.21 (0.14 – 0.29)	<0.001	**<0.001**	0.17 (0.06 – 0.28)	0.002	**0.026**
Diagnosis (ref = CN)		0.4	0.61		0.99	0.99		0.54	0.67
LMCI	0.01 (-0.05 – 0.06)			0.00 (-0.08 – 0.09)			0.06 (-0.06 – 0.18)		
AD	-0.03 (-0.09 – 0.03)			-0.01 (-0.11 – 0.09)			0.02 (-0.12 – 0.16)		
Observations	279			279			279		
R^2 / adjusted R^2	0.099 / 0.082			0.124 / 0.107			0.101 / 0.085		

Table 5. Pearson's correlations between plasma and CSF levels of the biomarkers analysed in the current study separetly in males and females. Differences in the correlations were determined using confidence intervals. Significant correlations and differences between correlations are P≤0.05 and trends are P≤0.08

	Correlation (r) in Males n = 149	P-value	Correlation (r) in Females n = 88	P-value	Difference (95% CI)	P-value
Cortisol	0.176	**0.032**	0.327	**0.002**	-0.151 (-0.388-0.101)	0.24
C reactive protein	0.793	**< 0.0001**	0.860	**< 0.0001**	-0.067 (-0.149-0.019)	0.12
CD40 antigen	0.374	**< 0.0001**	0.016	0.88	0.358 (0.103-0.606)	**0.01**
IL-16	0.156	**0.058**	0.290	**0.006**	-0.134 (-0.376-0.121)	0.30
IL-3	0.001	0.989	-0.246	**0.021**	0.247 (-0.016-0.493)	**0.06**
IL-6 receptor	0.459	**< 0.0001**	0.493	**< 0.0001**	-0.034 (-0.232-0.179)	0.75
IL-8	0.138	0.093	0.287	**0.007**	-0.149 (-0.392-0.107)	0.25
IgA	0.705	**< 0.0001**	0.529	**< 0.0001**	0.176 (0.012-0.361)	**0.03**
ICAM1	0.231	**0.005**	-0.021	0.849	0.252 (-0.011-0.507)	**0.06**

Table 6. Results of epigenetic age and epigenetic age acceleration calculation for all DNAme analyses, for both the larger DNAme cohort and the subset of samples with matched CSF biomarker data.

	DNAme Cohort				DNAme & CSF Biomarker Data Cohort			
		Sex				Sex		
	Total No. 640	Female No. 284	Male No. 356	P-value	Total No. 533	Female No. 243	Male No. 290	P-value
Age								
Mean (SD)	75.63 (±7.68)	74.78 (±8.03)	76.31 (±7.32)	<0.0001	75.01 (±7.61)	74.31 (±8.10)	75.61 (±7.11)	0.0019
Epigenetic Age (years)								
Mean (SD)	69.92 (±8.06)	67.45 (±8.15)	70.11 (±7.79)	<0.0001	68.47 (±8.17)	67.05 (±8.33)	69.72 (±7.82)	<0.0001
Epigenetic Age Acceleration (years)								
Mean (SD)	0.025 (±4.22)	-0.14 (±4.16)	0.16 (±4.26)	0.1	0.027 (±4.30)	-0.18 (±4.23)	0.20 (4.35)	0.057
Intrinsic Age Acceleration (years)								
Mean (SD)	0.026 (±4.11)	-0.19 (±4.06)	0.20 (±4.15)	0.019	0.020 (±4.18)	-0.25 (±4.14)	0.26 (±4.21)	0.021

P-values are from Wilcoxon rank sum tests for continuous variables and Fisher's exact tests for categorical variables

Table 7. Linear model for assessment of relationship of biochemical concentrations and presence of APOEε4 risk allele on universal epigenetic age acceleration. Intrinsic epigenetic age acceleration linear model not shown. CN, cognitively normal; EMCI, early mild cognitive impairment; LMCI, late mild cognitive impairment; AD, Alzheimer's disease.

Predictors	Estimates	CI	adjusted p
	Age Acceleration & CSF Biomarkers		
(Intercept)	-0.61	-1.67 – 0.46	0.61
Male (ref = Female)	0.6	-0.11 – 1.31	0.34
Diagnosis (ref = CN)			
EMCI	0.71	-0.17 – 1.60	0.39
LMCI	0.4	-0.55 – 1.36	0.64
AD	0.53	-1.07 – 2.14	0.64
APOE risk allele			
1 or 2 alleles (ref = 0 alleles)	-0.24	-1.01 – 1.31	0.813
CSF Tau	0.0001	-0.0077 – 0.023	0.95
Observations	533		
R^2 / adjusted R^2	0.0108/-0.000438		

Inflammation and epigenetic age in Alzheimer's disease: do sex and APOE matter?

Paula Duarte-Guterman[1*], Amy M. Inkster[2,3*], Arianne Y. Albert[4], Cindy K. Barha[5], Wendy P. Robinson[2,3], Liisa A.M. Galea[1], on behalf of the Alzheimer's Disease Neuroimaging Initiative[δ]

1. Djavad Mowafaghian Centre for Brain Health and Department of Psychology, University of British Columbia, Vancouver, BC, Canada
2. Department of Medical Genetics, University of British Columbia, Vancouver, BC, Canada.
3. BC Children's Hospital Research Institute, Vancouver, BC, Canada
4. Women's Health Research Institute of British Columbia, Vancouver, BC, Canada
5. Djavad Mowafaghian Centre for Brain Health and Department of Physical Therapy, University of British Columbia, Vancouver, BC, Canada

Address all correspondence and requests for reprints to:
L. A. M. Galea, PhD
Djavad Mowafaghian Centre for Brain Health
2215 Wesbrook Mall
Vancouver, British Columbia
V6T 1Z3, Canada
E-mail: lgalea@psych.ubc.ca.

* Considered co-first authors.

[δ] Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Table S1. Linear regression results for all CSF variables investigated by sex and APOE genotype (non-carriers or carriers of 1 or 2 APOEε4 alleles).

Predictors	Cortisol Cortisol ng ml			C Reactive Protein ug/ml			CD 40 antigen ng/ml			Interleukin 16 pg/ml			Interleukin 3 ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	1.43 (-7.00 – 9.87)			-3.03 (-3.84 – -2.23)			-1.10 (-1.27 – -0.93)			0.42 (0.15 – 0.68)			-3.03 (-3.49 – -2.57)		
AGE (years)	0.22 (0.12 – 0.32)			0.01 (-0.00 – 0.02)			0.01 (0.00 – 0.01)			0.01 (0.00 – 0.01)			0.01 (0.00 – 0.02)		
EDUCATION (years)	-0.21 (-0.45 – 0.03)			-0.01 (-0.03 – 0.02)			-0.00 (-0.01 – 0.00)			-0.01 (-0.01 – 0.00)			0.00 (-0.01 – 0.01)		
Male (ref = Female)	1.72 (0.25 – 3.19)	0.022	**0.07**	-0.11 (-0.25 – 0.03)	0.126	0.22	0.01 (-0.02 – 0.04)	0.432	0.55	0.13 (0.07 – 0.19)			0.08 (-0.00 – 0.16)	0.057	0.15
APOE4 1 or 2 alleles (ref = 0 alleles)	0.82 (-0.55 – 2.19)	0.241	0.35	-0.22 (-0.35 – -0.09)	0.001	**0.009**	0.01 (-0.02 – 0.04)	0.59	0.72	0.08 (0.01 – 0.14)			-0.04 (-0.12 – 0.03)	0.262	0.37
Interaction: Male by 1 or 2 alleles										-0.13 (-0.22 – -0.05)	0.003	**0.016**			
Observations	279			279			279			279			279		
R^2 / adjusted R^2	0.095 / 0.082			0.055 / 0.042			0.125 / 0.112			0.122 / 0.106			0.076 / 0.063		

Table S1. Continued

Predictors	Interleukin 6 receptor ng/ml			Interleukin 8 pg/ml			Immunoglobulin A mg/ml			Intercellular Adhesion Molecule ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	-0.27 (-0.48 – -0.06)			1.33 (1.11 – 1.54)			-2.91 (-3.35 – -2.48)			-0.23 (-0.83 – 0.38)		
AGE (years)	0.00 (0.00 – 0.01)			0.00 (0.00 – 0.01)			0.00 (-0.00 – 0.01)			0.01 (0.01 – 0.02)		
EDUCATION (years)	-0.00 (-0.01 – 0.00)			0.00 (-0.00 – 0.01)			0.00 (-0.01 – 0.01)			-0.00 (-0.02 – 0.02)		
Male (ref = Female)	0.02 (-0.02 – 0.05)	0.372	0.5	0.09 (0.04 – 0.14)	<0.001		0.21 (0.14 – 0.29)	<0.001	**<0.001**	0.18 (0.07 – 0.28)	0.001	**0.009**
APOE4 1 or 2 alleles (ref = 0 alleles)	0.04 (0.00 – 0.07)	0.025	**0.071**	0.03 (-0.02 – 0.09)	0.261		-0.00 (-0.07 – 0.07)	0.898	0.97	0.08 (-0.02 – 0.18)	0.128	0.22
Interaction: Male by 1 or 2 alleles				-0.09 (-0.16 – -0.02)	0.008	**0.035**						
Observations	279			279			279			279		
R^2 / adjusted R^2	0.046 / 0.032			0.092 / 0.076			0.123 / 0.111			0.105 / 0.092		

Table S2. Linear regression results for all CSF variables investigated by sex and baseline diagnosis (CN, cognitively normal; LMCI, late mild cognitive impairment; AD, Alzheimer's disease). There were no significant interactions between diagnosis and sex.

Predictors	Cortisol Cortisol ng ml			C Reactive Protein ug/ml			CD 40 antigen ng/ml			Interleukin 16 pg/ml			Interleukin 3 ng/ml		
	Estimates (CI)	*p*	*adjusted p*	*Estimates (CI)*	*p*	*adjusted p*	*Estimates (CI)*	*p*	*adjusted p*	*Estimates (CI)*	*p*	*adjusted p*	*Estimates (CI)*	*p*	*adjusted p*
(Intercept)	1.49 (-6.96 – 9.95)			-3.12 (-3.94 – -2.30)			-1.08 (-1.25 – -0.91)			0.51 (0.26 – 0.77)			-3.03 (-3.49 – -2.57)		
AGE (years)	0.22 (0.12 – 0.32)			0.01 (-0.00 – 0.02)			0.01 (0.00 – 0.01)			0.01 (0.00 – 0.01)			0.01 (0.00 – 0.02)		
EDUCATION (years)	-0.23 (-0.47 – 0.01)			-0.00 (-0.03 – 0.02)			-0.00 (-0.01 – 0.00)			-0.01 (-0.01 – 0.00)			0.00 (-0.01 – 0.01)		
Male (ref = Female)	1.55 (0.07 – 3.04)	0.041	0.18	-0.10 (-0.24 – 0.05)	0.19	0.42	0.01 (-0.02 – 0.04)	0.51	0.65	0.06 (0.02 – 0.11)	0.007	**0.055**	0.08 (-0.00 – 0.16)	0.052	0.18
Diagnosis (ref = CN)		0.24	0.49		0.15	0.36		0.067	0.2		0.4	0.61		0.5	0.65
LMCI	1.23 (-0.43 – 2.88)			-0.15 (-0.31 – 0.01)			0.01 (-0.03 – 0.04)			0.01 (-0.05 – 0.06)			-0.04 (-0.13 – 0.05)		
AD	0.10 (-1.83 – 2.02)			-0.15 (-0.34 – 0.03)			-0.03 (-0.07 – 0.00)			-0.03 (-0.09 – 0.03)			-0.06 (-0.17 – 0.04)		
Observations	279			279			279			279			279		
R^2 / adjusted R^2	0.100 / 0.084			0.030 / 0.012			0.141 / 0.125			0.099 / 0.082			0.077 / 0.060		

Table S2. Continued

Predictors	Interleukin 6 receptor ng/ml			Interleukin 8 pg/ml			Immunoglobulin A mg/ml			Intercellular Adhesion Molecule ng/ml		
	Estimates (CI)	*p*	*adjusted p*	*Estimates (CI)*	*p*	*adjusted p*	*Estimates (CI)*	*p*	*adjusted p*	*Estimates (CI)*	*p*	*adjusted p*
(Intercept)	-0.22 (-0.43 – -0.01)			1.35 (1.14 – 1.56)			-2.92 (-3.35 – -2.48)			-0.19 (-0.80 – 0.42)		
AGE (years)	0.00 (0.00 – 0.01)			0.00 (0.00 – 0.01)			0.00 (-0.00 – 0.01)			0.01 (0.01 – 0.02)		
EDUCATION (years)	-0.00 (-0.01 – 0.00)			0.00 (-0.00 – 0.01)			0.00 (-0.01 – 0.01)			-0.00 (-0.02 – 0.02)		
Male (ref = Female)	0.02 (-0.02 – 0.05)	0.42	0.61	0.04 (0.01 – 0.08)	0.019	0.12	0.21 (0.14 – 0.29)	<0.001	**<0.001**	0.17 (0.06 – 0.28)	0.002	**0.026**
Diagnosis (ref = CN)		0.46	0.65		0.85	0.96		0.99	0.99		0.54	0.67
LMCI	0.01 (-0.04 – 0.05)			0.01 (-0.03 – 0.05)			0.00 (-0.08 – 0.09)			0.06 (-0.06 – 0.18)		
AD	-0.02 (-0.07 – 0.03)			0.01 (-0.03 – 0.06)			-0.01 (-0.11 – 0.09)			0.02 (-0.12 – 0.16)		
Observations	279			279			279			279		
R^2 / adjusted R^2	0.034 / 0.016			0.062 / 0.045			0.124 / 0.107			0.101 / 0.085		

Table S3. Linear regression results for all plasma variables investigated by sex and APOE genotype (non-carriers or carriers of 1 or 2 APOEε4 alleles).

Predictors	Cortisol Cortisol ng ml			C Reactive Protein ug/ml			CD 40 antigen ng/ml			Interleukin 16 pg/ml			Interleukin 3 ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	2.06 (1.92 – 2.19)			0.46 (-0.05 – 0.97)			-0.52 (-0.65 – -0.39)			2.42 (2.27 – 2.58)			-1.86 (-2.16 – -1.57)		
AGE (years)	0.00 (-0.00 – 0.00)			0.00 (-0.00 – 0.01)			0.01 (0.00 – 0.01)			0.00 (0.00 – 0.00)			0.00 (-0.00 – 0.00)		
EDUCATION (years)	-0.00 (-0.00 – 0.00)			-0.03 (-0.04 – -0.01)			-0.00 (-0.01 – 0.00)			-0.01 (-0.01 – -0.00)			0.01 (0.00 – 0.02)		
Male (ref = Female)	0.01 (-0.01 – 0.04)	0.26	0.57	-0.13 (-0.22 – -0.04)	0.006	**0.048**	-0.04 (-0.07 – -0.01)			0.01 (-0.02 – 0.04)	0.50	0.79	-0.02 (-0.07 – 0.04)	0.57	0.8
APOE4 1 or 2 alleles (ref = 0 alleles)	0.02 (-0.00 – 0.04)	0.11	0.29	-0.31 (-0.39 – -0.22)	<0.001	**<0.001**	-0.03 (-0.07 – 0.00)			-0.01 (-0.04 – 0.01)	0.31	0.66	0.02 (-0.03 – 0.07)	0.43	0.67
Interaction: Male by 1 or 2 alleles							0.05 (0.01 – 0.10)	**0.022**	0.11						
Observations	527			526			527			527			527		
R² / adjusted R²	0.013 / 0.005			0.123 / 0.117			0.126 / 0.118			0.042 / 0.033			0.011 / 0.004		

Table S3. Continued

Predictors	Interleukin 6 receptor ng/ml			Interleukin 8 pg/ml			Immunoglobulin A mg/ml			Intercellular Adhesion Molecule ng/ml			Interleukin 18 pg/ml			Immunoglobulin E ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	1.43 (1.29 – 1.57)			0.79 (0.60 – 0.99)			0.48 (0.25 – 0.71)			1.89 (1.75 – 2.04)			2.42 (2.24 – 2.59)			1.42 (0.88 – 1.97)		
AGE (years)	0.00 (-0.00 – 0.00)			0.00 (0.00 – 0.01)			0.00 (-0.00 – 0.00)			0.00 (0.00 – 0.00)			-0.00 (-0.00 – 0.00)			0.00 (-0.00 – 0.01)		
EDUCATION (years)	-0.00 (-0.01 – 0.00)			0.00 (-0.00 – 0.01)			-0.00 (-0.01 – 0.01)			0.00 (-0.00 – 0.00)			-0.00 (-0.01 – 0.00)			0.01 (-0.01 – 0.02)		
Male (ref = Female)	-0.02 (-0.05 – -0.00)	0.049	0.14	-0.05 (-0.10 – 0.00)			0.02 (-0.02 – 0.06)	0.371	0.67	-0.04 (-0.06 – -0.01)	0.008	**0.051**	0.06 (0.03 – 0.09)	<0.001	**0.001**	0.24 (0.14 – 0.33)	<0.001	**<0.001**
APOE4 1 or 2 alleles (ref = 0 alleles)	0.00 (-0.02 – 0.03)	0.70	0.84	-0.08 (-0.13 – -0.02)			-0.02 (-0.06 – 0.02)	0.387	0.67	0.01 (-0.02 – 0.03)	0.586	0.80	-0.04 (-0.07 – -0.01)	0.019	0.10	0.01 (-0.09 – 0.10)	0.91	0.94
Interaction: Male by 1 or 2 alleles				0.07 (0.00 – 0.14)	0.046	0.14												
Observations	527			527			527			527			527			527		
R² / adjusted R²	0.011 / 0.004			0.036 / 0.026			0.019 / 0.012			0.008 / 0.000			0.038 / 0.031			0.052 / 0.045		

Table S4. Linear regression results for all plasma variables investigated by sex and baseline diagnosis (CN, cognitively normal; LMCI, late mild cognitive impairment; AD, Alzheimer's disease). There were no significant interactions between diagnosis and sex.

Predictors	Cortisol Cortisol ng ml			C Reactive Protein ug/ml			CD 40 antigen ng/ml			Interleukin 16 pg/ml			Interleukin 3 ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	2.09 (1.95 − 2.22)			0.33 (-0.21 − 0.88)			-0.55 (-0.68 − -0.42)			2.48 (2.31 − 2.65)			-1.78 (-2.08 − -1.47)		
AGE (years)	0.00 (-0.00 − 0.00)			0.01 (0.00 − 0.01)			0.01 (0.00 − 0.01)			0.00 (0.00 − 0.00)			0.00 (-0.00 − 0.00)		
EDUCATION (years)	-0.00 (-0.00 − 0.00)			-0.02 (-0.04 − -0.01)			-0.00 (-0.00 − 0.00)			-0.01 (-0.01 − -0.00)			0.01 (0.00 − 0.02)		
Male (ref = Female)	0.02 (-0.01 − 0.04)	0.18	0.44	-0.12 (-0.21 − -0.03)	0.012	**0.056**	-0.01 (-0.03 − 0.01)	0.38	0.59	0.01 (-0.02 − 0.04)	0.39	0.57	-0.02 (-0.07 − 0.04)	0.56	0.75
Diagnosis (ref = CN)		0.0009	**0.010**		0.007	**0.056**		0.016	**0.067**		0.009	**0.054**		0.044	0.17
LMCI	-0.02 (-0.07 − 0.02)			-0.27 (-0.44 − -0.10)			-0.02 (-0.06 − 0.02)			-0.08 (-0.13 − -0.03)			-0.04 (-0.13 − 0.06)		
AD	0.03 (-0.02 − 0.08)			-0.24 (-0.43 − -0.05)			0.02 (-0.02 − 0.07)			-0.08 (-0.14 − -0.02)			-0.11 (-0.21 − -0.00)		
Observations	527			526			527			526			527		
R^2 / adjusted R^2	0.034 / 0.024			0.061 / 0.052			0.131 / 0.123			0.048 / 0.039			0.022 / 0.013		

Table S4. Continued

Predictors	Interleukin 6 receptor ng/ml			Interleukin 8 pg/ml			Immunoglobulin A mg/ml			Intercellular Adhesion Molecule ng/ml			Interleukin 18 pg/ml			Immunoglobulin E ng/ml		
	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p	Estimates (CI)	p	adjusted p
(Intercept)	1.47 (1.33 − 1.61)			0.73 (0.54 − 0.93)			0.43 (0.20 − 0.67)			1.90 (1.75 − 2.05)			2.33 (2.15 − 2.51)			1.44 (0.89 − 2.00)		
AGE (years)	0.00 (-0.00 − 0.00)			0.00 (0.00 − 0.01)			0.00 (-0.00 − 0.00)			0.00 (-0.00 − 0.00)			0.00 (-0.00 − 0.00)			0.00 (-0.00 − 0.01)		
EDUCATION (years)	-0.00 (-0.01 − 0.00)			0.00 (-0.00 − 0.01)			-0.00 (-0.01 − 0.01)			0.00 (-0.00 − 0.01)			-0.00 (-0.01 − 0.00)			0.01 (-0.01 − 0.02)		
Male (ref = Female)	-0.02 (-0.05 − 0.00)	0.063	0.19	-0.01 (-0.04 − 0.03)	0.65	0.79	0.02 (-0.02 − 0.06)	0.42	0.59	-0.03 (-0.06 − -0.01)	0.011	**0.056**	0.06 (0.03 − 0.09)	<0.001	**0.004**	0.24 (0.14 − 0.33)	<0.001	**<0.001**
Diagnosis (ref = CN)		0.34	0.56		0.11	0.30		0.72	0.82		0.063	0.19		0.32	0.56		0.88	0.94
LMCI	-0.03 (-0.08 − 0.01)			-0.04 (-0.10 − 0.02)			0.03 (-0.05 − 0.10)			-0.01 (-0.06 − 0.03)			0.04 (-0.01 − 0.10)			-0.00 (-0.18 − 0.17)		
AD	-0.03 (-0.08 − 0.02)			-0.00 (-0.07 − 0.07)			0.02 (-0.07 − 0.10)			0.02 (-0.03 − 0.08)			0.03 (-0.03 − 0.10)			-0.03 (-0.23 − 0.16)		
Observations	527			527			527			527			527			527		
R^2 / adjusted R^2	0.015 / 0.006			0.029 / 0.020			0.008 / -0.002			0.029 / 0.019			0.032 / 0.023			0.053 / 0.043		

Table S5. Correlation matrix of biochemical variables for those ADNI samples with available DNAme and biochemical data (n=533). Correlation coefficients are Spearman's rho.

	Amyloid Beta	p value	Tau	p value	pTau	p value
Amyloid Beta	1.00	<0.0001	-0.18	<0.0001	-0.27	<0.0001
Tau	-0.18	<0.0001	1.00	<0.0001	0.98	<0.0001
pTau	-0.27	<0.0001	0.98	<0.0001	1.00	<0.0001

Table S6. Contingency table for the distribution of the APOEε4 alleles (0 or 1 and 2 ε4 alleles) in each of the diagnosis groups (CN, cognitively normal; LMCI, late mild cognitive impairment; AD, Alzheimer's disease).

	CN	LMCI	AD
0	55 (74.32%)	60 (43.48%)	19(28.36%)
1 and 2	19 (25.68%)	78 (56.52%)	48 (71.64%)