# Partitioning plant spectral diversity into alpha and beta components

Etienne Laliberté[1,2*], Anna K. Schweiger[1,2], Pierre Legendre[2]

[1]Institut de recherche en biologie végétale, Université de Montréal, 4101 Sherbrooke Est, Montréal, Québec H1X 2B2, Canada

[2]Département de sciences biologiques, Université de Montréal, C. P. 6128, succursale Centre-ville, Montréal, Québec H3C 3J7, Canada

*Corresponding author. Email: etienne.laliberte@umontreal.ca

Phone: +1 514 343 6132

**Running title:** Spectral alpha, beta, and gamma diversity

**Keywords:** remote sensing, imaging spectroscopy, spectral variation hypothesis, spectral heterogeneity, alpha diversity, beta diversity, gamma diversity, spectranomics

**Type of article:** Methods

**Number of words:** 152 (abstract), 5062 (main text, excluding abstract, acknowledgements references, table and figure legends).

**Number of references:** 49

**Number of figures:** 5

**Number of tables:** 1

**Statement of authorship:** EL conceived the ideas and performed the analyses. EL and AKS wrote the manuscript. AKS and PL contributed to the ideas, and PL contributed to the writing.

## ABSTRACT

Plant spectral diversity — how plants differentially interact with solar radiation — is an integrator of plant chemical, structural, and taxonomic diversity that can be remotely sensed. We propose to measure spectral diversity as spectral variance, which allows the partitioning of the spectral diversity of a region, called spectral *gamma* ($\gamma$) diversity, into additive alpha ($\alpha$; within communities) and beta ($\beta$; among communities) components. Our method calculates the contributions of individual bands or spectral features to spectral $\gamma$-, $\beta$-, and $\alpha$-diversity, as well as the contributions of individual plant communities to spectral diversity. We present two case studies illustrating how our approach can identify "hotspots" of spectral $\alpha$-diversity within a region, and discover spectrally unique areas that contribute strongly to $\beta$-diversity. Partitioning spectral diversity and mapping its spatial components has many applications for conservation since high local diversity and distinctiveness in composition are two key criteria used to determine the ecological value of ecosystems.

## INTRODUCTION

Major environmental changes, including land-use change, climate change, and invasive species are altering the Earth's biodiversity. The rapid rate and broad extent of those changes far exceed our capacity to monitor them via field-based sampling alone. This calls for the development of new remote sensing approaches that can provide rapid estimates of biodiversity over broad regions (Pereira *et al.* 2013; Turner 2014; Bush *et al.* 2017). For terrestrial plants, imaging spectroscopy is emerging as the most promising remote sensing method for estimating biodiversity (Féret & Asner 2014; Wang & Gamon 2019). This is because its high spectral resolution allows plant species to be discriminated from one another, while also enabling the determination of ecologically important foliar functional traits (Asner & Martin 2009; Ustin *et al.* 2009).

For every pixel of an aerial image, imaging spectroscopy measures reflected solar radiation in tens to hundreds of contiguous, narrow (~10 nm wide) wavelength bands, usually covering all or part of the visible to shortwave infrared range (400–2500 nm) of the electromagnetic spectrum. Leaf "spectral signatures" of plants provide unique expressions among species of how solar radiation interacts with photosynthetic pigments, water, proteins, as well as structural and chemical defense compounds, and thus represent the evolution of plant adaptations to different environmental conditions (Cavender-Bares *et al.* 2016; McManus *et al.* 2016). At the crown scale, these spectral signatures are further influenced by architectural traits due to scattering of photons within canopies (Asner 1998; Ollinger 2010). Therefore, plant spectral diversity is emerging as an integrator of plant chemical, structural and taxonomic diversity that can be remotely sensed (Cavender-Bares *et al.* 2017; Schweiger *et al.* 2018; see also Appendix S1 in Supporting Information).

One of the most influential conceptual developments in community ecology has been

Whittaker's (1960, 1972) suggestion to partition biodiversity across space into $\alpha$, $\beta$, and $\gamma$

components. Originally, $\alpha$ diversity was defined as the species diversity *within* communities, and

$\beta$ as the variation in species composition *among* communities; together, $\alpha$- and $\beta$-diversities

jointly determined $\gamma$-diversity, which is the species diversity across an entire region of interest. In

this paper, we transpose this foundational ecological concept from species diversity to spectral

diversity (Fig. 1). This requires that the spatial resolution of the imagery matches the size of the

object of interest (Woodcock & Strahler 1987), meaning that pixels should be approximately

equal or smaller than the size of an average canopy plant. At such fine spatial resolutions, the

relationship between spectral and taxonomic diversity is strongest (Wang *et al.* 2018a) and

imaging spectroscopy can provide direct, spatially explicit estimates of plant alpha ($\alpha$; within

community) diversity (Féret & Asner 2014; Wang *et al.* 2018b), and can detect changes in plant

community composition across landscapes (Draper *et al.* 2019). The ability to generate wall-to-

wall, high-resolution maps of canopy plant diversity across entire regions brings tremendous

benefits for biodiversity science and conservation (e.g., Asner *et al.* 2017); however, conceptual

and methodological challenges remain, especially with regard to $\beta$-diversity estimation (Rocchini

*et al.* 2010, 2018).

Spectral diversity is sometimes called spectral heterogeneity or spectral variability (Rocchini *et*

*al.* 2010), and has been defined as spatial variation in spectral reflectance (Rocchini *et al.* 2010;

Ustin & Gamon 2010; Gholizadeh *et al.* 2018; Wang & Gamon 2019). Intuitively, spectral

diversity can be conceptualized as multivariate dispersion, for which there are various statistical

measures highlighting different aspects of spectral diversity. For example, Wang et al. (2018a)

used the average coefficient of variation (CV) of each band for a set of pixels, whereas Rocchini

et al. (2010) used the mean distance from the spectral centroid; we note that the latter has also

been proposed as a measure of functional diversity in multivariate trait space (Laliberté &

Legendre 2010). However, none of the currently used metrics allow the partitioning of spectral

diversity into its $\alpha$ (within communities) and $\beta$ (among communities) components (Fig. 1).

Here we propose to use the *spectral variance* among image pixels as a measure of spectral

diversity. Our approach builds on that of Legendre and De Cáceres (2013) for species inventory

data, adapts it to spectral data, and extends it to jointly consider $\alpha$-, $\beta$-, and $\gamma$-diversity. Casting

spectral diversity as spectral variance has a number of benefits:

1.  the classical partitioning of sums of squares allows us to partition spectral $\gamma$-diversity into

additive spectral $\alpha$- and $\beta$-diversity components (Fig. 1), from which the relative

importance of local and regional processes regulating spectral diversity across a region

of interest can be inferred;

2.  it allows us to estimate the contributions of individual plots or communities to spectral $\beta$-

diversity, highlighting areas that are spectrally distinct within the broader region;

3.  it allows us to calculate the contributions of individual bands or spectral features to

spectral $\gamma$-, $\beta$- or $\alpha$-diversity (Fig. 2), providing information about the underlying biological

traits driving spectral diversity;

4.  it is easily implemented in software packages in a computationally efficient way, which is

important when dealing with high-volume image data;

5.  it provides a direct link to other statistical procedures based on least squares (e.g.,

MANOVA, multiple linear regression, canonical redundancy analysis, *K*-means

partitioning).

After describing the theory behind our spectral diversity partitioning approach, we illustrate it

using a simulation. We then apply our method to imaging spectroscopy data collected over the

Bartlett Experimental Forest by the National Ecological Observatory Network (NEON) Airborne

Observation Platform (AOP; Kampe *et al.* 2010). The R code and data for our analyses are

available online (https://github.com/elaliberte/specdiv).

# PARTITIONING SPECTRAL DIVERSITY

## Size and shape of spatial units

Partitioning spectral $\gamma$-diversity into its $\alpha$ and $\beta$ components first requires defining the extent of

the region of interest (Fig. 1). Delineating the region of interest is relatively straightforward since

it corresponds to the region over which imagery is acquired or a subset thereof. Delineating the

size and shape of communities across the region of interest, however, is more difficult. What

constitutes an ecological community has been the subject of considerable debate (see review

by Ricklefs 2008). Generally, a community is defined as "a group of organisms representing

multiple species living in a specified place and time" (Vellend 2010). This definition implies that

a community must be larger than the size of an individual organism, but how much larger will

depend on the objectives of the study. For the purpose of this work, we focus on communities of

canopy plants, because these are the organisms that can be seen in aerial images. We use

"community" in the sense of "sampling unit" in vegetation surveys, which can be defined as the

area in which the species composition of the vegetation type of interest is adequately

represented (Mueller-Dombois & Ellenberg 1974).

Setting the size of a community to the size of typical inventory plot for a given ecosystem type

facilitates interpretation as this is the sampling unit that field ecologists are familiar with. For

example, forest inventory plots often measure 20 m × 20 m (Fig. 1), which is large enough to

include several canopy trees. However, we recognize that setting fixed and regularly shaped

boundaries to delineate communities is artificial (Ricklefs 2008), and point out that community

size and shape can be changed in the analysis.

## Spectral gamma (*γ*) diversity

Let **Y** = [$y_{ij}$] be a matrix containing the positions, along the *p* axes defining the *spectral space* (column vectors $\mathbf{y}_1$, $\mathbf{y}_2$, … $\mathbf{y}_p$ of **Y**), of *n* pixels (row vectors $\mathbf{x}_1$, $\mathbf{x}_2$, … $\mathbf{x}_n$ of **Y**) in a region of interest (Fig. 2). We use indices *i* and *j* to denote rows (pixels) and columns (axes) of matrix **Y**, respectively. The *p* axes could be all or a subset of the original spectral bands, a set of vegetation indices calculated from selected spectral bands (Bannari *et al.* 1995), or a set of *p* uncorrelated spectral features extracted using dimensionality reduction methods such as principal component analysis (PCA). We use PCA in this section and in our case studies and point out the mathematical relationships between the principal components (PCs) and spectral variation below. We use the general term *variation* for sums of squares (SS, an abbreviation for "sum of the squared deviations from the mean"), and reserve the term *variance* when talking about spectral diversity (SD).

We refer to the total spectral diversity of the entire region as spectral *γ*-diversity ($SD_\gamma$). $SD_\gamma$ is measured by the total variance of **Y**, or Var(**Y**). This is done by first computing for every pixel and spectral feature $y_{ij}$ the squared deviations $s_{ij}$ from the average pixel (across the whole region) in terms of spectral reflectance, i.e. the column means of **Y**:

$$s_{ij} = \left( y_{ij} - \overline{y}_j \right)^2. \tag{1}$$

The total sum of squares (SS) of matrix **Y** is calculated by summing all $s_{ij}$:

$$SS_\gamma = \sum_{i=1}^{n} \sum_{j=1}^{p} s_{ij}. \tag{2}$$

Contrary to $SS_\gamma$, $SD_\gamma$ is scaled by the number of pixels in the region, such that $SD_\gamma$ of regions containing different numbers of pixels can be compared with one another:

$$SD_\gamma = \text{Var}(\mathbf{Y}) = SS_\gamma / (n-1). \tag{3}$$

We note that for calculating the joint $SD_\gamma$ of adjacent regions, their $SS_\gamma$ statistics can be added

and divided by the total number of pixels minus one, but their region-level $SD_\gamma$ statistics cannot

be added directly.

One might be interested in determining the individual contribution of the $j$th spectral feature to

$SS_\gamma$. We call this the *feature contribution to spectral γ-diversity* or $FCSD_{\gamma,j}$ (Fig. 2), which can be

calculated from the sum of squares of the $j$th feature:

$$SS_{\gamma,j} = \sum_{i=1}^{n} s_{ij}. \tag{4}$$

157    Dividing $SS_{\gamma,j}$ by $(n-1)$ gives the variance of the $j$th feature, or $Var(\mathbf{y}_j)$. $FCSD_{\gamma,j}$ can then be

158    calculated as:

$$FCSD_{\gamma,j} = Var(\mathbf{y}_j)/Var(\mathbf{Y}) = SS_{\gamma,j}/SS_\gamma. \tag{5}$$

If the $p$ features are principal components from PCA scaling type 1, then the $FCSD_{\gamma,j}$ values

correspond to their relative eigenvalues. We note that $FCSD_{\gamma,j}$ cannot be mapped because the

contribution of each spectral feature applies to the region as a whole.

Likewise, one might wish to estimate the individual contribution of the $i$th pixel within the region

to $SD_\gamma$. We refer to this as the *local contribution to spectral γ-diversity*, or $LCSD_{\gamma,i}$, which is

calculated as:

$$LCSD_{\gamma,i} = SS_{\gamma,i}/SS_\gamma \tag{6}$$

165    where

$$SS_{\gamma,i} = \sum_{j=1}^{p} s_{ij}. \tag{7}$$

We note that $LCSD_{\gamma,i}$ indices are important visual elements in PCA ordination: each $LCSD_{\gamma,i}$

value corresponds to the squared distance from one pixel to the centroid in the $p$-dimensional

PCA ordination plot. In addition, the $LCSD_{\gamma,i}$ can be plotted on maps since one value is

associated with every pixel in the image. Doing so indicates which pixels are most spectrally

dissimilar from the mean pixel of the region in spectral feature space. We note that the $SS_{\gamma,i}$ and

$LCSD_{\gamma,i}$ indices are additive. The indices from adjacent pixels within an area of interest, for

example an individual tree, can be added up, such that their sums represent the local

contributions of the area of interest to $SS_\gamma$ and $SD_\gamma$. LCSD indices are also useful when

computed at the community scale (i.e. $LCSD_\beta$), because they then correspond to the ecological

concept of $\beta$-diversity; see "Spectral beta ($\beta$) diversity" below.

## Partitioning the total sum of squares

177　Partitioning the sum of squares forms the basis of a series of classic statistical approaches

178　based on least squares, such as the analysis of variance (ANOVA). From these methods, it is

179　well known that the total sum of squares of a matrix **Y** ($SS_{total}$) can be partitioned into additive

180　among-group ($SS_{among}$) and within-group ($SS_{within}$) components:

$$SS_{total} = SS_{among} + SS_{within}. \tag{8}$$

181　In linear regression analysis, we talk about the SS explained by the regression equation and the

182　residual variation. These two components sum to the total sum of squares.

183　Using the same indices as in the previous section, the ANOVA relationship can be expressed

184　as:

$$\sum_{i=1}^{n} \sum_{j=1}^{p} \left(y_{ij} - \bar{y}_j\right)^2 = \sum_{k=1}^{q} \sum_{j=1}^{p} m \left(\hat{y}_{kj} - \bar{y}_j\right)^2 + \sum_{i=1}^{m} \sum_{k=1}^{q} \sum_{j=1}^{p} \left(y_{ij} - \hat{y}_{kj}\right)^2 \tag{9}$$

185　where $q$ is the number of groups, $\hat{y}_{kj}$ is the arithmetic mean of the $j$th variable (column) for the

186　$k$th group:

$$\hat{y}_{kj} = \left(\sum_{i=1}^{m} y_{ijk}\right)/m \tag{10}$$

and where $m$ is the number of samples (rows) in each group $k$; an important assumption here is

that $m$ is equal in each group. The proof of this theorem can be found in standard statistics

textbooks and is therefore not shown here.

In the next two sections, we apply Equation 9 to partition the total sum of squares of a region

$SS_Y$ into additive among- ($\beta$) and within-group ($\alpha$) components from which spectral $\beta$- and $\alpha$-

diversity can be calculated directly.

**Spectral beta ($\beta$) diversity**

Let us divide **Y** into $q$ groups of $m$ spatially contiguous pixels, where each group corresponds to

a local community (e.g., a vegetation survey plot); $n = q\,m$. Here, we assume that each of these

communities corresponds to a square of equal area, which is $\sqrt{m}$ pixels wide; with this setup,

each community is represented by the same number of pixels. We will present later our

suggestion to use a rarefaction procedure to handle situations where $m$ differs among groups.

*Spectral $\beta$-diversity*, or $SD_\beta$, represents the degree to which the $q$ communities within a region

differ from each other in terms of spectral composition. We note that $SD_\beta$ is a *non-directional*

measure of *$\beta$-diversity sensu* Anderson et al. (2011). To calculate $SD_\beta$, we first compute the

squared deviations $s_{kj}$ of the $k$th community from the average pixel of the region in terms of

spectral reflectance, i.e. the column means of **Y** across all variables $j$:

$$s_{kj} = \left(\hat{y}_{kj} - \bar{y}_j\right)^2 \tag{11}$$

204     where $\hat{y}_{kj}$ is the arithmetic mean of the $k$th community (i.e. the community centroid) for the $j$th

205     spectral feature (Equation 10).

206     The sum of squares associated with each community $k$ ($SS_{\beta,k}$) is:

$$SS_{\beta,k} = \Sigma_{j=1}^{p} m\, s_{kj}. \tag{12}$$

207     The total sum of squares of the $\beta$ component ($SS_\beta$) is:

$$SS_\beta = \sum_{k=1}^{q} SS_{\beta,k} \tag{13}$$

208     from which $SD_\beta$ is calculated as:

$$SD_\beta = SS_\beta/(n-1). \tag{14}$$

209     The contribution of each community $k$ to $SD_\beta$, which we call the *local contribution to spectral β-*

210     *diversity* ($LCSD_{\beta,k}$), can be computed by the following ratio of sum of squares:

$$LCSD_{\beta,k} = SS_{\beta,k}/SS_\beta. \tag{15}$$

211     Finally, one can compute the *feature contribution to spectral β-diversity* or $FCSD_{\beta,j}$ of the $j$th

212     spectral feature as:

$$FCSD_{\beta,j} = SS_{\beta,j}/SS_\beta \tag{16}$$

213     where

$$SS_{\beta,j} = \Sigma_{k=1}^{q} m\, s_{kj}. \tag{17}$$

214     We note here that $LCSD_{\beta,k}$ can be mapped because each community $k$ has its own $LCSD_\beta$

215     value. On the other hand, $FCSD_{\beta,j}$ or $SD_\beta$ cannot be mapped because they refer to the region as

216     a whole.

## Spectral alpha ($\alpha$) diversity

Spectral $\alpha$-diversity, or $SD_\alpha$, is the degree to which neighbouring pixels *within* a local community

differ spectrally from each other. Contrary to $SD_\beta$ and $SD_\gamma$, which apply to the entire region, $SD_\alpha$

is defined at the community level. Therefore, we denote $SD_\alpha$ by the index $k$, $SD_{\alpha,k}$, since it is

measured for each community $k$. To calculate $SD_{\alpha,k}$, we first compute for every pixel and

spectral feature per community $y_{ijk}$ the squared deviations $s_{ijk}$ from the mean pixel spectrum of

the $k$th community for each spectral feature or column of **Y**:

$$s_{ijk} = \left(y_{ijk} - \hat{y}_{kj}\right)^2.$$  (18)

224    The sum of squares associated with the $j$th spectral feature of community $k$ is:

$$SS_{\alpha,jk} = \sum_{i=1}^{m} s_{ijk},$$  (19)

225    and the total sum of squares for community $k$ is:

$$SS_{\alpha,k} = \sum_{j=1}^{p} \sum_{i=1}^{m} s_{ijk}.$$  (20)

226    $SD_{\alpha,k}$ is obtained by dividing $SS_{\alpha,k}$ by $(m-1)$, where $m$ is the number of pixels within one

227    community, to make it comparable with other communities differing in their numbers of pixels:

$$SD_{\alpha,k} = SS_{\alpha,k}/(m-1).$$  (21)

228    The total sum of squares of the $\alpha$-component for all $q$ communities within the entire region is:

$$SS_{\alpha} = \sum_{k=1}^{q} SS_{\alpha,k}.$$  (22)

229    Importantly, following Equations 8 and 9, $SS_{\alpha}$ and $SS_{\beta}$ are linked to $SS_{\gamma}$ by the relationship:

$$SS_{\gamma} = SS_{\beta} + SS_{\alpha}.$$  (23)

230    Therefore, $SS_{\alpha}$ and $SS_{\beta}$ can be used directly to determine the relative importance of the $\alpha$ and $\beta$

231    components to spectral $\gamma$-diversity.

232    Finally, the *contribution of the $j$th* feature *to the spectral $\alpha$-diversit*y of the $k$th community, which

233    we call FCSD$_{\alpha,jk}$, can be computed as:

$$FCSD_{\alpha,jk} = SS_{\alpha,jk}/SS_{\alpha,k}.$$  (24)

234    These FCSD$_{\alpha,jk}$ values can be mapped and give us useful information about the origin of

235    spectral $\alpha$-diversity across different communities.

## CASE STUDY 1: SIMULATED REGIONS

To illustrate our approach, we first use leaf spectra data to simulate imagery (Appendix S2).

This removes much of the complexity associated with real imagery, where one has to deal with

much higher numbers of pixels, varying illumination and sensor viewing geometry, and

presence of shaded and non-vegetated pixels. We simulated two regions with equal spectral $\gamma$-

diversity, but contrasting spectral $\beta$- and $\alpha$-diversities (Fig. 3). Each region is composed of 25 ×

25 pixels, populated with leaf-level spectra of three temperate tree species (i.e. *Populus*

*deltoides* W. Bartram ex Marshall subsp. *deltoides* Marsh, *P. tremuloides* Michaux, and *Betula*

*alleghaniensis* Britton) measured in the field on 15 individual plants (Fig. S1). These 25 × 25

pixels regions are equally split into 25 communities, each composed of 5 × 5 pixels.

For both scenarios, we calculated the SS across the entire region ($SS_\gamma$), partitioned $SS_\gamma$ into its

$\beta$ and $\alpha$ components, and calculated spectral $\gamma$- $\beta$-, and $\alpha$-diversity (Fig. 4a). As spectral

features (columns of **Y**) we used the first three PCs (using type-I scaling in PCA), which

together explained >97% of the total variation in spectral reflectance. As expected, spectral $\gamma$-

diversity was equal for both scenarios (Table 1), whether expressed as the total sum of squares

($SS_\gamma = 1.66$), or standardized by $n-1$ pixels ($SD_\gamma = 0.0027$). In addition, in the high $\beta$-diversity

scenario, spectral variation among communities ($SS_\beta$, ~84%) largely exceeded spectral

variation within communities ($SS_\alpha$, ~16%), whereas in the low $\beta$-diversity scenario $SS_\beta$ was

much lower (~5%) than $SS_\alpha$ (~95%) (Table 1).

Next, we determined the local contributions of individual communities to spectral $\beta$-diversity

($LCSD_\beta$). In the high $\beta$-diversity scenario (Fig. 4b, left panel), communities 12 and 21 (numbered

as in Fig. 3) contributed the most to spectral $\beta$-diversity. These were the only two plots (out of

25) containing spectra of *Populus tremuloides*. In other words, these two plots had the most

distinctive spectral composition compared to other communities. By contrast, in the low $\beta$-

diversity scenario, community 16 was the most spectrally distinct community, something that

could not be easily detected by examining this scenario visually (Fig. 3, right panel). As

illustrated here, it is important to note that a region with low $SD_\beta$ can still have individual

communities showing high $LCSD_\beta$ values, because $LCSD_\beta$ values are *proportions* of $SD_\beta$.

We then estimated the contributions of individual spectral features to spectral diversity (FCSD)

for each scenario. For spectral $\gamma$-diversity (total variance of the region), $FCSD_\gamma$ declined

progressively from the first to the third PC (Table 1). As mentioned previously, $FCSD_\gamma$ values are

equal to the relative PCA eigenvalues of the spectral feature. Likewise, for spectral $\beta$-diversity,

the contribution from the first to subsequent PCs decreased in both scenarios (Table 1). The

relative contributions of individual spectral features to $\beta$-diversity were fairly similar in both

regions, even though they differed considerably in spectral $\beta$-diversity. For spectral $\alpha$-diversity,

however, $FCSD_\alpha$ values differed noticeably among the two scenarios (Fig. 4c–d). In the low $\alpha$-

diversity scenario (Fig. 4c-d, left column), PC 2 contributed more strongly to the spectral $\alpha$-

diversity of most communities than PC 1, whereas the opposite was true for the high $\alpha$-diversity

scenario (Fig. 4c–d, right column). The $FCSD_\alpha$ values were not expected to decrease in a

monotonic way since $\alpha$-diversity is orthogonal to $\gamma$-diversity and the PCs are those of $\gamma$-, not of

$\alpha$-diversity.

We note that SS, SD and LCSD indices are exactly the same whether using the original spectral

bands or all PCs, because PCA type-I scaling preserves the Euclidean distance among objects

(e.g., image pixels in spectral space). Conversely, the equations developed in this paper hold

and can be used directly with the original band data. However, FCSD values would change

when using the original spectral bands instead of PCs, since this would then indicate the relative

contributions of individual spectral bands (instead of PCs) to spectral diversity.

## CASE STUDY 2: NEON IMAGERY

Next, we applied our method for partitioning spectral diversity to imaging spectroscopy data collected by NEON's Airborne Observation Platform (AOP; Kampe *et al.* 2010) over the Bartlett Experimental Forest (https://www.neonscience.org/field-sites/field-sites-map/BART). In this case study, we used a scene measuring 280 m (east-west) x 1000 m (north-south), acquired in August 2017. Spectral data were processed to surface reflectance and subsampled to 1-m pixel size by NEON. Our workflow is illustrated in Figure 2.

For spectral diversity calculations we selected a community (i.e. plot) size of 40 m × 40 m, which is the base plot size used by NEON. We used rarefaction to standardize the number of pixels per community used for analysis. We used a normalized difference vegetation index (NDVI) threshold of $\geq$0.8 to identify the minimum number of vegetated pixels across all plots in the image (termed $m_{min}$), which was 1474 (= 92% of the 1600 pixels per community). We randomly selected $m_{min}$ pixels per plot, and applied our spectral diversity partitioning approach to all selected pixels. The rarefaction was repeated 30 times and results were averaged across all 30 repeats (Fig. 2). Alternatively, one could take the median value instead of the mean if distributions are skewed.

Our analyses revealed that spectral $\alpha$-diversity in this forested landscape accounted for 77% of the spectral $\gamma$-diversity, whereas $\beta$-diversity accounted for the remaining 23% (Table 1). In other words, there is considerably more spectral diversity within individual 40 m × 40 m communities than among communities in this forest. Figure 5 illustrates how spectral diversity is spatially structured. Two areas contribute strongly to spectral $\beta$-diversity (LCSD$_\beta$, darker colours in Fig. 5). The tree communities in these areas are more spectrally dissimilar from the average community than communities with lower LCSD$_\beta$ (lighter colours in Fig. 5).

For completeness, we evaluated the effects of shadows and community size on spectral

diversity calculations (Appendix S3). We found that removing shadows had little influence on

spectral diversity (Figs. S4–S6) and that results remained remarkably stable for plots ranging

from 20 m × 20 m (400 m$^2$) to 140 m x 140 m (almost 2 ha) in size (Figs. S7–S9).

**DISCUSSION**

In this paper, we proposed a new method for partitioning plant spectral $\gamma$-diversity (i.e. the

spectral diversity of a region) into additive $\alpha$- (within community) and $\beta$-diversity (among

community) components. Our approach builds on a method for partitioning $\beta$-diversity initially

designed for community data (Legendre & Cáceres 2013), adapts it to spectral data and,

importantly, extends it to include $\alpha, \beta$ and $\gamma$ components. Partitioning spectral diversity can bring

new insights and generate new hypotheses about the origins and maintenance of plant spectral

diversity across regions. For instance, high spectral $\beta$-diversity could result from turnover in

plant species and/or functional trait composition across environmental gradients (e.g., soil

properties, hydrology), whereas high spectral $\alpha$-diversity might result from local biotic

interactions among co-occurring plants (e.g., resource partitioning, conspecific negative density

dependence). Mapping spectral indices such as LCSD$_\beta$ and SD$_\alpha$ could be used as a biodiversity

"discovery tool" to design targeted field sampling campaigns to test such hypotheses, e.g., by

comparing community composition and diversity in areas with high and low LCSD$_\beta$ and SD$_\alpha$

values, respectively (Fig. 5).

Partitioning spectral diversity allows the determination of the spectral features contributing most

strongly to spectral $\alpha$-, $\beta$- or $\gamma$-diversity (FCSD), which helps in understanding the underlying

biological traits driving spectral diversity at different spatial scales. In our case studies, the

spectral features were principal components (PCs), which are linear combinations of the original

wavelength bands. As such, the individual contributions of all wavelength bands to each

spectral feature can be retrieved. The bands, in turn, can be linked to specific plant properties,

since the biophysical and biological causes of spectral variation across spectral regions and for

specific absorption features of molecules are reasonably well understood (Gates *et al.* 1965;

Curran 1989; Asner 1998; Kokaly *et al.* 2009; Ustin *et al.* 2009). Identifying the traits contributing

most strongly to spectral $\alpha$-diversity might inform us about how co-occurring species are

partitioning resources at the local scale, whereas identifying the traits contributing most strongly

to spectral $\beta$-diversity might reveal important mechanisms driving changes in community

composition across environmental gradients.

Partitioning plant spectral diversity and mapping its spatial components has applications in

biodiversity management. Indeed, managers often need to estimate the ecological value of

different ecosystems over large regions, for example to prioritize conservation or restoration

efforts. However, access to field data might be limited. Using imaging spectroscopy data, our

approach of partitioning spectral diversity allows the identification of areas with high spectral $\alpha$-

diversity, which likely coincide with local "hotspots" of taxonomic and/or functional trait diversity.

Further, high LCSD$_\beta$ values indicates areas with rare spectral composition, i.e., containing

communities that are most spectrally dissimilar from the average community within the region of

interest. Given that species spectral dissimilarity is linked to their functional and phylogenetic

dissimilarity (Schweiger *et al.* 2018), spectrally rare communities can be expected to have rare

taxonomic and/or functional composition, either because they harbor uncommon species, or

rare combinations of common species.

Our approach measures spectral variance directly (Fig. 2), which is in contrast to other studies

that have prior to deriving biodiversity metrics first translated remotely-sensed spectra into plant

species (e.g., Féret & Asner 2013), "spectral species" (Féret & Asner 2014), or plant functional

traits (e.g., Dahlin *et al.* 2013; Schneider *et al.* 2017). While spectral diversity does not isolate

any particular facet of plant biodiversity (e.g., taxonomic, chemical, structural), it integrates all of

these facets (Schweiger *et al.* 2018; Appendix S1). From a practical perspective, casting

spectral diversity as spectral variance depends on fewer user decisions compared to other

approaches (e.g., selecting the number of clusters for classifying spectral species, selecting the

plant traits and modelling approach to predict traits from spectra). This makes spectral diversity

easily comparable across different regions. Therefore, maps of $SD_\alpha$ and $LCSD_\beta$ could be ideal

candidates for biodiversity products from remotely sensed spectral imagery.

**Comparison with other approaches**

Much of the interest in measuring spectral diversity from remote sensing data stems from the

spectral variation hypothesis (Palmer *et al.* 2002), stating that the spatial variation in spectral

reflectance expresses overall variation of the environment. As areas of high environmental

variation often harbour more species than areas with low environmental variation, spectral

variation across space can potentially uncover botanically interesting areas (Palmer *et al.* 2002).

However, spectral diversity has been predominantly used to investigate relationships between

plant spectra and taxonomic units at the $\alpha$- and $\gamma$-diversity scale, whereas the $\beta$ component has

received less attention (Rocchini *et al.* 2018).

Historically, Landsat satellites were instrumental for spurring large-scale biodiversity studies.

Early sensors contained few spectral bands; thus, a large body of literature deals with using

NDVI for predicting and mapping taxonomic diversity (Gould 2000; see review by Pettorelli *et al.*

2005). Recent advances in sensor technology, particularly increased spectral resolution, have

led to a variety of approaches to calculate spectral $\alpha$-diversity (Rocchini *et al.* 2010). This

includes metrics such as the standard deviation or coefficient of variation of spectral indices

(Oindo & Skidmore 2002), or spectral bands among pixels (Hall *et al.* 2010; Gholizadeh *et al.*

2018; Wang *et al.* 2018a), the convex hull volume of pixels in spectral feature space (Dahlin

2016), the mean distance of pixels from the spectral centroid (Rocchini et al. 2010), the number

of spectrally distinct clusters or spectral species in ordination space (Féret & Asner 2014), and

diversity metrics based on dissimilarity matrices among species spectra or image pixels

(Schweiger *et al.* 2018). Of these, our method is most similar to the mean distance to the

spectral centroid (Rocchini *et al.* 2010). The difference is that we square the individual distances

to the spectral centroid; doing so allows us to partition sums of squares into additive

components (Equation 9).

Fewer studies have considered spectral *β*-diversity (Rocchini *et al.* 2018). One approach for

studying *β*-diversity using spectra has been to combine ordination scores of species inventories

with spectral data in multivariate models to predict the positions of pixels with unknown species

composition in species-ordination-space (Schmidtlein *et al.* 2007). This method and some of its

variants (Rocchini *et al.* 2018) do not measure spectral *β*-diversity *per se*, but instead use

spectra to estimate changes in community composition across the landscape. Rao's quadratic

entropy has been suggested as a measure of spectral *β*-diversity, based on the dissimilarity

among image pixels within a moving window (Rocchini *et al.* 2018). However, a moving window

approach expresses spectral *β*-diversity for many small sub-regions independently from one

another and does not estimate the spectral *β*-diversity of the region as a whole. Another

approach for studying spectral *β*-diversity has been to measure the pairwise dissimilarity in the

composition of spectral species among mapping units, and to re-project those pairwise

dissimilarities onto an RGB colour space (Féret & Asner 2014). This method yields a useful map

showing changes in spectral composition across the region, similar to our mapping of the first

three PCs in Figure 5, but it does not calculate spectral diversity.

**Methodological considerations**

A number of methodological aspects should be considered before applying our approach to

imaging spectroscopy data. These include: (1) the choice of a brightness normalization

procedure, (2) whether all or a subset of the wavelength bands, or spectral features, should be

used, (3) masking non-vegetated pixels, or not, (4) determining community size, and (5)

deciding on the scaling type (i.e. type I or II; Legendre & Legendre 2012) if using PCA as a

spectral feature extraction method. We discuss these methodological points in detail in

Appendix S4.

**CONCLUSION**

Plant spectral diversity is emerging as an integrator of chemical, structural, and taxonomic

aspects of plant biodiversity, which can be remotely sensed (Cavender-Bares *et al.* 2017).

Partitioning plant spectral diversity using our approach can help us to better understand and

generate new hypotheses about the origins of, and the processes that drive, biodiversity

variation across regions. Given the rapid and broad extent of current environmental changes,

remote sensing of plant biodiversity over large regions is more important than ever (Turner

2014; Wang & Gamon 2019). Our approach can identify local $\alpha$-diversity hotspots as well as

unique areas contributing strongly to $\beta$-diversity – two central facets of biodiversity.

Our approach is timely since current technological developments in high-resolution UAV

imaging spectroscopy will make this technology more accessible to ecologists in the coming

years (Aasen *et al.* 2018; Arroyo-Mora *et al.* 2019). For example, the Canadian Airborne

Biodiversity Observatory (www.caboscience.org) is developing UAV spectroscopy to understand

how plant biodiversity is responding to major environmental changes across Canada. We

anticipate that a growing number of ecologists will embrace this transformative technology for

mapping plant biodiversity. In fact, a wealth of moderate-resolution imaging spectroscopy data

are already freely available for a wide range of ecosystems across the United States as part of

the NEON program (www.neonscience.org). Partitioning spectral diversity could become a

useful tool for remotely sensing plant biodiversity from these new data sources.

## ACKNOWLEDGEMENTS

EL was supported by a Discovery Grant from the Natural Sciences and Engineering Research Council of Canada (NSERC; Grant numbers RGPIN-2014-06106, RGPIN-2019-04537). PL was supported by NSERC Discovery Grant number 7738. Teaching relief to EL that facilitated the writing of this manuscript was provided by the Canada Research Chair (CRC) in Plant Functional Biodiversity, and by Université de Montréal (UdeM) in support of a Discovery Frontiers grant from NSERC (Grant number 509190-2017) to EL to establish the Canadian Airborne Biodiversity Observatory (CABO). AKS was supported by a postdoctoral fellowship from CABO. EL wishes to thank colleagues from CABO and graduate students from the Laboratory of Plant Functional Ecology (LEFO) at UdeM for stimulating discussions about remote sensing of plant biodiversity, which helped to shape some of the ideas presented here.

## REFERENCES

Aasen, H., Honkavaara, E., Lucieer, A., Zarco-Tejada, P., Aasen, H., Honkavaara, E., *et al.* (2018). Quantitative remote sensing at ultra-high resolution with UAV spectroscopy: a review of sensor technology, measurement procedures, and data correction workflows. *Remote Sens.*, 10, 1091.

Anderson, M.J., Crist, T.O., Chase, J.M., Vellend, M., Inouye, B.D., Freestone, A.L., *et al.* (2011). Navigating the multiple meanings of β diversity: a roadmap for the practicing ecologist. *Ecol. Lett.*, 14, 19–28.

Arroyo-Mora, J.P., Kalacska, M., Inamdar, D., Soffer, R., Lucanus, O., Gorman, J., *et al.* (2019). Implementation of a UAV–hyperspectral pushbroom imager for ecological monitoring. *Drones*, 3, 12.

Asner, G.P. (1998). Biophysical and biochemical sources of variability in canopy reflectance. *Remote Sens. Environ.*, 64, 234–253.

Asner, G.P. & Martin, R.E. (2009). Airborne spectranomics: mapping canopy chemical and taxonomic diversity in tropical forests. *Front. Ecol. Environ.*, 7, 269–276.

Asner, G.P., Martin, R.E., Knapp, D.E., Tupayachi, R., Anderson, C.B., Sinca, F., *et al.* (2017). Airborne laser-guided imaging spectroscopy to map forest trait diversity and guide conservation. *Science*, 355, 385–389.

Bannari, A., Morin, D., Bonn, F. & Huete, A.R. (1995). A review of vegetation indices. *Remote Sens. Rev.*, 13, 95–120.

Bush, A., Sollmann, R., Wilting, A., Bohmann, K., Cole, B., Balzter, H., *et al.* (2017). Connecting Earth observation to high-throughput biodiversity data. *Nat. Ecol. Evol.*, 1, s41559-017-0176–017.

Cavender-Bares, J., Gamon, J.A., Hobbie, S.E., Madritch, M.D., Meireles, J.E., Schweiger, A.K.,

*et al.* (2017). Harnessing plant spectra to integrate the biodiversity sciences across biological

and spatial scales. *Am. J. Bot.*, 104, 966–969.

Cavender-Bares, J., Meireles, J.E., Couture, J.J., Kaproth, M.A., Kingdon, C.C., Singh, A., *et al.*

(2016). Associations of leaf spectra with genetic and phylogenetic variation in oaks: prospects

for remote detection of biodiversity. *Remote Sens.*, 8, 221.

Curran, P.J. (1989). Remote sensing of foliar chemistry. *Remote Sens. Environ.*, 30, 271–278.

Dahlin, K.M. (2016). Spectral diversity area relationships for assessing biodiversity in a

wildland–agriculture matrix. *Ecol. Appl.*, 26, 2758–2768.

Dahlin, K.M., Asner, G.P. & Field, C.B. (2013). Environmental and community controls on plant

canopy chemistry in a Mediterranean-type ecosystem. *Proc. Natl. Acad. Sci.*, 110, 6895–6900.

Draper, F.C., Baraloto, C., Brodrick, P.G., Phillips, O.L., Martinez, R.V., Coronado, E.N.H., *et al.*

(2019). Imaging spectroscopy predicts variable distance decay across contrasting Amazonian

tree communities. *J. Ecol.*, 107, 696–710.

Feilhauer, H., Asner, G.P., Martin, R.E. & Schmidtlein, S. (2010). Brightness-normalized partial

least squares regression for hyperspectral data. *J. Quant. Spectrosc. Radiat. Transf.*, 111,

1947–1957.

Féret, J.B. & Asner, G.P. (2013). Tree species discrimination in tropical forests using airborne

imaging spectroscopy. *IEEE Trans. Geosci. Remote Sens.*, 51, 73–84.

Féret, J.-B. & Asner, G.P. (2014). Mapping tropical forest canopy diversity using high-fidelity

imaging spectroscopy. *Ecol. Appl.*, 24, 1289–1296.

Gates, D.M., Keegan, H.J., Schleter, J.C. & Weidner, V.R. (1965). Spectral properties of plants.

*Appl. Opt.*, 4, 11–20.

Gholizadeh, H., Gamon, J.A., Zygielbaum, A.I., Wang, R., Schweiger, A.K. & Cavender-Bares,

J. (2018). Remote sensing of biodiversity: Soil correction and data dimension reduction methods

improve assessment of α-diversity (species richness) in prairie ecosystems. *Remote Sens.*

*Environ.*, 206, 240–253.

Gould, W. (2000). Remote sensing of vegetation, plant species richness, and regional

biodiversity hotspots. *Ecol. Appl.*, 10, 1861–1870.

Hall, K., Johansson, L.J., Sykes, M.T., Reitalu, T., Larsson, K. & Prentice, H.C. (2010).

Inventorying management status and plant species richness in semi-natural grasslands using

high spatial resolution imagery. *Appl. Veg. Sci.*, 13, 221–233.

Kampe, T.U., Johnson, B.R., Kuester, M. & Keller, M. (2010). NEON: the first continental-scale

ecological observatory with airborne remote sensing of vegetation canopy biochemistry and

structure. *J. Appl. Remote Sens.*, 4, 043510-043510–24.

Kokaly, R.F., Asner, G.P., Ollinger, S.V., Martin, M.E. & Wessman, C.A. (2009). Characterizing

canopy biochemistry from imaging spectroscopy and its application to ecosystem studies.

*Remote Sens. Environ.*, 113, 78–91.

Laliberté, E. & Legendre, P. (2010). A distance-based framework for measuring functional

diversity from multiple traits. *Ecology*, 91, 299–305.

Legendre, P. & Cáceres, M.D. (2013). Beta diversity as the variance of community data:

dissimilarity coefficients and partitioning. *Ecol. Lett.*, 16, 951–963.

Legendre, P. & Legendre, L. (2012). *Numerical Ecology.* 3rd English edition. Elsevier Science,

Amsterdam, Netherlands.

McManus, K.M., Asner, G.P., Martin, R.E., Dexter, K.G., Kress, W.J. & Field, C.B. (2016).

Phylogenetic structure of foliar spectral traits in tropical forest canopies. *Remote Sens.*, 8, 196.

507 Mueller-Dombois, D. & Ellenberg, H. (1974). *Aims and Methods of Vegetation Ecology*. John

508 Wiley & Sons, Inc., New York, USA.

509 Oindo, B.O. & Skidmore, A.K. (2002). Interannual variability of NDVI and species richness in

510 Kenya. *Int. J. Remote Sens.*, 23, 285–298.

511 Ollinger, S.V. (2010). Sources of variability in canopy reflectance and the convergent properties

512 of plants. *New Phytol.*, 189, 375–394.

513 Palmer, M.W., Earls, P.G., Hoagland, B.W., White, P.S. & Wohlgemuth, T. (2002). Quantitative

514 tools for perfecting species lists. *Environmetrics*, 13, 121–137.

515 Pereira, H.M., Ferrier, S., Walters, M., Geller, G.N., Jongman, R.H.G., Scholes, R.J., *et al.*

516 (2013). Essential biodiversity variables. *Science*, 339, 277–278.

517 Pettorelli, N., Vik, J.O., Mysterud, A., Gaillard, J.-M., Tucker, C.J. & Stenseth, N.Chr. (2005).

518 Using the satellite-derived NDVI to assess ecological responses to environmental change.

519 *Trends Ecol. Evol.*, 20, 503–510.

520 Ricklefs, R.E. (2008). Disintegration of the ecological community. *Am. Nat.*, 172, 741–750.

521 Rocchini, D., Balkenhol, N., Carter, G.A., Foody, G.M., Gillespie, T.W., He, K.S., *et al.* (2010).

522 Remotely sensed spectral heterogeneity as a proxy of species diversity: Recent advances and

523 open challenges. *Ecol. Inform.*, 5, 318–329.

524 Rocchini, D., Luque, S., Pettorelli, N., Bastin, L., Doktor, D., Faedi, N., *et al.* (2018). Measuring

525 β-diversity by remote sensing: A challenge for biodiversity monitoring. *Methods Ecol. Evol.*, 9,

526 1787–1798.

527 Schmidtlein, S., Zimmermann, P., Schüpferling, R. & Weiß, C. (2007). Mapping the floristic

528 continuum: Ordination space position estimated from imaging spectroscopy. *J. Veg. Sci.*, 18,

529 131–140.

Schneider, F.D., Morsdorf, F., Schmid, B., Petchey, O.L., Hueni, A., Schimel, D.S., *et al.* (2017).

Mapping functional diversity from remotely sensed morphological and physiological forest traits.

*Nat. Commun.*, 8, 1441.

Schweiger, A.K., Cavender-Bares, J., Townsend, P.A., Hobbie, S.E., Madritch, M.D., Wang, R.,

*et al.* (2018). Plant spectral diversity integrates functional and phylogenetic components of

biodiversity and predicts ecosystem function. *Nat. Ecol. Evol.*, 2, 976–982.

Turner, W. (2014). Sensing biodiversity. *Science*, 346, 301–302.

Ustin, S.L. & Gamon, J.A. (2010). Remote sensing of plant functional types. *New Phytol.*, 186,

795–816.

Ustin, S.L., Gitelson, A.A., Jacquemoud, S., Schaepman, M., Asner, G.P., Gamon, J.A., *et al.*

(2009). Retrieval of foliar information about plant pigment systems from high resolution

spectroscopy. *Remote Sens. Environ.*, 113, 67–77.

Vellend, M. (2010). Conceptual synthesis in community ecology. *Q. Rev. Biol.*, 85, 183–206.

Wang, R. & Gamon, J.A. (2019). Remote sensing of terrestrial plant biodiversity. *Remote Sens.*

*Environ.*, 231, 111218.

Wang, R., Gamon, J.A., Cavender-Bares, J., Townsend, P.A. & Zygielbaum, A.I. (2018a). The

spatial sensitivity of the spectral diversity–biodiversity relationship: an experimental test in a

prairie grassland. *Ecol. Appl.*, 28, 541–556.

Wang, R., Gamon, J.A., Schweiger, A.K., Cavender-Bares, J., Townsend, P.A., Zygielbaum,

A.I., *et al.* (2018b). Influence of species richness, evenness, and composition on optical

diversity: A simulation study. *Remote Sens. Environ.*, 211, 218–228.

Whittaker, R.H. (1960). Vegetation of the Siskiyou Mountains, Oregon and California. *Ecol.*

*Monogr.*, 30, 279–338.

Whittaker, R.H. (1972). Evolution and measurement of species diversity. *Taxon*, 21, 213–251.

Woodcock, C.E. & Strahler, A.H. (1987). The factor of scale in remote sensing. *Remote Sens.*

*Environ.*, 21, 311–332.

# TABLES

557 **Table 1** Partitioning spectral diversity into additive components for the two simulated regions

558 (Figs. 3–4) and for the NEON imagery (Fig. 5).

| | | High $\beta$, low $\alpha$ | Low $\beta$, high $\alpha$ | NEON imagery |
|---|---|---|---|---|
| | |  |  |  |
| **Sums of squares (SS)** | | | | |
| $SS_\gamma$ | | 1.66 | 1.66 | 740.24 |
| $SS_\beta$ (% of $SS_\gamma$) | | 1.40 (84.4%) | 0.08 (4.8%) | 168.85 (22.8%) |
| $SS_\alpha$ (% of $SS_\gamma$) | | 0.26 (15.6%) | 1.58 (95.2%) | 571.39 (77.2%) |
| **Spectral diversity (SD)** | | | | |
| $SD_\gamma$ | | 0.0027 | 0.0027 | 0.0029 |
| $SD_\beta$ | | 0.0022 | 0.00013 | 0.00065 |
| $\overline{SD}_\alpha$ | | 0.00043 | 0.0026 | 0.0022 |
| **Feature contribution to SD (FCSD)** | | | | |
| $FCSD_\gamma$ | PC 1 | 0.800 | 0.800 | 0.690 |

| | | | | |
|---|---|---|---|---|
| | PC 2 | 0.157 | 0.157 | 0.154 |
| | PC 3 | 0.023 | 0.023 | 0.090 |
| $FCSD_\beta$ | PC 1 | 0.906 | 0.859 | 0.520 |
| | PC 2 | 0.079 | 0.108 | 0.329 |
| | PC 3 | 0.013 | 0.019 | 0.092 |
| $\overline{FCSD}_\alpha$ | PC 1 | 0.259 | 0.798 | 0.733 |
| | PC 2 | 0.513 | 0.158 | 0.108 |
| | PC 3 | 0.050 | 0.022 | 0.092 |

## FIGURE LEGENDS

**Figure 1** Partitioning plant spectral γ-diversity into additive *β* and *α* components. A region of interest is split into a number of communities of a specific size and shape (here, 20 m × 20 m squares, representing standard forest inventory plots). Spectral *γ*-diversity refers to the total spectral diversity in the entire region calculated from pixel-level reflectance. The *β* component corresponds to spectral diversity *among* communities, with similar colours sharing more similar spectral composition. The *α* component refers to spectral diversity *within* individual communities. The left-most panel is a true colour (red-green-blue, RGB) image of an area of Bartlett Experimental Forest; colours for the other panels were obtained using the reflectance of different wavelength bands (R = 779 nm, G = 639 nm, B = 2301 nm), followed by linear stretching.

**Figure 2** Overview of our proposed workflow for partitioning plant spectral diversity. In our NEON case study, spectral data pre-processing included removing atmospheric water absorption bands (wavelengths between 1340–1455 nm and 1790–1955 nm) and noisy regions of the spectrum (wavelengths <400 nm and >2400 nm), and applying a Savitzky-Golay filter (order = 3, size = 7) to every pixel in the image to remove high-frequency noise. We masked all pixels with normalized difference vegetation index (NDVI) values <0.8, and brightness-normalized all spectra (Feilhauer *et al.* 2010). Then, we performed a PCA with type I-scaling, and visually inspected the first 17 PCs which together accounted for >99% of the total spectral variation among all pixels (Fig. S3). Only the first five PCs showed meaningful biological spatial patterns and were retained for spectral diversity measurements; PCs 6–17 were excluded based on visual inspection as they expressed artefacts from image acquisition and processing (Fig. S3). For illustration purposes, in the diversity partitioning analysis (bottom panel) we show communities composed of only three pixels, whereas in fact we used a community size of 40 ×

40 pixels in our NEON case study. For abbreviations see text. *The shade mask is for illustrative

purposes and is not applied to the PCs shown in the middle panel.

**Figure 3** Two simulated landscapes of equal spectral $\gamma$-diversity, but with contrasting spectral $\beta$-

diversity and $\alpha$-diversity. **Left**: high spectral $\beta$-diversity but low $\alpha$-diversity. **Right**: low spectral $\beta$-

diversity but high $\alpha$-diversity. Each landscape is composed of 25 communities (numbered black

squares), each composed of 5 × 5 pixels (smaller coloured squares). The size of each pixel is

equivalent to the size of an individual plant and their colour corresponds to one of the 15 leaf

spectra (= 3 species × 5 individuals) shown in Figure S1. These colours were set by mapping

the scores of the first three principal components (PC) for each spectrum to a red-green-blue

(RGB) scale (PC 1 = green, PC 2 = red, PC 3 = blue). We generated the high spectral $\beta$-

diversity but low spectral $\alpha$-diversity scenario (left panel) by randomly assigning (with

replacement) pixels within each community with individual spectra from single species (Fig. S1,

bottom row). We selected species identity per community at random using the following

probabilities: 0.60 (*Betula alleghaniensis*, green hues), 0.35 (*Populus deltoides*, blue hues) and

0.05 (*Populus tremuloides*, red hues). In this scenario, spectral $\beta$-diversity was high and spectral

$\alpha$-diversity low because interspecific spectral variation (particularly between *Betula* and the two

*Populus* species) was higher than intraspecific spectral variation (Fig. S2). Next, to reduce

spectral $\beta$-diversity and increase $\alpha$-diversity while holding $\gamma$-diversity constant, we moved the

pixels of the left panel to randomly selected positions in the right panel.

**Figure 4** (a) Spectral $\alpha$-diversity (SD$_\alpha$), (b) local contribution to spectral $\beta$-diversity (LCSD$_\beta$), and

(c-d) feature contribution to spectral $\alpha$-diversity (FCSD$_\alpha$) for the first two spectral features (i.e.

first two principal components of the brightness-normalized reflectance data) of each community

in the two simulated regions. PC = principal component.

**Figure 5** Partitioning spectral diversity using imaging spectroscopy data acquired by the

National Ecological Observatory Network (NEON) over the Bartlett Experimental Forest site.

From left to right: (1) true colour (red-green-blue, RGB) image with 0.1 m ground resolution, (2)

false colour image with 1 m resolution based on the first three principal components (PCs) of

the spectral image cube (PC1 = red, PC2 = green, PC3 = blue), (3) local contribution to spectral

$\beta$-diversity (LCSD$_\beta$), and (4) spectral $\alpha$-diversity (SD$_\alpha$) of forest communities, each measuring 40

m × 40 m. For panels 3 and 4, light hues correspond to low, dark hues to high values of LCSD$_\beta$

coefficients and SD$_\alpha$ values, respectively.

**FIGURES**

**Figure 1**

**Figure 2**

**Figure 3**

**Figure 4**

627 **Figure 5**

RGB        PC 1,2,3        LCSD$_\beta$        SD$_\alpha$