Partitioning plant spectral diversity into alpha and beta components

Etienne Laliberté[1,2*], Anna K. Schweiger[1,2], Pierre Legendre[2]

[1]Institut de recherche en biologie végétale, Université de Montréal, 4101 Sherbrooke Est, Montréal, Québec H1X 2B2, Canada

[2]Département de sciences biologiques, Université de Montréal, C. P. 6128, succursale Centre-ville, Montréal, Québec H3C 3J7, Canada

*Corresponding author. Email: etienne.laliberte@umontreal.ca

Phone: +1 514 343 6132

Running title: Spectral alpha, beta, and gamma diversity

Keywords: remote sensing, imaging spectroscopy, spectral variation hypothesis, spectral heterogeneity, alpha diversity, beta diversity, gamma diversity, spectranomics

Type of article: Methods

Number of words: 152 (abstract), 5264 (main text, excluding abstract, acknowledgements references, table and figure legends).

Number of references: 49

Number of figures: 5

Number of tables: 1

Statement of authorship: EL conceived the ideas, performed the analyses, and wrote the manuscript. AKS and PL contributed to the ideas and to the writing of the manuscript.

Data statement: Should the manuscript be accepted, the data supporting the results will be archived in an appropriate public repository such as Dryad and the data DOI will be included at the end of the article.

ABSTRACT

Plant spectral diversity — how plants differentially interact with solar radiation — is a powerful

integrator of plant functional and phylogenetic diversity that can be remotely sensed. We

propose to measure spectral diversity as spectral variance, which allows the partitioning of the

spectral diversity of a region, called spectral *gamma* (γ) diversity, into additive alpha (α; within

communities) and beta (β; among communities) components. Our method calculates the

contributions of individual bands or spectral features to spectral γ-, β-, and α-diversity, as well

as the contributions of individual plant communities to spectral diversity. We present two case

studies illustrating how our approach can identify "hotspots" of spectral α-diversity within a

region, and discover spectrally unique areas that contribute strongly to β-diversity. Partitioning

spectral diversity and mapping its spatial components has many applications for conservation

since high local diversity and distinctiveness in composition are two key criteria used to

determine the ecological value of ecosystems.

INTRODUCTION

Major environmental changes, including land-use change, climate change, and invasive species are altering the Earth's biodiversity. The rapid rate and broad extent of those changes far exceed our capacity to monitor them via field-based sampling alone. This calls for the development of new remote sensing approaches that can provide rapid estimates of biodiversity over broad regions (Pereira *et al.* 2013; Turner 2014; Bush *et al.* 2017). For terrestrial plants, imaging spectroscopy is emerging as the most promising remote sensing method for estimating biodiversity (Féret & Asner 2014; Wang & Gamon 2019). This is because its high spectral resolution allows plant species to be discriminated from one another, while also enabling the determination of ecologically important foliar functional traits (Asner & Martin 2009; Ustin *et al.* 2009).

For every pixel of an aerial image, imaging spectroscopy measures reflected solar radiation in tens to hundreds of contiguous, narrow (~10 nm wide) wavelength bands, usually covering all or part of the optical range (400–2500 nm) of the electromagnetic spectrum. Leaf "spectral signatures" of plants provide unique expressions of how solar radiation interacts with photosynthetic pigments, water, proteins, as well as structural and chemical defense compounds, and thus represent the evolution of plant adaptations to different environmental conditions (Cavender-Bares *et al.* 2016; McManus *et al.* 2016). At the crown scale, these spectral signatures are further influenced by architectural traits due to scattering of photons within canopies (Asner 1998; Ollinger 2010). As a result, plant spectral diversity is emerging as a powerful integrator of plant functional, structural and phylogenetic diversity that can be remotely sensed (Cavender-Bares *et al.* 2017; Schweiger *et al.* 2018; see also Appendix S1 in Supporting Information).

One of the most influential conceptual developments in community ecology has been

Whittaker's (1960, 1972) suggestion to partition biodiversity across space into α, β, and γ

components. Originally, α diversity was defined as the species diversity *within* communities, and

β as the variation in species composition *among* communities; together, α and β diversity jointly

determined γ diversity, which is the total species diversity across an entire region of interest. In

this paper, we transpose this foundational ecological concept from species diversity to spectral

diversity (Fig. 1). Because of the tight link between spectral diversity and functional and

phylogenetic diversity, imaging spectroscopy can provide direct, spatially explicit estimates of

plant alpha (α; within community) diversity (Féret & Asner 2014; Wang *et al.* 2018b), and can

detect changes in plant community composition across landscapes (Draper *et al.* 2019). The

ability to generate wall-to-wall, high-resolution maps of canopy plant diversity across entire

regions brings tremendous benefits for biodiversity science and conservation (e.g., Asner *et al.*

2017); however, conceptual and methodological challenges remain, especially with regard to β-

diversity estimation (Rocchini *et al.* 2010, 2018).

Spectral diversity is sometimes called spectral heterogeneity or spectral variability (Rocchini *et*

al. 2010), and has been defined as spatial variation in spectral reflectance (Rocchini *et al.* 2010;

Ustin & Gamon 2010; Gholizadeh *et al.* 2018; Wang & Gamon 2019). Intuitively, spectral

diversity can be conceptualized as multivariate dispersion, for which there are various statistical

measures highlighting different aspects of spectral diversity. For example, Wang et al. (2018a)

used the average coefficient of variation (CV) of each band for a set of pixels, whereas Rocchini

et al. (2010) used the mean distance from the spectral centroid; we note that the latter has also

been proposed as a measure of functional diversity in multivariate trait space (Laliberté &

Legendre 2010). However, none of the currently used metrics allow the partitioning of spectral

diversity into its α- (within communities) and β- (among communities) components (Fig. 1).

Here we propose to use the *spectral variance* among image pixels in an image as a measure of

spectral diversity. Our approach builds on that of Legendre and De Cáceres (2013) for species

inventory data, adapting it to spectral data, and extends it to jointly consider α, β, and γ

diversity. Casting spectral diversity as spectral variance has a number of important benefits:

1. the classical partitioning of sums of squares allows us to partition spectral γ-diversity into

additive spectral α- and β-diversity components (Fig. 1), from which one can infer the

relative importance of local and regional processes regulating spectral diversity across a

region of interest;

2. it allows us to estimate the contributions of individual plots or communities to spectral β-

diversity, highlighting areas that are spectrally distinct within the broader region;

3. it allows us to calculate the contributions of individual bands or spectral features to

spectral γ-, β- or α-diversity (Fig. 2), providing information about the underlying biological

traits driving spectral diversity;

4. it is easily implemented in software packages in a computationally efficient way, which is

important when dealing with high-volume image data;

5. it provides a direct link to other statistical procedures based on least squares (e.g.,

MANOVA, multiple linear regression, canonical redundancy analysis, *K*-means

partitioning).

After describing the theory behind our spectral diversity partitioning approach, we illustrate it

using a simple simulation. We then apply our method to imaging spectroscopy data collected

over the Bartlett Experimental Forest by the National Ecological Observatory Network (NEON)

Airborne Observation Platform (AOP; Kampe *et al.* 2010). The R code and data for our analyses

are available online (https://github.com/elaliberte/specdiv).

PARTITIONING SPECTRAL DIVERSITY

Size and shape of spatial units

Partitioning spectral γ-diversity into its α and β components first requires defining the extent of

the region of interest (Fig. 1). Delineating the region of interest is relatively straightforward since

it corresponds to the spatial extent under study, e.g., the region over which imagery is acquired

or a subset thereof. Delineating the size and shape of communities across the region of interest,

however, is more difficult. What constitutes an ecological community has been the subject of

considerable debate (see review by Ricklefs 2008), and we adopt here the general definition of

a community as "a group of organisms representing multiple species living in a specified place

and time" (Vellend 2010). This definition implies that a community must be larger than the size

of an individual organism, but how much larger will depend on the objectives of the study. For

the purpose of this work, we focus on communities of canopy plants (e.g., trees), because these

are the organisms that can be seen in aerial images. We use "community" in the sense of

"sampling unit" in vegetation surveys, which can be defined as the area in which the species

composition of the vegetation type of interest is adequately represented (Mueller-Dombois &

Ellenberg 1974).

Setting the size of a community to the size of typical inventory plot for a given ecosystem type

facilitates interpretation as this is the sampling unit that field ecologists are familiar with. For

example, forest inventory plots often measure 20 m × 20 m (Fig. 1), which is large enough to

include several canopy trees. However, we recognize that setting fixed and regularly shaped

boundaries to delineate communities is artificial (Ricklefs 2008), and point out that community

size and shape can be changed in the analysis.

Spectral gamma (γ) diversity

Let $\mathbf{Y} = [y_{ij}]$ be a matrix containing the positions, along the p axes defining the *spectral space*

(column vectors $\mathbf{y}_1$, $\mathbf{y}_2$, … $\mathbf{y}_p$ of $\mathbf{Y}$), of the n pixels (row vectors $\mathbf{x}_1$, $\mathbf{x}_2$, … $\mathbf{x}_n$ of $\mathbf{Y}$) in a region of

interest (Fig. 2). We use indices i and j to denote rows (pixels) and columns (axes) of matrix $\mathbf{Y}$,

respectively. The p axes could be all or a subset of the original spectral bands, a set of

vegetation indices calculated from selected spectral bands (Bannari *et al.* 1995), or a set of p

uncorrelated spectral features extracted using dimensionality reduction methods such as

principal component analysis (PCA). Hereafter, we use PCA in this section and in our case

studies and point out the mathematical relationships between the principal components (PCs)

and spectral variation below. We use the general term *variation* for sums of squares (SS, an

abbreviation for "sum of the squared deviations from the mean"), and reserve the term *variance*

when talking about spectral diversity (SD).

We refer to the total spectral diversity of the entire region as spectral γ-diversity (SD$_\gamma$). SD$_\gamma$ is

measured by the total variance of $\mathbf{Y}$, or Var($\mathbf{Y}$). This is done by first computing for every pixel

and spectral feature y_{ij} the squared deviations s_{ij} from the average pixel (across the whole

region) in terms of spectral reflectance, i.e. the column means of $\mathbf{Y}$:

$$s_{ij} = \left(y_{ij} - \overline{y}_j\right)^2. \tag{1}$$

147 The total sum of squares (SS) of matrix $\mathbf{Y}$ is calculated by summing all s_{ij}:

$$SS_\gamma = \sum_{i=1}^{n}\sum_{j=1}^{p} s_{ij}. \tag{2}$$

148 Contrary to SS$_\gamma$, SD$_\gamma$ is scaled by the number of pixels in the region, such that SD$_\gamma$ of regions

149 containing different numbers of pixels can be compared with one another:

$$SD_\gamma = \text{Var}(\mathbf{Y}) = SS_\gamma/(n-1). \tag{3}$$

We note that for calculating the joint SD_γ of adjacent regions, their SS_γ statistics can be added

and divided by the total number of pixels minus one, but their region-level SD_γ statistics cannot

be added directly.

One might be interested in determining the individual contribution of the jth spectral feature to

SS_γ. We call this the *feature contribution to spectral γ-diversity* or $FCSD_{\gamma,j}$, which can be

calculated from the sum of squares of the jth feature:

$$SS_{\gamma,j} = \sum_{i=1}^{n} s_{ij}. \tag{4}$$

156   Dividing $SS_{\gamma,j}$ by $(n-1)$ gives the variance of the jth feature, or $Var(\mathbf{y}_j)$. $FCSD_{\gamma,j}$ can then be

157   calculated as:

$$FCSD_{\gamma,j} = Var(\mathbf{y}_j)/Var(\mathbf{Y}) = SS_{\gamma,j}/SS_\gamma. \tag{5}$$

If the p features are principal components from PCA scaling type 1, then the $FCSD_{\gamma,j}$ values

correspond to their relative eigenvalues. We note that $FCSD_{\gamma,j}$ cannot be mapped because the

contribution of each spectral feature applies to the region as a whole.

Likewise, one might wish to estimate the individual contribution of the ith pixel within the region

to SD_γ. We refer to this as the *local contribution to spectral γ-diversity*, or $LCSD_{\gamma,i}$, which is

calculated as:

$$LCSD_{\gamma,i} = SS_{\gamma,i}/SS_\gamma \tag{6}$$

164   where

$$SS_{\gamma,i} = \sum_{j=1}^{p} s_{ij}. \tag{7}$$

We note that $LCSD_{\gamma,i}$ indices are important visual elements in PCA ordination: each $LCSD_{\gamma,i}$

value corresponds to the squared distance from one pixel to the centroid in the p-dimensional

PCA ordination plot. In addition, the $LCSD_{\gamma,i}$ could be plotted on maps since one value is

associated with every pixel in the image. Doing so indicates which pixels are most spectrally

distinct from the mean pixel of the region in spectral feature space. We note that the $SS_{Y,i}$ and

$LCSD_{Y,i}$ indices are additive. The indices from adjacent pixels within an area of interest, for

example an individual tree, can be added up, such that their sums represent the local

contributions of the area of interest to SS_Y and SD_Y. LCSD indices are also useful when

computed at the community scale (i.e. $LCSD_\beta$), because they then correspond to the ecological

concept of β-diversity; see "Spectral beta (β) diversity" below.

Partitioning the total sum of squares

Partitioning the sum of squares forms the basis of a series of classic statistical approaches

based on least squares, such as the analysis of variance (ANOVA). From these methods, it is

well known that the total sum of squares of a matrix **Y** (SS_{total}) can be partitioned into additive

among-group (SS_{among}) and within-group (SS_{within}) components:

$$SS_{total} = SS_{among} + SS_{within}. \tag{8}$$

180   In linear regression analysis, we talk about the SS explained by the regression equation and the

181   residual variation. These two components sum to the total sum of squares.

182   Using the same indices as in the previous section, the ANOVA relationship can be expressed

183   as:

$$\sum_{i=1}^{n}\sum_{j=1}^{p}\left(y_{ij}-\bar{y}_j\right)^2 = \sum_{k=1}^{q}\sum_{j=1}^{p} m\left(\hat{y}_{kj}-\bar{y}_j\right)^2 + \sum_{i=1}^{m}\sum_{k=1}^{q}\sum_{j=1}^{p}\left(y_{ij}-\hat{y}_{kj}\right)^2 \tag{9}$$

184   where q is the number of groups, $\hat{y}_{kj}$ is the arithmetic mean of the jth variable (column) for the

185   kth group:

$$\hat{y}_{kj} = \left(\sum_{i=1}^{m} y_{ijk}\right)/m \tag{10}$$

and where m is the number of samples (rows) in each group k; an important assumption here is

that m is equal in each group. The proof of this theorem can be found in standard statistics

textbooks and is therefore not shown here.

In the next two sections, we apply Equation 9 to partition the total sum of squares of a region

SS_γ into additive among- (β) and within-group (α) components from which spectral β- and α-

diversity can be calculated directly.

Spectral beta (β) diversity

Let us divide **Y** into q groups of m spatially contiguous pixels, where each group corresponds to

a local community (e.g., a vegetation survey plot); $n = q\,m$. Here, we assume that each of these

communities corresponds to a square of equal area that is $\sqrt{m}$ pixels wide; with this setup, each

community is represented by the same number of pixels. We will present later our suggestion to

handle the situation where m differs among groups.

Spectral β-diversity, or SD_β, represents the degree to which the q communities within a region

differ from each other in terms of spectral composition. We note that SD_β is a *non-directional*

measure of *β-diversity sensu* Anderson et al. (2011). To calculate SD_β, we first compute the

squared deviations s_{kj} of the kth community from the average pixel of the region in terms of

spectral reflectance, i.e. the column means of **Y** across all variables j:

$$s_{kj} = \left(\hat{y}_{kj} - \bar{y}_j\right)^2 \tag{11}$$

203   where $\hat{y}_{kj}$ is the arithmetic mean of the kth community (i.e. the community centroid) for the jth

204   spectral feature (Equation 10).

205   The sum of squares associated with each community k ($SS_{\beta,k}$) is:

$$SS_{\beta,k} = \sum_{j=1}^{p} m\, s_{kj}. \tag{12}$$

206 The total sum of squares of the β component (SS_β) is:

$$SS_\beta = \sum_{k=1}^{q} SS_{\beta,k} \tag{13}$$

207 from which SD_β is calculated as:

$$SD_\beta = SS_\beta/(n-1). \tag{14}$$

208 The contribution of each community *k* to SD_β, which we call the *local contribution to spectral β-*

209 *diversity* ($LCSD_{\beta,k}$), can be computed by the following ratio of sum of squares:

$$LCSD_{\beta,k} = SS_{\beta,k}/SS_\beta. \tag{15}$$

210 Finally, one can compute the *feature contribution to spectral β-diversity* or $FCSD_{\beta,j}$ of the *j*th

211 spectral feature as:

$$FCSD_{\beta,j} = SS_{\beta,j}/SS_\beta \tag{16}$$

212 where

$$SS_{\beta,j} = \sum_{k=1}^{q} m\, s_{kj}. \tag{17}$$

We note here that $LCSD_{\beta,k}$ can be mapped because each community *k* has its own $LCSD_\beta$

value. On the other hand, $FCSD_{\beta,j}$ or SD_β cannot be mapped because they refer to the region as

a whole.

Spectral alpha (α) diversity

Spectral α-diversity, or SD_α, is the degree to which neighbouring pixels *within* a local community

differ spectrally from each other. Contrary to SD_β and SD_γ, which apply to the entire region, SD_α

is defined at the community level. Therefore, we denote SD_α by the index *k*, $SD_{\alpha,k}$, since it is

measured for each community *k*. To calculate $SD_{\alpha,k}$, we first compute for every pixel and

spectral feature per community y_{ijk} the squared deviations s_{ijk} from the mean pixel spectrum of

the kth community for each spectral feature or column of **Y**:

$$s_{ijk} = \left(y_{ijk} - \hat{y}_{kj}\right)^2. \tag{18}$$

223   The sum of squares associated with the jth spectral feature of community k is:

$$SS_{\alpha,jk} = \sum_{i=1}^{m} s_{ijk}, \tag{19}$$

224   and the total sum of squares for community k is:

$$SS_{\alpha,k} = \sum_{j=1}^{p} \sum_{i=1}^{m} s_{ijk}. \tag{20}$$

225     $SD_{\alpha,k}$ is obtained by dividing $SS_{\alpha,k}$ by $(m-1)$, where m is the number of pixels within one

226   community, to make it comparable with other communities differing in their numbers of pixels:

$$SD_{\alpha,k} = SS_{\alpha,k}/(m-1). \tag{21}$$

227   The total sum of squares of the α-component for all q communities within the entire region is:

$$SS_{\alpha} = \sum_{k=1}^{q} SS_{\alpha,k}. \tag{22}$$

228   Importantly, following Equations 8 and 9, SS_α and SS_β are linked to SS_γ by the relationship:

$$SS_\gamma = SS_\beta + SS_\alpha. \tag{23}$$

229   Therefore, SS_α and SS_β can be used directly to determine the relative importance of the α and β

230   components to spectral γ-diversity.

231   Finally, the *contribution of the jth feature to the spectral α-diversity* of the kth community, which

232   we call $FCSD_{\alpha,jk}$, can be computed as:

$$FCSD_{\alpha,jk} = SS_{\alpha,jk}/SS_{\alpha,k}. \tag{24}$$

233   These $FCSD_{\alpha,jk}$ values can be mapped and give us useful information about the origin of

234   spectral α-diversity across different communities.

CASE STUDY 1: SIMULATED REGIONS

To illustrate our approach, we first use leaf-level spectral data to simulate imagery (Appendix

S2). This removes much of the complexity associated with real imagery, where one has to deal

with much higher numbers of pixels, varying illumination and sensor viewing geometry, and

presence of shaded and non-vegetated pixels. We simulated two regions with equal spectral γ-

diversity, but contrasting spectral β- and α-diversities (Fig. 3). Each region is composed of 25 ×

25 pixels, populated with leaf-level spectra of three temperate tree species measured in the field

on 15 individual plants (Fig. S1). These 25 × 25 pixels regions are equally split into 25

communities, each composed of 5 × 5 pixels.

For both scenarios, we calculated the SS across the entire region (SS_γ), partitioned SS_γ into its

β and α components, and calculated spectral γ- β-, and α-diversity (Fig. 4a). As spectral

features (columns of **Y**) we used the first three PCs (using type-I scaling in PCA), which

together explained >97% of the total variation in spectral reflectance. As expected, spectral γ-

diversity was equal for both scenarios (Table 1), whether expressed as the total sum of squares

(SS_γ = 1.66), or standardized by $n - 1$ pixels (SD_γ = 0.0027). In addition, in the high β-diversity

scenario, spectral variation among communities (SS_β, ~84%) largely exceeded spectral

variation within communities (SS_α, ~16%), whereas in the low β-diversity scenario SS_β was

much lower (~5%) than SS_α (~95%) (Table 1).

Next, we determined the local contributions of individual communities to spectral β-diversity

($LCSD_\beta$). In the high β-diversity scenario (Fig. 4b, left panel), communities 12 and 21 (numbered

as in Fig. 3) contributed the most to spectral β-diversity. These were the only two plots (out of

25) containing spectra of *Populus tremuloides*. In other words, these two plots had the most

distinctive spectral composition compared to other communities. By contrast, in the low β-

diversity scenario, community 16 was the most spectrally distinct community, something that

could not be easily detected from visual examination of this scenario (Fig. 3, right panel). The

ability to map $LCSD_\beta$ (Fig. 4b) is especially useful for spatially-continuous imagery data since it

allows the identification of areas with rare spectral composition, possibly indicating unique

taxonomic and/or functional composition. As illustrated here, it is important to note that a region

with low SD_β can still have individual communities showing high $LCSD_\beta$ values, because $LCSD_\beta$

values are *proportions* of SD_β.

We then estimated the contributions of individual spectral features to spectral diversity (FCSD)

for each scenario. For spectral γ-diversity (total variance), $FCSD_\gamma$ declined progressively from

the first to the third PC (Table 1). As mentioned previously, the $FCSD_\gamma$ values are equal to the

relative PCA eigenvalues of the spectral feature. Likewise, for spectral β-diversity, the

contribution from the first to subsequent PCs decreased in both scenarios (Table 1). The

relative contributions of individual spectral features (e.g., PCs of the wavelength bands) to β-

diversity were fairly similar in both regions, even though they differed considerably in spectral β-

diversity. For spectral α-diversity, however, $FCSD_\alpha$ values differed noticeably among the two

scenarios (Fig. 4c–d). In the low α-diversity scenario (Fig. 4c-d, left column), PC 2 contributed

more strongly to the spectral α-diversity of most communities than PC 1, whereas the opposite

was true for the high α-diversity scenario (Fig. 4c–d, right column). The $FCSD_\alpha$ values were not

expected to decrease in a monotonic way since α-diversity is orthogonal to γ-diversity and the

PCs are those of γ, not of α diversity. Mapping those $FCSD_\alpha$ values (Fig. 4c–d) can provide

insights into the biological origins of spectral diversity (Gholizadeh *et al.* 2018).

We note that SS, SD and LCSD indices are exactly the same whether using the original spectral

bands or all PCs, because PCA type-I scaling preserves the Euclidean distance among objects

(e.g., image pixels in spectral space). However, FCSD values would change when using the

original spectral bands instead of PCs, since this would then indicate the relative contributions

of individual spectral bands (instead of PCs) to spectral diversity.

CASE STUDY 2: NEON IMAGERY

We then applied our method for partitioning spectral diversity to imaging spectroscopy data

collected by NEON's Airborne Observation Platform (AOP; Kampe *et al.* 2010) over the Bartlett

Experimental Forest (https://www.neonscience.org/field-sites/field-sites-map/BART). In this case

study, we used a scene measuring 280 m (east-west) x 1000 m (north-south), acquired in

August 2017. Spectral data were processed to surface reflectance and subsampled to 1-m pixel

size by NEON. Our entire workflow is illustrated in Figure 2.

For spectral diversity calculations we selected a community (i.e. plot) size of 40 m × 40 m, which

is the base plot size used by NEON. We used rarefaction to standardize the number of pixels

per community used for analysis. We used a normalized difference vegetation index (NDVI)

threshold of ≥0.8 to identify the minimum number of vegetated pixels across all plots in the

image (termed m_{min}), which was 1474 (= 92% of the 1600 pixels per community). We randomly

selected m_{min} pixels per plot, and applied our spectral diversity partitioning approach to all

selected pixels. The rarefaction was repeated 30 times and results were averaged across all 30

repeats (Fig. 2). Alternatively, we note that one could take the median value instead of the mean

if distributions are skewed.

Our analyses revealed that spectral α-diversity in this forested landscape accounted for 77% of

the spectral γ-diversity, whereas β-diversity accounted for the remaining 23% (Table 1). In other

words, there is considerably more spectral diversity within individual 40 m × 40 m communities

than among communities in this forest. Figure 5 illustrates how spectral diversity is spatially

structured. Two areas contribute strongly to spectral β-diversity (LCSD$_\beta$, darker colours in Fig.

5). The tree communities in these areas are more spectrally dissimilar from the average

community than communities with lower LCSD$_\beta$ (lighter colours in Fig. 5).

For completeness, we evaluated the potential effects of shadows in the image, as well as

community size, on spectral diversity calculations (Appendix S3). We found that removing

shadows had little influence (Figs. S4–S6) and that results remained remarkably stable for plots

ranging from 20 m × 20 m (400 m^2) to 140 m x 140 m (almost 2 ha) in size (Figs. S7–S9).

DISCUSSION

In this paper, we proposed a new method for partitioning plant spectral γ-diversity (i.e. the

spectral diversity of a region) into additive α- (within community) and β-diversity (among

community) components. Our approach builds on a method for partitioning β-diversity initially

designed for community data (Legendre & Cáceres 2013), adapts it to spectral data and,

importantly, extends it to include α, β and γ components. Casting spectral diversity as spectral

variance allows the determination of spectral α-, β- and γ-diversity within a single cohesive

framework, and makes it possible to calculate spectral β-diversity (SD statistics) independently

(i.e. in a non-additive and non-multiplicative way) from α and γ (Legendre & Cáceres 2013),

although the sum-of-square statistics (SS) are additive. In addition, the column-wise partitioning

of variance in spectral data matrices allows the determination of which spectral bands or PC

features contribute most strongly to the different components of spectral diversity (FCSD), while

the row-wise partitioning of spectral diversity allows the identification of local spectral α-diversity

"hotspots" and of spectrally distinct communities that contribute strongly to spectral β-diversity

(LCSD).

Benefits of our approach

Partitioning spectral diversity can bring new insights and generate new hypotheses about the

origins and maintenance of plant spectral diversity across regions. As illustrated in our

simulated example, two regions with similar levels of regional spectral γ-diversity may have

contrasting levels of spectral α- and β-diversity, suggesting that different processes could be

regulating diversity. For instance, high spectral β-diversity could result from strong turnover in

plant species and/or functional trait composition across pronounced environmental gradients

(e.g., soil properties, hydrology), whereas high spectral α-diversity might result from local biotic

interactions among co-occurring plants (e.g., resource partitioning, conspecific negative density

dependence). Mapping spectral indices such as SD_α and $LCSD_\beta$ could be used as a biodiversity

"discovery tool" to design targeted field sampling campaigns to test such hypotheses (Fig. 5).

Partitioning spectral diversity allows the determination of the spectral features contributing most

strongly to spectral α-, β- or γ-diversity (FCSD), which helps in understanding the underlying

biological traits driving spectral diversity at different spatial scales. In our case studies, the

spectral features were principal components (PCs), which are linear combinations of the original

wavelength bands. As such, the individual contributions of all wavelength bands to each

spectral feature can be retrieved. The bands, in turn, can be linked to specific plant properties,

since the biophysical and biological causes of spectral variation across spectral regions and for

specific absorption features of molecules are reasonably well understood (Gates *et al.* 1965;

Curran 1989; Asner 1998; Kokaly *et al.* 2009; Ustin *et al.* 2009). Identifying the traits contributing

most strongly to spectral α-diversity might inform us about how co-occurring species are

partitioning resources at the local scale, whereas identifying the traits contributing most strongly

to spectral β-diversity might reveal important mechanisms driving changes in community

composition across environmental gradients.

Partitioning plant spectral diversity and mapping its spatial components has applications in

biodiversity management. Indeed, managers often need to estimate the ecological value of

different ecosystems over large regions, for example to prioritize conservation or restoration

efforts. However, access to field data might be limited. Using imaging spectroscopy data, our

approach of partitioning spectral diversity allows the identification of areas with high spectral α-

diversity, which likely coincide with local "hotspots" of taxonomic and/or functional trait diversity.

Further, high LCSD$_\beta$ values indicate areas with rare spectral composition, i.e., containing

communities that are most spectrally dissimilar from the average community within the region of

interest. Given that species spectral dissimilarity is linked to their functional and phylogenetic

dissimilarity (Schweiger *et al.* 2018), spectrally rare communities can be expected to have rare

taxonomic and/or functional composition, either because they harbor uncommon species,

desirable (i.e. rare native species) or not (e.g., invasive exotic species), or rare combinations of

common species.

Our approach measures spectral variance directly (Fig. 2), which is in contrast to other studies

that have prior to deriving biodiversity metrics first translated remotely-sensed spectra into plant

species (e.g., Féret & Asner 2013), "spectral species" (Féret & Asner 2014), or plant functional

traits (e.g., Dahlin *et al.* 2013; Schneider *et al.* 2017). While spectral diversity does not isolate

any particular facet of plant biodiversity (e.g., taxonomic, phylogenetic, functional, structural), it

acts as a powerful integrator of all of these facets (Schweiger *et al.* 2018; Appendix S1). From a

practical perspective, casting spectral diversity as spectral variance depends on fewer user

decisions compared to other approaches (e.g., selecting the number of clusters for classifying

spectral species, selecting the plant traits and modelling approach to predict traits from spectra).

This makes spectral diversity easily comparable across different regions. Therefore, maps of

SD$_\alpha$ and LCSD$_\beta$ could be ideal candidates for biodiversity remote sensing products from

imaging spectroscopy data.

Comparison with other approaches

Much of the interest in measuring spectral diversity from remote sensing data stems from the

spectral variation hypothesis (Palmer *et al.* 2002), stating that the spatial variation in spectral

reflectance expresses overall variation of the environment. As areas of high environmental

variation often harbour more species than areas with low environmental variation, spectral

variation across space can potentially uncover botanically interesting areas (Palmer *et al.* 2002).

However, spectral diversity has been predominantly used to investigate relationships between

plant spectra and taxonomic units at the *α*- and *γ*-diversity scale, whereas the *β* component has

received less attention (Rocchini *et al.* 2018).

Historically, Landsat satellites were instrumental for spurring large-scale biodiversity studies.

Early sensors contained few spectral bands; thus, a large body of literature deals with using

NDVI for predicting and mapping taxonomic diversity (Gould 2000; see review by Pettorelli *et al.*

2005). Recent technological advances in sensor technology, particularly increased spectral

resolution, have led to a variety of approaches to calculate spectral *α*-diversity (Rocchini *et al.*

2010). This includes metrics such as the standard deviation or coefficient of variation of spectral

indices (Oindo & Skidmore 2002), or spectral bands among pixels (Hall *et al.* 2010; Gholizadeh

et al. 2018; Wang *et al.* 2018a), the convex hull volume of pixels in spectral feature space

(Dahlin 2016), the mean distance of pixels from the spectral centroid (Rocchini et al. 2010), the

number of spectrally distinct clusters or spectral species in ordination space (Féret & Asner

2014), and diversity measures based on dissimilarity matrices among species spectra or image

pixels (Schweiger *et al.* 2018). Of these, our method is most similar to the mean distance to the

spectral centroid (Rocchini *et al.* 2010). The difference is that we square the individual distances

to the spectral centroid; doing so allows us to partition sums of squares into additive

components (Equation 9).

Fewer studies have considered spectral *β*-diversity (Rocchini *et al.* 2018). One approach for

studying *β*-diversity using spectra has been to combine ordination scores of species inventories

with spectral data in multivariate models to predict the positions of pixels with unknown species

composition in species-ordination-space (Schmidtlein *et al.* 2007). This method and some of its

variants (Rocchini *et al.* 2018) do not measure spectral *β*-diversity *per se*, but instead use

spectra to estimate changes in community composition across the landscape. Rao's quadratic

entropy has been suggested as a measure of spectral *β*-diversity, based on the dissimilarity

among image pixels within a moving window (Rocchini *et al.* 2018). However, a moving window

approach expresses spectral *β*-diversity for many small sub-regions independently from one

another and does not estimate the spectral *β*-diversity of the region as a whole. Another

approach for studying spectral *β*-diversity has been to measure the pairwise dissimilarity in the

composition of spectral species among mapping units, and to re-project those pairwise

dissimilarities onto an RGB colour space (Féret & Asner 2014). This method yields a useful map

showing changes in spectral composition across the region, similar to our mapping of the first

three PCs in Figure 5, but it does not calculate spectral diversity.

Methodological considerations

A number of methodological aspects should be considered before applying our approach to

imaging spectroscopy data. These include: (1) the choice of a brightness normalization

procedure, (2) whether all or a subset of the wavelength bands, or spectral features, should be

used, (3) masking non-vegetated pixels, or not, (4) determining community size, and (5)

deciding on the scaling type (i.e. type I or II; Legendre & Legendre 2012) if using PCA as a

spectral feature extraction method. We discuss these methodological points in detail in

Appendix S4.

CONCLUSION

Plant spectral diversity is emerging as a powerful integrator of taxonomic, phylogenetic,

functional and structural aspects of biodiversity, which can be remotely sensed (Cavender-

Bares *et al.* 2017). Partitioning plant spectral diversity using our approach can help us to better

understand and generate new hypotheses about the origins of, and the processes that drive,

biodiversity variation across regions. Given the rapid and broad extent of current environmental

changes, remote sensing of plant biodiversity over large regions is now possible and more

important than ever (Turner 2014; Wang & Gamon 2019). Our approach can identify local α-

diversity hotspots as well as unique areas contributing strongly to β-diversity – two central

aspects of biodiversity.

We believe that our approach is timely since rapid technological developments in UAV imaging

spectroscopy will make this technology much more accessible to ecologists in the coming years

(Aasen *et al.* 2018; Arroyo-Mora *et al.* 2019). For example, the Canadian Airborne Biodiversity

Observatory (www.caboscience.org) is currently developing UAV-based imaging spectroscopy

to map, understand, and forecast how terrestrial plant biodiversity is responding to major

environmental changes across Canada. We anticipate that a growing number of ecologists will

embrace this transformative technology for mapping plant biodiversity. In fact, a wealth of

moderate-resolution imaging spectroscopy data are already freely available for a wide range of

terrestrial ecosystems across the United States as part of the NEON program

(www.neonscience.org). Moreover, upcoming hyperspectral satellite missions such as HyspIRI

(https://hyspiri.jpl.nasa.gov) and EnMAP (http://www.enmap.org) will bring new possibilities for

monitoring changes in plant biodiversity at global scale (Turner 2014; Jetz *et al.* 2016).

Partitioning spectral diversity could become a useful tool for remotely sensing plant biodiversity

from these new data sources.

ACKNOWLEDGEMENTS

EL was supported by a Discovery Grant from the Natural Sciences and Engineering Research

Council of Canada (NSERC; Grant number RGPIN-2014-06106). PL was supported by NSERC

Discovery Grant number 7738. Teaching relief to EL that facilitated the writing of this manuscript

was provided by the Canada Research Chair (CRC) in Plant Functional Biodiversity, and by

Université de Montréal (UdeM) in support of a Discovery Frontiers grant from NSERC (Grant

number 509190-2017) to EL to establish the Canadian Airborne Biodiversity Observatory

(CABO). AKS was supported by a postdoctoral fellowship from CABO. EL wishes to thank

colleagues from CABO and graduate students from the Laboratory of Plant Functional Ecology

(LEFO) at UdeM for stimulating discussions about remote sensing of plant biodiversity, which

helped to shape some of the ideas presented here.

REFERENCES

Aasen, H., Honkavaara, E., Lucieer, A., Zarco-Tejada, P., Aasen, H., Honkavaara, E., *et al.*

(2018). Quantitative remote sensing at ultra-high resolution with UAV spectroscopy: a review of

sensor technology, measurement procedures, and data correction workflows. *Remote Sens.*,

10, 1091.

Anderson, M.J., Crist, T.O., Chase, J.M., Vellend, M., Inouye, B.D., Freestone, A.L., *et al.*

(2011). Navigating the multiple meanings of β diversity: a roadmap for the practicing ecologist.

Ecol. Lett., 14, 19–28.

Arroyo-Mora, J.P., Kalacska, M., Inamdar, D., Soffer, R., Lucanus, O., Gorman, J., *et al.* (2019).

Implementation of a UAV–hyperspectral pushbroom imager for ecological monitoring. *Drones*,

3, 12.

Asner, G.P. (1998). Biophysical and biochemical sources of variability in canopy reflectance.

Remote Sens. Environ., 64, 234–253.

Asner, G.P. & Martin, R.E. (2009). Airborne spectranomics: mapping canopy chemical and

taxonomic diversity in tropical forests. *Front. Ecol. Environ.*, 7, 269–276.

Asner, G.P., Martin, R.E., Knapp, D.E., Tupayachi, R., Anderson, C.B., Sinca, F., *et al.* (2017).

Airborne laser-guided imaging spectroscopy to map forest trait diversity and guide conservation.

Science, 355, 385–389.

Bannari, A., Morin, D., Bonn, F. & Huete, A.R. (1995). A review of vegetation indices. *Remote*

Sens. Rev., 13, 95–120.

Bush, A., Sollmann, R., Wilting, A., Bohmann, K., Cole, B., Balzter, H., *et al.* (2017). Connecting

Earth observation to high-throughput biodiversity data. *Nat. Ecol. Evol.*, 1, s41559-017-0176–

017.

Cavender-Bares, J., Gamon, J.A., Hobbie, S.E., Madritch, M.D., Meireles, J.E., Schweiger, A.K.,

et al. (2017). Harnessing plant spectra to integrate the biodiversity sciences across biological

and spatial scales. *Am. J. Bot.*, 104, 966–969.

Cavender-Bares, J., Meireles, J.E., Couture, J.J., Kaproth, M.A., Kingdon, C.C., Singh, A., *et al.*

(2016). Associations of leaf spectra with genetic and phylogenetic variation in oaks: prospects

for remote detection of biodiversity. *Remote Sens.*, 8, 221.

Curran, P.J. (1989). Remote sensing of foliar chemistry. *Remote Sens. Environ.*, 30, 271–278.

Dahlin, K.M. (2016). Spectral diversity area relationships for assessing biodiversity in a

wildland–agriculture matrix. *Ecol. Appl.*, 26, 2758–2768.

Dahlin, K.M., Asner, G.P. & Field, C.B. (2013). Environmental and community controls on plant

canopy chemistry in a Mediterranean-type ecosystem. *Proc. Natl. Acad. Sci.*, 110, 6895–6900.

Draper, F.C., Baraloto, C., Brodrick, P.G., Phillips, O.L., Martinez, R.V., Coronado, E.N.H., *et al.*

(2019). Imaging spectroscopy predicts variable distance decay across contrasting Amazonian

tree communities. *J. Ecol.*, 107, 696–710.

Feilhauer, H., Asner, G.P., Martin, R.E. & Schmidtlein, S. (2010). Brightness-normalized partial

least squares regression for hyperspectral data. *J. Quant. Spectrosc. Radiat. Transf.*, 111,

1947–1957.

Féret, J.B. & Asner, G.P. (2013). Tree species discrimination in tropical forests using airborne

imaging spectroscopy. *IEEE Trans. Geosci. Remote Sens.*, 51, 73–84.

Féret, J.-B. & Asner, G.P. (2014). Mapping tropical forest canopy diversity using high-fidelity

imaging spectroscopy. *Ecol. Appl.*, 24, 1289–1296.

Gates, D.M., Keegan, H.J., Schleter, J.C. & Weidner, V.R. (1965). Spectral properties of plants.

Appl. Opt., 4, 11–20.

Gholizadeh, H., Gamon, J.A., Zygielbaum, A.I., Wang, R., Schweiger, A.K. & Cavender-Bares,

J. (2018). Remote sensing of biodiversity: Soil correction and data dimension reduction methods

improve assessment of α-diversity (species richness) in prairie ecosystems. *Remote Sens.*

Environ., 206, 240–253.

Gould, W. (2000). Remote sensing of vegetation, plant species richness, and regional

biodiversity hotspots. *Ecol. Appl.,* 10, 1861–1870.

Hall, K., Johansson, L.J., Sykes, M.T., Reitalu, T., Larsson, K. & Prentice, H.C. (2010).

Inventorying management status and plant species richness in semi-natural grasslands using

high spatial resolution imagery. *Appl. Veg. Sci.,* 13, 221–233.

Jetz, W., Cavender-Bares, J., Pavlick, R., Schimel, D., Davis, F.W., Asner, G.P., *et al.* (2016).

Monitoring plant functional diversity from space. *Nat. Plants*, 2, 16024.

Kampe, T.U., Johnson, B.R., Kuester, M. & Keller, M. (2010). NEON: the first continental-scale

ecological observatory with airborne remote sensing of vegetation canopy biochemistry and

structure. *J. Appl. Remote Sens.,* 4, 043510-043510–24.

Kokaly, R.F., Asner, G.P., Ollinger, S.V., Martin, M.E. & Wessman, C.A. (2009). Characterizing

canopy biochemistry from imaging spectroscopy and its application to ecosystem studies.

Remote Sens. Environ., 113, 78–91.

Laliberté, E. & Legendre, P. (2010). A distance-based framework for measuring functional

diversity from multiple traits. *Ecology*, 91, 299–305.

Legendre, P. & Cáceres, M.D. (2013). Beta diversity as the variance of community data:

dissimilarity coefficients and partitioning. *Ecol. Lett.*, 16, 951–963.

Legendre, P. & Legendre, L. (2012). *Numerical Ecology*. 3rd English edition. Elsevier Science,

Amsterdam, Netherlands.

McManus, K.M., Asner, G.P., Martin, R.E., Dexter, K.G., Kress, W.J. & Field, C.B. (2016).

Phylogenetic structure of foliar spectral traits in tropical forest canopies. *Remote Sens.*, 8, 196.

Mueller-Dombois, D. & Ellenberg, H. (1974). *Aims and Methods of Vegetation Ecology*. John

Wiley & Sons, Inc., New York, USA.

Oindo, B.O. & Skidmore, A.K. (2002). Interannual variability of NDVI and species richness in

Kenya. *Int. J. Remote Sens.*, 23, 285–298.

Ollinger, S.V. (2010). Sources of variability in canopy reflectance and the convergent properties

of plants. *New Phytol.*, 189, 375–394.

Palmer, M.W., Earls, P.G., Hoagland, B.W., White, P.S. & Wohlgemuth, T. (2002). Quantitative

tools for perfecting species lists. *Environmetrics*, 13, 121–137.

Pereira, H.M., Ferrier, S., Walters, M., Geller, G.N., Jongman, R.H.G., Scholes, R.J., *et al.*

(2013). Essential biodiversity variables. *Science*, 339, 277–278.

Pettorelli, N., Vik, J.O., Mysterud, A., Gaillard, J.-M., Tucker, C.J. & Stenseth, N.Chr. (2005).

Using the satellite-derived NDVI to assess ecological responses to environmental change.

Trends Ecol. Evol., 20, 503–510.

Ricklefs, R.E. (2008). Disintegration of the ecological community. *Am. Nat.*, 172, 741–750.

Rocchini, D., Balkenhol, N., Carter, G.A., Foody, G.M., Gillespie, T.W., He, K.S., *et al.* (2010).

Remotely sensed spectral heterogeneity as a proxy of species diversity: Recent advances and

open challenges. *Ecol. Inform.*, 5, 318–329.

Rocchini, D., Luque, S., Pettorelli, N., Bastin, L., Doktor, D., Faedi, N., *et al.* (2018). Measuring

β-diversity by remote sensing: A challenge for biodiversity monitoring. *Methods Ecol. Evol.*, 9,

1787–1798.

Schmidtlein, S., Zimmermann, P., Schüpferling, R. & Weiß, C. (2007). Mapping the floristic

continuum: Ordination space position estimated from imaging spectroscopy. *J. Veg. Sci.*, 18,

131–140.

Schneider, F.D., Morsdorf, F., Schmid, B., Petchey, O.L., Hueni, A., Schimel, D.S., *et al.* (2017).

Mapping functional diversity from remotely sensed morphological and physiological forest traits.

Nat. Commun., 8, 1441.

Schweiger, A.K., Cavender-Bares, J., Townsend, P.A., Hobbie, S.E., Madritch, M.D., Wang, R.,

et al. (2018). Plant spectral diversity integrates functional and phylogenetic components of

biodiversity and predicts ecosystem function. *Nat. Ecol. Evol.*, 2, 976–982.

Turner, W. (2014). Sensing biodiversity. *Science*, 346, 301–302.

Ustin, S.L. & Gamon, J.A. (2010). Remote sensing of plant functional types. *New Phytol.*, 186,

795–816.

Ustin, S.L., Gitelson, A.A., Jacquemoud, S., Schaepman, M., Asner, G.P., Gamon, J.A., *et al.*

(2009). Retrieval of foliar information about plant pigment systems from high resolution

spectroscopy. *Remote Sens. Environ.*, 113, 67–77.

Vellend, M. (2010). Conceptual synthesis in community ecology. *Q. Rev. Biol.*, 85, 183–206.

Wang, R. & Gamon, J.A. (2019). Remote sensing of terrestrial plant biodiversity. *Remote Sens.*

Environ., 231, 111218.

Wang, R., Gamon, J.A., Cavender-Bares, J., Townsend, P.A. & Zygielbaum, A.I. (2018a). The

spatial sensitivity of the spectral diversity–biodiversity relationship: an experimental test in a

prairie grassland. *Ecol. Appl.*, 28, 541–556.

Wang, R., Gamon, J.A., Schweiger, A.K., Cavender-Bares, J., Townsend, P.A., Zygielbaum,

A.I., *et al.* (2018b). Influence of species richness, evenness, and composition on optical

diversity: A simulation study. *Remote Sens. Environ.*, 211, 218–228.

Whittaker, R.H. (1960). Vegetation of the Siskiyou Mountains, Oregon and California. *Ecol.*

Monogr., 30, 279–338.

Whittaker, R.H. (1972). Evolution and measurement of species diversity. *Taxon*, 21, 213–251.

TABLES

576 **Table 1** Partitioning spectral diversity into additive components for the two simulated regions

577 (Figs. 3–4) and for the NEON imagery (Fig. 5).

		High β, low α	Low β, high α	NEON imagery
Sums of squares (SS)				
SS_γ		1.66	1.66	740.24
SS_β (% of SS_γ)		1.40 (84.4%)	0.08 (4.8%)	168.85 (22.8%)
SS_α (% of SS_γ)		0.26 (15.6%)	1.58 (95.2%)	571.39 (77.2%)
Spectral diversity (SD)				
SD_γ		0.0027	0.0027	0.0029
SD_β		0.0022	0.00013	0.00065
$\overline{SD}_\alpha$		0.00043	0.0026	0.0022
Feature contribution to SD (FCSD)				
$FCSD_\gamma$	PC 1	0.800	0.800	0.690
	PC 2	0.157	0.157	0.154
	PC 3	0.023	0.023	0.090
$FCSD_\beta$	PC 1	0.906	0.859	0.520
	PC 2	0.079	0.108	0.329

	PC 3	0.013	0.019	0.092
$\overline{FCSD}_\alpha$	PC 1	0.259	0.798	0.733
	PC 2	0.513	0.158	0.108
	PC 3	0.050	0.022	0.092

FIGURE LEGENDS

Figure 1 Partitioning plant spectral γ-diversity into additive $β$ and $α$ components. A region of interest is split into a number of communities of a specific size and shape (here, 20 m × 20 m squares, representing standard forest inventory plots). Spectral γ-diversity refers to the total spectral diversity in the entire region calculated from pixel-level reflectance. The $β$ component corresponds to spectral diversity *among* communities, with similar colours sharing more similar spectral composition. The $α$ component refers to spectral diversity *within* individual communities. The left-most panel is a true colour (red-green-blue, RGB) image of an area of Bartlett Experimental Forest; colours for the other panels were obtained using the reflectance of different wavelength bands (R = 779 nm, G = 639 nm, B = 2301 nm).

Figure 2 Overview of our proposed workflow for partitioning plant spectral diversity. In our NEON case study, spectral data pre-processing included removing atmospheric water absorption bands (wavelengths between 1340–1455 nm and 1790–1955 nm) and noisy regions of the spectrum (wavelengths <400 nm and >2400 nm), and applying a Savitzky-Golay filter (order = 3, size = 7) to every pixel in the image to remove high-frequency noise. We masked all pixels with normalized difference vegetation index (NDVI) values <0.8, and brightness-normalized all spectra (Feilhauer *et al.* 2010). Then, we performed a PCA with type I-scaling, and visually inspected the first 17 PCs which together accounted for >99% of the total spectral variation among all pixels (Fig. S3). Only the first five PCs showed meaningful biological spatial patterns and were retained for spectral diversity measurements; PCs 6–17 were excluded as they expressed artefacts from image acquisition and processing (Fig. S3). For illustration purposes, in the diversity partitioning analysis (bottom panel) we show communities composed of only three pixels, whereas in fact we used a community size of 40 × 40 pixels in our NEON case study. For abbreviations see text. *The shade mask is for illustrative purposes and is not applied to the PCs shown in the middle panel.

Figure 3 Two simulated landscapes of equal spectral γ-diversity, but with contrasting spectral β-

diversity and α-diversity. **Left**: high spectral β-diversity but low α-diversity. **Right**: low spectral β-

diversity but high α-diversity. Each landscape is composed of 25 communities (numbered black

squares), each composed of 5 × 5 pixels (smaller coloured squares). The size of each pixel is

equivalent to the size of an individual plant and their colour corresponds to one of the 15 leaf-

level spectra (= 3 species × 5 individuals) shown in Figure S1. These colours were set by

mapping the scores of the first three principal components (PC) for each spectrum to a red-

green-blue (RGB) scale (PC 1 = green, PC 2 = red, PC 3 = blue). We generated the high

spectral β-diversity but low spectral α-diversity scenario (left panel) by randomly assigning (with

replacement) pixels within each community with individual spectra from single species (Fig. S1,

bottom row). We selected species identity per community at random using the following

probabilities: 0.60 (*Betula alleghaniensis*, green hues), 0.35 (*Populus deltoides*, blue hues) and

0.05 (*Populus tremuloides*, red hues). In this scenario, spectral β-diversity was high and spectral

α-diversity low because interspecific spectral variation (particularly between *Betula* and the two

Populus species) was higher than intraspecific spectral variation (Fig. S2). Next, to reduce

spectral β-diversity and increase α-diversity while holding γ-diversity constant, we moved the

pixels of the left panel to randomly selected positions in the right panel.

Figure 4 (a) Spectral α-diversity (SD$_\alpha$), (b) local contribution to spectral β-diversity (LCSD$_\beta$), and

(c-d) feature contribution to spectral α-diversity (FCSD$_\alpha$) for the first two spectral features (i.e.

first two principal components of the brightness-normalized reflectance data), of each

community in the two simulated regions. PC = principal component.

Figure 5 Partitioning spectral diversity using imaging spectroscopy data acquired by the

National Ecological Observatory Network (NEON) over the Bartlett Experimental Forest site.

From left to right: (1) true colour (red-green-blue, RGB) image with 0.1 m ground resolution, (2)

false colour image with 1 m resolution based on the first three principal components (PCs) of

the spectral image cube (PC1 = red, PC2 = green, PC3 = blue), (3) local contribution to spectral

β-diversity (LCSD$_\beta$), and (4) spectral α-diversity (SD$_\alpha$) of forest communities, each measuring 40

m × 40 m. For panels 3 and 4, light hues correspond to low, dark hues to high values of LCSD$_\beta$

coefficients and SD$_\alpha$ values, respectively.

FIGURES

634 **Figure 1**

637   **Figure 2**

639   **Figure 3**

Figure 4

644   **Figure 5**

RGB   PC 1,2,3   LCSD$_\beta$ SD$_\alpha$