Cryptic Patterns of Speciation in Cryptic Primates:

Microendemic Mouse Lemurs and the Multispecies Coalescent

Running title: Cryptic speciation in mouse lemurs

Jelmer Poelstra[1#], Jordi Salmona[2#], George P. Tiley[1#], Dominik Schüßler[3], Marina B. Blanco[1,4],

Jean B. Andriambeloson[5], Sophie Manzi[2], C. Ryan Campbell[1†], Olivier Bouchez[6], Paul D. Etter[7],

Amaia Iribar[2], Paul A. Hohenlohe[8], Kelsie E. Hunnicutt[1@], Eric A. Johnson[7], Peter M. Kappeler[9],

Peter A. Larsen[1&], José M. Ralison[5], Blanchard Randrianambinina[10,11], Rodin M. Rasoloarison[9],

David W. Rasolofoson[10], Amanda R. Stahlke[8], David Weisrock[12], Rachel C. Williams[1,4], Lounès

Chikhi[2,13], Edward E Louis Jr.[14], Ute Radespiel[15]*, Anne D. Yoder[1]*

[1]: Department of Biology, Duke University, Durham, NC 27708, USA

[2]: CNRS, Université Paul Sabatier, IRD; UMR5174 EDB (Laboratoire Évolution & Diversité 11

Biologique), 118 route de Narbonne, 31062 Toulouse, France

[3]: Research Group Ecology and Environmental Education, Institute of Biology and Chemistry,

University of Hildesheim, Universitaetsplatz 1, 31141 Hildesheim, Germany

[4]: Duke Lemur Center, Duke University, Durham, NC 27705, USA

[5]: Zoology and Animal Biodiversity, University of Antananarivo, Antananarivo 101, Madagascar

[6]: INRA, US 1426, GeT-PlaGe, Genotoul, Castanet-Tolosan, France

[7]: *Institute of Molecular Biology, University of Oregon, Eugene, OR, USA*

[8]: *Institute for Bioinformatics and Evolutionary Studies, Department of Biological Sciences,*

University of Idaho, Moscow, ID 83844, USA

[9]: *Behavioral Ecology and Sociobiology Unit, German Primate Center, Kellnerweg 6, 37077*

Göttingen, Germany

[10]: *Groupe d'Etude et de Recherche sur les Primates de Madagascar (GERP); BP 779,*

Antananarivo 101, Madagascar

[11]: *Faculté des Sciences, University of Mahajanga, Mahajanga, Madagascar*

[12]: *Department of Biology, University of Kentucky, Lexington, KY, 40506, USA*

[13]: *Instituto Gulbenkian de Ciência, Rua da Quinta Grande, 6, 2780-156 Oeiras, Portugal*

[14]: *Grewcock Center for Conservation and Research, Omaha's Henry Doorly Zoo and Aquarium,*

Omaha, NE

[15]: *Institute of Zoology, University of Veterinary Medicine Hannover, Buenteweg 17, 30559*

Hannover, Germany

#: joint first authors

*: joint senior authors (corresponding authors)

@: current address: Department of Biological Sciences, University of Denver, Denver, CO

80208

&: current address: Department of Veterinary and Biomedical Sciences, University of Minnesota,

Saint Paul, MN 55108

†: current address: Department of Evolutionary Anthropology, Duke University, Durham, NC

27708, USA

Corresponding authors:

Ute Radespiel, Institute of Zoology, University of Veterinary Medicine Hannover, Germany,

email: ute.radespiel@tiho-hannover.de

Anne D. Yoder, Department of Biology, Duke University, Durham, NC, USA, email:

anne.yoder@duke.edu

Author contributions:

- Conception and design of study:

JP, GPT, DS, JS, LC, UR, ADY

- Data collection:

DS, MBB, JBA, EELJ, DWR, RMR, PK, JMR, BR collected samples in the field.

- Data analysis and interpretation:

DS, JS, LC, OB, PE, CRC, PAL, ARS, DW, AIP, PH, KEH, EJ, SM, RCW, EELJ, UR, ADY

generated sequencing data.

JP, GPT, DS, JS, LC, UR, ADY conducted population genetic and phylogenetic analyses.

- Drafting and revising manuscript:

JP, GPT, DS, JS, LC, UR, ADY drafted the manuscript.

All co-authors revised and agreed on the last version of the manuscript.

ABSTRACT

Mouse lemurs (*Microcebus*) are a radiation of morphologically cryptic primates distributed throughout Madagascar for which the number of recognized species has exploded in the past two decades. This taxonomic explosion has prompted understandable concern that there has been substantial oversplitting in the mouse lemur clade. Here, we take an integrative approach to investigate species diversity in two pairs of sister lineages that occur in a region in northeastern Madagascar with high levels of microendemism and predicted habitat loss. We analyzed RADseq data with multispecies coalescent (MSC) species delimitation methods for three named species and an undescribed lineage previously identified to have divergent mtDNA. Marked differences in effective population sizes, levels of gene flow, patterns of isolation-by-distance, and species delimitation results were found among them. Whereas all tests support the recognition of the presently undescribed lineage as a separate species, the species-level distinction of two previously described species, *M. mittermeieri* and *M. lehilahytsara* is not supported – a result that is particularly striking when using the genealogical discordance index (*gdi*). Non-sister lineages occur sympatrically in two of the localities sampled for this study, despite an estimated divergence time of less than 1 Ma. This suggests rapid evolution of reproductive isolation in the focal lineages, and in the mouse lemur clade generally. The divergence time estimates reported here are based on the MSC and calibrated with pedigree-based mutation rates and are considerably more recent than previously published fossil-

calibrated concatenated likelihood estimates, however. We discuss the possible explanations for

this discrepancy, noting that there are theoretical justifications for preferring the MSC estimates

in this case.

Keywords: effective population size, cryptic species, multispecies coalescent, species

delimitation, speciation, microendemism

INTRODUCTION

Mouse lemurs (*Microcebus* spp.) are small, nocturnal primates that are widespread in the

forests of Madagascar (Mittermeier et al. 2010), one of the world's most biodiverse environments

(Myers et al. 2000; Goodman and Benstead 2005; Estrada et al. 2017). Mouse lemur diversity

was long overlooked (Zimmermann and Radespiel 2014) until the introduction of genetic

analyses made it feasible to identify diverging lineages despite similar phenotypes and ecological

niches. This genetic perspective has led to the description of many new species, with 24 species

recognized at present. In one such study, Radespiel et al. (2008) surveyed the forests of the

Makira region (Fig. 1) in northeastern Madagascar and found evidence for three divergent

mitochondrial lineages occurring in sympatry. One of these was identified as *M. mittermeieri*

(Louis et al. 2006), while the second was newly described as *M. macarthurii*. A third lineage,

provisionally called M. sp. #3, was hypothesized to represent a new species closely related to *M.*

macarthurii but was not formally named because the data was limited to mtDNA sequence data

from one individual. Furthermore, two other species occur in the region, *M. lehilahytsara* (Roos

and Kappeler in Kappeler et al. 2005) at higher elevations, and *M. simmonsi* (Louis et al. 2006)

in lowland forests in the south (Fig. 1).

Figure 1: Sampling sites in northeastern Madagascar.

Size of the circles scales with the number of individuals sequenced for a given site. Green

background indicates forest cover as per Du Put & Moat (1998), with darker green indicating

"low altitude" and paler green indicating "mid altitude" evergreen humid forest. At Anjiahely

and Ambavala, two species were detected; in both cases, the leftmost site marker was slightly

Given that many previous taxonomic descriptions of mouse lemurs have relied strongly, if

not entirely, on mtDNA sequence divergence, there has been criticism that mouse lemurs (and

lemurs more generally) may have been oversplit (Tattersall 2007; Markolf et al. 2011). Species

delimitation using only mtDNA is now widely regarded as problematic, given that the

mitochondrial genome represents a single non-recombining locus whose gene tree may not

represent the underlying species tree (e.g., Pamilo and Nei 1988; Maddison 1997). Mitochondria

are also maternally inherited and therefore susceptible to effects of male-biased dispersal (e.g.,

Dávalos and Russell 2014), which is prevalent in mouse lemurs (reviewed in Radespiel 2016).

Moreover, previous attempts to resolve mouse lemur relationships using nuclear sequences have

been complicated by high gene tree discordance, consistent with strong incomplete lineage

sorting (e.g., Heckman et al. 2007; Weisrock et al. 2010). These issues can be overcome with

genomic approaches, which provide power for simultaneously resolving phylogenetic

relationships and estimating demographic parameters such as divergence times, effective

population sizes, and rates of gene flow — even among closely related species (e.g., Palkopoulou

et al. 2018; Pedersen et al. 2018).

Given that cryptic species are by definition difficult to identify based on phenotypic

characters (Bickford et al. 2007), recently developed methods for genomic species delimitation

have advanced our ability to recognize and quantify their species diversity. In the past decade,

both theory and methods for species delimitation have seen substantial progress, especially those

which leverage the multispecies coalescent (MSC) model (Pamilo and Nei 1988; Rannala and

Yang 2003). MSC-based species delimitation methods have been increasingly applied to

genomic data (e.g. Carstens and Dewey 2010; Yang and Rannala 2010; Grummer et al. 2014;

Dincă et al. 2019; Hundsdoerfer et al. 2019), though they have also been considered

controversial (Edwards and Knowles 2014; Sukumaran and Knowles 2017; Barley et al. 2018).

The controversy largely relates to the idea that strong population structure can be mistaken for

species boundaries, which may lead to oversplitting (Jackson et al. 2017; Sukumaran and

Knowles 2017; Luo et al. 2018; Leaché et al. 2019; Chambers and Hillis 2020). To overcome this

potential weakness, Jackson et al. (2017) proposed a heuristic criterion, the genealogical

divergence index (*gdi*), with Leaché et al. (2019) further suggesting that *gdi* helps to differentiate

between population structure and species-level divergence. In parallel, sophisticated statistical

approaches have been developed that can detect the presence and magnitude of gene flow during

or after speciation (Gronau et al. 2011; Payseur and Rieseberg 2016; Dalquen et al. 2017; Wen et

al. 2018). Taken together, these analytical developments are crucial to our ability to recognize the

patterns that characterize the speciation process, despite the challenge of identifying species

without universally agreed upon criteria (de Queiroz 2007).

In this study, we use a structured framework starting with phylogenetic placement of

lineages and culminating with the MSC to delimit species, estimate divergence times, identify

post-divergence gene flow, and to estimate both current and ancestral effective population sizes

(`Fig. S1`). We take advantage of increased geographic, population-level, and genomic sampling

to comparatively examine speciation dynamics for two pairs of closely related lineages in the

region (described below as Clades I and II) and perform MSC species delimitation methods with

Restriction-site Associated DNA sequencing (RADseq) data to infer divergence times, effective

population sizes, and rates of gene flow between these lineages. We also provide a novel whole-

genome assembly for the previously undescribed lineage and compare inferences of effective

population size (N_e) through time from whole-genome versus RADseq data. We find notably

different species delimitation results for the lineages in the two mouse lemur clades and believe

that the integrative analytic framework here used can be applied more generally to allow

investigators to test hypotheses of population- versus species-level differentiation.

MATERIALS AND METHODS

Summary of Analyses

We generated RADseq data for 63 individuals from 6 lineages, of which 48 were from

the two focal clades and passed quality control. First, we used maximum likelihood approaches

to infer relationships among lineages and to provide a framework for subsequent species

delimitation analyses (Fig. 2A and C). To delimit species, we performed clustering (Fig. 2B) and

PCA analyses (Fig. 3A-C), as well as formal MSC species delimitation analyses using SNAPP

and BPP. We also used the recently developed genealogical divergence index *gdi* based on BPP

parameter estimates (Fig. 3D) and performed an isolation-by-distance analysis (Fig. 4). To

determine to what extent ongoing and ancestral gene flow may have contributed to current

patterns of divergence, we used G-PhoCS and D-statistics (Fig. 5). Finally, we generated whole-

genome sequencing data for a single individual designated as *M.* sp. #3 comparing it to one for

M. mittermeieri from a previous study (Hunnicutt et al., 2020). The genomes were used to infer

N_e though time with Multiple Sequentially Markovian Coalescent (MSMC) analysis and

compared those estimated from G-PhoCS (Fig. 6). Below, we describe the methods in some

detail, while further details can be found in the Supplementary Material.

Study Sites and Sampling

Microcebus samples were obtained by taking ~2 mm² ear biopsies of captured (and

thereafter released) individuals between 2008 and 2017 at seven humid evergreen forest sites

(50-979 m a.s.l.) in the Analanjirofo and Sava regions of northeastern Madagascar (`Fig. 1;`

`Table S1`). Additional samples were used from Riamalandy, Zahamena National Park (NP),

Betampona Strict Nature Reserve (SNR) and Tampolo (Louis et al. 2006; Weisrock et al. 2010;

Louis and Lei 2016) (`Fig. 1`). With this sampling strategy, we expected to include all mouse

lemur species thought to occur in the region (from north to south): *M. mittermeieri*, *M.*

macarthurii, *M.* sp. #3, *M. lehilahytsara,* and *M. simmonsi* (`Fig. 1`). *Microcebus murinus,*

which occurs in western and southeastern Madagascar, was used as an outgroup.

Sequencing Data, Genotyping and Genome Assembly

We generated RADseq libraries using the *Sbf*I restriction enzyme, following three

protocols (Supplementary Methods, `Table S1`). Sequences were aligned to the *M.* sp. #3

nuclear genome generated by this study, and to the published *M. murinus* mitochondrial genome

(LeCompte et al. 2016). We used two genotyping approaches to ensure robustness of our results.

First, we estimated genotype likelihoods (GL) with `ANGSD` v0.92 (Nielsen et al. 2012;

Korneliussen et al. 2014), which retains information about uncertainty in base calls, thereby

alleviating some issues commonly associated with RADseq data such as unevenness in

sequencing depth and allele dropout (Lozier 2014; Pedersen et al. 2018; Warmuth and Ellegren

2019). Second, we called genotypes with `GATK` v4.0.7.0 (DePristo et al. 2011), and filtered `GATK`

genotypes following the "FS6" filter of O'Leary et al. (2018; their Table 2). We furthermore used

three mtDNA fragments [Cytochrome Oxidase II (COII), Cytochrome B (cytB), and d-loop] that

were amplified and Sanger sequenced for additional phylogenetic analyses.

The genome of the *M.* sp. #3 individual sampled in Mananara-Nord NP (`Table S3`) was

sequenced with a single 500bp insert library on a single lane of an Illumina HiSeq 3000 with

paired-end 150bp reads. We used `MaSuRCA` v3.2.2 (Zimin et al. 2013) for contig assembly and

`SSPACE` (Boetzer et al. 2011) for scaffolding. Scaffolds potentially containing mitochondrial or

X-chromosome sequence data were removed for downstream analyses.

Phylogenetic Analyses

We used three phylogenetic approaches to infer relationships among lineages: (1)

maximum likelihood using RAxML v8.2.11 (Stamatakis 2014), (2) SVDquartets, an MSC method

that uses phylogenetic invariants, implemented in PAUP v4a163 (Chifman and Kubatko 2014),

and (3) SNAPP, a full-likelihood MSC method for biallelic data that does not require joint gene

tree estimation (v1.3.0; Bryant et al. 2012). Analyses with RAxML and SVDquartets used all

available individuals, whereas SNAPP analyses were performed with subsets of 12 and 22

individuals for computational feasibility (see Supplementary Methods).

Species Delimitation

Clustering approaches and summary statistics. — Clustering analyses were performed

using corresponding methods based on ANGSD genotype likelihoods [clustering in NgsAdmix v32

(Skotte et al. 2013) and PCA in ngsTools va4d338d (Fumagalli et al. 2014)] and on GATK-called

genotypes [clustering in ADMIXTURE v1.3.0 (Alexander et al. 2009) and PCA using the glPca()

function in adegenet v2.1.1 (Jombart and Ahmed 2011)]. These analyses were run for Clade I

and II together and separately.

MSC-based approaches. — We used SNAPP to test if the two lineages each in Clade I and

II could be delimited using Bayes factors (Leaché et al. 2014), interpreting 2ln Bayes factors

greater than six as strong evidence for a given model (Kass and Raftery 1995). We also applied

guided species delimitation analyses with BPP (Yang and Rannala 2010; Rannala and Yang 2013)

using full-length fasta files for a subset of individuals based on the species tree estimated by

SVDquartets and SNAPP.

g*di.* — Coalescent node heights (τ) and ancestral effective population sizes (θ) estimated

by BPP were used to compute the genealogical divergence index (*gdi*; Jackson et al. 2017;

Leaché et al. 2019) for the lineages in Clade I and II. We calculated *gdi* as in Leaché et al.

(2019), using their equation 7 ($gdi = 1-e^{-2\tau/\theta}$), where $2\tau/\theta$ represents the population divergence

time between two taxa in coalescent units. θ is taken from one of the two taxa and therefore *gdi*

was calculated twice for each species pair, alternating the focal taxon. We computed *gdi* using τ

and θ parameter estimates for each posterior BPP sample to incorporate uncertainty in the

estimates. Jackson et al. (2017) suggested the following interpretation of *gdi* values: the taxon

pair (a) is unambiguously a single species for *gdi* < 0.2, (b) is unambiguously two separate

species for *gdi* > 0.7, and (c) falls in an ambiguous zone for 0.7 > *gdi* < 0.2.

Isolation-by-distance. — We tested for isolation-by-distance using the VCF file produced

by GATK with the gl.ibd() function in the R package dartR 1.1.11 (Gruber et al. 2018).

Inference of gene flow and divergence times

G-PhoCS v1.3 (Gronau et al. 2011), a Bayesian MSC approach that allows for the

estimation of periods of gene flow (*i.e.* "migration bands"), was used to jointly infer divergence

times, population sizes, and rates of gene flow between specific lineages. Based on the results of

exploratory models each containing a single "migration band" between two lineages, we ran a

final model with a migration band allowing gene flow from *mittermeieri* to *lehilahytsara*, and a

migration band allowing gene flow from *macarthurii* to *M.* sp. #3. Given the observed

mitonuclear discordance between *M.* sp. #3 and *M. macarthurii* (see Results)*,* we investigated

gene flow between them in more detail by running G-PhoCS using a dataset with only *M.* sp. #3,

M. macarthurii, and *M. lehilahytsara* individuals, wherein *M.* sp. #3 was divided into two

populations detected using clustering approaches.

The D-statistic and related formal statistics for admixture use phylogenetic invariants to

infer post-divergence gene flow between non-sister populations or taxa. We used the qpDstat tool

of admixtools v4.1 (Patterson et al. 2012) to compute four-taxon D-statistics for all possible

configurations in which gene flow between non-sister lineages among the five ingroup lineages

could be tested. We additionally tested for gene flow between *M. macarthurii* and *M.* sp. #3 by

separately treating (1) the two distinct *M.* sp. #3 populations detected by clustering approaches,

and (2) *M. macarthurii* individuals with and without "*M.* sp. #3-type" mtDNA (see Results). In

all tests, *M. murinus* was used as P4 (outgroup).

Effective Population Size Through Time

Studies have shown that population structure can generate spurious signals of population

size change (Beaumont 2004; Chikhi et al. 2010; Heller et al. 2013). For example, sequentially

Markovian coalescent approaches such as MSMC (Schiffels and Durbin 2014) actually estimate

the inverse instantaneous coalescence rate, which is only equivalent to an effective size in

panmictic models (Mazet et al. 2016; Rodríguez et al. 2018). We therefore inferred and

compared population size histories using two methods. We estimated N_e over time with MSMC

for two species, using the whole-genome data of *M.* sp. #3 and *M. mittermeieri* (Hunnicutt et al.

2020) mapped to the chromosome-level genome assembly of *M. murinus* (Larsen et al. 2017).

These estimates were compared to inferred changes in N_e over time based on θ estimates from G-

PhoCS for each predefined extant or ancestral population. Although the MSC was not expressly

developed to estimate change in N_e over time, this allowed us to examine broad demographic

trends for relatively small population-level sampling with RADseq data, and to explicitly

incorporate divergence events.

Mutation Rate and Generation Time

We used empirical estimates of mutation rate and generation time to convert coalescent

units from BPP, G-PhoCS and MSMC analyses into absolute times and population sizes. We

incorporated uncertainty by drawing from mutation rate and generation time distributions for

each sampled generation of the MCMC chains in BPP and G-PhoCS (MSMC parameter estimates

were converted using the point estimates). For the mutation rate, we used a gamma distribution

based on the mean (1.236×10^{-8}) and variance (0.107×10^{-8}) of seven pedigree-based mutation

rate estimates for primates (see Campbell et al. 2019, `Table S1`). For the generation time, we

used a lognormal distribution with a mean of ln(3.5) and standard deviation of ln(1.16) based on

estimates of 4.5 years calculated from survival data (Zohdy et al. 2014; Yoder et al. 2016) from

M. rufus, and 2.5 years from average parent age based on capture-mark-recapture and parentage

data in the wild (Radespiel et al. 2019) for *M. murinus*.

RESULTS

RADseq Data and Whole-Genome Assembly

We used three library generation protocols, two sequencing lengths, and a combination of

single and paired-end sequencing, yielding data for all 63 individuals in the study and

demonstrating the utility of cross-laboratory RAD sequencing, as previously shown in other taxa

(e.g., Gonen et al. 2015). From more than 447 million raw reads (`Table S1`), over 394 million

passed quality filters, with approximately 182 million successfully aligning to the *M.* sp. #3

reference genome. We obtained an average of 120,000 loci per individual with coverage ranging

from ~1 to ~22x (`Table S1`).

We assembled approximately 2.5 Gb of nuclear genome sequence data for *M*. sp. #3 with a

contig N50 around 36 Kb (Table S3). While the final assembly was fragmented, as expected

for a single Illumina library genome, only 6.4% of mammalian BUSCOs were found to be

missing. The genome sequence and associated gene annotations can be accessed through NCBI

(Bioproject PRJNA512515).

Phylogenetic Relationships

RAxML and SVDquartets recovered well-supported nDNA clades for *M. simmonsi*, *M.*

macarthurii, and *M*. sp. #3, the latter two as sister taxa with 100% bootstrap support (Fig. 2;

Fig. S2). SNAPP also supported *M*. sp. #3 as sister taxon to *M. macarthurii* (referred to as

Clade I) and placed *M. lehilahytsara* as sister taxon to *M. mittermeieri* (referred to as Clade II)

(Fig. S2). However, *M. lehilahytsara* was not monophyletic in RAxML analyses of nDNA

(Fig. 2C) or mtDNA (Fig. 2A), and a SVDquartets analysis of nDNA placed one *M.*

lehilahytsara individual from Ambavala as sister to all other *M. lehilahytsara* and *M.*

mittermeieri, and only weakly supported a monophyletic *M. mittermeieri* (Fig. S2A).

Figure 2: Phylogenetic relationships and ancestry proportions

a) Maximum-likelihood RAxML tree of 57 samples represented by 4,060 bp of mtDNA recovered

from RADseq and Sanger sequencing (Table S1). The gray shaded box highlights individuals of

M. macarthurii with *M.* sp. #3 mtDNA haplotypes. **b)** Clustering results for all species except the

outgroup *M. murinus*, using NgsAdmix at *K* = 5. **c)** Maximum-likelihood RAXML tree obtained

using RADseq nuclear data (nDNA). For all trees, *M. murinus* is used as the outgroup. In a) and

c), bootstrap support values >90% are indicated above each node as a red asterisk. Clades

Although mtDNA analyses placed several individuals from Anjiahely in a well-supported

clade with *M.* sp. #3, individuals from Ambavala (see `Fig. 1`), Mananara-Nord NP, and

Antanambe (`Fig. 2A`; see lower gray box), nuclear RADseq data placed them unambiguously

within the *M. macarthurii* clade (`Fig. 2B,C`). This suggests that individuals from Anjiahely are

in fact *M. macarthurii*, but carry two divergent mtDNA lineages, and that true *M.* sp. #3 are only

found between Ambavala and Antanambe (`Fig. 1`). The cause of this mitonuclear discordance

for *macarthurii* in Anjiahely was investigated further (see the section "Interspecific Gene

Flow").

Species Delimitation

Genetic structure. — A PCA with both pairs of sister lineages (Clade I: *M. macarthurii* and

M. sp. #3; Clade II: *M. mittermeieri* and *M. lehilahytsara*) distinguished the two clades along

PC1, and distinguished *M. macarthurii* and *M.* sp. #3 along PC2 (`Fig. 3B`). When restricting

clustering analyses to Clade I, K = 2 was the best-supported number of clusters with both

approaches, distinguishing *M. macarthurii* and *M.* sp. #3 (`Fig. S5; Fig. S7B`). At K = 3, *M.*

sp. #3 was divided into two clusters with individuals from Mananara-Nord NP and Antanambe

separated from Ambavala individuals (`Fig. S7B, Fig. S10`). A separate PCA analysis for

Clade I also distinguished these two groups along PC2 (`Fig. 3C`), which we hereafter refer to

as "southern *M.* sp. #3" (Mananara-Nord NP and Antanambe are south of the Mananara river)

and "northern *M.* sp. #3" (Ambavala is north of the river, and 24.0 km from Mananara-Nord NP

Figure 3: Population genetic structure and the gdi.

a) Genealogical divergence index (gdi) for *M. macarthurii* – *M.* sp. #3 and *M. lehilahytsara* – *M. mittermeieri*. *gdi* values > 0.7 suggest separate species, *gdi* values < 0.2 are below the lower threshold for species delimitation, and 0.2 < *gdi* < 0.7 is an "ambiguous" range (Jackson et al. 2017). **b-d)** PCA analyses for b) all four species in Clades I and II, c) Clade I only: *M.* sp. #3 and *M. macarthurii*, with the former showing a split into two population groups: "northern" (Ambavala) and "southern" *M.* sp. #3 (Antanambe and Mananara-Nord NP), d) Clade II only: *M. lehilahytsara* and *M. mittermeieri*.

and 35.2 km from Antanambe; Fig. 1). When restricting clustering analyses to Clade II,

ADMIXTURE and ngsAdmix suggested optimal values of 1 and 2, respectively; at K=2, *M.*

mittermeieri and *M. lehilahytsara* were largely but not entirely separated by both approaches

(`Fig. S5`, `Fig. S7C`, `Fig. S11`). A PCA distinguished *M. mittermeieri* and *M. lehilahytsara*

along PC1 but with little separation (`Fig. 3D`).

SNAPP and BPP. — SNAPP Bayes factors strongly favored splitting Clade I into two

species (2lnBF = 17,304 and 34,326 for two different datasets, `Table S6`), as well as splitting

Clade II, although with a smaller difference in marginal likelihood scores (2lnBF = 1,828 and

993). All putative species assignments were recovered by the guided delimitation analysis with

BPP (`Fig. S12`).

Genealogical divergence index (gdi). — For the Clade I sister pair, *gdi* was 0.727 (95%

HPD: 0.718-0.737) from the perspective of *M. macarthurii* (i.e. above the upper threshold for

species delimitation), and 0.500 (0.488-0.511) from the perspective of *M.* sp. #3 (i.e. in the upper

ambiguous zone for species delimitation; `Fig. 3A`). In contrast, *gdi* values for the Clade II

putative species pair were much lower and even below the lower threshold for species

delimitation: 0.080 (0.074-0.086) from the perspective of *M. lehilahytsara*, and 0.193 (0.187-

0.201) from the perspective of *M. mittermeieri* (`Fig. 3A`).

Isolation-by-distance (IBD). — While comparisons within and between lineages appeared

to follow a single isolation-by-distance pattern for *M. mittermeieri* and *M. lehilahytsara* (Clade

II, r=0.693, p=0.002, `Fig. 4B`), comparisons within versus between lineages differed strongly

for *M. macarthurii* and *M.* sp. #3 (Clade I, `Fig. 4A`). Specifically, genetic distances between *M.*

macarthurii and *M.* sp. #3 were much larger than within lineages and were also much larger than

between *M. mittermeieri* and *M. lehilahytsara*, despite similar geographic distances.

Figure 4: Patterns of isolation-by-distance in the two clades.

a) Clade I (*M. macarthurii* and *M. sp. #3*). **b)** Clade II (*M. mittermeieri* and *M. lehilahytsara*).

Population comparisons within lineages are shown as blue points, and comparisons between

lineages are shown as red points. Both panels have the same y-axis scale, while the inset in B has

a lower limit on the y-axis to better show the spread of points, given the smaller genetic distances

between *M. mittermeieri* and *M. lehilahytsara*.   24

Interspecific gene flow. — G-PhoCS inferred high levels of gene flow in Clade II, from *M.*

mittermeieri to *M. lehilahytsara* [population migration rate (2Nm) = 1.59 (95% HPD: 1.50-1.68),

migrants per generation: 0.18% (95% HPD: 0.09-0.27%)], and much lower levels of gene flow in

Clade I, from *M.* sp. #3 to *macarthurii* [2Nm = 0.08 (95% HPD: 0.07-0.09), migrants per

generation: 0.10% (0.05-0.15%)] (Fig. 5C). G-PhoCS also inferred low levels of gene flow

between the two clades, most likely between ancestral populations, but the timing and direction

of gene flow could not be determined (Supplementary Results; Fig. S13), and D-statistics

testing for gene flow between the clades were not significant (Fig. S14).

We further investigated gene flow between *M.* sp. #3 and *M. macarthurii* by taking the

strong population structure within *M.* sp. #3 into account. D-statistics suggested that northern *M.*

sp. #3 and *M. macarthurii* with "*M.* sp. #3-type" mtDNA share a slight excess of derived alleles

in relation to southern *M.* sp. #3, significantly deviating from 0, which indicates gene flow

(Fig. S15A). Using a G-PhoCS model with separate northern and southern groups for *M.* sp.

#3, we found that (1) gene flow with *M. macarthurii* took place before and after the onset of

divergence between northern and southern *M.* sp. #3, (2) gene flow between extant lineages

occurred or occurs only between northern (and not southern) *M.* sp. #3 and *M. macarthurii,* and

(3) gene flow is asymmetric, predominantly into *M. macarthurii* (Fig. S15B).

Divergence Times

We estimated divergence times under the MSC model using BPP and G-PhoCS both with

and without interspecific gene flow (Fig. 5; Fig. S17). Results were similar across these

approaches, with the exception of divergence times between sister lineages in G-PhoCS models

with versus without gene flow (Fig. 5). Specifically, the divergence time between *M.* sp. #3

and *M. macarthurii* (Clade I) without gene flow was estimated at 115 ka ago (95% HPD range:

52-190 ka across G-PhoCS and BPP models) (Fig. 5; Fig. S17), but at 193 ka ago (95%

HPD: 89-318 ka) when incorporating gene flow (Fig. 5C-D). In Clade II, this difference in

estimated divergence times was considerably larger: under an isolation model it was estimated to

be 103 ka ago (95% HPD: 49-171 ka; Fig. 5) and as much as 520 ka ago (95% HPD: 249-871

ka) when modeled with gene flow (Fig. 5). Deeper nodes were not as strongly affected:

divergence time between Clades I and II was estimated at 687 ka ago (95% HPD: 337-1126 ka)

across G-PhoCS and BPP isolation models, and at 796 ka ago (95% HPD: 360-1311 ka) in a G-

PhoCS model with gene flow (Fig. 5D).

Figure 5: Demographic histories inferred by G-PhoCS and BPP.

a-c) Divergence times (y-axis) and effective population sizes (x-axis) inferred with and without

migration. Migration bands representing the estimated magnitude of gene flow are illustrated in

(c). **d-e)** Comparison of divergence times and effective population sizes for each node and

lineage, respectively. The symbol "A" represents the lineage ancestral to *M. simmonsi*, *M.*

mittermeieri and *M. lehilahytsara*, "B" represents the lineage ancestral to *M.* sp. #3, *M.*

macarthurii, *M. simmonsi*, *M. mittermeieri* and *M. lehilahytsara*, and "root" represents the

lineage ancestral to all six species included.

Effective Population Sizes

We found large differences in N_e among lineages, with considerably larger N_e for the

lineages in Clade II, *M. lehilahytsara* (point estimate and 95% HPD range across the BPP and G-

PhoCS models with and without interspecific gene flow: 159 k; 58-265 k) and *M. mittermeieri*

(78 k; 36-140 k), than the lineages in Clade I, *M.* sp. #3 (24 k; 12-38 k) and *M. macarthurii* (12

k; 5-19 k) (Fig. 5A-C). Wide HPD intervals for *M. mittermeieri* and *lehilahytsara* are due to

differences between models with and without gene flow. Using the G-PhoCS model focused on

Clade I, fairly similar effective population sizes were estimated separately for northern (47 k; 17-

78 k), southern (23 k; 12-37 k), and ancestral (33 k; 17-53 k) *M.* sp. #3 lineages (Fig. S13).

Using the whole-genome data for one individual of *M.* sp. #3 (from the southern group)

and for *M. mittermeieri*, a comparison of MSMC analyses and G-PhoCS models with and without

gene flow (Fig. 6) showed highly similar and markedly declining estimates of population

sizes towards the present for *M.* sp. #3 (Fig. 6A). Estimates for *M. mittermeieri* were more

variable across analyses and did not show a consistent decline towards the present (Fig. 6B).

Figure 6: Estimates of effective population size through time for two species.

Effective population sizes through time as inferred by MSMC for whole-genome data from a single individual (green

lines), and by G-PhoCS for RADseq data without ("RAD: iso", orange lines) and with ("RAD: mig", blue lines) gene

flow. **(A)** *M.* sp. #3. G-PhoCS analyses are shown for the southern *M.* sp. #3 population group and its ancestral

lineages in the 3-species model given that the whole-genome individual was sampled from that population. **(B)** *M.*

mittermeieri. G-PhoCS analyses are shown for *M. mittermeieri* and its ancestral lineages in the 5-species model. The

sharp "jag" in the model with gene flow represents a small N_e estimate for the *M. mittermeieri-lehilahytsara*

ancestor, which exists for an extremely short time in this model (see Fig. 6B,C), likely preventing proper estimation

of N_e.

DISCUSSION

We used a MSC-based framework for genomic species delimitation and identified rapid

and recent diversification of mouse lemurs in a relatively small area in northeastern Madagascar.

The same region was previously identified to harbor high levels of lemur microendemism that is

vulnerable to the effects of climate change (Brown and Yoder 2015) and anthropogenic habitat

alteration (Schüßler et al. 2020), marking it as a region of conservation concern. Species-level

divergence was strongly supported for *M.* sp. #3 and its sister species *M. macarthurii* (Clade I,

`Fig. 2`), but not for the pair of *M. mittermeieri* and *M. lehilahytsara* (Clade II, `Fig. 2`),

despite our baseline assumption that the latter were distinct species (Olave et al., 2014). We

inferred that the focal species all diverged from their common ancestors within the past million

years and documented two cases of sympatric occurrence, each with one representative from

Clade I and one from Clade II. The combined findings of recent divergence and sympatric

overlap suggest that reproductive isolation can evolve rapidly in mouse lemurs.

Support for Separate Sister Species Differs Sharply Between the Two Clades

Evidence for distinguishing *M.* sp. #3 and *M. macarthurii* as separate species was strong

and consistent across analyses. They were reciprocally monophyletic across all phylogenetic

analyses of RADseq data (`Fig. 2C; Fig. S2; Fig. S3`), separated unambiguously in

clustering and PCA analyses (`Fig. 2B; Fig. 3BC; Fig. S4; Fig. S6-S10`), were

strongly supported as separate lineages using SNAPP Bayes factors (`Table 1`) and BPP (`Fig.`

S12), and passed the heuristic species delimitation criterion of *gdi* (Fig. 3A). A comparison of

genetic and geographic distances moreover showed a clear distinction between intra- and

interspecific genetic distances (Fig. 4). Finally, gene flow between these two lineages was

estimated to have occurred at very low levels (G-PhoCS migration band = 0.08; Fig. 5c).

In contrast, separate species status of *M. lehilahytsara* and *M. mittermeieri* (Clade II) was

not supported by genomic data. These species were paraphyletic in RAxML and SVDquartets

analyses (Fig. 2A,C; Fig. S2) and not as clearly separated in clustering and PCA analyses

(Fig. 2B; Fig. 3BD; Fig. S5, S7-S9; Fig. S10). Although the Bayes factor support

from SNAPP was strong by standard guidelines (Kass and Raftery 1995), the evidence was much

weaker relative to species in Clade I and decreased when more individuals were included

(Table S6). It is unsurprising that Bayes factors will support splitting lineages with genetic

structure (Sukumaran and Knowles 2017; Leaché et al. 2019) even with low levels of gene flow

(Barley et al. 2018). Therefore, standard guidelines for interpreting Bayes factors may be of

limited value for delimiting species, as informed by the lack of monophyly, high levels of

inferred gene flow, and failure of additional delimitation tests observed here. Guided delimitation

also separated *M. lehilahytsara* and *M. mittermeieri* (Fig. S11), but similar criticisms of

oversplitting (e.g. Barley et al. 2018) lead us to not interpret MSC delimitation results as

evidence of species status. Most strikingly, reciprocal *gdi* statistics for Clade II were <0.2, thus

falling in the range suggested to unambiguously indicate a single species (Jackson et al. 2017;

Leaché et al. 2019; Fig. 3A). Finally, comparing genetic and geographic distances within Clade

II showed that a single isolation-by-distance pattern fits both intra- and interspecific comparisons

(Fig. 4). While the range of *M. lehilahytsara* expands considerably further south than the

populations examined here, our results strongly suggest that *M. mittermeieri* and *M.*

lehilahytsara are best considered a single species. Sampling gaps are expected to cause false

positive species delimitations rather than false negatives (Barley et al. 2018; Chambers and Hillis

2020; Mason et al. 2020), therefore additional sampling of *M. lehilahytsara* populations farther

south should not affect our recommendation to synonymize *M. mittermeieri* as *M. lehilahytsara*.

Mitonuclear Discordance and Gene Flow

Mitonuclear discordance was observed for a subset of *M. macarthurii* individuals from

Anjiahely. These individuals carried mtDNA similar to that of *M.* sp. #3 (see Radespiel et al.

2008) but had nDNA indistinguishable from sympatric *M. macarthurii*. Although genealogical

discordance could be due to incomplete lineage sorting (e.g., Heckman et al. 2007; Weisrock et

al. 2010), mitochondrial introgression is supported by D-statistics (Fig. S15) and the inferred

low levels of gene flow from the northern *M.* sp. #3 population into *M. macarthurii* by G-PhoCS

(Fig. S13). Besides a possible case in Sgarlata et al. (2019), of mitochondrial introgression has

not previously been reported in mouse lemurs. Somewhat curiously, the discovery of a divergent

mtDNA lineage at Anjiahely (Radespiel et al. 2008), which prompted the current work, was

apparently the result of mtDNA introgression from an undescribed species into its sister species.

Population Size and Species Delimitation

The comparison of effective population sizes in Clades I and II reveals that they are

markedly different, which can affect species delimitation tests such as *gdi* (Leaché et al., 2019).

The *gdi* is calculated using population sizes and divergence times estimated under models with

no gene flow, and since divergence time estimates in these models were highly similar in both

clades (Fig. 5), differences in effective population sizes also appear to play a role in the stark

difference in *gdi*. Indeed, *gdi* aims to quantify the probability that two sequences from the focal

taxon coalesce more recently than the divergence time between the taxa, and larger effective

population sizes result in slower sorting of ancestral polymorphisms (Maddison 1997).

Assessing "progress" in speciation by quantifying rates of neutral coalescence, however,

implies that the magnitude of genetic drift is a good predictor of species limits. At least when

considering reproductive isolation (i.e., biological species), this can be problematic, given that

the role of drift in speciation is generally thought to be small (Rice and Hostert 1993; Coyne and

Orr 2004; Czekanski-Moir and Rundell 2019; but see Uyeda et al. 2009). Therefore, additional

measures of divergence should be taken into account, including those that do not depend on

population size, such as rates of gene flow and divergence time itself (Yang and Rannala 2010;

Leaché et al. 2019).

Sympatric Occurrence and the Tempo of Speciation in Mouse Lemurs

Sympatric *Microcebus* species were found at two study sites, with a representative of

each of the two focal clades in Anjiahely (*M. macarthurii* and *M. mittermeieri*) as well as in

Ambavala (*M.* sp. #3 and *M. lehilahytsara*; `Fig. 1`). These cases of sympatric occurrence, with

no evidence for recent admixture, imply that the two clades are reproductively isolated. Though

our methods cannot address the mechanisms underlying reproductive isolation, possible barriers

include male advertisement calls, which tend to differ strongly among species (Braune et al.

2008), and timing of reproduction, which has previously been found to differ among sympatric

mouse lemur species (Schmelting et al. 2000; Evasoa et al. 2018) including the focal species

(Schüßler et al., in revision). Only six other cases of sympatry among mouse lemur species are

known, five of which include *M. murinus* as one of the co-occurring species (Radespiel 2016;

Sgarlata et al. 2019).

Given that the sympatrically occurring species were estimated to have had a common

ancestor as recently as ~700-800 ka ago (i.e., the divergence time between Clade I and Clade II,

see `Fig. 5`), this suggests rapid evolution of reproductive isolation and a short time to sympatry

among mouse lemurs. By comparison, Pigot & Tobias (2015) estimated that after 5 Ma of

divergence, only 21–23% of primate species have attained sympatry. In fact, the one sympatric

pair within their dataset of 74 sister species pairs younger than 2.5 million years consisted of the

sympatric *Galago gallarum* and *G. senegalensis* (Pigot and Tobias 2015), which are also

Strepsirrhini. Moreover, Curnoe et al. (2006) compiled data for naturally hybridizing primate

species, and found the median estimated divergence time to be 2.9 Ma. More broadly, primate

speciation rates do not appear to be lower than those for other mammals or even vertebrates

(Curnoe et al. 2006, Upham et al. 2019). It should be noted, however, that the temporal estimates

reported in our study are based on MSC analyses using mutation rates estimated from pedigree

studies, whereas dates for other primate clades were largely calculated from fossil-calibrated

relaxed clock methods.

Complexities of Divergence Time Estimates

There are two noteworthy discrepancies in divergence time estimates highlighted by this

study. First, the age estimate between the *M. mittermeieri* and *M. lehilahytsara* lineages

increased from approximately 100 kya (`Fig. 5b`) to more than 500 kya (`Fig. 5c`) when the

MSC model allowed for gene flow. The substantial effect of incorporating or disregarding gene

flow on divergence time estimation has been previously noted (Leaché et al. 2014; Tseng et al.

2014) and we here reiterate its significance. Second, the coalescent-based estimates of

divergence times presented here differ drastically from estimates based on fossil-calibrated

relaxed-clock methods. In the present study, we estimated the mean age of the most recent

common ancestor (MRCA) of mouse lemurs to be under 1.5 Ma, with the highest upper bound of

95% HPDs across models at 2.40 Ma. This age estimate is in stark contrast to previous fossil-

calibrated estimates of 8 - 10 Ma (Yang and Yoder 2003; dos Reis et al. 2018).

Several factors likely contribute to this large difference. First, the MSC estimate uses a *de*

novo mutation rate sampled from a distribution based on available pedigree-based mutation rates

in primates, including mouse lemurs (Campbell et al., 2019). This rate is nearly two-fold higher

than the estimated substitution rate for *M. murinus* (dos Reis et al., 2018). Second, converting

coalescent units to absolute time also requires a generation time estimate. We attempted to

account for uncertainty in generation time by similarly drawing from a distribution based on

empirical parent age estimates (Zohdy et al., 2014; Radespiel et al., 2019) in mouse lemurs.

Thus, either overestimation of the mutation rate and/or underestimation of the generation time

would lead to divergence time estimates that are too recent. However, theoretical considerations

suggest that instead, mouse lemur divergence time estimates from fossil-calibrated clock models

are too old.

When incomplete lineage sorting is common, clock models that assume a single topology

underlies all loci can overestimate species divergences compared to MSC estimates that allow

gene trees to vary (Stange et al., 2018; Feng et al., 2020). This is likely to apply to mouse lemurs

given that high levels of incomplete lineage sorting have been previously documented (Heckman

et al., 2007; Weisrock et al., 2010; Hotaling et al., 2016). Moreover, due to the absence of a post-

K-Pg terrestrial fossil record for Madagascar, clock-model estimates of divergence times in

mouse lemurs have relied on fossil calibrations from the distantly-related African sister lineage

of lemurs, the Lorisiformes (Seiffert et al., 2003), as well as from anthropoid primates and other

mammals. This scenario - estimation of divergence times for younger, internal nodes with

calibrations placed on much older nodes - is expected to lead to overestimation of divergence

times (Angelis and dos Reis 2015). Therefore, it is likely that divergence times between mouse

lemur species have been overestimated by previous studies with fossil-calibrated clock models

(e.g. Yang and Yoder 2003; dos Reis et al., 2018), and we suggest that the mutation rate-

calibrated MSC divergence times presented here are more accurate.

Our estimates of divergence times imply that the entire mouse lemur radiation originated in

the Pleistocene, in turn suggesting that Pleistocene climatic oscillations represent a likely factor

leading to geographic isolation and subsequent genetic divergence. Periods of drought during

glacial maxima are hypothesized to have caused dramatic contraction of forest habitats (Burney

et al. 1997; Gasse and Van Campo 2001; Wilmé et al. 2006; Kiage and Liu 2016) and to isolation

of previously connected populations. Notably, the patterns of differentiation observed in this

study are consistent with the predictions of Wilmé et al. (2006) wherein Madagascar's river

drainage systems created high-elevation retreat-dispersal corridors during periods of climatic

oscillation. That is, whereas the lineages in Clade I (highly differentiated and low N_e) appear to

occur only in lowland forests, those in the Clade II (poorly differentiated and high N_e) occur at

both higher and lower elevations (Schüßler et al., in revision). Moreover, the Mananara river

runs between the fairly distinct northern and southern populations of *M*. sp. #3, further

emphasizing the potential of large rivers to act as phylogeographic barriers in lemurs (Martin

1972; Pastorini et al. 2003; Goodman and Ganzhorn 2004; Olivieri et al. 2007).

Population Size Dynamics

A long-term decline in population size was inferred for the lineage leading to *M*. sp. #3.

While changes in inferred N_e may be confounded by changes in population structure – especially

for single-population sequential Markovian coalescent (PSMC/MSMC) models that do not

explicitly consider population subdivision (Mazet et al. 2016; Chikhi et al. 2018) – we recovered

similar results in both MSMC and G-PhoCS analyses (Fig. 6A). This congruence is especially

persuasive given the underlying differences between the G-PhoCS and MSMC models and their

input data. Moreover, Markovian coalescent approaches have been shown to be robust to genome

assembly quality (Patton et al. 2019), yielding further confidence in the results. The inferred

decline and population subdivision of *M*. sp. #3 was initiated long before anthropogenic land use,

supporting the emerging consensus that human colonization in Madagascar alone does not

explain the occurrence of open habitats and isolated forest fragments (Quéméré et al. 2012;

Vorontsova et al. 2016; Yoder et al. 2016; Salmona et al. 2017, 2020; Hackel et al. 2018).

Conversely, results for the *M. mittermeieri* lineage do not indicate a declining N_e through time

(`Fig. 6B`). This latter result may well be a simple corollary of the evidence described above,

that this lineage is part of a single species complex represented by Clade II and thus occurs at

both higher and lower elevations in northeastern Madagascar.

Conclusions

We have shown that substantial mouse lemur diversity exists within a 130-km-wide stretch

in northeastern Madagascar, including two instances of sympatric occurrence between

representatives of two closely related clades. Within one of these clades, our integrative approach

indicates that the undescribed lineage *M.* sp. #3 represents a distinct species, while the two

named species in the other clade, *M. mittermeieri* and *M. lehilahytsara*, are better considered a

single, widespread species with significant population structure. Given that the original

description of *M. lehilahytsara* precedes that of *M. mittermeieri*, primate taxonomists should

synonymize the two as *M. lehilahytsara.*

The divergence times calculated here using pedigree-based mutation rate estimates with

the MSC are much younger than those of previous studies that used external fossil-based

calibrations with concatenated likelihood methods. The younger dates suggest rapid evolution of

reproductive isolation in mouse lemurs as well as a Pleistocene origin of the radiation, likely

following population isolation due to climatic oscillations. This departure from previous

hypotheses of mouse lemur antiquity emphasizes the need for future studies focused on resolving

discrepancies in divergence time estimates, both in mouse lemurs and in other recently evolved

organismal groups for which such comparisons have yet to be made.

ACKNOWLEDGMENTS

This study was conducted under the research permit No.

197/17/MEEF/SG/DGF/DSAP/SCB.Re (DS), 072/15/MEEMF/SG/DGF/DCB.SAP/SCB

(MBB), 137/13/MEF/SG/DGF/DCB.SAP/SCB (DWR), 175/14/MEF/SG/DGF/DCB.SAP/SCB

(A. Miller), kindly issued by the directeur du système des aires protégées, Antananarivo and the

regional authorities (Direction Régional de l'Environnement, de l'Ecologie et de Forêts). We are

endebted to J.H. Ratsimbazafy, N.V. Andriaholinirina, C. Misandeau, B. Le Pors and S.

Rasoloharijaona, for their help with administrative tasks, to A. Miller for sharing samples, and to

G. Besnard for facilitating this study. We thank our field assistants (T. Ralantoharijaona, I.

Sitrakarivo, C. Hanitriniaina and T. Ralantoharijaona), the Wildlife Conservation Society

Madagascar and the ADAFAM (Association Des Amis de la Forêt d'Ambodiriana-Manompana,

C. Misandeau in particular) for their valuable help during sample collection. We warmly thank

the many local guides and cooks for sharing their incomparable expertise and help in the field,

misaotra anareo jiaby.

Funding was granted by the Bauer Foundation and the Zempelin Foundation of the

"Deutsches Stiftungszentrum" under grant no. T237/22985/2012/kg and T0214/32083/2018/sm

to DS, Duke Tropical Conservation Initiative Grant to ADY, and Duke Lemur Center/SAVA

Conservation research funds to MBB, the School of Animal Biology at The University of

Western Australia to AM, the Fundação para a Ciência e a Tecnologia, Portugal (PTDC/BIA-

BEC/100176/2008, PTDC/BIA-BIC/4476/2012, and SFRH/BD/64875/2009), the Groupement

de Recherche International (GDRI) Biodiversité et développement durable – Madagascar, the

Laboratoire d'Excellence (LABEX) TULIP (ANR-10-LABX-41) and CEBA (ANR-10-LABX-

25-01, the Instituto Gulbenkian de Ciência, Portugal to LC and JS, the ERA-NET BiodivERsA

project: INFRAGECO (Inference, Fragmentation, Genomics, and Conservation, ANR-16-

EBI3-0014 & FCT-Biodiversa/0003/2015) the LIA BEEG-B (Laboratoire International Associé

– Bioinformatics, Ecology, Evolution, Genomics and Behaviour, CNRS) to LC and JS. Further

financial support came from the Institute of Zoology, University of Veterinary Medicine

Hannover and UR acknowledges the long-term support of the late Elke Zimmermann for her

research activities on Madagascar. The genomic data were generated with funds from NSF DEB-

1354610 to ADY and DWW and from the EDB Lab to JS. ADY also gratefully acknowledges

support from the John Simon Guggenheim Memorial Foundation and the Alexander von

Humboldt Foundation. EELJ would like to acknowledge support from the Ahmanson Foundation

for the data generation. This work was performed in collaboration with the GeT core facility,

Toulouse, France (http://get.genotoul.fr), and was supported by France Génomique National

infrastructure, funded as part of "Investissement d'avenir" program managed by Agence

Nationale pour la Recherche (contract ANR-10-INBS-09). JS, UR & LC also gratefully

acknowledge support from the Get-Plage sequencing and Genotoul bioinformatics

(BioinfoGenotoul) platforms Toulouse Midi-Pyrenees. This is DLC publication #XXXX.

REFERENCES

Alexander D.H., Novembre J., Lange K. 2009. Fast model-based estimation of ancestry in unrelated
individuals. Genome Res. 19:1655–1664.

Ali O.A., O'Rourke S.M., Amish S.J., Meek M.H., Luikart G., Jeffres C., Miller M.R. 2016. RAD capture
(Rapture): Flexible and efficient sequence-based genotyping. Genetics. 202:389–400.

Angelis K., Dos Reis M. 2015. The impact of ancestral population size and incomplete lineage sorting on
Bayesian estimation of species divergence times. Curr. Zool. 61:874–885.

Barley A.J., Brown J.M., Thomson R.C. 2018. Impact of model violations on the inference of species
boundaries under the multispecies coalescent. Syst. Biol. 67:269–284.

Beaumont M.A. 2004. Recent developments in genetic data analysis: what can they tell us about human
demographic history? Heredity. 92:365–379.

Bickford D., Lohman D.J., Sodhi N.S., Ng P.K.L., Meier R., Winker K., Ingram K.K., Das I. 2007.
Cryptic species as a window on diversity and conservation. Trends Ecol. Evol. 22:148–155.

Blanco M.B., Rasoazanabary E., Godfrey L.R. 2015. Unpredictable environments, opportunistic
responses: Reproduction and population turnover in two wild mouse lemur species (Microcebus
rufus and M. griseorufus) from eastern and western Madagascar. Am. J. Primatol. 77:936–947.

Boetzer M., Henkel C.V., Jansen H.J., Butler D., Pirovano W. 2011. Scaffolding pre-assembled contigs
using SSPACE. Bioinformatics. 27:578–579.

Braune P., Schmidt S., Zimmermann E. 2008. Acoustic divergence in the communication of cryptic
species of nocturnal primates (*Microcebus* ssp.). BMC Biology. 6:19.

Brown J.L., Yoder A.D. 2015. Shifting ranges and conservation challenges for lemurs in the face of
climate change. Ecology and Evolution. 5:1131–1142.

Bryant D., Bouckaert R., Felsenstein J., Rosenberg N.A., RoyChoudhury A. 2012. Inferring species trees
directly from biallelic genetic markers: bypassing gene trees in a full coalescent analysis. Mol.
Biol. Evol. 29:1917–1932.

Burney D., James H., Grady F., Rafamantanantsoa J.-G., Ramilisonina, Wright H., Cowart J. 1997.
Environmental change, extinction and human activity: evidence from caves in NW Madagascar.
Journal of Biogeography. 24:755–767.

Campbell C.R., Tiley G.P., Poelstra J.W., Hunnicutt K.E., Larsen P.A., Reis M. dos, Yoder A.D. 2019.
Pedigree-based measurement of the de novo mutation rate in the gray mouse lemur reveals a high
mutation rate, few mutations in CpG sites, and a weak sex bias. bioRxiv.:724880.

Carstens B.C., Dewey T.A. 2010. Species delimitation using a combined coalescent and information-
theoretic approach: an example from North American *Myotis* bats. Syst. Biol. 59:400–414.

Chambers E.A., Hillis D.M. 2020. The multispecies coalescent over-splits species in the case of
geographically widespread taxa. Syst. Biol. 69:184–193.

Chifman J., Kubatko L. 2014. Quartet inference from SNP data under the coalescent model.
Bioinformatics. 30:3317–3324.

Chikhi L., Rodríguez W., Grusea S., Santos P., Boitard S., Mazet O. 2018. The IICR (inverse
instantaneous coalescence rate) as a summary of genomic diversity: insights into demographic
inference and model choice. Heredity. 120:13–24.

Chikhi L., Sousa V.C., Luisi P., Goossens B., Beaumont M.A. 2010. The confounding effects of
population structure, genetic diversity and the sampling scheme on the detection and
quantification of population size change. Genetics. 186:983–995.

Curnoe D., Thorne A., Coate J.A. 2006. Timing and tempo of primate speciation. J. Evol. Biol. 19:59–65.

Czekanski-Moir J.E., Rundell R.J. 2019. The ecology of nonecological speciation and nonadaptive
radiations. Trends Ecol. Evol. 34:400–415.

Dalquen D.A., Zhu T., Yang Z. 2017. Maximum likelihood implementation of an isolation-with-migration
model for three species. Syst. Biol. 66:379–398.

Dávalos L.M., Russell A.L. 2014. Sex-biased dispersal produces high error rates in mitochondrial
distance-based and tree-based species delimitation. J. Mammal. 95:781–791.

DePristo M.A., Banks E., Poplin R., Garimella K.V., Maguire J.R., Hartl C., Philippakis A.A., del Angel
G., Rivas M.A., Hanna M., McKenna A., Fennell T.J., Kernytsky A.M., Sivachenko A.Y.,
Cibulskis K., Gabriel S.B., Altshuler D., Daly M.J. 2011. A framework for variation discovery
and genotyping using next-generation DNA sequencing data. Nature Genetics. 43:491–498.

Dincă V., Lee K.M., Vila R., Mutanen M. 2019. The conundrum of species delimitation: a genomic
perspective on a mitogenetically super-variable butterfly. Proc. R. Soc. Lond. B Biol. Sci.
286:20191311.

dos Reis M., Gunnell G.F., Barba-Montoya J., Wilkins A., Yang Z., Yoder A.D. 2018. Using
phylogenomic data to explore the effects of relaxed clocks and calibration strategies on
divergence time estimation: primates as a test case. Syst. Biol. 67:594–615.

Du Puy D.J., Moat J. 1998. Vegetation mapping and classification in Madagascar (using GIS):
implications and recommendations for the conservation of biodiversity. In: Huxley C., Lock J.,
Cutler D., editors. Chorology, Taxonomy and Ecology of the Floras of Africa and Madagascar.
Kew: Royal Botanical Gardens. p. 97–117.

Edwards D.L., Knowles L.L. 2014. Species detection and individual assignment in species delimitation:
can integrative data increase efficacy? Proc. R. Soc. Lond. B Biol. Sci. 281:20132765.

Eriksson A., Manica A. 2012. Effect of ancient population structure on the degree of polymorphism
687   shared between modern human populations and ancient hominins. Proc. Natl. Acad. Sci. U.S.A.
688   109:13956–13960.

689   Estrada A., Garber P.A., Rylands A.B., Roos C., Fernandez-Duque E., Fiore A.D., Nekaris K.A.-I.,
690   Nijman V., Heymann E.W., Lambert J.E., Rovero F., Barelli C., Setchell J.M., Gillespie T.R.,
691   Mittermeier R.A., Arregoitia L.V., Guinea M. de, Gouveia S., Dobrovolski R., Shanee S., Shanee
692   N., Boyle S.A., Fuentes A., MacKinnon K.C., Amato K.R., Meyer A.L.S., Wich S., Sussman
693   R.W., Pan R., Kone I., Li B. 2017. Impending extinction crisis of the world's primates: Why
694   primates matter. Science Advances. 3:e1600946.

695   Feder J.L., Egan S.P., Nosil P. 2012. The genomics of speciation-with-gene-flow. Trends in Genetics.
696   28:342–350.

697   Feng B, Merilä J, Matschiner M, Momigliano P. 2020. Estimating uncertainty in divergence times among
698   three-spined stickleback clades using the multispecies coalescent. Mol Phylogent Evol.
699   142:106646

700   Flouri T., Jiao X., Rannala B., Yang Z. 2018. Species tree inference with BPP using genomic sequences
701   and the multispecies coalescent. Mol. Biol. Evol. 35:2585–2593.

702   Fujita M.K., Leaché A.D., Burbrink F.T., McGuire J.A., Moritz C. 2012. Coalescent-based species
703   delimitation in an integrative taxonomy. Trends Ecol. Evol. 27:480–488.

704   Fumagalli M., Vieira F.G., Linderoth T., Nielsen R. 2014. ngsTools: methods for population genetics
705   analyses from next-generation sequencing data. Bioinformatics. 30:1486–1487.

706   Gasse F., Van Campo E. 2001. Late Quaternary environmental changes from a pollen and diatom record
707   in the southern tropics (Lake Tritrivakely, Madagascar). Palaeogeography, Palaeoclimatology,
708   Palaeoecology. 167:287–308.

Gavrilets S., Boake C.R. 1998. On the evolution of premating isolation after a founder event. Am. Nat.
152:706–716.

Gonen S., Bishop S.C., Houston R.D. 2015. Exploring the utility of cross-laboratory RAD-sequencing
datasets for phylogenetic analysis. BMC Research Notes. 8:299.

Goodman S.M., Benstead J.P. 2005. Updated estimates of biotic diversity and endemism for Madagascar.
Oryx. 39:73–77.

Goudet J. 2005. hierfstat, a package for r to compute and test hierarchical F-statistics. Mol. Ecol. Notes.
5:184–186.

Gronau I., Hubisz M.J., Gulko B., Danko C.G., Siepel A. 2011. Bayesian inference of ancient human
demography from individual genome sequences. Nat. Genet. 43:1031–1034.

Gruber B., Unmack P.J., Berry O.F., Georges A. 2018. dartr: An r package to facilitate analysis of SNP
data generated from reduced representation genome sequencing. Mol. Ecol. Res. 18:691–699.

Grummer J.A., Bryson R.W., Reeder T.W. 2014. Species delimitation using Bayes factors: simulations
and application to the *Sceloporus scalaris* species group (Squamata: Phrynosomatidae). Syst.
Biol. 63:119–133.

Hackel J., Vorontsova M.S., Nanjarisoa O.P., Hall R.C., Razanatsoa J., Malakasi P., Besnard G. 2018.
Grass diversification in Madagascar: In situ radiation of two large C3 shade clades and support
for a Miocene to Pliocene origin of C4 grassy biomes. Journal of Biogeography. 45:750–761.

Heckman K.L., Mariani C.L., Rasoloarison R., Yoder A.D. 2007. Multiple nuclear loci reveal patterns of
incomplete lineage sorting and complex species history within western mouse lemurs
(*Microcebus*). Mol. Phylogenet. Evol. 43:353–367.

Heller R., Chikhi L., Siegismund H.R. 2013. The confounding effect of population structure on Bayesian
skyline plot inferences of demographic history. PLoS ONE. 8:e62992.

Huang J.-P., Leavitt S.D., Lumbsch H.T. 2018. Testing the impact of effective population size on
speciation rates – a negative correlation or lack thereof in lichenized fungi. Scientific Reports.
8:1–6.

Hundsdoerfer A.K., Lee K.M., Kitching I.J., Mutanen M. 2019. Genome-wide SNP data reveal an
overestimation of species diversity in a group of hawkmoths. Genome Biol. Evol. 11:2136–2150.

Hunnicutt K.E., Tiley G.P., Williams R.C., Larsen P.A., Blanco M.B., Rasoloarison R.M., Campbell C.R.,
Zhu K., Weisrock D.W., Matsunami H., Yoder A.D. 2020. Comparative genomic analysis of the
pheromone receptor class 1 family (V1R) reveals extreme complexity in mouse lemurs (Genus,
Microcebus) and a chromosomal hotspot across mammals. Genome Biol. Evol. 12:3562–3579.

Jackson N.D., Carstens B.C., Morales A.E., O'Meara B.C. 2017. Species delimitation with gene flow.
Syst. Biol. 66:799–812.

Jombart T., Ahmed I. 2011. adegenet 1.3-1: new tools for the analysis of genome-wide SNP data.
Bioinformatics. 27:3070–3071.

Kappeler P.M., Rasoloarison R.M., Razafimanantsoa L., Walter L., Roos, C. 2005. Morphology,
behaviour and molecular evolution of giant mouse lemurs (Mirza spp.) Gray, 1870, with
description of a new species. Primate Rep. 71:3–26.

Kass R.E., Raftery A.E. 1995. Bayes Factors. J. Am. Stat. Assoc. 90:773–795.

Kiage L.M., Liu K. 2016. Late Quaternary paleoenvironmental changes in East Africa: a review of
multiproxy evidence from palynology, lake sediments, and associated records. Progress in
Physical Geography. 30:633–658.

Kim B.Y., Wei X., Fitz-Gibbon S., Lohmueller K.E., Ortego J., Gugger P.F., Sork V.L. 2018. RADseq data
reveal ancient, but not pervasive, introgression between Californian tree and scrub oak species
(*Quercus* sect. *Quercus*: *Fagaceae*). Mol. Ecol. 27:4556–4571.

Knaus B.J., Grünwald N.J. 2017. vcfr: a package to manipulate and visualize variant call format data in R.
Mol. Ecol. Res. 17:44–53.

Knoop S., Chikhi L., Salmona J. 2017. Mouse lemurs' and degraded habitat. bioRxiv.:216382.

Korneliussen T.S., Albrechtsen A., Nielsen R. 2014. ANGSD: Analysis of Next Generation Sequencing
Data. BMC Bioinformatics. 15:356.

Langmead B., Trapnell C., Pop M., Salzberg S.L. 2009. Ultrafast and memory-efficient alignment of short
DNA sequences to the human genome. Genome Biology. 10:R25.

Leaché A.D., Fujita M.K., Minin V.N., Bouckaert R.R. 2014. Species delimitation using genome-wide
SNP data. Syst. Biol. 63:534–542.

Leaché A.D., Zhu T., Rannala B., Yang Z. 2019. The spectre of too many species. Syst Biol. 68:168–181.

LeCompte E., Crouau-Roy B., Aujard F., Holota H., Murienne J. 2016. Complete mitochondrial genome
of the gray mouse lemur, *Microcebus murinus* (Primates, Cheirogaleidae). Mitochondrial DNA A
DNA Mapp. Seq. Anal. 27:3514–3516.

Linck E., Epperly K., Van Els P., Spellman G.M., Bryson R.W., McCormack J.E., Canales-Del-Castillo
R., Klicka J. 2019. Dense geographic and genomic sampling reveals paraphyly and a cryptic
lineage in a classic sibling species complex. Syst. Biol. 68:956–966.

Louis E.E., Coles M.S., Andriantompohavana R., Sommer J.A., Engberg S.E., Zaonarivelo J.R., Mayor
M.I., Brenneman R.A. 2006. Revision of the mouse lemurs (*Microcebus*) of eastern Madagascar.
Int. J. Primatol. 27:347–389.

Louis E.E.Jr., Engberg S.E., McGuire S.M., McCormick M.J., Randriamampionona R., Ranaivoarisoa
J.F., Bailey C.A., Mittermeier R.A., Lei R. 2008. Revision of the mouse lemurs, *Microcebus*
(Primates, Lemuriformes), of northern and northwestern Madagascar with descriptions of two

new species at Montagne d'Ambre National Park and Antafondro Classified Forest. Primate

Conservation. 23:19–38.

Louis E.E.Jr., Lei R. 2016. Mitogenomics of the family Cheirogaleidae and relationships to taxonomy and

biogeography in Madagascar. In: Lehman S., Radespiel U., Zimmermann E., editors. The Dwarf

and Mouse Lemurs of Madagascar: Biology, Behavior and Conservation Biogeography of the

Cheirogaleidae. Cambridge, UK: Cambridge University Press. p. 54–93.

Lozier J.D. 2014. Revisiting comparisons of genetic diversity in stable and declining species: assessing

genome-wide polymorphism in North American bumble bees using RAD sequencing. Mol. Ecol.

23:788–801.

Luo A., Ling C., Ho S.Y.W., Zhu C.-D. 2018. Comparison of methods for molecular species delimitation

across a range of speciation scenarios. Syst. Biol. 67:830–846.

Maddison W.P. 1997. Gene trees in species trees. Syst. Biol. 46:523–536.

Mallet J., Besansky N., Hahn M.W. 2016. How reticulated are species? Bioessays. 38:140–149.

Markolf M., Brameier M., Kappeler P.M. 2011. On species delimitation: Yet another lemur species or just

genetic variation? BMC Evol. Biol. 11:216.

Mason NA, Fletcher NK, Gill BA, Funk WC, Zamudio KR. 2020. Coalescent-based species delimitation

is sensitive to geographic sampling and isolation by distance. Systematics and Biodiversity,

18:269-280.

Matute D.R. 2013. The role of founder effects on the evolution of reproductive isolation. J. Evol. Biol.

26:2299–2311.

Mazet O., Rodríguez W., Grusea S., Boitard S., Chikhi L. 2016. On the importance of being structured:

instantaneous coalescence rates and human evolution--lessons for ancestral population size

inference? Heredity. 116:362–371.

McLaughlin J.F., Winker K. 2020. An empirical examination of sample size effects on population

demographic estimates in birds using single nucleotide polymorphism (SNP) data.

bioRxiv.:2020.03.10.986463.

Mittermeier R. A., Louis E. E. Jr., Richardson M., Schwitzer C. , Langrand O., Rylands A. B., Hawkins F.,

Rajaobelina S., Ratsimbazafy J. , Rasoloarison R. , Roos C. , Kappeler P. M., Mackinnon J..

2010. Lemurs of Madagascar, 3rd Edition. Conservation International Tropical Field Guide Series,

Washington, USA.

Myers N., Mittermeier R.A., Mittermeier C.G., da Fonseca G.A.B., Kent J. 2000. Biodiversity hotspots

for conservation priorities. Nature. 403:853–858.

Nielsen R., Korneliussen T., Albrechtsen A., Li Y., Wang J. 2012. SNP calling, genotype calling, and

sample allele frequency estimation from new-generation sequencing data. PLoS ONE. 7:e37558.

O'Leary S.J., Puritz J.B., Willis S.C., Hollenbeck C.M., Portnoy D.S. 2018. These aren't the loci you're

looking for: Principles of effective SNP filtering for molecular ecologists. Mol. Ecol. 27:3193–

3206.

Olave M, Sola E, Knowles LL. 2014. Upstream analyses create problems with DNA-based species

delimitation. Syst Biol, 63:263-271.

Olivieri G, Zimmermann E, Randrianambinina B, Rasoloharijaona S, Rakotondravony D, Guschanski K,

Radespiel U. 2007. The ever-increasing diversity in mouse lemurs: Three new species in north

and northwestern Madagascar. Mol. Phy. Evol., 43:309-327.

Palkopoulou E., Lipson M., Mallick S., Nielsen S., Rohland N., Baleka S., Karpinski E., Ivancevic A.M.,

To T.-H., Kortschak R.D., Raison J.M., Qu Z., Chin T.-J., Alt K.W., Claesson S., Dalén L.,

MacPhee R.D.E., Meller H., Roca A.L., Ryder O.A., Heiman D., Young S., Breen M., Williams

C., Aken B.L., Ruffier M., Karlsson E., Johnson J., Palma F.D., Alfoldi J., Adelson D.L., Mailund

T., Munch K., Lindblad-Toh K., Hofreiter M., Poinar H., Reich D. 2018. A comprehensive
genomic history of extinct and living elephants. Proc. Natl. Acad. Sci. U.S.A. 115:E2566–E2574.

Pamilo P., Nei M. 1988. Relationships between gene trees and species trees. Mol. Biol. Evol. 5:568–583.

Patton AH, Margres MJ, Stahlke AR, Hendricks S, Lewallen K, Hamede RK, Ruiz-Aravena M, Ryder O,
McCallum HI, Jones ME, et al. 2019. Contemporary Demographic Reconstruction Methods Are
Robust to Genome Assembly Quality: A Case Study in Tasmanian Devils. Mol Biol Evol,
36:2906-2921.

Pastorini J., Thalmann U., Martin R.D. 2003. A molecular approach to comparative phylogeography of
extant Malagasy lemurs. Proc. Natl. Acad. Sci. U.S.A. 100:5879–5884.

Patterson N., Moorjani P., Luo Y., Mallick S., Rohland N., Zhan Y., Genschoreck T., Webster T., Reich D.
2012. Ancient admixture in human history. Genetics. 192:1065–1093.

Pedersen C.-E.T., Albrechtsen A., Etter P.D., Johnson E.A., Orlando L., Chikhi L., Siegismund H.R.,
Heller R. 2018. A southern African origin and cryptic structure in the highly mobile plains zebra.
Nat. Ecol. Evol. 2:491–498.

Pigot A.L., Tobias J.A. 2015. Dispersal and the transition to sympatry in vertebrates. Proc. R. Soc. Lond.
B Biol. Sci. 282:20141929.

de Queiroz K. 2007. Species concepts and species delimitation. Syst. Biol. 56:879–886.

Quéméré E., Amelot X., Pierson J., Crouau-Roy B., Chikhi L. 2012. Genetic data suggest a natural
prehuman origin of open habitats in northern Madagascar and question the deforestation narrative
in this region. Proc. Natl. Acad. Sci. U.S.A. 109:13028–13033.

R Core Development Team. 2013. R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing. Vienna, Austria.

Radespiel U. 2016. Can behavioral ecology help to understand the divergent geographic range sizes of
846   mouse lemurs? In: Lehman S., Radespiel U., Zimmermann E., editors. The Dwarf and Mouse
847   Lemurs of Madagascar: Biology, Behavior and Conservation Biogeography of the
848   Cheirogaleidae. Cambridge, UK: Cambridge University Press. p. 498–519.

849   Radespiel U., Lutermann H., Schmelting B., Zimmermann E. 2019. An empirical estimate of the
850   generation time of mouse lemurs. Am. J. Primatol. 81:e23062.

851   Radespiel U., Olivieri G., Rasolofoson D.W., Rakotondratsimba G., Rakotonirainy O., Rasoloharijaona
852   S., Randrianambinina B., Ratsimbazafy J.H., Ratelolahy F., Randriamboavonjy T.,
853   Rasolofoharivelo T., Craul M., Rakotozafy L., Randrianarison R.M. 2008. Exceptional diversity
854   of mouse lemurs (*Microcebus* spp.) in the Makira region with the description of one new species.
855   Am. J. Primatol. 70:1033–1046.

856   Radespiel U., Sarikaya Z., Zimmermann E., Bruford M.W. 2001. Sociogenetic structure in a free-living
857   nocturnal primate population: sex-specific differences in the grey mouse lemur (*Microcebus*
858   *murinus*). Behav. Ecol. Sociobiol. 50:493–502.

859   Rannala B., Yang Z. 2003. Bayes estimation of species divergence times and ancestral population sizes
860   using DNA sequences from multiple loci. Genetics. 164:1645–1656.

861   Rannala B., Yang Z. 2013. Improved reversible jump algorithms for Bayesian species delimitation.
862   Genetics. 194:245–253.

863   Rasolooarison R.M., Goodman S.M., Ganzhorn J.U. 2000. Taxonomic revision of mouse lemurs
864   (*Microcebus*) in the western portions of Madagascar. Int. J. Primatol. 21:963–1019.

865   Rice W.R., Hostert E.E. 1993. Laboratory experiments on speciation: what have we learned in 40 years?
866   Evolution. 47:1637–1653.

Evasoa M., Radespiel U., Hasiniaina A.F., Rasoloharijaona S., Randrianambinina B., Rakotondravony R.,
Zimmermann E. 2018. Variation in reproduction of the smallest-bodied primate radiation, the
mouse lemurs (*Microcebus* spp.): A synopsis. Am. J. Primatol. 80:e22874.

Rodríguez W., Mazet O., Grusea S., Arredondo A., Corujo J.M., Boitard S., Chikhi L. 2018. The IICR and
the non-stationary structured coalescent: towards demographic inference with arbitrary changes in
population structure. Heredity. 121:663–678.

Salmona J., Heller R., Quéméré E., Chikhi L. 2017. Climate change and human colonization triggered
habitat loss and fragmentation in Madagascar. Mol. Ecol. 26:5203–5222.

Salmona J., Olofsson J.K., Hong-Wa C., Razanatsoa J., Rakotonasolo F., Ralimanana H.,
Randriamboavonjy T., Suescun U., Vorontsova M.S., Besnard G. 2020. Late Miocene origin and
recent population collapse of the Malagasy savanna olive tree (*Noronhia lowryi*). Biol. J. Linn.
Soc. 129:227–243.

Schmelting B., Ehresmann P., Lutermann H., Randrianambinina B., Zimmermann, E. 2000. Reproduction
of two sympatric mouse lemur species (*Microcebus murinus* and *M. ravelobensis*) in northwest
Madagascar: first results of a long term study. In: Lourenço W.R., Goodman S.M. editors.
Diversité et Endémisme à Madagascar. Paris: Société de Biogéographie. p. 165-175.

Schüßler D., Mantilla-Contreras J., Stadtmann R., Ratsimbazafy J.H., Radespiel U. 2020. Identification of
crucial stepping stone habitats for biodiversity conservation in northeastern Madagascar using
remote sensing and comparative predictive modeling. Biodivers. Conserv.
https://doi.org/10.1007/s10531-020-01965-z.

Seiffert ER, Simons EL, Attia Y. 2003. Fossil evidence for an ancient divergence of lorises and galagos.
Nature, 422:421-424.

Sgarlata G.M., Salmona J., Pors B.L., Rasolondraibe E., Jan F., Ralantoharijaona T., Rakotonanahary A.,
Randriamaroson J., Marques A.J., Aleixo-Pais I., Zoeten T. de, Ousseni D.S.A., Knoop S.B.,

Teixeira H., Gabillaud V., Miller A., Ibouroi M.T., Rasoloharijaona S., Zaonarivelo J.R.,
892   Andriaholinirina N.V., Chikhi L. 2019. Genetic and morphological diversity of mouse lemurs
893   (*Microcebus* spp.) in northern Madagascar: The discovery of a putative new species? Am. J.
894   Primatol. 81:e23070.

895   Skotte L., Korneliussen T.S., Albrechtsen A. 2013. Estimating individual admixture proportions from next
896   generation sequencing data. Genetics. 195:693–702.

897   Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
898   phylogenies. Bioinformatics. 30:1312–1313.

899   Stange M, Sánchez-Villagra MR, Salzburger W, Matschiner M. 2018. Bayesian divergence-time
900   estimation with genome-wide single-nucleotide polymorphism data of sea catfishes (Ariidae)
901   supports Miocene closure of the Panamanian Isthmus. Syst Biol. 67:681-699.

902   Sukumaran J., Knowles L.L. 2017. Multispecies coalescent delimits structure, not species. Proc. Natl.
903   Acad. Sci. U.S.A. 114:1607–1612.

904   Tattersall I. 2007. Madagascar's lemurs: Cryptic diversity or taxonomic inflation? Evolutionary
905   Anthropology: Issues, News, and Reviews. 16:12–23.

906   Tseng S.-P., Li S.-H., Hsieh C.-H., Wang H.-Y., Lin S.-M. 2014. Influence of gene flow on divergence
907   dating - implications for the speciation history of *Takydromus* grass lizards. Mol. Ecol. 23:4770–
908   4784.

909   Upham N.S., Esselstyn J.A., Jetz W. 2019. Inferring the mammal tree: Species-level sets of phylogenies
910   for questions in ecology, evolution, and conservation. PLOS Biology 17(12): e3000494.

911   Uyeda J.C., Arnold S.J., Hohenlohe P.A., Mead L.S. 2009. Drift promotes speciation by sexual selection.
912   Evolution. 63:583–594.

Vorontsova M.S., Besnard G., Forest F., Malakasi P., Moat J., Clayton W.D., Ficinski P., Savva G.M.,
Nanjarisoa O.P., Razanatsoa J., Randriatsara F.O., Kimeu J.M., Luke W.R.Q., Kayombo C.,
Linder H.P. 2016. Madagascar's grasses and grasslands: anthropogenic or natural? Proc. R. Soc.
Lond. B Biol. Sci. 283:20152262.

Wang K., Mathieson I., O'Connell J., Schiffels S. 2020. Tracking human population structure through
time from whole genome sequences. PLoS Genetics. 16:e1008552.

Warmuth V.M., Ellegren H. 2019. Genotype-free estimation of allele frequencies reduces bias and
improves demographic inference from RADSeq data. Mol. Ecol. Res. 19:586–596.

Weisrock D.W., Rasoloarison R.M., Fiorentino I., Ralison J.M., Goodman S.M., Kappeler P.M., Yoder
A.D. 2010. Delimiting species without nuclear monophyly in Madagascar's mouse lemurs. PLoS
ONE. 5:e9883.

Wen D., Nakhleh L., Kubatko L. 2018. Coestimating reticulate phylogenies and gene trees from
multilocus sequence data. Syst. Biol. 67:439–457.

Wilmé L., Goodman S.M., Ganzhorn J.U. 2006. Biogeographic evolution of Madagascar's microendemic
biota. Science. 312:1063–1065.

Yang Z., Rannala B. 2010. Bayesian species delimitation using multilocus sequence data. Proc. Natl.
Acad. Sci. U.S.A. 107:9264–9269.

Yang Z., Yoder A.D. 2003. Comparison of likelihood and Bayesian methods for estimating divergence
times using multiple gene Loci and calibration points, with application to a radiation of cute-
looking mouse lemur species. Syst. Biol. 52:705–716.

Yoder A.D., Campbell C.R., Blanco M.B., Reis M. dos, Ganzhorn J.U., Goodman S.M., Hunnicutt K.E.,
Larsen P.A., Kappeler P.M., Rasoloarison R.M., Ralison J.M., Swofford D.L., Weisrock D.W.

2016. Geogenetic patterns in mouse lemurs (genus *Microcebus*) reveal the ghosts of
936   Madagascar's forests past. Proc. Natl. Acad. Sci. U.S.A. 113:8049–8056.

937   Yoder A.D., Rasoloarison R.M., Goodman S.M., Irwin J.A., Atsalis S., Ravosa M.J., Ganzhorn J.U. 2000.
938   Remarkable species diversity in Malagasy mouse lemurs (primates, *Microcebus*). Proc. Natl.
939   Acad. Sci. U.S.A. 97:11325–11330.

940   Zimin A.V., Marçais G., Puiu D., Roberts M., Salzberg S.L., Yorke J.A. 2013. The MaSuRCA genome
941   assembler. Bioinformatics. 29:2669–2677.

942   Zimmermann E., Cepok S., Rakotoarison N., Zietemann V., Radespiel U. 1998. Sympatric mouse lemurs
943   in north-west Madagascar: a new rufous mouse lemur species (*Microcebus ravelobensis*). Folia
944   Primatol. 69:106–114.

945   Zimmermann E., Radespiel U. 2014. Species concepts, diversity, and evolution in primates: Lessons to be
946   learned from mouse lemurs. Evolutionary Anthropology: Issues, News, and Reviews. 23:11–14.

947   Zohdy S., Gerber B.D., Tecot S., Blanco M.B., Winchester J.M., Wright P.C., Jernvall J. 2014. Teeth, sex,
948   and testosterone: aging in the world's smallest primate. PLoS ONE. 9:e109528.