Alejandro Sosnovsky[1*], Mailén Elizabeth Lallement[1], Magalí Rechencq[1], Eduardo Enrique Zattara[1], María

Valeria Fernández[1], María José Suárez [3], Sergio Leiva[2] and Rodolfo Juan Carlos Cantet[3]

Title: Topography and Land Use Modulate Hydrological and Nutrient Export Dynamics in

Andean Streams

Short Title: Hydrology and Nutrient exports in Andean streams

[1]Grupo de Evaluación y Manejo de Recursos Ícticos, INIBIOMA, CONICET-Universidad Nacional del

Comahue, Quintral 1250, R8400FRF San Carlos de Bariloche, Argentina

[2] Grupo de Nuevos Materiales y Dispositivos, Comisión Nacional de Energía Atómica Av. Ezequiel Bustillo Km

9.5 (8400) San Carlos de Bariloche, Argentina

[3]Departamento de Producción Animal, Facultad de Agronomía, Universidad de Buenos Aires, e INPA,

CONICET. Av. San Martín 4453, C1417DSQ Ciudad Autónoma de Buenos Aires, Argentina

[*] Corresponding autor e-mail address: alejandro.sosnovsky@comahue-conicet.gob.ar

Telephone number: +54 294 423374 (452), Fax number: 0054-2944-422111

ACKNOWLEDGMENTS

We thank Rio Negro's province Water Department and Argentina's National Weather Service for providing

hydrological and meteorological data; and D.S. Ahearn for his interesting comments during the data analysis and

writing periods. This study was funded by *Agencia Nacional de Promoción Científica y Tecnológica* (PICT

2959). Lallement, M. E. has a National Scientific and Technical Research Council (CONICET) postdoctoral

fellowship. Sosnovsky, A, Cantet, R.J., Rechenq, M., Fernandez, M.V. and Zattara E.E. are CONICET

researchers.

ABSTRACT

Waterways act as integrators of ecosystems, their dynamics driven by multiple environmental factors and scales.

To elucidate factors affecting hydrological and nutrient dynamics in Andean streams, we characterized two

adjacent North Patagonian streams with contrasting drainage basins: Casa de Piedra (CP), originating in a small

mountain lake and running through a pristine landscape, and Gutiérrez (G), originating in a large piedmont lake

and running through an anthropized landscape. The drainage basins share both temperature and precipitation

regimes; however, the streams presented contrasting hydrological dynamics and nutrient export values. CP had

higher discharge flashiness with shorter response delays to precipitation, while G showed more stable

hydrograms, a difference explained by buffering from a large upstream lake in G's basin. Streams showed

differences in the quality and timing of coarse particulate organic matter export and basal export levels of

phosphorus and nitrogen that could be explained by human activities affecting G but not CP. Moreover,

nitrogen:phosphorus ratio indicates a possible shift to phosphorus as the limiting nutrient in the future in G

drainage basin. In summary, our results show that even under a common climatic regime, dynamics of adjacent

basins can be strongly driven by topographic and land use factors.

INTRODUCTION:

River ecosystems have traditionally been studied as isolated entities; however, over time it has become clear that they are deeply intertwined with their surroundings (Likens 1985). Climate, topography, geology and land cover are environmental factors that control spatial patterns in these ecosystems through often hierarchical influence (Snelder & Biggs 2002). In other words, a river is the product of its landscape (Wiens 2002) and thus riverine hydrological, chemical and biological processes should be studied in relation to the river's overall drainage basin (Hynes 1975; Vannote et al. 1980).

Regional climate and drainage basin topography are factors acting at a macro and meso spatial scales upon the river ecosystem, strongly driving hydrological and chemical dynamics (Snelder & Biggs 2002). Climate directly drives the hydrological cycle, determining stream discharge, a key ecosystem variable. Discharge is fed by rain, snowmelt and groundwater, the relative importance of these factors varying over time and space (Brown et al. 2003). After a rain episode, surface and sub-surface run-off are favored at mountainous landscapes, while water infiltration and evapotranspiration are favored at highly vegetated lowlands with permeable soils. Snowfall at high altitudes contributes to formation of a snowpack, which will store water until the melt period, when this water flows down into a stream (Bailey 1995). Lakes are also topographic features of a drainage basin that can attenuate run-off response and sediment transport (Snelder & Biggs 2002). Thus, headwater lakes influence directly the hydrograph and sediment dynamic of downstream fluvial ecosystems (Gordon et al. 2004). These hydrographs tend to be stable over time, furthermore, considering lakes as sediment and nutrient traps, low values of nutrient concentration and export could be expected in these streams (Parker et al. 2009).

At a lower spatial scale, drainage basin traits also affect fluvial dynamics, since nutrients and contaminants generated at the basin eventually reach the river ecosystem (Allan 2004; Dodds & Smith 2016). Non-point source pollution, associated with changing land use patterns and practices, has resulted in increased impacts on water bodies (Jordan et al. 1997; Carpenter et al. 1998). For example, growth of urban areas favors nutrient export (Howarth et al. 1996; Meyer et al. 2005; Walsh et al. 2005). Moreover, direct dumping of sewage water to a stream greatly increases nutrient load (Pieterse et al. 2003) and can also change nutrient stoichiometry (Merseburger et al. 2005). Thus, increases of nutrient load leading to eutrophication of running waters and carbon dynamics largely depend on the drainage basin state.

In Northern Patagonia, climate and topography are largely determined by the Andes mountain range (Paruelo et al. 1998). According to the updated world map of the Köppen-Geiger climate classification, Northern

Patagonia is characterized by a temperate climate with warm and dry summers (Peel et al. 2007). The Andes

represent an important barrier for humid air masses brought from the Pacific Ocean by westerly winds. Most of

the humidity in these maritime air masses precipitates on the west side of the Andes as they are blown to higher,

colder altitudes across the mountain range. Upon crossing the Andes, the air descends along their east side and

becomes hotter and drier through adiabatic warming. East of the Andes, the amount of rainfall follows a steep

west-east gradient: annual precipitation drops from 3500 mm to 700 mm in less than 60 km. Moreover, the

precipitation regime shows marked seasonality, with most rain falling during fall and winter (Paruelo et al.

1998). Thus, in this mountain region, the hydrological regime of fluvial ecosystems can be divided into three

contrasting periods (Sosnovsky et al. 2019): 1) a storm period, in which stream discharge variability and water

turbidity are high due to high run-off and the interaction between land and river ecosystems; 2) a snowmelt

period, with highest stream discharge and electrical conductivity (EC) dilution values; and 3) a dry period, with

overall low stream discharge values and stronger influx from groundwater.

86   North Patagonian drainage basins along the East side of the Andes are in a relatively pristine state. In

this region, many river systems originate from high-altitude wetlands and small lakes, or large piedmont lakes.

Human population density is low, and native forest occupies the western, most humid zone while steppe

vegetation occupies the eastern zone (Mermoz & Martín 1987). Forests are dominated by the genus *Nothofagus*

(southern beeches) and Andean cypress species *Austrocedrus chilensis*, but other deciduous and evergreen

species of high ecological and commercial value are present (Veblen et al. 1996). As in other South American

temperate forests, N deposition is remarkably low (Holland et al. 1999), and vegetation is limited mainly by N

(Diehl et al. 2003). Soils are classified as Andisols, which are characterized by high capacity to stabilize organic

matter, store water and retain phosphorous (Satti et al. 2003). They also have a high pH buffering capacity. Work

carried out in these drainage basins has revealed the oligotrophic character of the fluvial ecosystems (Pedrozo et

al. 1993). Diverse authors have studied anthropic (Miserendino et al. 2016) and natural (Lallement et al. 2016;

Williams Subiza & Brand 2018) elements that have affected Andean drainage basins, and consequently, their

fluvial ecosystems. Some of these effects are sustained on chemical and biological structure over time (Carrillo

et al. 2018). Recent research has revealed the dynamics of organic matter, which is intimately associated with the

stream-basin interaction (García et al. 2015b; Díaz Villanueva et al. 2016; García et al. 2018). Nevertheless,

studies focusing on stream discharge (Masiokas et al. 2008; Barros & Camilloni 2016; Sosnovsky et al. 2019)

and, nutrient load or export (Oyarzún et al. 1998; Oyarzún et al. 2004; Temporetti 2006) are scarce; despite the

hydrological, ecological and human importance of these variables. Even scarcer are studies where natural

processes and undisturbed nutrient cycling can be assessed in pristine ecosystems (Hedin et al. 1995) like the

ones can be found in North Patagonian region.

The main hypothesis of our study is that climate is the principal driver of stream ecosystems dynamics,

which are in turn modulated by topography and land use. As a consequence, we expect precipitation, air

temperature and stream discharge to be intimately connected. In turn, adjacent drainage basins with contrasting

topography and land use should present differences in their hydrological, physical and chemical dynamics. To

test this hypothesis, we selected two streams located in adjacent yet contrasting drainage basins: Casa de Piedra

(CP) and Gutiérrez (G). CP drainage basin is pristine, and the stream originates in a high-mountain lake. G

drainage basin suffers from significant anthropization, and the stream originates from a large piedmont lake. On

one hand, we expect to find buffered hydrological and chemical dynamics in G, relative to CP. On the other,

higher anthropization of G´s basin could impact the stream ecosystem, potentially causing elevated or

unpredictable variability in discharge regime along with increase nutrient concentration and export. Our first

goal is to compare discharge variability of CP and G, and study how discharge relates to climatic variables (basin

precipitation and air temperature). Our second goal is to determine the trophic state of these two streams, by

measuring basal concentration, export and stoichiometry of nitrogen (TN: total N) and phosphorous (TP: total P),

and estimate basal export of coarse particulate organic matter (CPOM, organic matter > 1mm). Our third

objective is to identify seasonal patterns of key physical and chemical variables.

MATERIALS Y METHODS

Study area

Research was conducted at the drainage basin of the streams Casa de Piedra (CP; 41° 07' 30.11" S 71° 27'

13.16"W) and Gutiérrez (G; 41° 09' 36.18" S 71° 24' 37.19"W), both located in Nahuel Huapi National Park,

Rio Negro, Argentina (Fig. 1). These drainage basins are very different from each other, despite of their

proximity (Table 1). CP's basin covers 63 km^2 and the stream originates in lake Jakob, a small (0.15 km^2), ultra-

oligotrophic (1-3 μg l^{-1} TP), high altitude water body situated at 1550 m a.s.l. (García et al. 2015a). CP's only

perennial affluent is the Rucaco stream, which originates at a high-altitude wetland. CP flows 19.7 km through a

steep V-shaped valley (average slope: 33.9 m km^{-1}) from most of its length. CP's drainage basin is mostly

pristine, with human activities restricted to hiking and infrequent camping; human population (less than 1000

inhabitants, 2010 National Census) is restricted to lowland areas. Forest below the tree line includes deciduous

Nothofagus pumilio and evergreens *N. dombeyi* and *Austrocedrus chilensis*. In contrast, G's drainage basin is

larger (160 km^2), and the stream originates in lake Gutiérrez, a deep (111 m maximum depth) and large (17 km^2)

water body that occupies ~11% of the basin's area. The seasonal concentration of TP is 3.4 µg l^{-1} on average

(Diaz et al. 2007). G's only perennial affluent is the Cascada stream, whose catchment covers 12.5 km^2 and

includes a major summer and winter sports resort. G flows 9 km through a wide, gently sloped valley (5.9 m km^{-1}

^{1}). Its riverbanks are extensively colonized by the exotic crack willow, *Salix fragilis*, and pass through a series of

populated areas (approximately 4000 inhabitants, 2010 National Census) belonging to the city of San Carlos de

Bariloche (approximately 108000 inhabitants, 2010 National Census). In addition, two trout farms with a total

production of ~3 tons year^{-1} use G's waters. In summary, CP's basin is largely free of impact from human

activities, while G's basin is not.

Fig. 1 a) Nahuel Huapi National Park; the study site is shaded in black. The city marker indicates San Carlos de

Bariloche downtown. b) Study area, Casa de Piedra and Gutiérrez drainage basins, Jakob lake and Gutiérrez lake

sub-catchments are also delimited. Curves represent 100 m slope on the terrain. Populated zones are shown in

grey, crosses indicate sampling sites, black circles indicate the Limnigraphic stations and white circle indicate

the rain meter location. R Str. and C Str., Rucaco and Cascada streams (modified from Sosnovsky et al. 2019).

Data collection

The present study was carried out over a period of one year, from March 2013 to March 2014. Data on rainfall

and discharge was kindly provided by the Rio Negro province's Water Department. Daily precipitation was

recorded with a rain meter (Fig. 1), and did not differentiate between rain and snow. Discharge data were

average daily records from limnigraphic stations. CP's limnigraphic station is located close to its mouth, while

G's station is located at its source (Fig. 1). Values of discharge at our G sampling site were estimated as

discharge as measured at the station plus discharge for the Cascada stream, determined using the Drainage-Area

Ratio method (Emerson et al. 2005); based on the specific daily values of discharge for the adjacent CP stream.

Specific discharge (Q^{sp}) was defined as discharge per unit area. Daily average air temperature records were

obtained from the National Weather Service.

	Casa de Piedra	Gutiérrez
Drainage area (km2)	63.9 (60.3)	160.4 (30.1)
Lake area (km2)	0.15	17.21
Native Forest (%)	60.2 (61.7)	61.1 (56.3)
Bare ground (%)	(27.7) 29.7	(9.5) 17.2
Exotic Forest (%)	(10.5) 9.9	10.8 (34.2)
Habitants [N]	< 1000 (< 1000)	4900 (4000)
Urban area [km2]	0.65 (0.65)	11.35 (5.08)

Table 1 Drainage basins characteristics. The values are expressed based on the entire area of the drainage basins,

values in brackets exclude the sub-catchment area of the Jakob lake and Gutiérrez lake.

Baseline levels of nutrients and relevant physical and chemical variables were obtained from monthly sampling

at both streams (Fig. 2). Water samples were taken to analyze total phosphorus (TP) and total nitrogen (TN)

concentrations. Water turbidity was measured with a Velp turbidimeter. Temperature, electrical conductivity

(EC) at 25 °C and pH were measured with an Oakton probe. Coarse particulate organic matter (CPOM) was

sampled with 2 drift nets (1-mm, 20 X 20-cm opening at the mouth) that were staked in the streambed and left to

collect material for intervals of 10 to 200 minutes, depending on the amount of CPOM present in the stream.

During October, high and fast stream discharge prevented us from placing Surber sampler nets at G, so we

performed linear interpolation for this value. Data were taken exclusively on days without precipitation.

Fig. 2 Temperature and precipitation during the study period. Black dots and numbers indicate sampling

dates

Data processing and analysis

Discharge

To understand the dynamics of discharge, we used an autocorrelation analysis (Mangin 1984) and calculated

Richard- Baker (R-B) index (Baker et al. 2004). The autocorrelation analysis examines how a value depends on

preceding values over a period of time. The correlations are computed for a given lag time k to obtain an

autocorrelation coefficient $r(k)$. For example, the correlation between today´s and yesterday´s stream discharges

would be a lag-1 autocorrelation coefficient, $r(1)$. The autocorrelation function is represented with a

correlogram, where slope is determined by the response of the system to an event. If the event has only a short-

term influence on the response of the discharge, the slope of the correlogram will decrease steeply. In contrast, if

the system is influenced by an event for a long time, the slope of the correlogram will decrease slowly. Generally

the length of the influence of an event is given by the "memory effect" which is according to Mangin (1984) the

lag number k when $r(k)$ reaches a value of 0.2. For this purpose, data on the average daily discharge were

transformed to normality by the Box-Cox procedure as suggested by Peltier et al. (1998). The formula for

autocorrelation is (Mangin 1984; Larocque et al. 1998):

$$r(k) \frac{C(k)}{C(O)}$$

with

$$C(k) = \frac{1}{n} \sum_{t=1}^{n-k} (x_t - \bar{x})(x_{t+k} - \bar{x})$$

Where k is the time lag and varies from 0 to n. According to Mangin (1984), n has to be taken as 1/3 of the

whole dataset to avoid stability problems.

The annual variability in stream discharge (flashiness) was measured by the R-B index as follows:

$$R - B \ Index = \frac{\sum_{i=n}^{n} |q_i - q_{i-1}|}{\sum_{i=1}^{n} q_i}$$

The variable *q* is the mean daily discharge whereas *n* is the number of observations. This index measures

oscillations in discharge relative to total discharge. Thus, it is a dimensionless measure ranging between 0 and 2

(Fongers et al. 2012). A value of 0 represents an absolutely constant discharge; increased R-B index values

indicate increased flashiness of streamflow. As such, the index appears to provide a useful characterization of the

way drainage basins process hydrological inputs into their streamflow outputs.

In addition, we studied the relationship between discharge, precipitation and temperature by cross-

correlation analysis. This analysis is widely used to analyze the linear relationship between input and output

signals in hydrology (Larocque et al. 1998). In this case, input signals were precipitation and temperature, and

output signal was river discharge of CP and G streams. Cross-correlations are represented by a cross-

correlogram. The maximum amplitude and the lag value of the cross-correlogram provide information on the

delay, which indicates the time of the pulse transfer to the stream. According to Larocque *et al.* (1998) the

formula for cross-correlation is:

$$r_{xy} = \frac{C_{xy}\ (k)}{\sigma_x \sigma_y}$$

211   with

$$C_{xy}(k) = \frac{1}{n} \sum_{t=1}^{n-k} (x_t - \bar{x})\,(y_{t+k} - \bar{y})$$

Where σ_x and σ_y are the standard deviation of the two times series.

Physical and chemical analysis

We estimated concentration and export of nutrients (TN and TP) and CPOM. TN was analyzed through digestion

and subsequent reduction in cadmium column (Grasshoff et al. 1983). TP was analyzed using acid digestion

followed by evaluation of solubilized phosphorus as soluble reactive phosphorus (A.P.H.A. 2005). We estimated

Ash Free Dry Weight of the CPOM using the Pozo et al. (2009) methodology. The annual export of TN, TP and

CPOM were calculated by the following formula:

$$Nutrient\ export = \frac{[N]\ Q\ t}{A}$$

Where $[N]$ is nutrient concentration, Q is discharge, t is time and A is drainage basin area.

We performed a paired T-test to compare nutrient concentration and export between streams; to this end the data

were transformed to comply with normality and homoscedasticity assumptions. Last, we performed a Principal

Component Analysis (PCA) that included 12 variables: turbidity, temperature, EC, pH, discharge, TP, TN, TN

TP^{-1}, CPOM, TP export, TN export and CPOM export. PCs were calculated based on standardized correlation

matrix, using components with eigenvalues larger than 1. Number Cruncher (Hintze 1998) and InfoStat (Di

Rienzo et al. 2016) where employed for statistical analysis, GraphPad Prism version 6.01 was employed for

graphics editing.

RESULTS

Climate and hydrology of the region

Air temperature and precipitation in the study area agreed with a temperate climate. Air temperature was low

between May and August and increased monotonically after September, reaching its highest values from

December to February. Annual precipitation was 1164 mm and showed strong seasonality, with 80% falling

between mid-May and mid-September. The first two (#1-2) and last three (#11-13) samples thus took place

during a drier period; samples #3-6 were taken during the precipitation period; the remaining samples (#7-10)

fell within the period of monotonic air temperature warming (Fig. 2). Thus, samples were evenly distributed

across three climatically contrasting periods.

The flow regime was contrasting between the streams. CP had a lower annual average discharge value

than G (Discharge $_{CP}$= 3.357 m^3/s; Discharge $_G$= 4.627 m^3/s; t =-7.4437; $P < 0.0001$; d.f. = 704), but its Q^{sp} was

higher (Q^{sp}_{CP}= 0.051 m^3/s; Q^{sp}_G= 0.029 m^3/s; t=28.9156; $P < 0.0001$; d.f. = 487). CP presented the most extreme

discharge values (Fig. 3a): the R-B index was 0.143 for CP, compared to 0.031 for G, indicating that the average

day-to-day fluctuations of streamflow were 14.3 % and 3.1 %, respectively. Furthermore, CP showed a steep

autocorrelation function with a memory effect at $k = 38$, while G revealed a more gently sloped function with a

memory effect at $k = 61$ (Fig 3b). Thus, CP showed higher flashiness and lower memory (i.e., temporal

persistence) than G.

Fig. 3 a) Hydrographs of Casa de Piedra (CP) and Gutiérrez (G) streams, black dots and numbers indicate sampling dates. b) Autocorrelation functions and memory effects (r (k) = 0.2) of the discharge. c) Cross-correlations between discharge and precipitation, three different periods are shown, and d) cross-correlations between discharge and temperature.

Cross-correlation of discharge with precipitation and temperature showed opposing patterns. Discharge was

positively correlated with precipitation, but negatively correlated with air temperature (Fig. 3c, d and Table S1).

The discharge-precipitation correlogram shows three periods that coincide with peaks of the correlation function

(Fig. 3c). Both streams show differences at each of these peaks: CP had higher correlation values (r_{max}) and

lower lag times than G. Indeed, at CP the correlation reached its maximum the same day precipitation took place

(lag = 0) during the first period, while r was not even significant for G at the time (Table S1). During the second

period, correlation at CP reached an r_{max}=0.36 with a lag time of 34 days, while at G, correlation was lower

(r_{max}=0.19) and the lag higher (lag = 40). The third, longest period presented a series of peaks and valleys, and

lapsed from day 57 to day 110 after the precipitation event at CP and from day 62 to 142 at G. In contrast, the

discharge-temperature correlogram did not show similar periods; cross-correlation was significant from day 0 to

around day 90 for both streams (Fig. 3d, Table S1).

Physical and chemical characteristics of the streams studied

Although streams differed in the specific values, both had very low EC, turbidity, nutrient concentration and nutrient export (Table 2). G showed the highest TP and TN concentrations (Fig. 4a, c) and exports (Fig. 4b, d): CP exported 8.9 kg TP km^{-2} $year^{-1}$ and 59.0 kg TN km^{-2} $year^{-1}$, while G exported 11.7 kg TP km^{-2} $year^{-1}$ and 78.8 kg TN km^{-2} $year^{-1}$ (approximately 35 % more N). No significant differences in TN:TP average ratios were found (~15; Fig. 4e), although the median value of TN:TP was 11 for CP and 16 for G. As to the amount of CPOM exported, no significant difference between streams was found. There were differences, however, in the quality of exported material; native *N. pumilio* leaves dominated in CP, while introduced *S. fragilis* leaves dominated in G. The peaks of CPOM export also differed among streams: highest export at CP was found during June and July (samples #3-4), while at G this peak occurred earlier, in April (sample #2). In summary, the drainage basin showing more anthropic land uses also showed higher nutrient export, with most CPOM export represented by introduced tree species.

	Casa de Piedra	Gutiérrez
Temperature (°C)	7.8 (3.7-12.5)	10.6 (6.0-16.8)
Electrical Conductivity (μs cm^{-1})	43 (27-61)	74 (71-76)
Turbidity (NTU)	0.2 (0.0-0.8)	0.7 (0.0-2.0)
pH	7.8 (7.4-8.0)	7.9 (7.3-8.1)
Total Phosphorus (mg m^{-3})	5 (2-9)	14 (10-18)
Total Nitrogen (mg m^{-3})	29 (18-44)	94 (75-134)
TN TP^{-1} (molar)	15 (6-31)	15 (11-21)
Coarse Particulate Organic Matter (AFDW m^{-3})*10	0.46 (0.04-1.86)	0.91 (0.02-8.76)
Total Phosphorus export (kg km^{-2} y^{-1})	8.9 (1.7-26.4)	11.7 (3.8-22.6)
Total Nitrogen export (kg km^{-2} y^{-1})	59.0 (8.2-134.2)	78.8 (27.4-186.0)
Coarse Particulate Organic Matter export (kg km^{-2} y^{-1})	131.3 (1.3-778.2)	43.3 (1.5-320.2)

Table 2 Average values, maximum and minimum in brackets, of the physical and chemical variables in the streams (n=13, except for the Coarse particulate organic matter where n=12).

Fig. 4 Box plot showing annual differences in nutrient concentration, ratio and export in Casa de Piedra (CP) and Gutiérrez (G) streams (Paired T Test, $*$ $P \le 0.05$; $**$ $P \le 0.01$; $***$ $P \le 0.001$; ns: not significant). f) The Y axis is in logarithmic scale. Total Phosphorus (TP), Total Nitrogen (TN) and Coarse Particulate Organic Matter (CPOM).

Our PCA determined three Principal Component, FI, FII and FIII that respectively explain 33%, 24% and 13% of the variance, accounting for 70% of the total variance (Table 3). FI was mainly related to discharge and nutrient export (TP_{exp} and TN_{exp}) (Fig. 5a). Positive FI values corresponded to samples with high discharge, high nutrient export and high turbidity. The FII was mainly related to the EC of the water, and to a lesser extent with water temperature and nutrient concentration (Fig. 5a). The FIII was mainly related to the TN:TP ratio, and to a lesser extent with CPOM variables.

Variable	First Factor (33%)	Second Factor (24%)	Third Factor (13%)
Temperature		0.31	
EC		0.47	
Turbidity	0.35		
Q	0.38		
TP	0.35	0.41	
TN	0.36	0.32	
TN TP^{-1}			0.55
CPOM			-0.47
TP$_{exp}$	0.42		
TN$_{exp}$	0.42		
CPOM$_{exp}$		-0.35	-0.55

Table 3 Principal Component Analysis on physical and chemical variables. Factors are a linear combination of

the different variables. Percent of variance explained and factor loadings (> 0.30) are shown. Abbreviations are

as in Figure 5.

Both streams are clearly discriminated by PCA ordering, highlighting seasonal differences in each

stream (Fig. 5b). Besides separating samples from each stream, plotting FI versus FII shows that each stream is

temporally heterogeneous, allowing clustering in 3 groups for CP (CP$_A$, CP$_B$ and CP$_C$) and 2 for G (G$_A$ and G$_B$)

(Fig. 5b). CP$_A$ had low discharge and export rate values and high temperatures, taking place at the warmest

period (Fig. 2). CP$_B$ had higher nutrient concentration and export values, matching the period with more

abundant precipitation (Fig. 2). CP$_C$ was mainly characterized by low EC, nutrient concentration and turbidity

values, and coincides with the period of diminishing rainfall and monotonic air temperature warming (Fig. 2).

G$_A$ had similar characteristics to CP$_A$ and spans the same period. The remaining samples fall within G$_B$. Thus, the

values of physical and chemical variables sampled monthly through the year show that both streams have

marked seasonal differences that follow precipitation and air temperature patterns.

Fig 5 Principal Component Analysis (PCA) of physical and chemical variables in the study streams. a) The

weight of variables in each of the axes is shown. b) First factorial plane of the PCA, Casa de Piedra sampling

dates are represented by triangles and Gutiérrez ones are represented by rectangules, the numbers correspond to

the sampling dates and they are grouped by climatic characteristics of the region (a, b and c). Electrical

conductivity (EC), Discharge (Q), Total Phosphorus (TP), Total Nitrogen (TN), Coarse Particulate Organic

Matter (CPOM), Export (exp).

DISCUSSION AND CONCLUSIONS

We surveyed discharge and nutrient dynamics in two adjacent North Patagonian streams with contrasting

drainage basins. Precipitation had a direct influence on the magnitude of stream discharge at both streams, but

the response time varied between them. CP stream was flashy, showed a tighter relation between precipitation at

the basin and discharge response and stronger seasonality of physical and chemical parameter values. G stream,

in contrast, showed a more buffered discharge response to precipitation and higher nutrient exports. In general,

similarities seen between streams can be related to regional climate, while differences are better explained by

contrasting topography and land use of the respective drainage basins.

Precipitation and temperature have opposite influence in river discharge in the Andean region

(Masiokas et al. 2008). In this way, precipitation increased stream discharge of our study sites, while elevated air

temperature periods produced the opposite effect. The different pathways taken by precipitated water to reach

the streams (run-off, groundwater and snowmelt) (Brown et al. 2003) and their relative importance are evidenced

by the three periods found in the discharge-precipitation correlogram. Surface run-off was more relevant at CP, a

stream that feeds from run-off from a basin characterized by steep slopes and the lack of buffering from a large

upstream lake. In this stream, discharge increased drastically during the same day precipitation occurred, and the

following days. Water that does not reach the stream immediately as run-off, infiltrates into the soil and moves

to the stream as sub-surface run-off or groundwater; these pathways usually results in a delayed stream discharge

response (Kalff 2002). This delay time was close to a month for both streams; this similarity suggests that both

basins share the same subsurface geological traits, and thus that geology drives groundwater dynamics. The sub-

surface run-off and groundwater pathways were also more relevant at CP, which showed a higher *r(max)* and

shorter delay times. In colder regions, snow precipitation is stored during the winter months and released as

snowmelt in spring and summer (Bailey 1995). This results in a delay time between precipitation and discharge

response of about 2-4 months, as we found at both study sites during spring. Such discharge dynamics and its

relationship with the fractions of precipitation fallen as either rain or snow match the storm and melt periods

described at other mountainous regions (Ahearn et al. 2004).

The presence of a headwater lake in the drainage basin regulates the flow regime of a stream (Baker et

al. 2004). This was made evident by the contrasting values of discharge autocorrelation and flashiness R-B

indices between the streams in our study. The autocorrelation function fell sharply for CP but sloped more gently

for G, showing a shorter memory effect (Mangin 1984) in CP, and agreeing with differences in annual values of

the R-B index. Flashiness values seen at CP (R-B = 0.14) are similar to those found for streams at Austria (R-B

= 0.18) and Slovakia (R-B = 0.15) that flow through basins with steep slopes and small area (Holko et al. 2011).

In contrast, flashiness at G was much lower (R-B=0.03), and more akin the values found for streams with

regulated flows at Indiana, USA (Baker et al. 2004). Thus, the difference in flashiness between the adjacent

streams we studied is explained by the contrasting topography of their drainage basins, including the presence of

a large lake in one but not the other.

Leaves of riparian trees are generally the largest component of allochthonous inputs to stream located in

forested drainage basins and are often the major source of energy to heterotrophic organisms (Vannote et al.

1980; Webster et al. 1999)**.** In the case of Andean streams, *N. pumilio* leaves cover large areas of the waterway

during autumn (Albariño et al. 2009). A substantial amount of these may be transported downstream by spates

occurring from late autumn to winter and eventually reaches the lake (Modenutti et al. 2010). We observed no

significant differences in the amount of CPOM exported by the two streams studied. However, we highlight the

fact that in stream CP the CPOM was mainly made up of native *N. pumilio* leaves, while in stream G introduced

S. fragilis leaves dominated. This corresponds to the tree species present in the different catchments, and could

explain the temporal difference between streams in the surge of CPOM; the peak in CPOM export occurred

during June and July in stream CP, and during April in stream G. This supports the claim that afforestation or

invasion of exotic plants with different phenology, chemical and physical characteristics to those of the native

riparian species changes the timing, quantity and quality of leaf litter standing stock in streams (Naiman et al.

2005).

Streams in the north Patagonian region are oligotrophic (Pedrozo et al. 1993; García et al. 2015b). The

concentration and the annual export rate of the nutrients from CP stream reflected this characteristic. Values of N

and P concentrations were lower to those measured in a forested nearby sub-catchment (TN = 110.6 µg l^{-1} and

TP = 9.6 µg l^{-1}) while they were much smaller than those found in a burned sub-catchment of this region (TN =

675.1 µg l^{-1} and TP = 12.0 µg l^{-1}) (Temporetti 2006). Values of N export were lower to those measured in areas

of low anthropogenic impact in Patagonia (average TN export = 100 kg km^{-2} $year^{-1}$) (Little et al. 2008), and

values of P export were between the lowest measured in South America (TP export range from 4 to 450 kg TP

km^{-2} $year^{-1}$) (Álvarez-Cobelas & Angeler 2007). These results indicate that nutrient export in CP river ecosystem

is among the lowest measured for any basin worldwide (Álvarez-Cobelas et al. 2008; Álvarez-Cobelas et al.

2009). However, it should be taken into account that we have estimated basal nutrient export rates because we

measured them in days without precipitation, when their export tend to be low (Tate & Singer 2013). In any

case, it is interesting to note that these rates were higher for stream G than for CP, even though the opposite

result could have been expected, given that headwater lakes act as nutrient and sediment traps (Parker et al.

2009). The higher nutrient concentration seen at G is thus not explained by natural causes, but by the effects of

land use in its drainage basin, namely the presence of human settlements and trout farms.

Increases in eutrophication levels of river ecosystems is due to both point and diffuse sources

(Carpenter et al. 1998). While we did not formally weigh the relative impacts of fish farming and urbanization,

we suspect that a growing residential settlement of the Gutiérrez stream drainage basin, often times without

adequate wastewater treating infrastructure, is expected to have a larger impact than the detritus from fish farms

producing ~3 tons of fish per year. This is partially supported by nutrient measurements at G stream during the

low water period where dissolved inorganic nitrogen export (DIN_{exp}) was measured downstream the larger fish

farm (DIN_{exp} = 7 kg km^{-2} $year^{-1}$) and at its mouth (DIN_{exp} = 108 kg km^{-2} $year^{-1}$) (Sosnovsky, unpublished results).

Moreover, the additional nutrient sources of fish food and feces from salmonid production decrease the TN:TP

ratio of the water column in Patagonia reservoirs (Diaz et al. 2001), a fact we did not observe in G stream.

However, a more detailed spatial study in G drainage basin aimed to determine the sources and sinks of total and

dissolved nutrients associated to eutrophication processes is still necessary.

The molecular N:P ratio is said to distinguish nitrogen-limited systems from those constrained by

phosphorus (Rhee & Gotham 1980). Kahlert (1998), in a review of the literature, found that the optimal N:P ratio

of freshwater benthic algae was 18:1 (molar), which deviates from the phytoplankton optimal stoichiometric N:P

ratio of 16:1 (Redfield ratio) (Goldman et al. 1979). CP and G streams had an arithmetic mean N:P molar ratio

close to 15, which would indicate nitrogen deficit in these ecosystems, similar to what has been observed in

other ecosystems of Northern Patagonia, either forests (Satti et al. 2003) or lakes (Diaz et al. 2007). However,

considering that median values of N:P were 11 for CP and 16 for G, it becomes evident that nitrogen limitation is

being lifted at G, most likely due to wastewater leaching from human settlements. An increase in the N load is

expected to result in increased eutrophication of the G stream and downstream water bodies. Thus, it would be

advisable to build adequate wastewater treatment plants to avoid driving these, up until now, low impact

drainage basins towards ecosystem eutrophication. Our findings at the drainage basin level are in line with

observations at the global scale suggesting human actions generate a clear unbalance favoring N over P

(Peñuelas et al. 2013).

By studying two Andean streams located at adjacent, yet contrasting drainage basins, we were able to

investigate the influence of environmental factors acting at different spatial scales over these freshwater

ecosystems. At a regional scale, climate determined seasonality of water flow for both streams. At a landscape

scale, we found that the presence of a large, deep lake buffered the hydrological dynamic of its effluent stream.

At a lower spatial scale, we found that more intense human impact at G's drainage basin correlated with higher

nutrient export rates by the stream. Considering the strong demographic pressure at G's drainage basin, there is a

high risk of change in N:P stoichiometry and ecosystem eutrophication. Gutiérrez basin situation thus shows a

stark contrast to Casa de Piedra basin, which instead has the hydrological, physical and chemical dynamics of a

pristine drainage basin characteristics of Andean streams originating at high altitude wetlands and small lakes

(Sosnovsky et al. 2019). Besides geomorphology, the main differences between both basins are land use policies:

CP's basin is mostly within protected areas, while most of G's lower basin is within the limits of San Carlos de

Bariloche city and subject to several, potentially high impact uses, like residential neighborhoods, winter sports

and fish farming, among others. Local and global environmental problems are currently burning issues in stream

ecosystem processes and will be even more so in the future. Considering that lakes and streams are sentinels and

integrators of environmental change in the surrounding terrestrial landscape (Williamson et al. 2008), it is really

important to monitor the key variables of these aquatic ecosystems (Lovett et al. 2007) and emphasize the

preservation and restoration of their drainage basins.

428
Author contributions

Alejandro Sosnovsky performed the study conception and design. All authors contributed to the material

preparation, data collection and analysis. The first draft was written by Sosnovsky, Alejandro and all authors

commented on previous versions of the manuscript. All authors read and approved the final manuscript.

REFERENCES

A.P.H.A., 2005. Standard methods for examination of water and wastewater, 21 edn. American Public Health

Association, Washington DC.

Ahearn, D. S., R. W. Sheibley, R. A. Dahlgren & K. E. Keller, 2004. Temporal dynamics of stream water

chemistry in the last free-flowing river draining the western Sierra Nevada, California. Journal of Hydrology

295: 47-63.

Albariño, R. J., V. Díaz Villanueva & L. Buria, 2009. Leaf litter dynamics in a forested small Andean catchment,

northern Patagonia, Argentina. In Oyarzún, C., N. E. C. Verhoest, P. Boeckx & R. Godoy (eds), Ecological

advances on Chilean temperate rainforest. Academia Press, 183-211.

Álvarez-Cobelas, M. & D. G. Angeler, 2007. Exportación de nutrientes en las cuencas hidrográficas de

Latinoamérica: una recopilación. Revista Latinoamericana de Recursos Naturales 3: 31-43.

Álvarez-Cobelas, M., D. G. Angeler & S. Sánchez-Carrillo, 2008. Export of nitrogen from catchments: A

worldwide analysis. environmental Pollution 156: 261-269.

Álvarez-Cobelas, M., S. Sánchez-Carrillo & D. G. Angeler, 2009. Phosphorus export from catchments: a global

view. Journal of American Benthological Society 28: 805-820.

Allan, J. D., 2004. Landscapes and Riverscapes: The influence of land use on stream ecosystems. Annual

Review of Ecology, Evolution and Systematics 35: 257-284.

Bailey, R. G., 1995. Ecosystem Geography. Springer-Verlag, New York, New York.

Baker, D. B., R. P. Richards, T. T. Loftus & J. W. Kramer, 2004. A new flashiness index: Characteristics and

applications to midwestern rivers and streams. Journal of the American Water Resources Association 40:

503-522.

455 Barros, V. & I. Camilloni, 2016. La Argentina y el cambio climático De la física a la política, 1[st] edn. EUDEBA,

456  Ciudad Autónoma de Buenos Aires.

457 Brown, L. E., D. M. Hannah & A. M. Milner, 2003. Alpine Stream Habitat Classification: An Alternative

458  Approach Incorporating the Role of Dynamic Water Source Contributions. Artic, Antartic, and Alpine

459  Research 35: 313-322.

460 Carpenter, S. R., N. F. Caraco, D. L. Correll, R. W. Howarth, A. N. Sharpley & V. H. Smith, 1998. Nonpoint

461  Pollution of Surface Waters with Phosphorus and Nitrogen. Ecological Applications 8: 559-568.

462 Carrillo, U., V. Díaz Villanueva & B. E. Modenutti, 2018. Sustained effects of volcanic ash on biofilm

463  stoichiometry, enzyme activity and community composition in North-Patagonia streams. Science of the Total

464  Environment 621: 235-244.

465 Di Rienzo, J. A., F. Casanoves, M. G. Balzarini, M. González, M. Tablada & C. W. Robledo, 2016. InfoStat.

466  2016 edn. Universidad Nacional de Córdoba.

467 Diaz, M. M., F. L. Pedrozo, C. S. Reynolds & P. F. Temporetti, 2007. Chemical composition and the nitrogen-

468  regulated trophic state of Patagonian lakes. Limnologica 37: 17-27.

469 Diaz, M. M., P. F. Temporetti & F. L. Pedrozo, 2001. Response of phytoplankton to enrichment from cage fish

470  farm waste in Alicura Reservoir (Patagonia, Argentina). Lake & Reservoir Management 6: 151-158.

471 Díaz Villanueva, V., M. Bastidas Navarro & R. Albariño, 2016. Seasonal patterns of organic matter

472  stoichiometry along a mountain catchment. Hydrobiologia.

473 Diehl, L., M. J. Mazzarino, F. Funes, S. Fontenla, M. Gobbi & J. Ferrari, 2003. Nutrient conservation strategies

474  in native Andean-Patagonian forests. Journal of Vegetation Science 14: 63-70.

475 Dodds, W. K. & V. H. Smith, 2016. Nitrogen, phosphorus, and eutrophication in streams. Inland Waters 6: 155-

476  164.

477 Emerson, D. G., A. V. Vecchia & A. L. Dahl, 2005. Evaluation of Drainage-Area Ratio Method Used to

478  Estimate Streamflow for the Red River of the North Basin, North Dakota and Minnesota. In: 5017, S. I.-.

479  (ed). U.S. Geological Survey, Reston, Virginia, 18.

480 Fongers, D., R. Day & J. Rathbun, 2012. Application of the Richards-Baker Flashiness Index to gaged Michigan

481  rivers and Streams. Water Resources Division, Michigan Department of Environmental Quality, Lansing, 1-

482  141.

García, D. R., M. C. Diéguez, M. Gerea & P. E. García, 2018. Characterisation and reactivty continuum of

dissolved organic matter in forested headwater catchments of Andean Patagonia. Freshwater Biology 63:

1049-1062.

García, P., M. C. Diéguez & C. P. Queimaliños, 2015a. Landscape integration of North Patagonian mountain

lakes: a first approach using characterization of dissolved organic matter. Lakes & Reservoirs: Research and

Management 20: 1-14.

García, R. D., M. Reissig, C. P. Queimaliños, P. E. García & M. C. Diéguez, 2015b. Climate-driven terrestrial

inputs in ultraoligotrophic mountain streams of Andean Patagonia revealed through chromophoric and

fluorescent dissolved organic matter. Science of the Total Environment 521-522: 280-292.

Goldman, J. C., J. J. Mc Carthy & D. G. Peavey, 1979. Growth rate influence on the chemical composition of

phytoplankton in oceanic waters. Nature 279: 210-215.

Gordon, N. D., T. A. McMahon, C. J. Gippel & R. J. Natha, 2004. Stream Hydrology An Introduction for

Ecologist. John Wiley & Sons LTD, Chichester, England.

Grasshoff, M., K. Ehrhardt & K. Kremling, 1983. Methods of seawater analysis. Wiley, New York.

Hedin, L. O., J. J. Armesto & A. H. Johnson, 1995. Patterns of Nutrient Loss from Unpolluted, Old-Growth

Temperate Forests: Evaluation of biogeochemical theory. Ecology 76: 493-509.

Hintze, J. L., 1998. Number Cruncher Statistical System (NCSS). Version 2000. User´s Guide. Kaysville, Utah.

Holko, L., J. Parajka, Z. Kostka, P. Skoda & G. Blöschl, 2011. Flashiness of mountain streams in Slovakia and

Austria. Journal of Hydrology 405: 392-401.

Holland, E. A., F. J. Dentener, B. H. Braswell & J. M. Sulzman, 1999. Contemporary and pre-industrial global

reactive nitrogen budgets. Biogeochemistry 46: 7-43.

Howarth, R. W., G. Billen, D. Swaney, A. Townsend, N. A. Jaworski, K. Lajtha, J. A. Downing, R. Elmgren, N.

Caraco, T. Jordan, F. Berendse, J. Freney, V. Kudeyarov, P. Murdoch & Z. L. Zhu, 1996. Regional nitrogen

budgets and riverine N & P fluxes for the drainages to the North Atlantic Ocean: natural and human

influences. Biogeochemistry 35: 75-139.

Hynes, H. B. N., 1975. The stream and its valley. SIL, Proceedings, 1922-2010 19: 1-15.

Jordan, T. E., D. L. Correll & D. E. Weller, 1997. Effects of agriculture on discharges of nutrients from coastal

plain watersheds of Chesapeake Bay. Journal of Environmental Quality 26: 836-848.

Kahlert, M., 1998. C:N:P ratios of freshwater benthic algae. Archiv für Hydrobiologie, Special Issues, Advances

in Limnology 51: 105-114.

Kalff, J., 2002. Hidrology and Climate. In Ryu, T. (ed) Limnology Inland Water Ecosystems. Prentice Hall,

Upper Saddle River, 53-71.

Lallement, M., P. J. Macchi, P. H. Vigliano, S. Juarez, M. Rechencq, M. Baker, N. Bouwes & T. A. Crowl,

2016. Rising from the ashes: Change in salmonid fish assemblages after 30 months of the Puyehue-Cordon

Caulle volcanic eruption. Science of the Total Environment 541: 1041-1051.

Larocque, M., A. Mangin, M. Razack & O. Banton, 1998. Contribution of correlation and spectral analyses to

the regional study of a large karst aquifer (Charente, France). Journal of Hydrology 205: 217-231.

Likens, G. E., 1985. An Ecosystem Approach to Aquatic Ecology. Mirror Lake and Its Environment. Springer-

Verlag, New York.

Little, C., D. Soto, A. Lara & J. G. Cuevas, 2008. Nitrogen exports at multiple-scales in a southern Chilean

watershed (Patagonian Lakes district). Biogeochemistry 87: 297:309.

Lovett, G. M., D. A. Burns, C. T. Driscoll, J. C. Jenkins, M. J. Mitchell, L. Rustad, J. B. Shanley, G. E. Likens &

R. Haeuber, 2007. Who need environmental monitoring? Frontiers in Ecology and the Envirnonment 5: 253-

260.

Mangin, A., 1984. Pour une meilleure connaissance des systèmes hydrologgiques à partir des analyses

corrélatoire et spectrale. Journal of Hydrology 67: 25-43.

Masiokas, M. H., R. Villalba, B. H. Luckman, M. E. Lascano, S. Delgado & P. Stepanek, 2008. 20th-century

recession and regional hydroclimatic changes in northwestern Patagonia. Global and Planetary Change 60:

85-100.

Mermoz, M. & C. Martín, 1987. Mapa de vegetación de la Reserva y del Parque Nacional Nahuel Huapi.

Administración de Parques Nacionales, Delegación Técnica Regional Patagonia, San Carlos de Bariloche,

23.

Merseburger, G. C., E. Martí & F. Sabater, 2005. Net changes in nutrient concentrations below a point source

input in two streams draining catchments with contrasting land uses. Science of the Total Environment 347:

217-229.

Meyer, J. L., M. J. Paul & W. K. Taulbee, 2005. Stream ecosystem function in urbanizing landscapes. Journal of

the North American Benthological Society 24: 602-612.

Miserendino, M. L., M. A. Kutschker, C. Brand, L. La Manna, Y. C. Di Prinzio, G. Papazian & J. Bava, 2016.

Ecological Status of a Patagonian Mountain River: Usefulness of Environmental and Biotic Metrics for

Rehabilitation Assessment. Environmental Management 57: 1166-1187.

Modenutti, B. E., R. J. Albariño, M. Bastidas Navarro, V. D. Villanueva, A. F. Souza, C. Trochine, C.

Laspoumaderes, F. Cuassolo, G. Mariluán, L. Buria & E. G. Balseiro, 2010. Structure and dynamic of food

webs in Andean North Patagonian freshwater systems: organic matter, light and nutrient relationships.

Ecología Austral 20: 95-114.

Naiman, R. J., H. Décamps & M. E. McClain, 2005. Riparia: ecology, conservation and management of

streamside communities. Elsevier Academic Press, London, UK.

Oyarzún, C. E., R. Godoy, A. Schrijver De, J. Staelens & N. Lust, 2004. Water chemistry and nutrient budgets in

an undisturbed evergreen rainforest of souther Chile. Biogeochemistry 71: 107-123.

Oyarzún, C. E., R. Godoy & A. Sepulveda, 1998. Water and nutrient fluxes in a cool temperate rainforest at the

Cordillera de la Costa in southern Chile. Hydrological Processes 12: 1067-1077.

Parker, B. P., D. E. Schindler, K. G. Beaty, M. P. Stainton & S. E. M. Kasian, 2009. Long-term changes in

climate, streamflow, and nutrient budgets for first-order catchments at the Experimental Lakes Area (Ontario,

Canada). Canadian Journal of Fisheries and Aquatic Sciences 66: 1848-1863.

Paruelo, J. M., A. Beltran, E. Jobbágy, O. Sala & R. Golluscio, 1998. The climate of Patagonia: general patterns

and controls on biotic processes. Ecología Austral 8: 85-101.

Pedrozo, F. L., S. Chillrud, P. F. Temporetti & M. M. Diaz, 1993. Chemical composition and nutrient limitation

in rivers and lakes of northern Patagonian Andes (39.5°-42° S; 71° W) (Rep. Argentina). SIL Proceedings,

1922-2010 25: 207-214.

Peel, M. C., B. L. Finlayson & T. A. McMahon, 2007. Updated world map of the Köppen-Geiger climate

classification. Hydrology and Earth System Sciences 4: 439-473.

Peltier, M. R., C. J. Wilcox & D. C. Sharp, 1998. Technical Note: Application of the Box-Cox data

transformation to animal science experiments. Journal of Animal Science 78: 847-849.

Peñuelas, J., B. Poulter, J. Sardans, P. Ciais, M. van der Velde, L. Bopp, O. Boucher, Y. Godderis, P. Hinsinger,

J. Llusia, E. Nardin, S. Vicca, M. Obersteiner & I. A. Janssens, 2013. Human-induced nitrogen–phosphorus

imbalances alter natural and managed ecosystems across the globe. Nature Communications 4: 2934.

Pieterse, N. M., W. Bleuten & S. E. Jørgensen, 2003. Contribution of point sources and diffuse sources to

nitrogen and phosphorus loads in lowland rivers tributaries. Journal of Hydrology 271: 213-225.

Pozo, J., A. Elosegi, J. Díez & J. Molinero, 2009. Dinámica y relevancia de la materia orgánica. In Elosegi, A. &

S. Sabater (eds), Conceptos y técnicas en ecología fluvial. Fundación BBVA, Bilbao, España, 141-166.

Rhee, G. Y. & I. J. Gotham, 1980. Optimum N:P ratios and co-existence of planktonic algae. Journal of

Phycology 16: 486-489.

Satti, P., M. J. Mazzarino & M. Gobbi, 2003. Soil N dynamics in relation to leaf litter quality and soil fertility in

north-western Patagonian forests. Journal of Ecology 91: 173-181.

Snelder, T. H. & B. J. F. Biggs, 2002. Multiscale river environment classification for water resources

management. Journal of the American Water Resources Association 38: 1225-1239.

Sosnovsky, A., M. Rechencq, V. Fernández, M. J. Suarez & R. J. C. Cantet, 2019. Hydrological and Physico-

chemical dynamics in two Andean streams. Limnetica in press.

Tate, K. W. & M. J. Singer, 2013. Timing, Frecuency of Sampling Affect Accuracy of Water-Quality

Monitoring. California Agriculture 53: 44-48.

Temporetti, P. F., 2006. Efecto a largo plazo de los incendios forestales en la calidad del agua de dos arroyos en

la sub-región Andino-Patagónica, Argentina. Ecología Austral 16: 157-166.

Vannote, R. L., G. W. Minshall, K. W. Cummins, J. R. Sedell & C. E. Cushing, 1980. The River Continuum

Concept. Canadian Journal of Fisheries and Aquatic Sciences 37: 130-137.

Veblen, T. T., R. S. Hill & J. Read (eds), 1996. The ecology and biogeography on Nothofagus forests. Yale

University Press, London, UK.

Walsh, C. J., T. D. Fletcher & A. R. Ladson, 2005. Stream restoration in urban catchments through redesigning

stormwater systems: looking to the catchment to save the stream. Journal of the North American

Benthological Society 24.

Webster, J. R., E. F. Benfield, T. P. Ehrman, M. A. Schaeffer, J. L. Tank, J. J. Hutchens & D. J. D´angelo, 1999.

What happens to allochthonous material that falls into streams? A synthesis of new and published

information from Coweeta. Freshwater Biology 41: 687-705.

Wiens, J. A., 2002. Riverine landscapes: taking landscape ecology into the water. Freshwater Biology 47: 501-

515.

Williams Subiza, E. A. & C. Brand, 2018. Short-term effect of wildfire on Patagonian headwater streams.

International Journal of Wildland Fire 27: 457-470.

Williamson, C. E., W. K. Dodds, T. K. Kratz & M. A. Palmer, 2008. Lakes and streams as sentinels of

environmental change in terrestrial and atmospheric processes. Frontiers in Ecology and the Envirnonment 6:

247-254.