Topography and Land Use Modulate Hydrological and Nutrient

Export Dynamics in two Andean Streams

Alejandro Sosnovsky[1], Mailén Elizabeth Lallement[1], Magalí Rechencq[1],

Eduardo Enrique Zattara[1], María Valeria Fernández[1], Sergio Leiva[2], María

José Suárez[3], Romina Daga[4] and Rodolfo Juan Carlos Cantet[3]

[1]Grupo de Evaluación y Manejo de Recursos Ícticos, INIBIOMA, CONICET-

Universidad Nacional del Comahue, Quintral 1250, R8400FRF San Carlos de

Bariloche, Argentina

[2] Grupo de Nuevos Materiales y Dispositivos, Comisión Nacional de Energía Atómica

Av. Ezequiel Bustillo Km 9.5 (8400) San Carlos de Bariloche, Argentina

[3]Departamento de Producción Animal, Facultad de Agronomía, Universidad de Buenos

Aires, e INPA, CONICET. Av. San Martín 4453, C1417DSQ Ciudad Autónoma de

Buenos Aires, Argentina

[4]Laboratorio de Análisis por Activación Neutrónica, Centro Atómico Bariloche,

Comisión Nacional de Energía Atómica, Av. Bustillo 9500, 8400 Bariloche, Argentina.

Centro Científico Tecnológico CONICET. Patagonia Norte.

Corresponding author: Alejandro Sosnovsky

e-mail: alejandro.sosnovsky@comahue-conicet.gob.ar

Telephone number: +54 294 423374 (452), Fax number: +54-2944-422111

Topography and Land Use Modulate Hydrological and Nutrient

Export Dynamics in two Andean Streams

Abstract: Fluvial dynamics are driven by multiple environmental factors and

scales. Studies coupling hydrological and nutrient dynamics of Andean streams

are almost nonexistent. We characterized two adjacent streams with contrasting

drainage basins: Casa de Piedra, originating in a small mountain lake and running

through a pristine landscape, and Gutiérrez, originating in a large piedmont lake

and running through an anthropized landscape. Despite both drainage basins

sharing geology and climate, we found that the streams presented contrasting

hydrological and nutrient dynamics. Casa de Piedra had higher discharge

flashiness with shorter response delays to precipitation. Interestingly, Gutiérrez's

hydrology was buffered by the upstream lake, but its nutrient exports were not.

Differences observed in quality and timing of coarse particulate organic matter

export and basal export levels of phosphorus and nitrogen could be explained by

human activities affecting Gutiérrez but not Casa de Piedra. Moreover,

nitrogen:phosphorus ratio indicates a possible future shift to phosphorus as the

limiting nutrient as Andean basins become more densely populated. In summary,

our annual basis study shows that even under a common geology and climate,

stream dynamics of adjacent basins can be starkly different due to differences in

topography and land use.

INTRODUCTION

River ecosystems have traditionally been studied as isolated entities; however, it has

recently become clear that they are deeply intertwined with their surroundings. Climate,

topography, geology and land cover are environmental factors that show a hierarchical

influence on spatial patterns of these ecosystems (Snelder and Biggs 2002). In other

words, a river is the product of its landscape (Wiens 2002) and thus riverine

hydrological, chemical and biological processes should be studied in relation to the

river's overall drainage basin (Hynes 1975).

Regional climate and drainage basin topography are factors acting at a macro

and meso spatial scales upon the river ecosystem (Snelder and Biggs 2002). Climate

directly drives the hydrological cycle, determining the flow regime of a stream, a key

ecosystem variable (Poff et al. 1997). Rain, snowmelt and groundwater feed stream

discharge, the relative importance of these factors varying over time and space (Brown

et al. 2003). After a rain episode, surface and sub-surface run-off are favored at

mountainous landscapes, while water infiltration and evapotranspiration are favored at

highly vegetated lowlands with permeable soils. Snowfall at high altitudes contributes

to formation of a snowpack, which will store water until the melt period, when this

water flows down into a stream (Bailey 1995). Lakes are topographic features of a

drainage basin that can buffer run-off response and sediment transport (Snelder and

Biggs 2002). Thus, headwater lakes influence directly the hydrograph and nutrient

dynamic of downstream fluvial ecosystems (Gordon et al. 2004). These hydrographs

tend to be stable over time, furthermore, considering lakes as sediment and nutrient

traps, low values of nutrient concentration and export could be expected in these

streams (Little et al. 2008; Parker et al. 2009).

At a lower spatial scale, drainage basin traits also affect fluvial dynamics, since

nutrients and contaminants generated at the basin eventually reach the river ecosystem

(Allan 2004; Dodds and Smith 2016). Non-point source pollution, associated with

changing land use patterns and practices, has resulted in increased impacts on water

bodies (Jordan et al. 1997; Carpenter et al. 1998). For example, growth of urban areas

favors nutrient export (Howarth et al. 1996; Meyer et al. 2005; Walsh et al. 2005).

Moreover, direct dumping of sewage water to a stream greatly increases nutrient load

(Pieterse et al. 2003) and can also change nutrient stoichiometry (Merseburger et al.

2005). Thus, increases of nutrient load leading to eutrophication of running waters and

carbon dynamics largely depend on the drainage basin state.

In Northern Patagonia, characteristics of Andean ecosystems are largely

determined by the Andes mountain range. This region is characterized by a temperate-

cold climate with warm and dry summers (Peel et al. 2007). The Andes represent an

important barrier for humid air masses brought from the Pacific Ocean by westerly

winds. Most of the humidity in these maritime air masses precipitates on the west side

of the Andes as they are blown to higher, colder altitudes across the mountain range.

Upon crossing the Andes, the air descends along the east slopes and becomes hotter and

drier through adiabatic warming. East of the Andes, the amount of rainfall follows a

steep west-east gradient: annual precipitation drops from 3500 mm to 700 mm in less

than 60 km. Moreover, the precipitation regime shows marked seasonality, with most

rain falling during fall and winter (Paruelo et al. 1998). As in other South American

watersheds, N deposition is remarkably low (Holland et al. 1999), and forest (Diehl et

al. 2003) and lake ecosystems (Diaz et al. 2007) are limited mainly by N. Moreover, a

large fraction of drainage basins along the East side of the Andes are in a relatively

pristine state because human population density is low. Diverse authors have studied

anthropic (Miserendino et al. 2016) and natural (Lallement et al. 2016; Carrillo et al.

2018; Williams Subiza and Brand 2018) impacts on Andean drainage basins and their

fluvial ecosystems. Recent research has revealed that dynamics of organic matter are

intimately associated with the stream-basin interaction (García RD et al. 2015; Díaz

Villanueva et al. 2016; García et al. 2018). Andean streams originate mainly from

snowmelt, glacial melt and piedmont lakes and are predominantly oligotrophic

(Pedrozo et al. 1993). Sosnovsky et al (2020) described their discharge and physico-

chemical dynamics during the three contrasting hydrological periods characteristics of

mountain regions, storm, snowmelt and basal periods. Oyarzún et al (2004) and

Temporetti (2006) estimated the nutrient load of pristine and burned watersheds,

respectively. Nevertheless, studies coupling hydrological and nutrient dynamics in the

region are lacking, despite the hydrological, ecological and human importance of these

variables and their possible interactions.

The main hypothesis of our study is that climate is the main driver of stream

ecosystems dynamics, which are in turn modulated by topography and land use.

Therefore, we expect precipitation, air temperature and stream discharge to be

intimately related. Under this hypothesis, adjacent drainage basins with contrasting

topography and land use should present differences in their hydrological, physical and

chemical dynamics. To test this expectation, we selected two streams located in

adjacent yet contrasting drainage basins: Casa de Piedra (CP) and Gutiérrez (G). CP

stream originates in a small high-mountain lake and its drainage basin is pristine

throughout its length. In contrast, G stream originates from a large piedmont lake which

buffers it hydrological dynamic (Sosnovsky et al. 2020), and its drainage basin suffers

from significant anthropization. While the large headwater lake is expected to buffer

chemical dynamics in G, higher anthropization of G's basin could impact the stream

ecosystem, potentially causing increase nutrient concentration and export. Our first goal

was to characterize discharge dynamic within a year in these fluvial ecosystems. To this

end, we compare discharge magnitude and variability between CP and G, and study

how it relates to climatic variables (basin precipitation and air temperature). Our second

goal is to determine the trophic state of both streams, measuring basal concentration,

export and stoichiometry of nitrogen (TN: total N) and phosphorous (TP: total P), and

estimating basal export of coarse particulate organic matter (CPOM, organic matter >

1mm). Our third goal is to identify the seasonal patterns of key physical and chemical

variables.

MATERIALS Y METHODS

Study area

Regional landscapes of Andean North Patagonia are the result mainly of

glaciation/deglaciation, fluvial and mass movement processes over the previous

structural-tectonic landscape. In the region, most fluvial courses have clear structural

control. Highest mountains show mainly the result of glacial erosion processes while

both erosive and depositional geoforms are present in the hillsides corresponding to

ancient lateral glacial valleys. The alluvial plains of streams and tributaries in the area

run mainly on moraine systems and stream mouths form alluvial fans over glacifluvials

plains (Pereyra et al. 2005).

The lithology of the study site is mainly composed of rocks of Paleozoic

metamorphic and granitic basement, Cretacic and Miocene granite porphyry, and

limited outcrops of Jurassic volcanic and sedimentary rocks covering previous

granitoids. Oligocene volcanic and marine sedimentary rocks are also present (Giacosa

et al. 2001). In general, the rocky outcrops constitute the hydrogeologic basement with

almost absent primary permeability and secondary permeability acquired by folding and

fracturing of the rocks. Headwaters are dominated by rocky outcrops and poor soil

development is associated mainly to presence of vegetation and lower slopes on the

sides of glacial valleys. On the other hand, the lower parts of the landscape are

represented by Pliocene-Pleistocene glacial material (moraines, glaciofluvial-fluvial and

glaciolacustrine deposits) with poorly to moderately developed soils (Andisols reaching

more than 80 cm in some places) under well drained conditions, well provided of

superficial organic matter and phosphorus retention >85% (Satti et al. 2003). The most

recent sedimentary deposits correspond to the present alluvial materials (gravel, sand,

silt) in the flood plains surrounding the Gutiérrez lake and streams, with poorly

developed soils due to active morphodynamics and coarse materials (Giacosa et al.

2001; Pereyra et al. 2005). Associated to these deposits are unconfined aquifers of

variable permeability and discontinuous character due to the heterogeneity on the grain

size of the deposits (Pereyra et al. 2005).

Research was conducted at the drainage basin of the streams Casa de Piedra

(CP; 41° 07' 30.11" S 71° 27' 13.16"W) and Gutiérrez (G; 41° 09' 36.18" S 71° 24'

37.19"W), both located in Nahuel Huapi National Park, Río Negro, Argentina (Figure

1). These drainage basins are very different from each other, despite of their proximity

(Table 1). CP's basin covers 65.5 km^2 and the stream originates in lake Jakob, a small

(0.15 km^2), ultra-oligotrophic (1-3 μg l^{-1} TP), high altitude water body situated at 1550

m above sea level (García P et al. 2015). CP's only perennial affluent is the Rucaco

stream, which originates at a high-altitude wetland. CP flows 19.7 km through a steep

V-shaped valley (average slope: 33.9 m km^{-1}) for most of its length. CP's drainage

basin is mostly pristine, with human activities restricted to hiking and infrequent

camping; human population (less than 1000 inhabitants, 2010 National Census) is

restricted to lowland areas. Forest below the tree line includes deciduous *Nothofagus*

pumilio and evergreens *N. dombeyi* and *Austrocedrus chilensis*. In contrast, G's

drainage basin is larger (161.6 km^2), and the stream originates in lake Gutiérrez, a deep

(111 m maximum depth) and large (17 km^2) water body that occupies ~10 % of the

basin's area. The seasonal concentration of TP is 3.4 μg l^{-1} on average (Diaz et al.

2007). G's only perennial affluent is the Cascada stream, whose catchment covers 13.0

km^2 and includes a major winter sports resort. G flows 9 km through a wide, gently

sloped valley (5.9 m km^{-1}). Its riverbanks are extensively colonized by the exotic crack

willow, *Salix fragilis*, and pass through a series of populated areas (approximately 4500

inhabitants, 2010 National Census) belonging to the city of San Carlos de Bariloche

(approximately 108000 inhabitants, 2010 National Census). In addition, one trout farm

with a total production of ~10 tons year^{-1} use G's waters. In summary, CP's basin is

largely free of impact from human activities, while G's basin is not.

Figure 1. a) Nahuel Huapi National Park; the study site is shaded in black. The city
marker indicates San Carlos de Bariloche downtown. b) Study area, Casa de Piedra and
Gutiérrez drainage basins, Jakob lake and Gutiérrez lake sub-catchments are also
delimitated. Curves represent 100 m slope on the terrain. Populated zones are shown in
dark grey, crosses indicate sampling sites, black circles indicate the limnigraphic
stations and the white circle indicate the rain meter location. R Str: Rucaco stream; C
Str: Cascada stream (with permission from Sosnovsky et al. 2020).
194

Data collection

The present study was carried out over a period of one year, from March 2013 to March

2014. Daily precipitation was recorded with a rain meter (Figure 1) and did not

differentiate between rain and snow. Discharge data were taken from average daily

records from limnigraphic stations. CP's limnigraphic station is located close to its

mouth, while G's station is located at its source (Figure 1). Values of discharge at our G

sampling site were estimated as discharge as measured at the station plus discharge for

the Cascada stream, determined using the Drainage-Area Ratio method (Emerson et al.

2005); based on the specific daily values of discharge for the adjacent CP stream.

Specific discharge (Q^{sp}) was defined as discharge per unit area. Daily average air

temperature records were obtained from the National Weather Service.

	CP: Casa de Piedra	**G:** Gutiérrez
Drainage area (km^2)	65.5 (60.3)	161.6 (30.1)
Lake area (km^2)	0.15	17.21
Native forest (%)	60.2 (61.7)	61.1 (56.3)
Bare ground (%)	29.7 (27.7)	17.2 (9.5)
Exotic forest (%)	9.9 (10.5)	10.8 (34.2)
Population (N)	< 1000 (< 1000)	4528 (3580)
Urban area (km^2)	0.65 (0.65)	11.35 (5.08)

Table 1. Drainage basins characteristics. Values are expressed based on the entire area
of the drainage basins, values between parentheses exclude the sub-catchment areas
draining into Jakob lake and Gutiérrez lake.

Baseline levels of nutrients and relevant physical and chemical variables were

obtained from monthly samples at both streams (Figure 2). Water samples were taken to

analyze total phosphorus (TP) and total nitrogen (TN) concentrations. Water turbidity

was measured with a Velp turbidimeter. Temperature, electrical conductivity (EC) at 25

°C and pH were measured with an Oakton probe. Coarse particulate organic matter

(CPOM) was sampled with 2 drift nets (1-mm, 20 X 20-cm opening at the mouth) that

were staked in the streambed and left to collect material for intervals of 10 to 200

minutes, depending on the amount of CPOM present in the stream. During October,

high and fast stream discharge prevented us from placing Surber sampler nets at G, so

we performed linear interpolation for this value. Data were taken exclusively on days

without precipitation.

Figure 2. Temperature (solid line) and precipitation (dotted line) during the study
period. Black dots and numbers indicate sampling dates.
225

Data processing and analysis

Discharge

Differences in stream discharge were evaluated using a heterogeneous variance model.

For this purpose, data on the average daily discharge and average daily specific

discharge were transformed to normality using the Box-Cox procedure as suggested by

Peltier et al. (1998). To understand the dynamics of discharge, we used an

autocorrelation analysis (Mangin 1984). This analysis examines how a value depends

on preceding values over a period of time. The correlations are computed for a given lag

time k to obtain an autocorrelation coefficient r(k). For example, the correlation

between today's and yesterday's stream discharges would be a lag-1 autocorrelation

coefficient, r(1). The autocorrelation function is represented with a correlogram, where

slope is determined by the response of the system to an event. If the event has only a

short-term influence on the response of the discharge, the slope of the correlogram will

decrease steeply. In contrast, if the system is influenced by an event for a long time, the

slope of the correlogram will decrease slowly. Generally the length of the influence of

an event is given by the "memory effect" which is according to Mangin (1984) the lag

number k when r(k) reaches a value of 0.2. The formula for autocorrelation is (Larocque

et al. 1998):

$$r(k) = \frac{C(k)}{C(O)}$$

245 with

$$C(k) = \frac{1}{n} \sum_{t=1}^{n-k} (x_t - \bar{x})(x_{t+k} - \bar{x})$$

Where k is the time lag and varies from 0 to n. According to Mangin (1984), n has to be

taken as 1/3 of the whole dataset to avoid stability problems.

In addition, we studied the relationship between discharge, precipitation and

temperature by cross-correlation analysis. This analysis is widely used to analyze the

linear relationship between input and output signals in hydrology (Larocque et al.

1998). In this case, input signals were precipitation and temperature, and output signal

was river discharge of CP and G streams. Cross-correlations are represented by a cross-

correlogram. The maximum amplitude and the lag value of the cross-correlogram

provide information on the delay, which indicates the time of the pulse transfer to the

stream. According to Larocque et al. (1998) the formula for cross-correlation is:

$$r_{xy} = \frac{C_{xy}\,(k)}{\sigma_x \sigma_y}$$

259   with

$$C_{xy}(k) = \frac{1}{n} \sum_{t=1}^{n-k} (x_t - \bar{x})\,(y_{t+k} - \bar{y})$$

Where σx and σy are the standard deviation of the two times series.

Physical and chemical analysis

We estimated concentration and export of nutrients (TN and TP) and CPOM. TN was

analyzed through digestion and subsequent reduction in cadmium column (Grasshoff et

al. 1983). TP was analyzed using acid digestion followed by evaluation of solubilized

phosphorus as soluble reactive phosphorus (A.P.H.A. 2005). We estimated Ash Free

Dry Weight of the CPOM using the Pozo et al. (2009) methodology. The export of TN,

TP and CPOM were calculated for each sample date by the following formula:

$$Nutrient\ export = \frac{[N]\,Q\,t}{A}$$

Where [N] is instantaneous nutrient concentration (mg m^{-3}), Q is instantaneous

discharge (m^3 s^{-1}), t is time (yr) and A is drainage basin area (km^2). The annual nutrient

export was an average of each data set (n=13).

We performed a paired T-test to compare nutrient concentration and export

between streams; to this end the data were transformed to comply with normality and

homoscedasticity assumptions. Last, we performed a Principal Component Analysis

(PCA) that included 12 variables: turbidity, temperature, EC, pH, discharge, TP, TN,

TN:TP, CPOM, TP export, TN export and CPOM export. PCs were calculated based on

standardized correlation matrix, using components with eigenvalues larger than 1.

Number Cruncher (Hintze 1998) was employed for statistical analysis and GraphPad

Prism version 6.01 was employed for graphics editing.

RESULTS

Climate and hydrology of the region

Air temperature was low between May and August and increased monotonically after

September, reaching its highest values from December to February. Annual

precipitation was 1164 mm and showed strong seasonality, with 80% falling between

mid-May and mid-September. The first two (#1-2) and last three (#11-13) samples thus

took place during a drier period; samples #3-6 were taken during the precipitation

period; the remaining samples (#7-10) fell within the period of monotonic air

temperature warming (Figure 2). Thus, samples were evenly distributed across three

climatically contrasting periods.

Cross-correlation of discharge with precipitation and temperature showed

opposing patterns. Discharge was positively correlated with precipitation, but negatively

correlated with air temperature (Figure 3c, d, Appendix S1). The discharge-precipitation

correlogram shows three periods that coincide with peaks of the correlation function

(Figure 3c). Both streams show differences at each of these peaks: CP had higher

correlation values (r max) and lower lag times than G. Indeed, at CP the correlation

reached its maximum the same day precipitation took place (lag = 0) during the first

period, while r was not even significant for G at the time (Appendix S1). During the

second period, correlation at CP reached an r max=0.36 with a lag time of 34 days,

while at G, correlation was lower (r max=0.19) and the lag higher (lag = 40). The third,

longest period presented a series of peaks and valleys, and lapsed from day 57 to day

113 after the precipitation event at CP and from day 62 to 145 at G. In contrast, the

discharge-temperature correlogram did not show similar periods; cross-correlation was

significant from day 0 to around day 90 for both streams (Figure 3d, Appendix S1).

The flow regime was contrasting between the streams. CP had a lower annual

average discharge value than G (Discharge CP= 3.364 m^3 s^{-1}; Discharge G= 4.635 m^3 s^-

^{1}; t =7.4437; $P < 0.0001$; d.f. = 728), but its Q^{sp} was higher (Q^{sp} CP= 0.051 m^3 s^{-1}; Q^{sp}

G= 0.029 m^3 s^{-1}; t=-24.7917; $P < 0.0001$; d.f. = 728). CP presented the most extreme

discharge values (Figure 3a) and a steep autocorrelation function with a memory effect

at k = 38, while G revealed a more gently sloped function with a memory effect at k =

61 (Figure 3b). Thus, CP showed lower memory effect than G, indicating a higher

flashiness of stream discharge.

Figure 3. a) Hydrographs of Casa de Piedra (CP) and Gutiérrez (G) streams, black dots
and numbers indicate sampling dates. b) Autocorrelation functions and memory effects
(r (k) = 0.2) of the discharge. c) Cross-correlations between discharge and precipitation,
three different periods are shown, and d) cross-correlations between discharge and
temperature.

 Physical and chemical characteristics of the streams studied

Although streams differed in the specific values, both had very low EC, turbidity,

nutrient concentration and nutrient export (Table 2). G showed the highest TP and TN

concentrations (Figure 4a, c) and exports (Figure 4b, d): CP exported 8.6 kg TP km^{-2} yr^{-1}

and 57.5 kg TN km^{-2} yr^{-1}, while G exported 11.6 kg TP km^{-2} yr^{-1} and 78.4 kg TN km^{-2}

yr^{-1} (approximately 35 % more P and N). No significant differences in TN:TP average

ratios were found (~15; Figure 4e), although the median value of TN:TP was 11 for CP

and 16 for G. As to the amount of CPOM exported, no significant difference between

streams was found. There were differences, however, in the quality of exported

material; native *N. pumilio* leaves dominated in CP, while introduced *S. fragilis* leaves

dominated in G. The peaks of CPOM export also differed among streams: highest

export at CP was found during June and July (samples #3-4), while at G this peak

occurred earlier, in April (sample #2). In summary, the drainage basin showing more

anthropic land uses also showed higher nutrient export, with most CPOM export

represented by introduced tree species.

	Casa de Piedra	Gutiérrez
Temperature (°C)	7.8 (3.7-12.5)	10.6 (6.0-16.8)
Electrical Conductivity (μs cm^{-1})	43 (27-61)	74 (71-76)
Turbidity (NTU)	0.2 (0.0-0.8)	0.7 (0.0-2.0)
pH	7.8 (7.4-8.0)	7.9 (7.3-8.1)
Total Phosphorus (mg m^{-3})	5 (2-9)	14 (10-18)
Total Nitrogen (mg m^{-3})	29 (18-44)	94 (75-134)
TN:TP (molar)	15 (6-31)	15 (11-21)
Coarse Particulate Organic Matter (AFDW m^{-3})*10	0.46 (0.04-1.86)	0.91 (0.02-8.76)
Total Phosphorus export (kg km^{-2} yr^{-1})	8.6 (1.7-25.4)	11.6 (3.8-22.5)
Total Nitrogen export (kg km^{-2} yr^{-1})	57.5 (7.9-129.6)	78.4 (27.3-184.8)
Coarse Particulate Organic Matter export (kg km^{-2} yr^{-1})	126.4 (1.3-750.4)	43.1 (1.5-317.9)

341 Table 2. Average values, maximum and minimum in brackets, of the physical and
342 chemical variables in the streams (n=13, except for the Coarse particulate organic
343 matter where n=12).

Figure 4. Box plot showing annual differences in nutrient concentration, ratio and
export in Casa de Piedra (CP) and Gutiérrez (G) streams (Paired T Test, * P ≤ 0.05; * *
P ≤ 0.01; * ** P ≤ 0.001; ns: not significant). f) The Y axis is in logarithmic scale. Total
Phosphorus (TP), Total Nitrogen (TN) and Coarse Particulate Organic Matter (CPOM).

Our PCA determined three Principal Component, FI, FII and FIII that

respectively explain 33%, 24% and 13% of the variance, accounting for 70% of the total

variance (Table 3). FI was mainly related to discharge and nutrient export (TP$_{exp}$ and

TN$_{exp}$) (Figure 5a). Positive FI values corresponded to samples with high discharge,

high nutrient export and high turbidity. The FII was mainly related to the EC of the

water, and to a lesser extent with water temperature and nutrient concentration (Figure

5a). The FIII was mainly related to the TN:TP ratio, and to a lesser extent with CPOM

variables.

Variable	First Factor (33%)	Second Factor (24%)	Third Factor (13%)
Temperature		0.31	
EC		0.47	
Turbidity	0.35		
Q	0.38		
TP	0.35	0.41	
TN	0.36	0.32	
TN:TP			0.55
CPOM			-0.47
TP_{exp}	0.42		
TN_{exp}	0.42		
$CPOM_{exp}$		-0.35	-0.55

361 Table 3. Principal Component Analysis on physical and chemical variables. Factors are
362 a linear combination of the different variables. Percent of variance explained and factor
363 loadings (> 0.30) are shown. Abbreviations are as in Figure 5.
364

Both streams are clearly discriminated by PCA ordering, highlighting seasonal

differences in each stream (Figure 5b). Besides separating samples from each stream,

plotting FI versus FII shows that each stream is temporally heterogeneous, allowing

clustering in 3 groups for CP (CPA, CPB and CPC) and 2 for G (GA and GB) (Figure

5b). CPA had low discharge and export rate values and high temperatures, taking place

at the warmest period (Figure 2). CPB had higher nutrient concentration and export

values, matching the period with more abundant precipitation (Figure 2). CPC was

mainly characterized by low EC, nutrient concentration and turbidity values, and

coincides with the period of diminishing rainfall and monotonic air temperature

warming (Figure 2). GA had similar characteristics to CPA and spans the same period.

The remaining samples fall within GB. Thus, the values of physical and chemical

variables sampled monthly through the year show that both streams have marked

seasonal differences that follow precipitation and air temperature patterns.

Figure 5. Principal Component Analysis (PCA) of physical and chemical variables in
the study streams. a) The weight of variables in each of the axes is shown. b) First
factorial plane of the PCA, Casa de Piedra sampling dates are represented by triangles
and Gutiérrez ones are represented by rectangles, the numbers correspond to the
sampling dates and they are grouped by climatic characteristics of the region (a, b and
c). Electrical conductivity (EC), Discharge (Q), Total Phosphorus (TP), Total Nitrogen
(TN), Coarse Particulate Organic Matter (CPOM), Export (exp).
387

DISCUSSION AND CONCLUSIONS

We surveyed discharge and nutrient dynamics in two adjacent North Patagonian streams

with contrasting drainage basins. Precipitation had a direct influence on stream

discharge at both streams, but response time and amplitude varied between them. CP

stream was flashy, showed a tighter relation between precipitation at the basin and

discharge response and stronger seasonality of physical and chemical parameter values.

G stream, in contrast, showed a more buffered discharge response to precipitation and

higher nutrient exports. In general, similarities seen between streams can be related to

regional climate and geology, while differences are better explained by contrasting

topography and land use of the respective drainage basins.

Precipitation and temperature have opposite influence in river discharge in the

Andean region (Masiokas et al. 2008). Precipitation increased stream discharge of our

study sites. Run-off, groundwater and snowmelt, the different pathways taken by

precipitated water to reach the streams and their relative importance are evidenced by

the three periods found in the discharge-precipitation correlogram. Surface run-off was

relevant at CP, its basin characterized by steep slopes. In this stream, discharge

increased drastically during the same day precipitation occurred, and then during the

following days. On the other hand, surface run-off was not relevant at G stream, likely

due to buffering by the large upstream lake. However, at a shorter temporal scale,

Sosnovsky et al. (2020) reported a positive relation between G stream discharge and

precipitation with a 4-days lag during the storm season. Water not reaching the stream

immediately as run-off infiltrates into the soil and moves to the stream as sub-surface

run-off or groundwater; these pathways usually result in delayed stream discharge

responses (Kalff 2002). Sub-surface run-off and groundwater pathways were more

relevant at CP, which showed a higher r(max). However, delay time was similar for

both streams; this is likely due to shared basin characteristics driving groundwater

dynamics: lithology, geomorphology and soil development (Giacosa et al. 2001). In

colder regions, snow precipitation is stored during the winter months and released as

snowmelt in spring and summer (Bailey 1995). This results in a delay time between

precipitation and discharge response of about 2-4 months, as we found at both study

sites during spring. The observed discharge dynamics and its relationship with

precipitation (either rain or snow) agree with the different water pathways to streams

described at other mountainous regions (Brown et al. 2003; Ahearn et al. 2004). On the

other hand, air temperature decreased stream discharge of our study sites, suggesting

that direct evaporation and plant transpiration across the basin might have a sizable

effect on the amount of water reaching the streams. Interestingly, Sosnovsky et al.

(2020) reported a positive correlation between air temperature and discharge at shorter

timescales during the melting season, suggesting that the influence of air temperature on

discharge is both scale and context dependent. Altogether, the same meteorological

variable (air temperature) might modulate stream dynamics at several different points

along the precipitation/runoff/discharge cycle, sometimes even in opposite direction

(e.g., warm temperatures increasing snowmelt but decreasing runoff into stream through

evapotranspiration).

Nutrient concentration and annual export rate from CP reflect the usual

oligotrophy of north Patagonian streams (Pedrozo et al. 1993; García RD et al. 2015).

Values of N and P concentrations were much smaller than those found in either a sub-

catchment that had suffered a wildfire (TN = 675.1 µg l^{-1} and TP = 12.0 µg l^{-1})

(Temporetti 2006) or in areas of low anthropogenic impact in Patagonia (average TN

export = 100 kg km^{-2} yr^{-1}) (Little et al. 2008). Indeed, values of P export were between

the lowest measured in South America (TP export range from 4 to 450 kg TP km^{-2} yr^{-1})

(Álvarez-Cobelas and Angeler 2007). These results indicate that nutrient export in CP

stream and similar ecosystems are among the lowest measured for any basin in the

Southern hemisphere (Quinn and Stroud 2002; Temporetti 2006; Álvarez-Cobelas and

Angeler 2007; Little et al. 2008) and worldwide (Álvarez-Cobelas et al. 2008; Álvarez-

Cobelas et al. 2009). Such low export values could be explained by low atmospheric N

deposition (Holland 1996) and relatively light land use in this region. It must be noted,

however, that nutrient load was measured on a monthly basis, in the absence of

precipitation, when overall nutrient export is lower (Abell et al. 2013; Tate and Singer

2013). This implies that our estimates should be considered as baseline values. In any

case, it is interesting to note that nutrient export values were higher for G than for CP,

even though the opposite result could have been expected, given that headwater lakes

act as nutrient and sediment traps (Little et al. 2008; Parker et al. 2009).

Topography and land cover are environmental factors that control spatial

patterns in river ecosystems (Snelder and Biggs 2002). For example, the presence of a

headwater lake in the drainage basin regulates the flow regime of a stream (Baker et al.

2004). This was made evident by the contrasting values and dynamics of discharge

autocorrelation between CP and G streams. The autocorrelation function fell sharply for

CP but sloped more gently for G. On the other hand, nutrient exports were higher in G

despite the expected trapping effect of its headwater lake. This could be explained by

land use of its drainage basin, namely the presence of human settlements and a trout

farm. While we did not formally weigh the relative impacts of fish farming and

urbanization, we suspect that a growing residential settlement of the Gutiérrez stream

drainage basin, often times without adequate wastewater treating infrastructure, is

expected to have a larger impact than the detritus from a fish farm producing ~10 tons

of fish per year. Moreover, the additional nutrient sources of fish food and feces from

salmonid production would be expected to decrease the TN:TP ratio of the water

column in Patagonia reservoirs (Diaz et al. 2001), yet we saw an opposite effect at G

stream. Our results suggest that while topography is the main driver of G stream

hydrology (Sosnovsky et al. 2020), nutrient dynamics are mainly driven by land use.

The molecular N:P ratio is said to distinguish nitrogen-limited systems from

those constrained by phosphorus (Rhee and Gotham 1980). Kahlert (1998, 2001) found

that the optimal N:P ratio of freshwater benthic algae was 18:1 (molar), which deviates

from the phytoplankton optimal stoichiometric N:P ratio of 16:1 (Redfield ratio)

(Goldman et al. 1979). CP and G streams had an arithmetic mean N:P molar ratio close

to 15, which would indicate nitrogen deficit in these ecosystems, similar to what has

been observed in other ecosystems of Northern Patagonia, either forests (Satti et al.

2003) or lakes (Diaz et al. 2007). However, considering that median values of N:P were

11 for CP and 16 for G, it becomes evident that nitrogen limitation is being lifted at G,

most likely due to wastewater from human settlements. Wastewater releases phosphorus

and nitrogen to the soil. While Andisols have high capacity in phosphorus retention

(Satti et al. 2003), nitrate leaching would reach the stream through the groundwater,

priming an eutrophication process of the G stream and downstream water bodies. This

finding at the drainage basin level is in line with observations at the global scale

suggesting human actions generate a clear unbalance favoring N over P which may

induce significant alterations at organism, community and ecosystem levels (Peñuelas et

al. 2013).

Leaves of riparian trees are generally the largest component of allochthonous

inputs to stream located in forested drainage basins and are often the major source of

energy to heterotrophic organisms (Vannote et al. 1980; Webster et al. 1999). In the

case of Andean streams, *N. pumilio* leaves cover large areas of the waterway during

autumn (Albariño et al. 2009). A substantial amount of these may be transported

downstream by spates occurring from late autumn to winter and eventually reaches the

lake (Modenutti et al. 2010). We observed no significant differences in the amount of

CPOM exported by the two streams studied. However, we highlight the fact that in

stream CP the CPOM was mainly made up of native *N. pumilio* leaves, while in stream

G introduced *S. fragilis* leaves dominated. This corresponds to the tree species present

in the different catchments, and could explain the temporal difference between streams

in the surge of CPOM; the peak in CPOM export occurred during June and July in

stream CP, and during April in stream G. This supports the claim that afforestation or

invasion of exotic plants with different phenology, chemical and physical characteristics

to those of the native riparian species changes the timing, quantity and quality of leaf

litter standing stock in streams (Naiman et al. 2005).

By studying two Andean streams located at adjacent, yet contrasting drainage

basins, we were able to investigate the influence of environmental factors acting at

different spatial scales over these freshwater ecosystems. At a regional scale, climate

determined seasonality of water flow for both streams. At a landscape scale, we found

that the presence of a large, deep lake buffered the hydrological dynamic of its effluent

stream. At a lower spatial scale, we found that more intense human impact at G's

drainage basin correlated with higher nutrient export rates by the stream. We could

propose, therefore, that the hydrological and nutrients dynamics are decoupled in this

ecosystem. Considering the strong demographic pressure at G's drainage basin, there is

a high risk of change in N:P stoichiometry and ecosystem eutrophication. Gutiérrez

basin situation thus shows a stark contrast to Casa de Piedra basin, which instead has

the hydrological, physical and chemical dynamics of a pristine drainage basin

characteristics of Andean streams originating at high altitude wetlands and lakes.

Besides geomorphology, the main differences between both basins are land use policies:

CP's basin is mostly within protected areas, while most of G's lower basin is within the

limits of San Carlos de Bariloche city and subject to several, potentially high impact

land uses, like residential neighbourhoods, winter sports and fish farming, among

others. Local and global environmental problems are currently burning issues in stream

ecosystem processes and will be even more so in the future. Considering that lakes and

streams are sentinels and integrators of environmental change in the surrounding

terrestrial landscape (Williamson et al. 2008), it is really important to implement

monitoring schemes for key variables of these aquatic ecosystems (Lovett et al. 2007)

and encourage restoration and sustainable management of their drainage basins.

ACKNOWLEDGMENTS

We thank Rio Negro's province Water Department and Argentina's National Weather

Service for providing hydrological and meteorological data; and Ahearn D.S. for his

interesting comments during the data analysis and writing periods. This study was

supported by the Agencia Nacional de Promoción Científica y Tecnológica under Grant

number 2959.

REFERENCES

A.P.H.A. 2005. Standard methods for examination of water and wastewater. 21 ed.
Washington DC: American Public Health Association.
Abell JM, Hamilton DP, Rutherfor JC. 2013. Quantifying temporal and spatial
variations in sediment, nitrogen and phosphorus transport in stream inflows to a
large eutrophic lake. Envirnomental Science Processes and Impacts.
Ahearn DS, Sheibley RW, Dahlgren RA, Keller KE. 2004. Temporal dynamics of
stream water chemistry in the last free-flowing river draining the western Sierra
Nevada, California. Journal of Hydrology. 295:47-63.
Albariño RJ, Díaz Villanueva V, Buria L. 2009. Leaf litter dynamics in a forested small
Andean catchment, northern Patagonia, Argentina. In: Oyarzún C, Verhoest
NEC, Boeckx P et al., editors. Ecological advances on Chilean temperate
rainforest. Academia Press; p. 183-211.
Álvarez-Cobelas M, Angeler DG. 2007. Exportación de nutrientes en las cuencas
hidrográficas de Latinoamérica: una recopilación. Revista Latinoamericana de
Recursos Naturales. 3(1):31-43.
Álvarez-Cobelas M, Angeler DG, Sánchez-Carrillo S. 2008. Export of nitrogen from
catchments: A worldwide analysis. environmental Pollution. 156:261-269.
Álvarez-Cobelas M, Sánchez-Carrillo S, Angeler DG. 2009. Phosphorus export from
catchments: a global view. Journal of American Benthological Society.
28(4):805-820.
Allan JD. 2004. Landscapes and Riverscapes: The influence of land use on stream
ecosystems. Annual Review of Ecology, Evolution and Systematics. 35:257-
284.
Bailey RG. 1995. Ecosystem Geography. New York, New York: Springer-Verlag.
Baker DB, Richards RP, Loftus TT, Kramer JW. 2004. A new flashiness index:
Characteristics and applications to midwestern rivers and streams. Journal of the
American Water Resources Association. 40(2):503-522.
Brown LE, Hannah DM, Milner AM. 2003. Alpine Stream Habitat Classification: An
Alternative Approach Incorporating the Role of Dynamic Water Source
Contributions. Artic, Antartic, and Alpine Research. 35(3):313-322.
Carpenter SR, Caraco NF, Correll DL, Howarth RW, Sharpley AN, Smith VH. 1998.
Nonpoint Pollution of Surface Waters with Phosphorus and Nitrogen. Ecological
Applications. 8(3):559-568.
Carrillo U, Díaz Villanueva V, Modenutti BE. 2018. Sustained effects of volcanic ash
on biofilm stoichiometry, enzyme activity and community composition in
North-Patagonia streams. Science of the Total Environment. 621:235-244.
Diaz MM, Pedrozo FL, Reynolds CS, Temporetti PF. 2007. Chemical composition and
the nitrogen-regulated trophic state of Patagonian lakes. Limnologica. 37:17-27.
Diaz MM, Temporetti PF, Pedrozo FL. 2001. Response of phytoplankton to enrichment
from cage fish farm waste in Alicura Reservoir (Patagonia, Argentina). Lake &
Reservoir Management. 6:151-158.
Díaz Villanueva V, Bastidas Navarro M, Albariño R. 2016. Seasonal patterns of organic
matter stoichiometry along a mountain catchment. Hydrobiologia.
Diehl L, Mazzarino MJ, Funes F, Fontenla S, Gobbi M, Ferrari J. 2003. Nutrient
conservation strategies in native Andean-Patagonian forests. Journal of
Vegetation Science. 14:63-70.
Dodds WK, Smith VH. 2016. Nitrogen, phosphorus, and eutrophication in streams.
Inland Waters. 6:155-164.

Emerson DG, Vecchia AV, Dahl AL. 2005. Evaluation of Drainage-Area Ratio Method
Used to Estimate Streamflow for the Red River of the North Basin, North
Dakota and Minnesota. Reston, Virginia: U.S. Geological Survey.
García DR, Diéguez MC, Gerea M, García PE. 2018. Characterisation and reactivty
continuum of dissolved organic matter in forested headwater catchments of
Andean Patagonia. Freshwater Biology. 63(9):1049-1062.
García P, Diéguez MC, Queimaliños CP. 2015. Landscape integration of North
Patagonian mountain lakes: a first approach using characterization of dissolved
organic matter. Lakes & Reservoirs: Research and Management. 20:1-14.
García RD, Reissig M, Queimaliños CP, García PE, Diéguez MC. 2015. Climate-driven
terrestrial inputs in ultraoligotrophic mountain streams of Andean Patagonia
revealed through chromophoric and fluorescent dissolved organic matter.
Science of the Total Environment. 521-522:280-292.
Giacosa R, Heredia N, González R, Faroux A, Césari O, Franchi M. 2001. Hoja
Geológica 4172-IV, San Carlos de Bariloche. Provincias de Río Negro y
Neuquén.
Goldman JC, Mc Carthy JJ, Peavey DG. 1979. Growth rate influence on the chemical
composition of phytoplankton in oceanic waters. Nature. 279:210-215.
Gordon ND, McMahon TA, Gippel CJ, Natha RJ. 2004. Stream Hydrology An
Introduction for Ecologist. Chichester, England: John Wiley & Sons LTD.
Grasshoff M, Ehrhardt K, Kremling K. 1983. Methods of seawater analysis. New York:
Wiley.
Hintze JL. 1998. Number Cruncher Statistical System (NCSS). Version 2000. User´s
Guide. Utah: Kaysville.
Holland EA, Dentener FJ, Braswell BH, Sulzman JM. 1999. Contemporary and pre-
industrial global reactive nitrogen budgets. Biogeochemistry. 46:7-43.
Howarth RW, Billen G, Swaney D, Townsend A, Jaworski NA, Lajtha K, Downing JA,
Elmgren R, Caraco N, Jordan T et al. 1996. Regional nitrogen budgets and
riverine N & P fluxes for the drainages to the North Atlantic Ocean: natural and
human influences. Biogeochemistry. 35:75-139.
Hynes HBN. 1975. The stream and its valley. Verhandlungen der Internationale
Vereingung für Limnologie. 19:1-15.
Jordan TE, Correll DL, Weller DE. 1997. Effects of agriculture on discharges of
nutrients from coastal plain watersheds of Chesapeake Bay. Journal of
Environmental Quality. 26:836-848.
Kahlert M. 1998. C:N:P ratios of freshwater benthic algae. Archiv für Hydrobiologie,
Special Issues, Advances in Limnology. 51:105-114.
Kahlert M. 2001. Biomass and Nutrient Status of Benthic Algae in Lakes. Acta
Universitatis Upsaliensis: Uppsala.
Kalff J. 2002. Chapter 5, Hidrology and Climate. In: Ryu T, editor. Limnology Inland
Water Ecosystems. Upper Saddle River: Prentice Hall; p. 53-71.
Lallement M, Macchi PJ, Vigliano PH, Juarez S, Rechencq M, Baker M, Bouwes N,
Crowl TA. 2016. Rising from the ashes: Change in salmonid fish assemblages
after 30 months of the Puyehue-Cordon Caulle volcanic eruption. Science of the
Total Environment. 541:1041-1051.
Larocque M, Mangin A, Razack M, Banton O. 1998. Contribution of correlation and
spectral analyses to the regional study of a large karst aquifer (Charente,
France). Journal of Hydrology. 205:217-231.

Little C, Soto D, Lara A, Cuevas JG. 2008. Nitrogen exports at multiple-scales in a
southern Chilean watershed (Patagonian Lakes district). Biogeochemistry.
87:297-309.
Lovett GM, Burns DA, Driscoll CT, Jenkins JC, Mitchell MJ, Rustad L, Shanley JB,
Likens GE, Haeuber R. 2007. Who need environmental monitoring? Frontiers in
Ecology and the Envirnonment. 5(5):253-260.
Mangin A. 1984. Pour une meilleure connaissance des systèmes hydrologgiques à partir
des analyses corrélatoire et spectrale. Journal of Hydrology. 67:25-43.
Masiokas MH, Villalba R, Luckman BH, Lascano ME, Delgado S, Stepanek P. 2008.
20th-century recession and regional hydroclimatic changes in northwestern
Patagonia. Global and Planetary Change. 60:85-100.
Merseburger GC, Martí E, Sabater F. 2005. Net changes in nutrient concentrations
below a point source input in two streams draining catchments with contrasting
land uses. Science of the Total Environment. 347:217-229.
Meyer JL, Paul MJ, Taulbee WK. 2005. Stream ecosystem function in urbanizing
landscapes. Journal of the North American Benthological Society. 24:602-612.
Miserendino ML, Kutschker MA, Brand C, La Manna L, Di Prinzio YC, Papazian G,
Bava J. 2016. Ecological Status of a Patagonian Mountain River: Usefulness of
Environmental and Biotic Metrics for Rehabilitation Assessment. Environmental
Management. 57(6):1166-1187.
Modenutti BE, Albariño RJ, Bastidas Navarro M, Villanueva VD, Souza AF, Trochine
C, Laspoumaderes C, Cuassolo F, Mariluán G, Buria L et al. 2010. Structure and
dynamic of food webs in Andean North Patagonian freshwater systems: organic
matter, light and nutrient relationships. Ecología Austral. 20(2):95-114.
Naiman RJ, Décamps H, McClain ME. 2005. Riparia: ecology, conservation and
management of streamside communities. London, UK: Elsevier Academic Press.
Oyarzún CE, Godoy R, Schrijver De A, Staelens J, Lust N. 2004. Water chemistry and
nutrient budgets in an undisturbed evergreen rainforest of souther Chile.
Biogeochemistry. 71:107-123.
Parker BP, Schindler DE, Beaty KG, Stainton MP, Kasian SEM. 2009. Long-term
changes in climate, streamflow, and nutrient budgets for first-order catchments
at the Experimental Lakes Area (Ontario, Canada). Canadian Journal of
Fisheries and Aquatic Sciences. 66:1848-1863.
Paruelo JM, Beltran A, Jobbágy E, Sala O, Golluscio R. 1998. The climate of
Patagonia: general patterns and controls on biotic processes. Ecología Austral.
8(2):85-101.
Pedrozo FL, Chillrud S, Temporetti PF, Diaz MM. 1993. Chemical composition and
nutrient limitation in rivers and lakes of northern Patagonian Andes (39.5º-42º S;
71º W) (Rep. Argentina). SIL Proceedings, 1922-2010. 25:207-214.
Peel MC, Finlayson BL, McMahon TA. 2007. Updated world map of the Köppen-
Geiger climate classification. Hydrology and Earth System Sciences. 4(2):439-
473.
Peltier MR, Wilcox CJ, Sharp DC. 1998. Technical Note: Application of the Box-Cox
data transformation to animal science experiments. Journal of Animal Science.
78:847-849.
Peñuelas J, Poulter B, Sardans J, Ciais P, van der Velde M, Bopp L, Boucher O,
Godderis Y, Hinsinger P, Llusia J et al. 2013. Human-induced nitrogen–
phosphorus imbalances alter natural and managed ecosystems across the globe.
Nature Communications. 4:2934.

Pereyra FX, Albertoni J, Bréard C, Cavaliaro S, Coccia M, Ducós E, Dzendoletas M,
Fookes S, Getino E, Helms F et al. 2005. Estudio geocientífico aplicado al
ordenamiento territorial, San Carlos de Bariloche.
Pieterse NM, Bleuten W, Jørgensen SE. 2003. Contribution of point sources and diffuse
sources to nitrogen and phosphorus loads in lowland rivers tributaries. Journal of
Hydrology. 271:213-225.
Poff NL, Allan JD, Bain MB, Karr JR, Prestegaard KL, Richter BD, Sparks RE,
Stromberg JC. 1997. The Natural Flow Regime. Bioscience. 47(11):769-784.
Pozo J, Elosegi A, Díez J, Molinero J. 2009. Chapter 10, Dinámica y relevancia de la
materia orgánica. In: Elosegi A, Sabater S, editors. Conceptos y técnicas en
ecología fluvial. Bilbao, España: Fundación BBVA; p. 141-166.
Quinn JM, Stroud MJ. 2002. Water quality and sediment and nutrient export from New
Zealand hill-land. New Zealand Journal of Marine and Freshwater Research.
36(2):409-429.
Rhee GY, Gotham IJ. 1980. Optimum N:P ratios and co-existence of planktonic algae.
Journal of Phycology. 16:486-489.
Satti P, Mazzarino MJ, Gobbi M. 2003. Soil N dynamics in relation to leaf litter quality
and soil fertility in north-western Patagonian forests. Journal of Ecology.
91:173-181.
Snelder TH, Biggs BJF. 2002. Multiscale river environment classification for water
resources management. Journal of the American Water Resources Association.
38:1225-1239.
Sosnovsky A, Rechencq M, Fernández V, Suarez MJ, Cantet RJC. 2020. Hydrological
and Physico-chemical dynamics in two Andean streams. Limnetica. 39(1):17-33.
Tate KW, Singer MJ. 2013. Timing, Frecuency of Sampling Affect Accuracy of Water-
Quality Monitoring. California Agriculture. 53(6):44-48.
Temporetti PF. 2006. Efecto a largo plazo de los incendios forestales en la calidad del
agua de dos arroyos en la sub-región Andino-Patagónica, Argentina. Ecología
Austral. 16(2):157-166.
Vannote RL, Minshall GW, Cummins KW, Sedell JR, Cushing CE. 1980. The River
Continuum Concept. Canadian Journal of Fisheries and Aquatic Sciences.
37:130-137.
Walsh CJ, Fletcher TD, Ladson AR. 2005. Stream restoration in urban catchments
through redesigning stormwater systems: looking to the catchment to save the
stream. Journal of the North American Benthological Society. 24(690-705).
Webster JR, Benfield EF, Ehrman TP, Schaeffer MA, Tank JL, Hutchens JJ, D´angelo
DJ. 1999. What happens to allochthonous material that falls into streams? A
synthesis of new and published information from Coweeta. Freshwater Biology.
41:687-705.
Wiens JA. 2002. Riverine landscapes: taking landscape ecology into the water.
Freshwater Biology. 47:501-515.
Williams Subiza EA, Brand C. 2018. Short-term effect of wildfire on Patagonian
headwater streams. International Journal of Wildland Fire. 27:457-470.
Williamson CE, Dodds WK, Kratz TK, Palmer MA. 2008. Lakes and streams as
sentinels of environmental change in terrestrial and atmospheric processes.
Frontiers in Ecology and the Envirnonment. 6(5):247-254.

Appendix S1. Cross-correlation between climatic variables and discharge (* P ≤ 0.05; * * P ≤ 0.01; * ** P ≤ 0.001; ns: not significant).

Lag (days)	Discharge-Precipitation				Lag (days)	Discharge-Temperature			
	r_{CP}	P_{CP}	r_G	P_G		r_{CP}	P_{CP}	r_G	P_G
0	0.31	***	-0.04	NS	0	-0.38	***	-0.13	**
1	0.30	***	-0.03	NS	1	-0.37	***	-0.13	**
2	0.27	***	-0.03	NS	2	-0.38	***	-0.14	***
3	0.25	***	-0.02	NS	3	-0.39	***	-0.15	***
4	0.26	***	-0.01	NS	4	-0.40	***	-0.16	***
5	0.26	***	0.01	NS	5	-0.42	***	-0.17	***
6	0.25	***	0.01	NS	6	-0.44	***	-0.19	***
7	0.23	***	0.01	NS	7	-0.44	***	-0.20	***
8	0.24	***	0.02	NS	8	-0.44	***	-0.20	***
9	0.23	***	0.02	NS	9	-0.44	***	-0.21	***
10	0.19	***	0.01	NS	10	-0.44	***	-0.22	***
11	0.15	***	0.00	NS	11	-0.44	***	-0.22	***
12	0.13	**	-0.01	NS	12	-0.46	***	-0.24	***
13	0.14	**	-0.00	NS	13	-0.49	***	-0.25	***
14	0.14	**	-0.00	NS	14	-0.51	***	-0.27	***
15	0.14	**	0.00	NS	15	-0.53	***	-0.29	***
16	0.12	**	-0.00	NS	16	-0.55	***	-0.31	***
17	0.13	**	0.00	NS	17	-0.55	***	-0.32	***
18	0.11	**	0.00	NS	18	-0.55	***	-0.34	***
19	0.10	**	-0.00	NS	19	-0.59	***	-0.35	***
20	0.11	**	0.01	NS	20	-0.57	***	-0.36	***
21	0.14	**	0.02	NS	21	-0.57	***	-0.36	***
22	0.16	**	0.03	NS	22	-0.57	***	-0.37	***
23	0.17	**	0.03	NS	23	-0.57	***	-0.38	***
24	0.17	**	0.03	NS	24	-0.57	***	-0.39	***
25	0.19	***	0.05	NS	25	-0.58	***	-0.40	***
26	0.20	***	0.06	NS	26	-0.59	***	-0.41	***
27	0.19	***	0.06	NS	27	-0.60	***	-0.43	***
28	0.20	***	0.07	NS	28	-0.61	***	-0.44	***
29	0.20	***	0.08	NS	29	-0.61	***	-0.45	***
30	0.22	***	0.10	*	30	-0.61	***	-0.45	***
31	0.22	***	0.11	*	31	-0.61	***	-0.46	***
32	0.23	***	0.11	*	32	-0.61	***	-0.47	***
33	0.24	***	0.13	**	33	-0.60	***	-0.48	***
34	0.24	***	0.14	**	34	-0.60	***	-0.48	***
35	0.24	***	0.14	**	35	-0.59	***	-0.49	***
36	0.22	***	0.15	**	36	-0.59	***	-0.49	***
37	0.20	**	0.14	**	37	-0.58	**	-0.50	***
38	0.18	**	0.14	**	38	-0.58	**	-0.51	***
39	0.15	*	0.13	**	39	-0.57	**	-0.51	***
40	0.14	*	0.13	**	40	-0.58	**	-0.52	***
41	0.11	NS	0.12	*	41	-0.58	**	-0.52	***
42	0.10	NS	0.12	*	42	-0.58	**	-0.53	***
43	0.09	NS	0.12	*	43	-0.58	**	-0.54	***
44	0.09	NS	0.12	*	44	-0.59	**	-0.55	***
45	0.10	NS	0.12	*	45	-0.60	**	-0.56	***

46	0.11	NS	0.12	*	46	-0.61	**	-0.58	***
47	0.10	NS	0.11	*	47	-0.62	**	-0.59	***
48	0.10	NS	0.10	NS	48	-0.64	**	-0.60	***
49	0.11	NS	0.11	*	49	-0.66	**	-0.62	***
50	0.13	NS	0.12	*	50	-0.67	**	-0.63	***
51	0.12	NS	0.11	NS	51	-0.67	**	-0.64	***
52	0.13	NS	0.10	NS	52	-0.68	**	-0.65	***
53	0.13	NS	0.00	NS	53	-0.68	**	-0.66	***
54	0.16	NS	0.00	NS	54	-0.67	**	-0.66	***
55	0.17	*	0.00	NS	55	-0.66	**	-0.67	***
56	0.18	*	0.00	NS	56	-0.66	**	-0.67	***
57	0.20	*	0.00	*	57	-0.66	**	-0.68	***
58	0.23	**	0.00	*	58	-0.67	**	-0.69	***
59	0.25	**	0.00	*	59	-0.68	**	-0.69	***
60	0.24	**	0.00	*	60	-0.68	**	-0.70	***
61	0.25	**	0.00	**	61	-0.67	**	-0.69	**
62	0.25	**	0.00	**	62	-0.66	**	-0.69	**
63	0.24	**	0.00	**	63	-0.66	**	-0.69	**
64	0.23	*	0.00	**	64	-0.67	**	-0.69	**
65	0.23	*	0.00	**	65	-0.67	**	-0.69	**
66	0.23	*	0.00	**	66	-0.65	**	-0.69	**
67	0.24	**	0.00	**	67	-0.64	**	-0.69	**
68	0.27	**	0.00	**	68	-0.63	**	-0.68	**
69	0.30	**	0.00	***	69	-0.63	**	-0.68	**
70	0.31	**	0.00	***	70	-0.63	**	-0.68	**
71	0.33	***	0.00	***	71	-0.64	**	-0.69	**
72	0.33	***	0.00	***	72	-0.64	*	-0.69	**
73	0.33	***	0.00	***	73	-0.64	*	-0.69	**
74	0.32	**	0.00	***	74	-0.63	*	-0.69	*
75	0.32	**	0.00	***	75	-0.62	*	-0.69	*
76	0.31	**	0.00	***	76	-0.61	*	-0.69	*
77	0.30	**	0.00	***	77	-0.62	*	-0.69	*
78	0.27	**	0.00	***	78	-0.62	*	-0.70	*
79	0.25	*	0.00	***	79	-0.62	*	-0.70	*
80	0.26	*	0.00	***	80	-0.63	*	-0.71	*
81	0.25	*	0.00	***	81	-0.64	*	-0.72	*
82	0.26	*	0.00	***	82	-0.65	*	-0.73	*
83	0.30	**	0.00	***	83	-0.65	*	-0.73	*
84	0.30	**	0.00	***	84	-0.65	*	-0.73	*
85	0.30	**	0.00	***	85	-0.64	*	-0.73	*
86	0.29	**	0.00	***	86	-0.62	*	-0.72	*
87	0.29	**	0.00	***	87	-0.60	*	-0.71	*
88	0.29	**	0.00	***	88	-0.59	*	-0.71	*
89	0.30	**	0.00	***	89	-0.58	*	-0.70	*
90	0.30	**	0.00	***	90	-0.57	*	-0.69	*
91	0.30	**	0.00	***	91	-0.55	*	-0.68	NS
92	0.30	**	0.00	***	92	-0.54	*	-0.67	NS
93	0.31	**	0.00	***	93	-0.53	NS	-0.67	NS
94	0.31	**	0.00	***	94	-0.50	NS	-0.65	NS
95	0.30	**	0.00	***	95	-0.47	NS	-0.64	NS
96	0.31	**	0.00	***	96	-0.45	NS	-0.62	NS
97	0.33	**	0.00	***	97	-0.43	NS	-0.61	NS
98	0.34	**	0.00	***	98	-0.41	NS	-0.60	NS
99	0.35	**	0.00	***	99	-0.40	NS	-0.59	NS

100	0.37	**	0.00	***	100	-0.40	NS	-0.58	NS
101	0.37	**	0.00	***	101	-0.41	NS	-0.58	NS
102	0.38	**	0.00	***	102	-0.41	NS	-0.57	NS
103	0.37	**	0.00	***	103	-0.41	NS	-0.57	NS
104	0.34	**	0.00	***	104	-0.40	NS	-0.56	NS
105	0.34	**	0.00	***	105	-0.40	NS	-0.55	NS
106	0.33	**	0.00	***	106	-0.39	NS	-0.55	NS
107	0.31	**	0.00	***	107	-0.38	NS	-0.54	NS
108	0.30	*	0.00	***	108	-0.38	NS	-0.54	NS
109	0.29	*	0.00	***	109	-0.37	NS	-0.53	NS
110	0.28	*	0.41	***	110	-0.35	NS	-0.52	NS
111	0.27	*	0.41	***	111	-0.34	NS	-0.51	NS
112	0.26	*	0.40	**	112	-0.33	NS	-0.51	NS
113	0.24	*	0.39	**	113	-0.33	NS	-0.50	NS
114	0.22	NS	0.38	**	114	-0.33	NS	-0.50	NS
115	0.21	NS	0.37	**	115	-0.33	NS	-0.50	NS
116	0.22	NS	0.36	**	116	-0.34	NS	-0.51	NS
117	0.21	NS	0.35	**	117	-0.34	NS	-0.51	NS
118	0.22	NS	0.35	**	118	-0.34	NS	-0.51	NS
119	0.23	*	0.35	**	119	-0.32	NS	-0.50	NS
120	0.24	*	0.35	**	120	-0.29	NS	-0.49	NS
121	0.22	NS	0.34	**	121	-0.28	NS	-0.48	NS
122	0.24	*	0.34	*	122	-0.27	NS	-0.48	NS
123	0.24	*	0.34	*	123	-0.25	NS	-0.47	NS
124	0.26	*	0.34	*	124	-0.24	NS	-0.46	NS
125	0.25	*	0.34	*	125	-0.22	NS	-0.45	NS
126	0.23	NS	0.33	*	126	-0.21	NS	-0.43	NS
127	0.25	*	0.35	*	127	-0.20	NS	-0.43	NS
128	0.24	*	0.35	*	128	-0.21	NS	-0.42	NS
129	0.25	*	0.36	**	129	-0.21	NS	-0.41	NS
130	0.25	*	0.36	**	130	-0.21	NS	-0.40	NS
131	0.25	*	0.36	**	131	-0.20	NS	-0.40	NS
132	0.24	*	0.36	**	132	-0.19	NS	-0.39	NS
133	0.26	*	0.38	**	133	-0.17	NS	-0.38	NS
134	0.25	*	0.38	**	134	-0.15	NS	-0.36	NS
135	0.27	*	0.39	**	135	-0.12	NS	-0.35	NS
136	0.27	*	0.39	**	136	-0.10	NS	-0.33	NS
137	0.25	*	0.38	**	137	-0.08	NS	-0.32	NS
138	0.26	*	0.37	*	138	-0.06	NS	-0.30	NS
139	0.26	*	0.37	*	139	-0.06	NS	-0.29	NS
140	0.26	*	0.37	*	140	-0.06	NS	-0.28	NS
141	0.25	*	0.36	*	141	-0.06	NS	-0.27	NS
142	0.24	NS	0.35	*	142	-0.04	NS	-0.25	NS
143	0.22	NS	0.35	*	143	-0.01	NS	-0.23	NS
144	0.20	NS	0.33	*	144	0.02	NS	-0.21	NS
145	0.15	NS	0.31	*	145	0.04	NS	-0.20	NS