Whole-genome sequencing of five proteobacteria isolated from Lechuguilla Cave, New Mexico, USA:
Insights into taxonomy and quorum-sensing

Han Ming Gan[1,2*], Peter C. Wengert[4], Hazel A. Barton[3], André O. Hudson[4], Michael A. Savka[4]

[1] Centre for Integrative Ecology, School of Life and Environmental Sciences, Deakin University, Geelong,
Victoria, Australia

[2] Deakin Genomics Centre, Deakin University, Geelong, Victoria, Australia

[3] Department of Biology, University of Akron, Akron, Ohio, USA

[4] The Thomas H. Gosnell School of Life Sciences, Rochester Institute of Technology, Rochester, New York,
USA

*Corresponding author

Name: Han Ming Gan

Email: han.gan@deakin.edu.au

Abstract. Genomic resources remain scarce for bacteria isolated from oligotrophic caves. We
sequenced the genomes of five proteobacteria isolated from Lechuguilla Cave in New Mexico, USA.
Genome-based phylogeny indicates that each strain belongs to a distinct genus. Two *Rhizobiaceae*
isolates possess the genomic potential for the biosynthesis of acyl-homoserine lactone.

Main Text:

Adequate genomic resources are crucial for the understanding of adaptation for culturable
microbes in nutrient-limited cave environments (1). To date, bacterial genomes isolated from caves are
poorly represented in the public databases (2-5). Here, we sequenced the genomes of five non-clonal
Lechuguilla cave isolates and used genome-based phylogeny to refine their taxonomic assignment that
was previously based on the variable regions 3 and 4 of the 16S rRNA gene. We also identified genomic
potential for the biosynthesis of cell-cell communication signal in two *Rhizobiaeae* isolates and
confirmed the predicted phenotype using an *Agrobacterium tumefaciens* reporter assay (6).

Bacteria were grown in half strength tryptic soy broth with shaking at $30^{o}C$ for 2 days. DNA
extraction was performed using the GenElute bacterial genomic DNA kit (MilliporeSigma, St Louis, MO).
The gDNA was processed with the Illumina Nextera XT DNA sample prep kit (Illumina, San Diego, CA)
according to the manufacturer's instructions and sequenced on an Illumina MiSeq (run configuration of
2×150 bp). The paired-end reads were adapter-trimmed with Trimmomatic v0.36 (7) and assembled

using Unicycler v0.4.7 (8). The assembled genomes range from 4.5 to 6.5 Mb with N_{50} length of more

than 100 kb (Table 1).

The genome of strain LC387 was assembled into a single contig (Table 1). Its full-length 16S rRNA

sequence is 100% identical to that of *Afipia massiliensis* CCUG 45153 [T] (RefSeq Accession: NR_025646.1),

making LC387 the first member of *A. massiliensis* with genome availability. Using phylogenomic

approach, GToTree v1.2.1 (9), confident taxonomic assignment to the genus level was obtained for three

of the four remaining strains (Figure 1A). The basal placement of LC148 in the *Neorhizobium* clade

suggests that it is a divergent member within the genus or a member of an undescribed genus closely

related to *Neorhizobium*. Therefore, strain LC148 was classified as *Rhizobacteriaceae* sp. pending future

taxonomic investigation.

We used a previously described Hidden Markov Model (HMM) approach (10) to identify the

genomic potential for biosynthesis of acyl-homoserine lactone (AHL) molecules involved in the

regulation of gene expression in response to cell density (11). AHL synthase homologs (LuxI) were

identified in strains LC34 (GenBank Protein ID: TKT57503.1) and LC148 (GenBank Protein IDs:

TKT46115.1 and TKT66962.1). The genes coding for these proteins were also classified as *luxI* homologs

with AntiSMASH 4 (12). To confirm the predicted phenotype, LC34 and LC148 strains were grown in YM

media and ethyl acetate extracts from the cultures were tested for the presence of AHL signals using the

AHL-dependent biosensor *Agrobacterium tumefaciens* NTL4(pZLR4) as previously described(13). Under

the described culturing and detection setting, LC34 produced one type of AHL signal with retardation

factor (RF) that is to similar 3-oxo-C8. LC148 produced two distinct AHLs, one with RF similar to C6 and

the other with RF smaller than the included AHL standards (Figure 1B). Future work investigating the

role of quorum sensing in these cave isolates through transposon mutagenesis (14), targeted gene

deletion (15) or transcriptome sequencing (16) will be informative in understanding bacteria from cave

environments.

Data availability. The raw Illumina paired-end reads and genome assemblies have been deposited in

GenBank under the BioProject numbers as listed in Table 1. Bacterial strains can be requested from

Professor Michael A. Savka, Rochester Institute of Technology, New York, USA.

Acknowledgements. The authors acknowledge the Thomas H. Gosnell School of Life Sciences (GSoLS)

and the College of Science (COS) at the Rochester Institute of Technology (RIT) for ongoing support.

PCW was supported by a 2019 COS Summer Undergraduate Research Fellowship from RIT.

Competing Interests. The authors have declared that no competing interests exist.

68   Table 1. Genome assembly metrics and data availability

BioProject	Assembly Accession	Organism	Strain	Genome Size (bp)	GC content (%)	No. of contigs (>500 bp)	N_{50} length (bp)
PRJNA281560	LBCQ02	*Aquamicrobium* sp.	LC103	6,044,022	64.06	34	330,476
PRJNA532460	SZVV01	*Rhizobiaceae* sp.	LC148	5,822,494	61.45	86	124,816
PRJNA281572	LBHX02	*Agrobacterium* sp.	LC34	5,303,297	59.35	36	377,718
PRJNA281679	LBIA02	*Afipia massiliensis*	LC387	4,495,198	61.35	1	4,495,198
PRJNA281681	LBHW02	*Achromobacter* sp.	LC458	6,547,009	64.55	102	143,492

Figure Legends

Figure 1. (A) Maximum likelihood tree constructed based on the concatenated alignment of 119

conserved proteobacteria single-copy gene sets. Branch lengths indicate the number of substitutions per

site while node labels are Shimodaira-Hasegawa (SH) local support values computed in FastTree2 (17).

(B) Separation of AHL molecules from the LC34 and LC148 extracts by C_{18} reversed-phase thin-layer plate

developed with methanol/water (60:40, vol/vol). Standards include: C6, N-hexanoyl-L-homoserine

lactone; C6-O, N-(3-oxohexanoyl)-L-homoserine lactone; C8-O, N-(3-oxo-octanoyl)-L-homoserine

lactone.

References

1.   Barton MD, Petronio M, Giarrizzo JG, Bowling BV, Barton HA. 2013. The genome of

Pseudomonas fluorescens strain R124 demonstrates phenotypic adaptation to the mineral

environment. J Bacteriol 195:4793-803.

2.   Gosse JT, Ghosh S, Sproule A, Overy D, Cheeptham N, Boddy CN. 2019. Whole Genome

Sequencing and Metabolomic Study of Cave Streptomyces Isolates ICC1 and ICC4. Frontiers in

Microbiology 10.

3.   Pawlowski AC, Westman EL, Koteva K, Waglechner N, Wright GD. 2018. The complex resistomes

of Paenibacillaceae reflect diverse antibiotic chemical ecologies. The ISME Journal 12:885-897.

4.   Pawlowski AC, Wang W, Koteva K, Barton HA, McArthur AG, Wright GD. 2016. A diverse intrinsic

antibiotic resistome from a cave bacterium. Nature Communications 7:13803.

5.   Gan HY, Gan HM, Tarasco AM, Busairi NI, Barton HA, Hudson AO, Savka MA. 2014. Whole-

genome sequences of five oligotrophic bacteria isolated from deep within Lechuguilla Cave, New

Mexico. Genome Announc 2:e01133-14.

6.   Scott RA, Weil J, Le PT, Williams P, Fray RG, von Bodman SB, Savka MA. 2006. Long- and Short-

Chain Plant-Produced Bacterial N-Acyl-Homoserine Lactones Become Components of

Phyllosphere, Rhizosphere, and Soil. Molecular Plant-Microbe Interactions 19:227-239.

7.   Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence

data. Bioinformatics 30:2114-2120.

8.   Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies

from short and long sequencing reads. PLoS computational biology 13:e1005595.

9.   Lee MD. 2019. GToTree: a user-friendly workflow for phylogenomics. Bioinformatics
doi:10.1093/bioinformatics/btz188.

10.   Gan HM, Gan HY, Ahmad NH, Aziz NA, Hudson AO, Savka MA. 2015. Whole genome sequencing
and analysis reveal insights into the genetic structure, diversity and evolutionary relatedness of
luxI and luxR homologs in bacteria belonging to the Sphingomonadaceae family. Frontiers in
Cellular and Infection Microbiology 4.

11.   Miller MB, Bassler BL. 2001. Quorum sensing in bacteria. Annual Reviews in Microbiology
55:165-199.

12.   Blin K, Wolf T, Chevrette MG, Lu X, Schwalen CJ, Kautsar SA, Suarez Duran HG, de Los Santos EL,
Kim HU, Nave M. 2017. antiSMASH 4.0—improvements in chemistry prediction and gene cluster
boundary identification. Nucleic acids research 45:W36-W41.

13.   Lowe N, Gan HM, Chakravartty V, Scott R, Szegedi E, Burr TJ, Savka MA. 2009. Quorum-sensing
signal production by *Agrobacterium vitis* strains and their tumor-inducing and tartrate-catabolic
plasmids. FEMS microbiology letters 296:102-109.

14.   Gan HM, Ibrahim Z, Shahir S, Yahya A. 2011. Identification of genes involved in the 4-
aminobenzenesulfonate degradation pathway of *Hydrogenophaga* sp. PBC via transposon
mutagenesis. FEMS microbiology letters 318:108-114.

15.   Marx CJ. 2008. Development of a broad-host-range *sacB*-based vector for unmarked allelic
exchange. BMC research notes 1:1-1.

16.   Wagner VE, Gillis RJ, Iglewski BH. 2004. Transcriptome analysis of quorum-sensing regulation
and virulence factor expression in *Pseudomonas aeruginosa*. Vaccine 22:S15-S20.

17.   Price MN, Dehal PS, Arkin AP. 2010. FastTree 2–approximately maximum-likelihood trees for
large alignments. PloS one 5:e9490.