Draft genome sequences of five Proteobacteria isolated from Lechuguilla Cave, New Mexico, USA:
Insights into taxonomy and quorum-sensing

Han Ming Gan[1,2*], Peter C. Wengert[4], Hazel A. Barton[3], André O. Hudson[4], Michael A. Savka[4]

[1] Centre for Integrative Ecology, School of Life and Environmental Sciences, Deakin University, Geelong,
Victoria, Australia

[2] Deakin Genomics Centre, Deakin University, Geelong, Victoria, Australia

[3] Department of Biology, University of Akron, Akron, Ohio, USA

[4] The Thomas H. Gosnell School of Life Sciences, Rochester Institute of Technology, Rochester, New York,
USA

*Corresponding author

Name: Han Ming Gan

Email: han.gan@deakin.edu.au

Abstract. Genomic resources remain scarce for bacteria isolated from oligotrophic caves. We
sequenced the genomes of five Proteobacteria isolated from Lechuguilla Cave in New Mexico, USA.
Genome-based phylogeny indicates that each strain belongs to a distinct genus. Two *Rhizobiaceae*
isolates possess the genomic potential for the biosynthesis of acyl-homoserine lactone.

Main Text:

Adequate genomic resources are crucial for the understanding of adaptation for culturable
microbes in nutrient-limited cave environments (1). To date, bacterial genomes isolated from caves are
poorly represented in the public databases (2-5). Here, we report the genomes of five Lechuguilla cave
isolates and used genome-based phylogeny to refine their taxonomic assignment. We also identified
genomic potential for the biosynthesis of cell-cell communication signal in two *Rhizobiaeae* isolates and
confirmed the predicted phenotype using an *Agrobacterium tumefaciens* reporter assay (6).

Initial isolation of the bacterial strains from remote sample sites in Lechuguilla Cave was
previously described by Bhullar *et al* (7). Strains were grown in half strength tryptic soy broth with
shaking at 30°C for 2 days. DNA extraction used the GenElute bacterial genomic DNA kit (MilliporeSigma,
St Louis, MO). Sequencing library was generated using the tagmentation-based Illumina Nextera XT DNA
sample prep kit (Illumina, San Diego, CA) and sequenced on an Illumina MiSeq (run configuration of 2 ×

150 bp). The paired-end reads were adapter-trimmed and assembled with Trimmomatic v0.36 (8) and
Unicycler v0.4.7 (9), respectively, using the default setting.

The genome of LC387 was assembled into a single contig (Table 1). Based on BLASTN alignment
(10), its full-length 16S rRNA sequence is 100% identical to that of *Afipia massiliensis* CCUG 45153 [T]
(RefSeq Accession: NR_025646.1). For the generation of genome-based phylogeny using GToTree v1.2.1
(11), genome assemblies of bacterial type strains exhibiting high 16S rRNA gene sequence similarity to
the cave isolates were downloaded and included in the pipeline. Based on phylogenomic clustering,
confident taxonomic assignment to the genus level was obtained for LC34, LC103 and LC458 (Figure 1A).
The basal placement of LC148 in the *Neorhizobium* clade suggests that it is a divergent member within
the genus or a member of an undescribed genus. Therefore, strain LC148 was classified as
Rhizobacteriaceae sp. pending future taxonomic investigation.

We used a previously described Hidden Markov Model (HMM) approach (12) to identify the
genomic potential for biosynthesis of acyl-homoserine lactone (AHL) molecules involved in the
regulation of gene expression in response to cell density (13). AHL synthase homologs (LuxI) were
identified in strains LC34 (GenBank Protein ID: TKT57503.1) and LC148 (GenBank Protein IDs:
TKT46115.1 and TKT66962.1). The genes coding for these proteins were similarly classified as *luxI*
homologs by AntiSMASH 4 and the NCBI prokaryotic genome annotation pipeline (14, 15). To confirm
the predicted phenotype, LC34 and LC148 were grown in YM media and ethyl acetate extracts from the
cultures were tested for the presence of AHL signals using the AHL-dependent biosensor *Agrobacterium*
tumefaciens NTL4(pZLR4) (16). LC34 produced one type of AHL signal with retardation factor (RF) that is
similar to 3-oxo-C8. LC148 produced two distinct AHLs, one with RF similar to C6 and the other with RF
smaller than the included AHL standards (Figure 1B). Future work investigating the role of quorum
sensing in these cave isolates through transposon mutagenesis (17), targeted gene deletion (18) or
transcriptome sequencing (19) will be informative in understanding bacteria from cave environments.

Data availability. The raw Illumina paired-end reads and genome assemblies have been deposited in
GenBank under the BioProject numbers as listed in Table 1. Bacterial strains can be requested from
Professor Michael A. Savka, Rochester Institute of Technology, New York, USA.

Acknowledgements. The authors acknowledge the Thomas H. Gosnell School of Life Sciences (GSoLS)
and the College of Science (COS) at the Rochester Institute of Technology (RIT) for ongoing support.
PCW was supported by a 2019 COS Summer Undergraduate Research Fellowship from RIT.

Competing Interests. The authors have declared that no competing interests exist.

67   Table 1. Genome assembly metrics and data availability

BioProject	Assembly Accession	Organism	Strain	Number of reads	Number of bases (Mb)	Estimated genome coverage (×)	Genome Size (bp)	GC content (%)	No. of contigs (>500 bp)	N_{50} length (bp)
PRJNA281560	LBCQ02	*Aquamicrobium* sp.	LC103	1,246,649	417	69.5	6,044,022	64.06	34	330,476
PRJNA532460	SZVV01	*Rhizobiaceae* sp.	LC148	399,832	110	18.8	5,822,494	61.45	86	124,816
PRJNA281572	LBHX02	*Agrobacterium* sp.	LC34	843,163	247	46.6	5,303,297	59.35	36	377,718
PRJNA281679	LBIA02	*Afipia massiliensis*	LC387	777,637	216	48.0	4,495,198	61.35	1	4,495,198
PRJNA281681	LBHW02	*Achromobacter* sp.	LC458	1,286,382	448	68.4	6,547,009	64.55	102	143,492

Figure Legends

Figure 1. (A) Maximum likelihood tree based on the concatenated alignment of 119 conserved

proteobacteria single-copy gene sets generated using the default setting of GToTree v1.2.1. Branch

lengths indicate the number of substitutions per site while node labels are Shimodaira-Hasegawa (SH)

local support values computed in FastTree2 (20). (B) Separation of AHL molecules from the LC34 and

LC148 extracts by C_{18} reversed-phase thin-layer plate developed with methanol/water (60:40, vol/vol).

Standards include: C6, N-hexanoyl-L-homoserine lactone; C6-O, N-(3-oxohexanoyl)-L-homoserine

lactone; C8-O, N-(3-oxo-octanoyl)-L-homoserine lactone.

References

1.   Barton MD, Petronio M, Giarrizzo JG, Bowling BV, Barton HA. 2013. The genome of
Pseudomonas fluorescens strain R124 demonstrates phenotypic adaptation to the mineral
environment. J Bacteriol 195:4793-803.
2.   Gosse JT, Ghosh S, Sproule A, Overy D, Cheeptham N, Boddy CN. 2019. Whole Genome
Sequencing and Metabolomic Study of Cave Streptomyces Isolates ICC1 and ICC4. Frontiers in
Microbiology 10.
3.   Pawlowski AC, Westman EL, Koteva K, Waglechner N, Wright GD. 2018. The complex resistomes
of Paenibacillaceae reflect diverse antibiotic chemical ecologies. The ISME Journal 12:885-897.
4.   Pawlowski AC, Wang W, Koteva K, Barton HA, McArthur AG, Wright GD. 2016. A diverse intrinsic
antibiotic resistome from a cave bacterium. Nature Communications 7:13803.
5.   Gan HY, Gan HM, Tarasco AM, Busairi NI, Barton HA, Hudson AO, Savka MA. 2014. Whole-
genome sequences of five oligotrophic bacteria isolated from deep within Lechuguilla Cave, New
Mexico. Genome Announc 2:e01133-14.
6.   Scott RA, Weil J, Le PT, Williams P, Fray RG, von Bodman SB, Savka MA. 2006. Long- and Short-
Chain Plant-Produced Bacterial N-Acyl-Homoserine Lactones Become Components of
Phyllosphere, Rhizosphere, and Soil. Molecular Plant-Microbe Interactions 19:227-239.
7.   Bhullar K, Waglechner N, Pawlowski A, Koteva K, Banks ED, Johnston MD, Barton HA, Wright GD.
2012. Antibiotic resistance is prevalent in an isolated cave microbiome. PloS one 7:e34953-
e34953.
8.   Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics 30:2114-2120.
9.   Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies
from short and long sequencing reads. PLoS computational biology 13:e1005595.
10.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. J
Mol Biol 215:403-10.
11.   Lee MD. 2019. GToTree: a user-friendly workflow for phylogenomics. Bioinformatics
doi:10.1093/bioinformatics/btz188.
12.   Gan HM, Gan HY, Ahmad NH, Aziz NA, Hudson AO, Savka MA. 2015. Whole genome sequencing
and analysis reveal insights into the genetic structure, diversity and evolutionary relatedness of
luxI and luxR homologs in bacteria belonging to the Sphingomonadaceae family. Frontiers in
Cellular and Infection Microbiology 4.
13.   Miller MB, Bassler BL. 2001. Quorum sensing in bacteria. Annual Reviews in Microbiology
55:165-199.

14.   Blin K, Wolf T, Chevrette MG, Lu X, Schwalen CJ, Kautsar SA, Suarez Duran HG, de Los Santos EL,
Kim HU, Nave M. 2017. antiSMASH 4.0—improvements in chemistry prediction and gene cluster
boundary identification. Nucleic acids research 45:W36-W41.
15.   Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt
KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic genome annotation pipeline. Nucleic Acids
Res 44:6614-24.
16.   Lowe N, Gan HM, Chakravartty V, Scott R, Szegedi E, Burr TJ, Savka MA. 2009. Quorum-sensing
signal production by *Agrobacterium vitis* strains and their tumor-inducing and tartrate-catabolic
plasmids. FEMS microbiology letters 296:102-109.
17.   Gan HM, Ibrahim Z, Shahir S, Yahya A. 2011. Identification of genes involved in the 4-
aminobenzenesulfonate degradation pathway of *Hydrogenophaga* sp. PBC via transposon
mutagenesis. FEMS microbiology letters 318:108-114.
18.   Marx CJ. 2008. Development of a broad-host-range *sacB*-based vector for unmarked allelic
exchange. BMC research notes 1:1-1.
19.   Wagner VE, Gillis RJ, Iglewski BH. 2004. Transcriptome analysis of quorum-sensing regulation
and virulence factor expression in *Pseudomonas aeruginosa*. Vaccine 22:S15-S20.
20.   Price MN, Dehal PS, Arkin AP. 2010. FastTree 2—approximately maximum-likelihood trees for
large alignments. PloS one 5:e9490.