RNA-seq analysis of a zebrafish *tlr2* mutant shows a broad function of this Toll-like

receptor in transcriptional and metabolic control and defense to *Mycobacterium*

***marinum* infection**

Shuxin Yang[1,6*] Wanbin Hu[1*], Yasuhito Shimada[1,2], Magnus Münch [3,4], Rubén Marín-Juez[1,5],

Annemarie H. Meijer[1], Herman P. Spaink[1**]

[1]Institute of Biology, Leiden University, P.O. Box 9502, 2300 RA Leiden, the Netherlands

[2]Department of Integrative Pharmacology, Mie University Graduate School of Medicine,

Japan.

[3] Mathematical Institute, Leiden University, Leiden

[4] Department of Epidemiology & Biostatistics, Amsterdam Public Health research institute,

Amsterdam University medical center, Amsterdam

[5] Department of Developmental Genetics, Max Planck Institute for Heart and Lung Research,

Ludwigstrasse 43, 61231 Bad Nauheim, Germany.

[6]Institute of Synthetic Biology, Shenzhen Institutes of Advanced Technology, Chinese

Academy of Sciences, Shenzhen 518055, China.

*These authors contributed equally

**Corresponding author email: h.p.spaink@biology.leidenuniv.nl

Abstract

Background

The function of Toll-like receptor 2 (TLR2) in host defense against pathogens, especially

Mycobacterium tuberculosis (Mtb) is poorly understood. To investigate the role of TLR2

during mycobacterial infection, we analyzed the response of *tlr2* zebrafish mutant larvae to

infection with *Mycobacterium marinum* (Mm), a close relative to Mtb, as a model for

tuberculosis. We measured infection burdens and transcriptome responses using RNA deep

sequencing in mutant and control larvae.

Results

tlr2 mutant embryos at 2 dpf do not show morphological alterations or differences in the

number of macrophages and neutrophils when compared to control embryos. However, we

found substantial changes in gene expression in these mutants, particularly in developmental

and metabolic pathways, when compared with the heterozygote $tlr2^{+/-}$ control. After Mm

infection, bacterial burden was six to ten fold higher in $tlr2^{-/-}$ larvae than in $tlr2^{+/-}$, or $tlr2^{+/+}$

larvae, indicating that Tlr2 acts as a protective factor in zebrafish host defense. RNAseq

analysis of infected $tlr2^{-/-}$ versus $tlr2^{+/-}$ shows that the number of up-regulated and down-

regulated genes in response to infection was greatly diminished in *tlr2* mutants by at least 2

fold and 10 fold, respectively. Analysis of the transcriptome data and qPCR validation shows

that Mm infection of *tlr2* mutants leads to decreased mRNA levels of genes involved in

inflammation and immune responses, including *il1b*, *tnfb*, *cxcl11aa/ac*, *fosl1a*, and *cebpb*.

Furthermore, RNAseq analyses revealed that the expression of genes for Maf family

transcription factors, vitamin D receptors, and Dicps proteins is significantly altered in *tlr2*

mutants with or without infection. In addition, the data indicate a function of Tlr2 in the

control of induction of cytokines and chemokines, such as the CXCR3-CXCL11 signaling

axis.

Conclusion

The transcriptome and infection burden analyses give support for a function of TLR2 in host

defense against mycobacteria. Transcriptome analysis revealed *tlr2*-specific pathways

involved in Mm infection, which are related to responses to Mtb infection in human

macrophages. Considering its dominant function in control of transcriptional processes that

govern defense responses and metabolism, the TLR2 protein can be expected to be also of

importance for other infectious diseases and interactions with the microbiome.

Background

Mycobacterium tuberculosis (Mtb) is the causative agent of tuberculosis (TB), which infects

nearly 23% of the world's population, and kills about 1.6 million people annually (WHO

Global Tuberculosis Report 2018). TB is characterized by the formation of granulomas,

aggregates of infected macrophages and other immune cells, not only in the lung but also in

other tissues and organs[1]. The formation of granulomas is the result of a concerted action of

host innate and adaptive immunity [2-4].

Innate immune responses play a critical role in defense against TB infection in the host, and

for a major part these processes are mediated by Toll-like receptors (TLRs), a conserved

family of pattern recognition receptors. TLR2 is one of the most widely reported members of

the TLR family to be involved in defense against Mtb by virtue of its recognition of cell wall-

associated components associated with this pathogen [5-7]. Following mycobacterial infection

in human cell cultures, TLR2 dimerizes with TLR1 or TLR6, and recognizes mycobacterial

components such as cell wall glycolipids LAM and LM [8], 38-kDa and 19-kDa

mycobacterial glycoprotein (LpqH) [9-11], phosphatidylinositol mannoside (PIM) [12], and

triacylated (TLR2/TLR1) [13, 14] or diacylated (TLR2/TLR6) lipoproteins [15, 16]. Then,

these heterodimers recruit the MYD88 and TIRAP (MAL) proteins to activate the IRAK (1

and 4)/TRAF6/IKK (α or β) cascade, which subsequently leads to the ubiquitination of IκBα

and the activation of transcription factor NF-κB or AP-1 to induce the expression of host

defense genes and cytokine and chemokine responses [17, 18]. Once released after further

processing, these cytokines and chemokines attract migration of macrophages and neutrophils

to the infection site and activate the microbicidal functions of these cells.

TLR2 has been shown to be important for granuloma formation in Mtb infection in mouse

infection model [19]. Other studies have reported that *Tlr2*[-/-] mice lose control to high dose

infection of Mtb delivered by aerosol administration and show higher susceptibility to Mtb

infection compared to the wild type [20, 21]. However, using lower doses of infection of 100

bacteria, there are controversial results as to the function of TLR2 in defense to Mtb in rodent

models [20-22]. Furthermore, the function of TLR2 in susceptibility to Mtb is still unclear

because several independent studies have reported polymorphisms of TLR2 (TLR1 and TLR6)

in humans that have been linked with TB susceptibility, whereas others have been unable to

find such links [23, 24].

Several studies suggest that the interaction between TLR2 and Mtb or other pathogens does

not always promote the killing of bacteria, but can in fact be part of the pathogens's strategy

to evade the immune system [25-27]. TLR2 has been reported to inhibit MHC-II expression

on the surface of murine macrophages, thereby preventing presentation of Mtb antigens,

which may allow intracellular Mtb to evade immune surveillance and maintain chronic

infection [11, 28]. Furthermore, LprG from Mtb was reported to inhibit human macrophage

class II MHC antigen processing through TLR signaling [25]. In other infection systems, *Tlr2*

mutant mice have an increased resistance against infection of *Candida albicans* [29] and

Yersinia pestis [30]. It has been proposed that this phenotype is the result of Tlr2-dependent

induction of the anti-inflammatory cytokine IL-10 [30]. Furthermore, TLR2 activation

inhibits the release of IL-12 via activation of the cFos transcription factor. It also has been

shown that IFN-γ or IFN-γ-induced signals [31] are inhibited after infection of murine

macrophages in a Tlr2-dependent fashion [32]. These effects show that TLR2 activation can

yield a bias to T helper Type 2 (Th2) cells [33], and by breaking the Th1/Th2 balance can

lead to less Th1 type responses and reducing the killing of intracellular pathogens. However,

most of the molecular mechanisms underlying TLR2 functions remain unknown and a better

understanding of the TLR2-mediated immune response and immune evasion can help in

planning prevention and therapy strategies against Mtb infection.

Animal models have shown their power in studies of the mechanisms of interaction of host

and TB pathogens, and discovering new anti-TB drugs. Zebrafish adult and larvae models

have become a useful complement for rodent studies, for three important reasons. First,

zebrafish have a 3-4 weeks separation stage between development of innate and adaptive

immunity after fertilization [34, 35], which gives the possibility to study the host innate

immune response to infection in the absence of adaptive immune responses. Second, zebrafish

can be infected by *Mycobacterium marinum* (Mm), a natural pathogen of cold-blooded

vertebrates and a close relative of Mtb, which can induce granuloma formation in zebrafish,

similar to human TB [36, 37]. Third, the transparent larvae are ideal for imaging the early

steps of the infection process in real time. Hence, zebrafish has earned its place of being a

versatile tuberculosis model [1, 38, 39].

In our previous study, we demonstrated that the mammalian TLR2 ligand Pam3CSK4, a

synthetic triacylated lipopeptide that mimics the triacylated lipoprotein of mycobacteria,

could also specifically activate the zebrafish Tlr2 pathway, inducing *fosl1a* and *cebpb* gene

upregulation [40]. In the current study, to further explore the involvement of Tlr2 in Mm

infection, we conducted infection studies using *tlr2* mutant zebrafish. We found that a *tlr2*

mutation led to an overall increased Mm bacterial burden, corresponding with reduced

presence of macrophages in the granulomatous aggregates and more extracellular growth.

These results indicate that Tlr2 plays an important role in protecting the host during the early

stage of mycobacterial infection. In addition, we performed RNA deep sequencing (RNAseq)

and determined a Tlr2-specific gene list for the response to Mm infection. This analysis

revealed that most of transcriptional downregulation caused by Mm infection in control

animals was abrogated by *tlr2* mutation, in addition to a dampening effect on the upregulation

of transcription factors and inflammatory genes.

Results

1. Characterisation of a *tlr2* mutant zebrafish line

The *tlr2*[sa19423] mutant (*tlr2*[-/-]) carries a thymine to adenine point mutation that creates a

premature stop codon (Fig. 1a), which is located in the sequence coding for the C-terminus of

the leucine-rich repeat (LRR) domain. This leads to a truncated protein without the Toll/IL-1

receptor (TIR) domain, which is required for the interaction with Myd88 and Tirap (Mal) [41,

42].

To confirm whether *tlr2* mutation blocks its downstream pathway, we analyzed the gene

expression profiles of zebrafish treated with the TLR2 agonist, Pam3CSK4 similarly as in our

previous work [43], now also including the heterozygote mutant (*tlr2*[+/-]). Pam3CSK4 was

injected into the blood island of zebrafish embryos at 27 hours post fertilization (hpf). One

hour after injection (hpi), we collected samples and performed qPCR to analyze the

expression levels of CCAAT/enhancer-binding protein beta (*cebpb*) and FOS Like Antigen 1a

(*fosl1a*), previously shown to be specific targets of Tlr2 signaling [40]. In wild-type siblings

and *tlr2*[+/ sa19423] heterozygotes (*tlr2*[+/-]), the expression levels of *cebpb* and *fosl1a*, as well as

the inflammatory gene *il1b* were significantly induced upon Pam3CSK4 injection, whereas

tlr2[-/-] showed no significant response (Fig. 1b-d). *tlr2*[+/-] animals showed a lower induction of

these marker genes than the wild-types, indicating that there is an effect of the *tlr2* mutation

even in the heterozygote. To confirm that these results are specific to the Tlr2 pathway, we

injected flagellin, a Tlr5 agonist, into 27 hpf embryos. Flagellin induced *il1b* expression, but

not *cebpb* and *fosl1a* expression in wild-type siblings, *tlr2*[+/-] and *tlr2*[-/-] larvae (Fig. 1e-g).

Overall, these data show that *tlr2* mutation specifically blocks the Tlr2 downstream pathways.

To determine if immune cell development was affected by *tlr2* mutation, we used the double-

transgenic   line   *tlr2*[+/+]   *Tg(mpeg1:mCherry-F);TgBAC(mpx:EGFP)*   and   *tlr2*[-/-]

Tg(mpeg1:mCherry-F);TgBAC(mpx:EGFP) to count the number of macrophages and

neutrophils at 2 dpf. The results show that there is no significant difference in the number of

macrophages and neutrophils at 2 dpf between the wild type and mutant (Fig. 1j, k).

2 Comparison of gene expression profiles of *tlr2* homozygote and heterozygote mutants

in the absence of infection

In order to investigate the systemic effects of the *tlr2* mutation we compared basal levels of

gene expression in the absence of infection between *tlr2* homozygote and heterozygote

mutants (Fig. 2). We chose to use the heterozygote *tlr2*$^{+/-}$ line as a control since these are

genetically as comparable as possible, considering the fact that random ENU mutations could

still contaminate our background even after 3 generations of outcrossing and the use of sibling

lines. To further minimize the effect of polymorphisms, we pooled 10 larvae in each of our

samples. These results show that there is a large group of genes that are expressed differently

even at extremely stringent false discovery rate (FDR) adjusted *p*-value of 10^{-10} (Fig. 2). Since

the results of the DEseq2 analyses showed such a surprisingly large number of significant

differences we also used another statistical method for analyses, called edgeR, that differs in

normalization and estimation of the dispersion parameters [44]. EdgeR analyses confirmed

the statistical significance of the differences between the homozygote and heterozygote

mutants (Fig. 2a and 2b). While no differences in numbers of mpeg-positive macrophages

were detected (Fig. 1j, k), the RNAseq analysis showed that the *mpeg1* gene is expressed

approximately 2 fold higher in the *tlr2*$^{+/-}$ control than in the *tlr2*$^{-/-}$ mutant (Supplementary

Table 1). This was confirmed by pixel count measurements in the transgenic line *tlr2*$^{+/+}$

Tg(mpeg1:mCherry-F);TgBAC(mpx:EGFP) and *tlr2*$^{-/-}$ *Tg(mpeg1:mCherry-*

F);TgBAC(mpx:EGFP) (Fig. S1 a, b). These results showed lower fluorescence of the

mCherry *mpeg1* reporter as compared to the wild-type siblings at 2 dpf (Fig. S1 a) whereas no

difference was detected for the eGFP *mpx* reporter (Fig. S1 b). Therefore these *in vivo* results

confirm the RNAseq data. We also examined the expression levels of the *tlr2* gene in the

homozygote and heterozygote mutants in the RNAseq data (Fig. S2). The results show that,

although *tlr2* is very lowly expressed, there is no difference in expression levels between the

homozygote and heterozygote mutants, indicating that there is no non-sense mediated mRNA

decay (NMD).

To investigate whether the basal expression level differences of genes might be relevant to the

Tlr2 pathway, we performed GO analysis of a set of genes that were expressed differently

with the very low FDR adjust *P* value of 10^{-10} (Supplementary Table 2). Gene ontology

analyses of the gene sets that differ with an FDR of 10^{-10} in both analyses show that there is a

large enrichment of genes belonging to the GO terms related to neural development

(Supplementary Table 2). In addition, we have also analyzed differences with a fold change

criterion of two and FDR of 0.05. These analyses indicated that under the GO category

"Transcription factor genes" only two categories of genes including the c-Maf transcription

factor were present (Supplementary Table 3). Pathway analysis shows that genes that function

in glucose metabolism are differentially expressed in the *tlr2* homozygote versus heterozygote

mutant (Fig. S3).

3 Tlr2-specific gene expression profiles after Mm infection

To study the role of Tlr2 in Mm infection we tested the *tlr2* mutant line in comparison with

the heterozygote and wild-type sibling controls (Fig. 3). No differences in bacterial burden

were observed at 3 dpi (Fig. 3d). However, we found that bacterial burden was significantly

higher in $tlr2^{-/-}$ than in $tlr2^{+/-}$ and wild-type larvae at 4 dpi (Fig. 3e). There was no significant

difference in bacterial infection burden between the heterozygote mutant strain ($tlr2^{+/-}$) and

the wild-type siblings ($tlr2^{+/+}$).

Next, we set out to assess the general inflammation and specific immune responses in *tlr2*

mutants. We first studied by qPCR analysis the expression of *il1b*, *tnfa*, *tnfb*, *irg1l*, and *tlr2*

specific transcription factors *fosl1a* and *cebpb* in $tlr2^{-/-}$ and $tlr2^{+/-}$ larvae upon Mm (strain

Mma20) infection at 4 dpi. Our results show that the induction levels of *il1b*, *tnfb*, *fosl1a* and

cebpb in $tlr2^{-/-}$ larvae were significantly reduced when compared to the heterozygotes in the

infected condition (Fig. 4). $tlr2^{-/-}$ larvae failed to upregulate *fosl1a* and *cebpb* expression in

response to Mm administration (Fig. 4e, f). In our previous work, we showed that Cxcl11-like

chemokines play a crucial role in macrophage-mediated granuloma formation upon Mm

infection [45, 46]. We therefore conducted qPCR to assess the expression levels of *cxcl11-*

like genes, previously shown to be induced by infection [45], including *cxcl11aa* and

cxcl11ac (Fig. 4g, h). The expression levels of *cxcl11aa* and *cxcl11ac* was significantly higher

in $tlr2^{+/-}$ larvae than in $tlr2^{-/-}$ upon Mm infection (Fig. 4g, h). These results indicate that *tlr2*

mutation results in a defective immune or inflammatory response to Mm infection.

To further study the function of *tlr2* in defense against mycobacterial infection, we performed

RNAseq of $tlr2^{+/-}$ and $tlr2^{-/-}$ larvae at 4 dpi with *M. marinum* strain (strain Mma20) and PBS

as control. We summarized the number of differential expressed genes (DEGs) according to

P-value (Fig. 5a, b) and in volcano plots (Fig. S4). The number of DEGs in $tlr2^{+/-}$ infected

with Mm was higher than those in $tlr2^{-/-}$ at any given *P*-value or false discovery rate less than

0.05, or any given fold-change with a *P*-value less than 0.05. These data also show that most

of the genes downregulated by the Mm infection in the control remained unchanged in *tlr2*

mutants (Fig. 5a, b). To further analyze these RNAseq data, we chose the genes with a

threshold of a *P*-value less than 0.05 in $tlr2^{+/-}$ with Mm infection (1102 up- and 827 down-

regulated genes, Fig. 5a). Then, for these genes, we calculated the fold-change ratio of $tlr2^{+/-}$

versus *tlr2^{-/-}*, and genes with ratios greater than 2 or less than 0.5 were selected for further

analysis (Fig. 5c). As a result, 97 and 92 genes were scored as *tlr2* specific up- and down-

regulated genes, respectively. Next, we conducted GO analysis (Fig. 5d, e) showing that

genes grouped into the immune system category are the most prominently deregulated (36%)

in the whole *tlr2* up-regulated 97-gene set (Fig. 5d). Within this category we found genes

involved in lysosome, chemotaxis, transcription regulation, diverse immunoglobulin domain-

containing proteins (*dicps*) and other immune processes (Fig. 5d and 6a-e). For other

categories, many up-regulated genes fell in the categories oxidation-reduction process, DNA

repair, transcription regulation and apoptotic process regulation (Fig. S5). In the *tlr2* down-

regulated 92-gene set, the immune related genes also were the largest portion (15%; Fig. 5e,

6f). Many of these genes are poorly annotated and include genes encoding cysteine proteases,

a *nitr* gene, a *mafb* transcription factor gene and *hsp70*. Categories encompassing non-

immune related genes are listed in Fig. S6. These results show that, upon infection with Mm,

tlr2 mutants show a dampened response of immune genes.

To show the relationship between DEGs, we constructed networks based on common

expression targets in the 97 up- and 92 down-regulated genes [47]. The involved networks

with the up-regulated genes contain *lipc*, *nr0b2*, *hmgcr*, *itln1*, *fgf23*, *vdr*, *irak3* and *tlr8* (Fig.

S7). In *tlr2^{+/-}* controls we observed positively regulated *hmgcr*, *fgf23*, *vdr* and *tlr8*, and

negatively regulated *nr0b2*, whereas *tlr2^{-/-}* showed an opposite regulation for most of these

genes. However, *lipc* and *irak3*, were positively regulated in both *tlr2^{+/-}* and *tlr2^{-/-}*. For the 92

downregulated genes, a network containing *nr0b2* and *mafb* was constructed (Fig. S8). *nr0b2*

and *mafb* were positively regulated in *tlr2^{-/-}* zebrafish and down regulated in *tlr2^{+/-}*.

4 Enrichment analysis of Tlr2 specific genes after Mm infection

To link our data to gene sets defined based on prior biological knowledge (including GO), we

conducted Gene-Set Enrichment Analysis (GSEA) of the differently expressed genes. This

method derives its power by focusing on gene sets, that is, groups of genes that share common

biological function, chromosomal location, or regulation [48]. Because the number of

predicted gene sets were too large for our analysis method (more than 1,000 with $P < 0.05$),

we focused on the gene sets related to metabolic, immunological and inflammation pathways.

GSEA predicted 61 pathways in $tlr2^{+/-}$ and 67 pathways in $tlr2^{-/-}$ zebrafish responsive to Mm

infection (Supplementary Table 4A and 4B). Interestingly, most of these pathways are

common in $tlr2^{+/-}$ and $tlr2^{-/-}$, however some of them are detected as being anti-correlated in

regulation, including some pathways underlying natural killer cell functions and omega-6-

fatty acid metabolism (Supplementary Table 4C). We also performed Sub-Network

Enrichment Analysis (SNEA) [49] to identify possible key genes that are responsible for the

difference in response of the $tlr2^{+/-}$ and $tlr2^{-/-}$ group to Mm infection ($P < 0.05$). SNEA

predicted 565 and 503 pathways for $tlr2^{+/-}$ and $tlr2^{-/-}$ zebrafish that are linked to the response

to infection, respectively (Supplementary Table 4D and 4E), and 264 and 202 of them are

specific for the response in $tlr2^{+/-}$ and $tlr2^{-/-}$ fish, respectively (Supplementary Table 4F).

Since the RNAseq was conducted with the total RNA from whole body of zebrafish, it is not

possible to define which pathways are involved in macrophage functions related to $tlr2$

expression. Therefore, we analyzed previously published DNA microarray data (GDS4781) of

human macrophages transfected with *Mtb* from Gene Expression Omnibus [50]. SNEA

analysis of human macrophages shows that 659 pathways are linked to Mtb infection

(Supplementary Table 4G). By comparing the human SNEA result to $tlr2^{+/-}$ specific pathways

in zebrafish, 56 pathways were defined as $tlr2^{+/-}$-specific (Supplementary Table 4H). Of these,

the pathway of TLR8, which has the lowest *P*-value in both zebrafish and human enrichment

(Supplementary Table 4H), and its network is depicted as example in Figure S9. Overall,

these analyses show that $tlr2^{-/-}$ mutants have a strongly altered immune response of which

many pathways that are linked to human tuberculosis are differently responding.

Discussion

TLR2 has been shown to play a role in host defense against Mtb in several rodent studies but

its role in host innate immunity during infection is still not clear. Moreover, little is known

about the systemic regulation of down-stream signaling of TLR receptors in animal models.

As part of this study we generated a *tlr2* zebrafish mutant to study Tlr2 function in the innate

immune system during mycobacterial infection. Our results show a clear effect of Tlr2 in

defense against mycobacterial infection. The deep sequencing data provided the possibility to

study the whole transcriptome profile in our mycobacterial infection model at the systems

level. There are only a few RNA sequencing results of tuberculosis studies in rodents [51, 52],

human serum [53], human pulmonary epithelial cells [54], and bovine systems.

To characterize the effect of *tlr2* mutation in the absence of infection, we compared the

transcriptome of homozygous mutant larvae with that of heterozygote larvae, thereby

excluding the effect of non-dominant background polymorphisms that might have resulted

from ENU mutagenesis. The results show large differences between these genotypes, for

instance in genes involved in metabolism (Fig. S3). In agreement, our previous results

suggested a role in metabolism also for Myd88, the adaptor in TLR signaling [43]. The largest

category of genes that was significantly affected in *tlr2* homozygous mutants was

"neurological system process" (Supplementary table 2). Many recent studies show that a

mutation in Tlr2 in mice resulted in effects of neuronal development and responses to injury

[55, 56]. Some of these studies show a connection of Tlr2 deficiency to neuronal defects that

could be related to Il10 function and autophagy [57, 58]. When focusing on the signaling

pathways that could be involved, we observed that there was a significant effect in the GO

category of transcription factors, namely the c-Maf factors that totals up to 546

representatives that were affected (Supplementary Table 3). The members of the Maf family

of transcription factors, c-Maf and Mafb are specifically expressed in monocyte and

macrophage lineages [59, 60], and in addition, c-Maf is also expressed in T helper cells [60].

c-Maf was also reported to directly regulate IL-10 expression induced by LPS in macrophages

[61]. Double deficiency of Mafb/c-Maf promotes self-renewal of differentiated macrophages

[62]. We did not detect differences in macrophage numbers between *tlr2* mutants and controls,

but it remains to be further investigated if macrophage differentiation is altered as a result of

Tlr2 deficiency.

Recently, we have shown that Tlr2 and its adaptor MyD88 are important for the response of

the zebrafish host to the microbiome [43]. Therefore, it is very well possible that the

transcriptional differences we find in the *tlr2* mutant versus the controls are due to an aberrant

response to the microbiome. Similarly, we have found that mutation in *myd88* leads to a large

difference in the zebrafish gene expression profile, even in the absence of pathogenic

infection[43]. This difference appeared to be dependent on the presence of a microbiome. In

future work we therefore aim to study the transcriptome response of the *tlr2* mutant under

germfree conditions and in the presence of a microbiome under gnotobiotic conditions and

investigate whether the dysregulation of neural development pathways in *tlr2* mutants might

be linked to control of the gut-brain axis [63-67]. Such studies could also show whether the

composition of the microbiome affects the progression of mycobacterial infection.

The analysis of differential gene expression during infection showed that there is a very

pronounced effect of mutation of the *tlr2* gene. With respect to the number of genes affected,

the strongest effect was observed in genes that are down regulated during infection, since this

category was strongly diminished in the *tlr2* mutant (Fig. 5a, b and Fig. S4). This observation

suggests that Tlr2 has an important function in anti-inflammatory responses, in line with

previous reports of studies in mice that showed a strongly decreased anti-inflammatory

response in *Tlr2* knockouts [68, 69]. In a further analysis of the quantitative effects on the

differences in expression levels of genes in both the up regulated and down regulated groups,

we selected a number of genes that were most significantly affected in GO analysis (Fig. 5d,

e). This GO analysis showed many groups to be affected with the immune response as the

biggest group. We also performed GSEA and SNEA analyses (Fig. S7, S8, S9 and

Supplementary Table 4) showing that the *tlr2* mutant has a very different immune response

than heterozygote controls. Furthermore, we show that many signaling pathways that have

been reported to be linked to tuberculosis infection in humans are differentially regulated in

our data set. Most significantly, activation of the Tlr8 pathway was strongly affected (Fig. S9).

This evidence suggests that the Tlr2 signaling is strongly connected with Tlr8 function. *TLR8*

mutations (polymorphisms) increase susceptibility to mycobacteria in the human population

[70, 71]. In addition, our analysis revealed differential expression of three other interesting

categories of immune genes, discussed below.

The vitamin D receptor pathway genes that are normally up-regulated during infection in

zebrafish larvae were down regulated in the *tlr2* mutant. Furthermore, pathway analysis (Fig.

S7 and S9) also implicated the expression of the Tlr8 pathway connected to vitamin D

signaling as being strongly affected in the *tlr2* mutant. Vitamin D has been shown to be an

important regulatory factor during tuberculosis [72] and has been linked previously to TLR2

function in studies in cell cultures [73]. Therefore, aberrant vitamin D signaling could be a

major contributing factor to the hyper-susceptibility phenotype of *tlr2* mutants in Mm

infection.

Tlr2 has been shown to be essential for the up-regulation of a group of genes that encode the

Diverse Immunoglobulin Domain-Containing Proteins (DICPS) (Fig. 5d). This group is as

novel multigene family encoding diversified immune receptors. Haire *et al*. [74] reported that

recombinant DICP Ig domains bind lipids and lipid extracts of different bacteria, including

Mtb and Mm, a property shared by mammalian CD300 and TREM family members. In the

down-regulated set also several DICP members appear to be dependent on Tlr2, such as

dicp1.17, *dicp3.3*, that are linked to the GO term insulin-like growth factor binding. These

correlations might relate to functions of Tlr2 in other processes such as the control of diabetes

type II by gut microbiota. However, the DICP gene family lacks easily recognizable genetic

homologs in mammals, making a translation to a function in mammalian tuberculosis and

other diseases currently not yet possible [75].

Another highly relevant category of genes of which the induction or repression during

infection is dependent on Tlr2 includes the chemokines. In a previous study of our laboratory,

Torraca *et al*. demonstrated the function of the Cxcr3-Cxcl11 axis in macrophage recruitment

to infection foci and showed that disruption of this axis by *cxcr3.2* mutation increases the

resistance to mycobacterial infection [45]. Furthermore, by Fluorescence-activated cell

sorting (FACS) of Mm-infected cells, we recently showed that *cxcl11a* is a robust marker of

infected macrophages [46]. The infection induced expression of this chemokine is dependent

on Myd88, the common adaptor of the majority of Tlrs, including *tlr2* [46]. In agreement, the

tlr2 mutant shows a significant lower expression of *cxcl11aa* and also of an related

chemokine, *cxcl11ac,* during Mm infection (Fig. 4). Considering the large number of other

chemokines that are controlled by Tlr2 during infection, it is clear that the integrative network

of connections cannot yet be understood from these expression studies and need more detailed

functional analyses, e.g. by combinations of different mutations or directed studies on

responses to chemokines as shown by Torraca *et al* [45]. However, we can state that the

phenotype of the *tlr2* mutant, at the infection level, and the level of transcriptional control

such as the mentioned effects on regulation of MafB/c-Maf and chemokines shows a clear

connection with macrophage chemotaxis. These evidences do not exclude that Tlr2 has many

other functions during infection such as phagocytosis. For instance, Blander *et al*. [76] and

Rahman *et al*. [77] showed that phagocytosis of bacteria and phagosome maturation are

impaired in the absence of TLR signaling. Therefore, the large number of unannotated genes

of which the expression during infection is dependent on Tlr2 is also worth studying in more

detail in future studies.

Conclusion

Our study shows that Tlr2, as a part of innate immunity, plays an important role in controlling

mycobacterial infection as observed on the transcriptome and infection level. This function

may be mediated by several mechanisms, including a general attenuation of the inflammatory

response, reduced mycobacterial dissemination by dampening of CXCR3-CXCL11 signaling,

and anti-mycobacterial effects like vitamin D signaling. Our results show that Tlr2 is a major

Tlr family member upstream of Myd88 that activates the CXCR3-CXCL11 signaling axis.

The *tlr2* mutant is therefore a valuable model for further studies using published infection

models for other pathogens and the study of the interactions with gut microbiota in zebrafish

larvae.

Methods

Zebrafish husbandry

All zebrafish were handled in compliance with the local animal welfare regulations and

maintained according to standard protocols (zfin.org). Larvae were raised in egg water (60

g/ml Instant Ocean sea salts) at 28.5 °C. For the duration of bacterial injections, larvae were

kept under anesthesia in egg water containing 0.02% buffered 3-aminobenzoic acid ethyl ester

(Tricaine, Sigma-Aldrich, the Netherlands). The culture of zebrafish with mutations in

immune genes was approved by the local animal welfare committee (DEC) of the University

of Leiden (protocol 14198). All protocols adhered to the international guidelines specified by

the EU Animal Protection Directive 2010/63/EU.

The *tlr2*^{sa19423} mutant line (ENU-mutagenized) was obtained from the Sanger Institute

Zebrafish Mutation Resource (Hinxton, Cambridge, UK) and shipped by the Zebrafish

Resource Center of the Karlsruhe Institute of Technology. The mutant allele was identified by

sequencing. Heterozygous carriers of the mutation were outcrossed three times against wild

type (*AB* strain), and were subsequently incrossed three times. Heterozygous fish of the

resulting family were used to produce embryos. Homozygous mutants were outcrossed to the

Tg(mpeg1:mCherry-F);TgBAC(mpx:EGFP) double transgenic line [78, 79], and the offspring

with GFP and mCherry fluorescence were subsequently incrossed to produce the

Tg(mpeg1:mCherry-F);TgBAC(mpx:EGFP) line.

Bacterial strain preparation

The bacterial strain, *Mycobacterium marinum* m20 (Mma20) expressing mCherry fluorescent

protein [80], was used in this study. For the infection to zebrafish larvae, the bacteria were

prepared as previously described [81]. The infection inoculum was prepared in 2%

polyvinylpyrrolidone40 solution (Calbiochem, the Netherlands), and 150 colony-forming

units (CFU) of bacteria were injected into the blood stream at 28 hours post fertilization (hpf)

as previously described [82].

Ligands injection

Purified Pam3CSK4 (InvivoGen, France) and flagellin from *S. typhimurium* (Flagellin FliC

VacciGrade™, Invitrogen, France) were diluted in 1 mg/ml and 100 µg/ml in sterile water,

respectively. For injection, 1 nl of the ligand solutions were injected into the blood stream at

28hpf. Sterile water was injected as a control experiment. Injections were performed using a

FemtoJet microinjector (Eppendorf, the Netherlands) equipped with a capillary glass needle.

Bacterial burden imaging and quantification

425 Pools of 20 larvae were collected at 3- and 4-day post infection (dpi) and imaged by using the

426 Leica MZ16FA Fluorescence Stereo Microscope (Leica Microsystems, Wetzlar Germany)

427 equipped with the DFC420C color camera (Leica Microsystems). Bacterial loads were

428 analyzed using dedicated pixel counting software as previously described [83].

Confocal laser scanning microscopy imaging and image quantification

Larvae (2 dpf) were embedded in 1% low melting point agarose (Sigma Aldrich), and image

acquisition was performed by using a Leica TCS SP8 confocal microscope (Leica

Microsystems) with 10 times objectives (N.A. 0.4). Acquisition settings and area of imaging

(in the caudal vein region) were kept the same across the groups for macrophages and

neutrophils number counting (Fig. 1 j, k) and pixel counting (Fig. S1). Experiments were

performed in two independent series. Double fluorescent lines *tlr2*$^{+/+}$ *Tg*(*mpeg1:mCherry-*

F);TgBAC(mpx:EGFP) and *tlr2*$^{-/-}$ *Tg*(*mpeg1:mCherry-F);TgBAC(mpx:EGFP)* were used for

macrophages and neutrophils number counting. Macrophage and neutrophil cell counting was

either performed manually or by using the plugin Find Maxima in Fiji

(http://imagej.nih.gov/ij/docs/menus/ process.html#find-maxima) (Fig. 1 j, k) using

projections of the z-series. 25 individual larvae for each group were used for counting and

representative images are given in Fig. 1h, i. With manual counting the z-series data was

examined in cases when it was unclear whether the fluorescent pixels presented one or

multiple cells. The result of manual cell counting (Fig. S10) is comparable with the result of

automated cell counting using Fiji. Pixel counting of the double transgenic lines (Fig. 1S a, b)

was performed using dedicated pixel counting software as well as previously described [83].

RNA isolation, cDNA synthesis and qPCR

Total RNAs were extracted using TRIzol Reagent (Life Technologies) and purified using

RNeasy MinElute Cleanup Kit (Qiagen, the Netherlands). The concentration and quality of

RNAs were evaluated by NanoDrop 2000 (Thermo Scientific, the Netherlands). cDNAs were

synthesized from 1 µg total RNAs and qPCR were performed by using the iScript™ cDNA

Synthesis Kit (BioRad, the Netherlands) and iQ™ SYBR Green Supermix (BioRad) and

normalized against the expression of *ppial* as a housekeeping gene [84]. Results were

analyzed using the ΔΔCt method [85]. Primer sequences are described in Supplementary

Table 5.

Deep sequencing and data analysis

Triplicates of 10 larvae of $tlr2^{+/-}$ and $tlr2^{-/-}$ with PBS (as control) or Mma20 injection, were

homogenized in 300ul of TRIzol reagent, and total RNAs were purified as described above.

RNAseq was performed using Illumina Hi-Seq 2500 as previously described [86]. The raw

data is available in the NCBI GEO database under accession number GSE102766. The

RNAseq data were mapped on the zebrafish genome (version GRCz10) and tag counts were

performed by Bowtie 2 using GeneTiles software (http://www. genetiles. com) [87]. Then, we

performed normalization and gene expression analysis using the R package and DESeq2 [88].

After statistical tests, we performed further bioinformatics analyses Gene-Set Enrichment

Analysis [48], Sub-Network Enrichment Analysis [49] and Pathway Enrichment Analysis

[89]. For creating gene networks based on common regulatory targets, we used Pathway

Studio 9.0 (Elsevier, Amsterdam, the Netherlands) as previously described [47].

Comparison of edgeR and DEseq2: EdgeR and DEseq2 differ mainly in the aspects of

normalization and estimation of the dispersion parameters. Normalization in edgeR is done

via the trimmed mean of M values, while in DEseq2 this is done by comparing each library

with a virtual library based on the relative log expressions. The dispersion parameters in

edgeR are estimated by empirical Bayes and are therefore shrunken towards the overall mean

of the estimates. Dispersion in DEseq2 is estimated by taking the maximum of the individual

dispersions and the mean trend of the dispersions. As a consequence, edgeR tends to be more

sensitive to outliers, while DEseq2 is less powerful [44].

Statistical analyses

Graphpad Prism software (Version 7.00; GraphPad Software, San Diego, CA, USA) was used

for statistical analysis. All experiment data are shown as mean ± SEM. In immune gene

expression in $tlr2^{+/-}$ and $tlr2^{-/-}$ fish lines infected with Mm (Fig. 4), statistical significance of

differences was determined by two-way ANOVA with Šidák Multiple Comparison test as a

post-hoc test. The other experiments were analyzed by using unpaired, two-tailed t-tests for

comparisons between two groups and one-way ANOVA with Tukey's multiple comparison

methods as a post-hoc test for comparisons between more than two groups. (ns, no significant

difference; $*p<0.05$; $**p<0.01$; $***p<0.001$; $****p<0.0001$).

List of abbreviations

CFU : colony-forming units

CLSM: confocal laser scanning microscopy

DEGs: differential expressed genes

Dpf: days post fertilization

Dpi: days post injection

FDR: false discovery rate

FACS: Fluorescence-activated cell sorting

GSEA: Gene-Set Enrichment Analysis

Hpf: hours post fertilization

Hpi: hours post injection

Mm: *Mycobacterium marinum*

Mtb: *Mycobacterium tuberculosis*

NMD: non-sense mediated mRNA decay

qPCR: quantitative polymerase chain reaction

RNAseq: Deep sequencing of cDNA derived from polyA RNA

SNEA: Sub-Network Enrichment Analysis

TB: tuberculosis

Declarations

Acknowledgments

We thank all members of the fish facility team for fish caretaking. We would like to thank

Lanpeng Chen and Gerda E.M. Lamers for assistance with confocal laser scanning imaging.

We also thank Vincenzo Torraca for helpful discussions and the qPCR primers of *cxcl11*-like

genes. The zebrafish *tlr2* mutant was obtained from the Sanger Institute Zebrafish Mutation

Resource (ZF-MODELS Integrated Project funded by the European Commission; contract

number LSHG-CT-2003-503496), also sponsored by the Welcome Trust [grant number WT

077047/Z/05/Z].

515 Funding

516 S. Y. and W.H were supported by a grant from the China Scholarship Council (CSC). This

517 funding body had no role in the design of the study and collection, analysis, and interpretation

518 of data and in writing the manuscript

520 Availability of data and materials

521 The raw data of the RNAseq experiments is available in the NCBI GEO database under

522 accession number GSE102766.

524 Authors' contributions

SY and WH carried out all the experimental operations. SY, YS, and HPS performed RNAseq

data analysis. SY, WH and HPS wrote the first draft of the manuscript. SY, WH and HPS

participated in the design of the study and coordination. AHM and RMJ helped to revise the

final manuscript. HPS conceived the study. MM assisted in statistical analysis of RNAseq
data. All authors read and approved the final manuscript.

Ethics approval and consent to participate

No animals were used for experimentation. Larvae for experiments were obtained from
zebrafish lines that were handled in compliance with the local animal welfare regulations and
maintained according to standard protocols (zfin.org). The breeding of adult fish was
approved by the local animal welfare committee (DEC) of the University of Leiden. All
protocols adhered to the international guidelines specified by the EU Animal Protection
Directive 2010/63/EU for which larvae under the age of 5 days post fertilization are not
considered test animals.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Ramakrishnan L: **The zebrafish guide to tuberculosis immunity and treatment**. *Cold Spring*
Harbor symposia on quantitative biology 2013, **78**:179-192.

2.   Co DO, Hogan LH, Kim SI, Sandor M: **Mycobacterial granulomas: keys to a long-lasting host-**
pathogen relationship. *Clin Immunol* 2004, **113**(2):130-136.

3.   Salgame P: **Host innate and Th1 responses and the bacterial factors that control**
Mycobacterium tuberculosis infection. *Curr Opin Immunol* 2005, **17**(4):374-380.

4.   Ramakrishnan L: **Revisiting the role of the granuloma in tuberculosis**. *Nature reviews*
Immunology 2012, **12**(5):352-366.

5.   Jo EK, Yang CS, Choi CH, Harding CV: **Intracellular signalling cascades regulating innate**
immune responses to Mycobacteria: branching out from Toll-like receptors. *Cell Microbiol*
2007, **9**(5):1087-1098.

6.   Tapping RI, Tobias PS: **Mycobacterial lipoarabinomannan mediates physical interactions**
between TLR1 and TLR2 to induce signaling. *J Endotoxin Res* 2003, **9**(4):264-268.

7.   Means TK, Jones BW, Schromm AB, Shurtleff BA, Smith JA, Keane J, Golenbock DT, Vogel SN,
Fenton MJ: **Differential effects of a Toll-like receptor antagonist on Mycobacterium**
tuberculosis-induced macrophage responses. *J Immunol* 2001, **166**(6):4074-4082.

8.   Quesniaux VJ, Nicolle DM, Torres D, Kremer L, Guerardel Y, Nigou J, Puzo G, Erard F, Ryffel B:
Toll-like receptor 2 (TLR2)-dependent-positive and TLR2-independent-negative regulation
of proinflammatory cytokines by mycobacterial lipomannans. *J Immunol* 2004, **172**(7):4425-
4434.

9.   Jung SB, Yang CS, Lee JS, Shin AR, Jung SS, Son JW, Harding CV, Kim HJ, Park JK, Paik TH *et al*:
The mycobacterial 38-kilodalton glycolipoprotein antigen activates the mitogen-activated
protein kinase pathway and release of proinflammatory cytokines through Toll-like
receptors 2 and 4 in human monocytes. *Infect Immun* 2006, **74**(5):2686-2696.

10.   Brightbill HD, Libraty DH, Krutzik SR, Yang RB, Belisle JT, Bleharski JR, Maitland M, Norgard
MV, Plevy SE, Smale ST *et al*: **Host defense mechanisms triggered by microbial lipoproteins**
through toll-like receptors. *Science* 1999, **285**(5428):732-736.

11.   Noss EH, Pai RK, Sellati TJ, Radolf JD, Belisle J, Golenbock DT, Boom WH, Harding CV: **Toll-like**
receptor 2-dependent inhibition of macrophage class II MHC expression and antigen
processing by 19-kDa lipoprotein of Mycobacterium tuberculosis. *J Immunol* 2001,
167(2):910-918.

12.   Gilleron M, Quesniaux VF, Puzo G: **Acylation state of the phosphatidylinositol**
hexamannosides from Mycobacterium bovis bacillus Calmette Guerin and mycobacterium
tuberculosis H37Rv and its implication in Toll-like receptor response. *J Biol Chem* 2003,
278(32):29880-29889.

13.   Gilleron M, Nigou J, Nicolle D, Quesniaux V, Puzo G: **The acylation state of mycobacterial**
lipomannans modulates innate immunity response through toll-like receptor 2. *Chem Biol*
2006, **13**(1):39-47.

14.   Takeuchi O, Sato S, Horiuchi T, Hoshino K, Takeda K, Dong Z, Modlin RL, Akira S: **Cutting edge:**
role of Toll-like receptor 1 in mediating immune response to microbial lipoproteins. *J*
Immunol 2002, **169**(1):10-14.

15.   Takeuchi O, Kawai T, Mühlradt PF, Morr M, Radolf JD, Zychlinsky A, Takeda K, Akira S:
Discrimination of bacterial lipoproteins by Toll-like receptor 6. *International Immunology*
2001, **13**(7):933-940.

16.   Marinho FAV, de Paula RR, Mendes AC, de Almeida LA, Gomes MTR, Carvalho NB, Oliveira FS,
Caliari MV, Oliveira SC: **Toll-like receptor 6 senses Mycobacterium avium and is required for**
efficient control of mycobacterial infection. *European Journal of Immunology* 2013,
43(9):2373-2385.

17.   Basu J, Shin DM, Jo EK: **Mycobacterial signaling through toll-like receptors**. *Front Cell Infect*
Microbiol 2012, **2**:145.

18.   Underhill DM, Ozinsky A, Smith KD, Aderem A: **Toll-like receptor-2 mediates mycobacteria-**
induced proinflammatory signaling in macrophages. *Proc Natl Acad Sci U S A* 1999,
96(25):14459-14463.

19. Carlos D, Frantz FG, Souza-Junior DA, Jamur MC, Oliver C, Ramos SG, Quesniaux VF, Ryffel B,
Silva CL, Bozza MT *et al*: **TLR2-dependent mast cell activation contributes to the control of**
Mycobacterium tuberculosis infection. *Microbes Infect* 2009, **11**(8-9):770-778.

20. Reiling N, Holscher C, Fehrenbach A, Kroger S, Kirschning CJ, Goyert S, Ehlers S: **Cutting edge:**
Toll-like receptor (TLR)2- and TLR4-mediated pathogen recognition in resistance to
airborne infection with Mycobacterium tuberculosis. *J Immunol* 2002, **169**(7):3480-3484.

21. Drennan MB, Nicolle D, Quesniaux VJ, Jacobs M, Allie N, Mpagi J, Fremond C, Wagner H,
Kirschning C, Ryffel B: **Toll-like receptor 2-deficient mice succumb to Mycobacterium**
tuberculosis infection. *Am J Pathol* 2004, **164**(1):49-57.

22. McBride A, Konowich J, Salgame P: **Host defense and recruitment of Foxp3(+) T regulatory**
cells to the lungs in chronic Mycobacterium tuberculosis infection requires toll-like
receptor 2. *PLoS Pathog* 2013, **9**(6):e1003397.

23. Ogus AC, Yoldas B, Ozdemir T, Uguz A, Olcen S, Keser I, Coskun M, Cilli A, Yegin O: **The**
Arg753GLn polymorphism of the human toll-like receptor 2 gene in tuberculosis disease.
Eur Respir J 2004, **23**(2):219-223.

24. Biswas D, Gupta SK, Sindhwani G, Patras A: **TLR2 polymorphisms, Arg753Gln and Arg677Trp,**
are not associated with increased burden of tuberculosis in Indian patients. *BMC research*
notes 2009, **2**:162.

25. Gehring AJ, Dobos KM, Belisle JT, Harding CV, Boom WH: **Mycobacterium tuberculosis LprG**
(Rv1411c): a novel TLR-2 ligand that inhibits human macrophage class II MHC antigen
processing. *J Immunol* 2004, **173**(4):2660-2668.

26. Saraav I, Singh S, Sharma S: **Outcome of Mycobacterium tuberculosis and Toll-like receptor**
interaction: immune response or immune evasion? *Immunol Cell Biol* 2014, **92**(9):741-746.

27. Netea MG, Van der Meer JW, Kullberg BJ: **Toll-like receptors as an escape mechanism from**
the host defense. *Trends Microbiol* 2004, **12**(11):484-488.

28. Noss EH, Harding CV, Boom WH: **Mycobacterium tuberculosis inhibits MHC class II antigen**
processing in murine bone marrow macrophages. *Cell Immunol* 2000, **201**(1):63-74.

29. Netea MG, Sutmuller R, Hermann C, Van der Graaf CA, Van der Meer JW, van Krieken JH,
Hartung T, Adema G, Kullberg BJ: **Toll-like receptor 2 suppresses immunity against Candida**
albicans through induction of IL-10 and regulatory T cells. *J Immunol* 2004, **172**(6):3712-
3718.

30.   Sing A, Rost D, Tvardovskaia N, Roggenkamp A, Wiedemann A, Kirschning CJ, Aepfelbacher M,
Heesemann J: **Yersinia V-antigen exploits toll-like receptor 2 and CD14 for interleukin 10-**
mediated immunosuppression. *J Exp Med* 2002, **196**(8):1017-1024.

31. Gehring AJ, Rojas RE, Canaday DH, Lakey DL, Harding CV, Boom WH: **The Mycobacterium**
tuberculosis 19-kilodalton lipoprotein inhibits gamma interferon-regulated HLA-DR and Fc
gamma R1 on human macrophages through Toll-like receptor 2. *Infect Immun* 2003,
71(8):4487-4497.

32. Alvarez GR, Zwilling BS, Lafuse WP: **Mycobacterium avium inhibition of IFN-gamma**
signaling in mouse macrophages: Toll-like receptor 2 stimulation increases expression of
dominant-negative STAT1 beta by mRNA stabilization. *J Immunol* 2003, **171**(12):6766-6773.

33. Agrawal S, Agrawal A, Doughty B, Gerwitz A, Blenis J, Van Dyke T, Pulendran B: **Cutting edge:**
different Toll-like receptor agonists instruct dendritic cells to induce distinct Th responses
via differential modulation of extracellular signal-regulated kinase-mitogen-activated
protein kinase and c-Fos. *J Immunol* 2003, **171**(10):4984-4989.

34. Lam SH, Chua HL, Gong Z, Lam TJ, Sin YM: **Development and maturation of the immune**
system in zebrafish, Danio rerio: a gene expression profiling, in situ hybridization and
immunological study. *Dev Comp Immunol* 2004, **28**(1):9-28.

35. Willett CE, Cortes A, Zuasti A, Zapata AG: **Early hematopoiesis and developing lymphoid**
organs in the zebrafish. *Developmental dynamics : an official publication of the American*
Association of Anatomists 1999, **214**(4):323-336.

36.   Swaim LE, Connolly LE, Volkman HE, Humbert O, Born DE, Ramakrishnan L: **Mycobacterium**
marinum Infection of Adult Zebrafish Causes Caseating Granulomatous Tuberculosis and Is
Moderated by Adaptive Immunity. *Infection and Immunity* 2006, **74**(11):6108-6117.
37.   Davis JM, Clay H, Lewis JL, Ghori N, Herbomel P, Ramakrishnan L: **Real-time visualization of**
mycobacterium-macrophage interactions leading to initiation of granuloma formation in
zebrafish embryos. *Immunity* 2002, **17**(6):693-702.
38.   Meijer AH: **Protection and pathology in TB: learning from the zebrafish model**. *Semin*
Immunopathol 2016, **38**(2):261-273.
39.   Cronan MR, Tobin DM: **Fit for consumption: zebrafish as a model for tuberculosis**. *Disease*
models & mechanisms 2014, **7**(7):777-784.
40.   Yang S, Marin-Juez R, Meijer AH, Spaink HP: **Common and specific downstream signaling**
targets controlled by Tlr2 and Tlr5 innate immune signaling in zebrafish. *BMC Genomics*
2015, **16**:547.
41.   Medzhitov R, Preston-Hurlburt P, Kopp E, Stadlen A, Chen C, Ghosh S, Janeway CA, Jr.:
MyD88 is an adaptor protein in the hToll/IL-1 receptor family signaling pathways. *Mol Cell*
1998, **2**(2):253-258.
42.   Fitzgerald KA, Palsson-McDermott EM, Bowie AG, Jefferies CA, Mansell AS, Brady G, Brint E,
Dunne A, Gray P, Harte MT *et al*: **Mal (MyD88-adapter-like) is required for Toll-like**
receptor-4 signal transduction. *Nature* 2001, **413**(6851):78-83.
43.   Bjørn E. V. Koch SY, Gerda Lamers, Jens Stougaard & Herman P. Spaink: **Intestinal**
microbiome adjusts the innate immune setpoint during colonization through negative
regulation of MyD88. *nature comunication* 2018, **9**(1):4099.
44.   Anders S, McCarthy DJ, Chen Y, Okoniewski M, Smyth GK, Huber W, Robinson MD: **Count-**
based differential expression analysis of RNA sequencing data using R and Bioconductor.
Nature protocols 2013, **8**(9):1765-1786.
45.   Torraca V, Cui C, Boland R, Bebelman JP, van der Sar AM, Smit MJ, Siderius M, Spaink HP,
Meijer AH: **The CXCR3-CXCL11 signaling axis mediates macrophage recruitment and**
dissemination of mycobacterial infection. *Disease models & mechanisms* 2015, **8**(3):253-269.
46.   Rougeot J, Torraca V, Zakrzewska A, Kanwal Z, Jansen HJ, Sommer F, Spaink HP, Meijer AH:
RNAseq Profiling of Leukocyte Populations in Zebrafish Larvae Reveals a cxcl11 Chemokine
Gene as a Marker of Macrophage Polarization During Mycobacterial Infection. *Frontiers in*
Immunology 2019, **10**.
47.   Nikitin A, Egorov S, Daraselia N, Mazo I: **Pathway studio--the analysis and navigation of**
molecular networks. *Bioinformatics* 2003, **19**(16):2155-2157.
48.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A,
Pomeroy SL, Golub TR, Lander ES *et al*: **Gene set enrichment analysis: a knowledge-based**
approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* 2005,
102(43):15545-15550.
49.   Kotelnikova E, Yuryev A, Mazo I, Daraselia N: **Computational approaches for drug**
repositioning and combination therapy design. *J Bioinform Comput Biol* 2010, **8**(3):593-606.
50.   Verway M, Bouttier M, Wang TT, Carrier M, Calderon M, An BS, Devemy E, McIntosh F,
Divangahi M, Behr MA *et al*: **Vitamin D induces interleukin-1beta expression: paracrine**
macrophage epithelial signaling controls M. tuberculosis infection. *PLoS Pathog* 2013,
9(6):e1003407.
51.   Leisching G, Pietersen RD, Mpongoshe V, van Heerden C, van Helden P, Wiid I, Baker B: **The**
Host Response to a Clinical MDR Mycobacterial Strain Cultured in a Detergent-Free
Environment: A Global Transcriptomics Approach. *PLoS One* 2016, **11**(4):e0153079.
52.   Kubler A, Larsson C, Luna B, Andrade BB, Amaral EP, Urbanowski M, Orandle M, Bock K,
Ammerman NC, Cheung LS *et al*: **Cathepsin K Contributes to Cavitation and Collagen**
Turnover in Pulmonary Tuberculosis. *J Infect Dis* 2016, **213**(4):618-627.

53.   Zhang H, Sun Z, Wei W, Liu Z, Fleming J, Zhang S, Lin N, Wang M, Chen M, Xu Y *et al*:
Identification of serum microRNA biomarkers for tuberculosis using RNA-seq. *PLoS One*
2014, **9**(2):e88909.

54.   Mvubu NE, Pillay B, Gamieldien J, Bishai W, Pillay M: **Canonical pathways, networks and**
transcriptional factor regulation by clinical strains of Mycobacterium tuberculosis in
pulmonary alveolar epithelial cells. *Tuberculosis (Edinb)* 2016, **97**:73-85.

55.   Ziegler G, Freyer D, Harhausen D, Khojasteh U, Nietfeld W, Trendelenburg G: **Blocking TLR2**
in vivo protects against accumulation of inflammatory cells and neuronal injury in
experimental stroke. *J Cereb Blood Flow Metab* 2011, **31**(2):757-766.

56.   Hoffmann O, Braun JS, Becker D, Halle A, Freyer D, Dagand E, Lehnardt S, Weber JR: **TLR2**
mediates neuroinflammation and neuronal damage. *J Immunol* 2007, **178**(10):6476-6481.

57.   Gu Y, Zhang Y, Bi Y, Liu J, Tan B, Gong M, Li T, Chen J: **Mesenchymal stem cells suppress**
neuronal apoptosis and decrease IL-10 release via the TLR2/NFkappaB pathway in rats with
hypoxic-ischemic brain damage. *Mol Brain* 2015, **8**(1):65.

58.   Kim C, Rockenstein E, Spencer B, Kim HK, Adame A, Trejo M, Stafa K, Lee HJ, Lee SJ, Masliah E:
Antagonizing Neuronal Toll-like Receptor 2 Prevents Synucleinopathy by Activating
Autophagy. *Cell Rep* 2015, **13**(4):771-782.

59.   Moriguchi T, Hamada M, Morito N, Terunuma T, Hasegawa K, Zhang C, Yokomizo T, Esaki R,
Kuroda E, Yoh K *et al*: **MafB is essential for renal development and F4/80 expression in**
macrophages. *Mol Cell Biol* 2006, **26**(15):5715-5727.

60.   Bauquet AT, Jin H, Paterson AM, Mitsdoerffer M, Ho IC, Sharpe AH, Kuchroo VK: **The**
costimulatory molecule ICOS regulates the expression of c-Maf and IL-21 in the
development of follicular T helper cells and TH-17 cells. *Nat Immunol* 2009, **10**(2):167-175.

61.   Cao S, Liu J, Song L, Ma X: **The protooncogene c-Maf is an essential transcription factor for**
IL-10 gene expression in macrophages. *J Immunol* 2005, **174**(6):3484-3492.

62.   Aziz A, Soucie E, Sarrazin S, Sieweke MH: **MafB/c-Maf deficiency enables self-renewal of**
differentiated functional macrophages. *Science* 2009, **326**(5954):867-871.

63.   Smith PA: **The tantalizing links between gut microbes and the brain**. *Nature* 2015,
526(7573):312-314.

64.   Davis DJ, Bryda EC, Gillespie CH, Ericsson AC: **Microbial modulation of behavior and stress**
responses in zebrafish larvae. *Behav Brain Res* 2016, **311**:219-227.

65.   de Abreu MS, Giacomini A, Zanandrea R, Dos Santos BE, Genario R, de Oliveira GG, Friend AJ,
Amstislavskaya TG, Kalueff AV: **Psychoneuroimmunology and immunopsychiatry of**
zebrafish. *Psychoneuroendocrinology* 2018, **92**:1-12.

66.   Phelps D, Brinkman NE, Keely SP, Anneken EM, Catron TR, Betancourt D, Wood CE,
Espenschied ST, Rawls JF, Tal T: **Microbial colonization is required for normal**
neurobehavioral development in zebrafish. *Sci Rep* 2017, **7**(1):11244.

67.   Borrelli L, Aceto S, Agnisola C, De Paolo S, Dipineto L, Stilling RM, Dinan TG, Cryan JF, Menna
LF, Fioretti A: **Probiotic modulation of the microbiota-gut-brain axis and behaviour in**
zebrafish. *Sci Rep* 2016, **6**:30046.

68.   Richardson ET, Shukla S, Sweet DR, Wearsch PA, Tsichlis PN, Boom WH, Harding CV: **Toll-like**
receptor 2-dependent extracellular signal-regulated kinase signaling in Mycobacterium
tuberculosis-infected macrophages drives anti-inflammatory responses and inhibits Th1
polarization of responding T cells. *Infect Immun* 2015, **83**(6):2242-2254.

69.   Piermattei A, Migliara G, Di Sante G, Foti M, Hayrabedyan SB, Papagna A, Geloso MC, Corbi
M, Valentini M, Sgambato A *et al*: **Toll-Like Receptor 2 Mediates In Vivo Pro- and Anti-**
inflammatory Effects of Mycobacterium Tuberculosis and Modulates Autoimmune
Encephalomyelitis. *Front Immunol* 2016, **7**:191.

70.   Davila S, Hibberd ML, Hari Dass R, Wong HE, Sahiratmadja E, Bonnard C, Alisjahbana B,
Szeszko JS, Balabanova Y, Drobniewski F *et al*: **Genetic association and expression studies**
indicate a role of toll-like receptor 8 in pulmonary tuberculosis. *PLoS Genet* 2008,
4(10):e1000218.

71. Lai YF, Lin TM, Wang CH, Su PY, Wu JT, Lin MC, Eng HL: **Functional polymorphisms of the**
TLR7 and TLR8 genes contribute to Mycobacterium tuberculosis infection. *Tuberculosis*
(Edinb) 2016, **98**:125-131.
72. Reeme AE, Robinson RT: **Dietary Vitamin D3 Suppresses Pulmonary Immunopathology**
Associated with Late-Stage Tuberculosis in C3HeB/FeJ Mice. *J Immunol* 2016, **196**(3):1293-
1304.
73. Liu PT, Stenger S, Li H, Wenzel L, Tan BH, Krutzik SR, Ochoa MT, Schauber J, Wu K, Meinken C
et al: **Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response**.
Science 2006, **311**(5768):1770-1773.
74. Haire RN, Cannon JP, O'Driscoll ML, Ostrov DA, Mueller MG, Turner PM, Litman RT, Litman
GW, Yoder JA: **Genomic and functional characterization of the diverse immunoglobulin**
domain-containing protein (DICP) family. *Genomics* 2012, **99**(5):282-291.
75. Rodriguez-Nunez I, Wcisel DJ, Litman GW, Yoder JA: **Multigene families of immunoglobulin**
domain-containing innate immune receptors in zebrafish: deciphering the differences. *Dev*
Comp Immunol 2014, **46**(1):24-34.
76. Blander JM, Medzhitov R: **Regulation of Phagosome Maturation by Signals from Toll-Like**
Receptors. *Science* 2004, **304**(5673):1014-1018.
77. Rahman MJ, Chuquimia OD, Petursdottir DH, Periolo N, Singh M, Fernandez C: **Impact of toll-**
like receptor 2 deficiency on immune responses to mycobacterial antigens. *Infect Immun*
2011, **79**(11):4649-4656.
78. Bernut A, Herrmann JL, Kissa K, Dubremetz JF, Gaillard JL, Lutfalla G, Kremer L:
Mycobacterium abscessus cording prevents phagocytosis and promotes abscess formation.
Proc Natl Acad Sci U S A 2014, **111**(10):E943-952.
79. Renshaw SA, Loynes CA, Trushell DM, Elworthy S, Ingham PW, Whyte MK: **A transgenic**
zebrafish model of neutrophilic inflammation. *Blood* 2006, **108**(13):3976-3978.
80. van der Sar AM, Spaink HP, Zakrzewska A, Bitter W, Meijer AH: **Specificity of the zebrafish**
host transcriptome response to acute and chronic mycobacterial infection and the role of
innate and adaptive immune components. *Molecular immunology* 2009, **46**(11–12):2317-
2332.
81. Benard EL, van der Sar AM, Ellett F, Lieschke GJ, Spaink HP, Meijer AH: **Infection of zebrafish**
embryos with intracellular bacterial pathogens. *Journal of visualized experiments : JoVE*
2012(61):2.
82. Stockhammer OW, Zakrzewska A, Hegedus Z, Spaink HP, Meijer AH: **Transcriptome profiling**
and functional analyses of the zebrafish embryonic innate immune response to Salmonella
infection. *J Immunol* 2009, **182**(9):5641-5653.
83. Stoop EJM, Schipper T, Rosendahl Huber SK, Nezhinsky AE, Verbeek FJ, Gurcha SS, Besra GS,
Vandenbroucke-Grauls CMJE, Bitter W, van der Sar AM: **Zebrafish embryo screen for**
mycobacterial genes involved in the initiation of granuloma formation reveals a newly
identified ESX-1 component. *Disease models & mechanisms* 2011, **4**(4):526-536.
84. van Soest JJ, Stockhammer OW, Ordas A, Bloemberg GV, Spaink HP, Meijer AH: **Comparison**
of static immersion and intravenous injection systems for exposure of zebrafish embryos
to the natural pathogen Edwardsiella tarda. *BMC Immunol* 2011, **12**:58.
85. Livak KJ, Schmittgen TD: **Analysis of relative gene expression data using real-time**
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 2001, **25**(4):402-408.
86. Marin-Juez R, Jong-Raadsen S, Yang S, Spaink HP: **Hyperinsulinemia induces insulin**
resistance and immune suppression via Ptpn6/Shp1 in zebrafish. *The Journal of*
endocrinology 2014.
87. Veneman W, de Sonneville J, van der Kolk KJ, Ordas A, Al-Ars Z, Meijer A, Spaink H: **Analysis**
of RNAseq datasets from a comparative infectious disease zebrafish model using GeneTiles
bioinformatics. *Immunogenetics* 2015, **67**(3):135-147.
88. Love MI, Huber W, Anders S: **Moderated estimation of fold change and dispersion for RNA-**
seq data with DESeq2. *Genome Biol* 2014, **15**(12):550.

89.   Kotelnikova E, Shkrob MA, Pyatnitskiy MA, Ferlini A, Daraselia N: **Novel approach to meta-**
analysis of microarray datasets reveals muscle remodeling-related drug targets and
biomarkers in Duchenne muscular dystrophy. *PLoS Comput Biol* 2012, **8**(2):e1002365.

Figure legends

Figure 1 Characterization of the Tlr2 mutant. (a) mutant DNA and protein sequence. A

point mutation (T to A) in the C-terminal of the second LRR domain of zebrafish Tlr2

introduces a premature stop codon. The predicted truncated protein lacks the whole TIR

domain. Nucleotide and amino acid positions are indicated with respect to the translation start

codon. $tlr2^{+/+}$, $tlr2^{+/-}$ and $tlr2^{-/-}$ embryos were injected at 27 hpf with 1 ng Pam3CSK4 (b-d) or

0.1ng flagellin (e-g) and expression levels of *il1b*, *fosl1a* and *cebpb* were determined at 1hour

post injection by qPCR. Data (mean ± SEM) are combined from three biological replicates

(n=10 embryos per group) and expressed relative to their corresponding mock injection (water)

control, which set at 1. Statistical significance of differences between mock and Pam3CSK4

groups was determined by one-way ANOVA with Tukey's Multiple Comparison method as a

post-hoc test (b-g), ns, non-significant; *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$; ****, $P<0.0001$.

(h, i). Representative Stereo images of the caudal hematopoietic tissue of $tlr2^{+/+}$

Tg(mpeg1:mCherry-F);TgBAC(mpx:EGFP)   (h)   and $tlr2^{-/-}$ *Tg(mpeg1:mCherry-*

F);TgBAC(mpx:EGFP) (i) were taken at 2 dpf for quantification of macrophages and

neutrophils numbers. At 2 dpf, numbers of mCherry-labeled macrophages (j) and GFP-

labeled neutrophils (k) were counted using Leica TCS SP8 confocal laser scanning

microscopy (CLSM) of transgenic lines. Data (mean ± SEM) were combined from two

independent experiments. No significant differences (ns) in the number of macrophages (j)

and neutrophils (k) was detected with a t-test.

Figure 2 Gene expression in the absence of infection between Tlr2 mutants. (a) a number

of down- and up-regulated genes per method (edgeR and DEseq2) and (b) a Venn diagram

that compares the down-, up- and non-differentially expressed genes per method at a

significance level of 10^{-10} in the edgeR and DEseq2 analyses methods. Down-regulated means

that a gene is less expressed in the $tlr2^{+/-}$ compared to the $tlr2^{-/-}$ strain.

Figure 3 Quantification of bacterial burden. $tlr2^{+/+}$ (a), $tlr2^{+/-}$ (b) and $tlr2^{-/-}$ (c) embryos

were infected with mCherry-labeled *M. marinum* strain Mma 20 at a dose of ~150 CFU by

caudal vein infection at 28 hpf. Representative images for bacterial pixel count in $tlr2^{+/+}$ (a),

$tlr2^{+/-}$ (b) and $tlr2^{-/-}$ (c) were taken at 4 dpi. Bacterial burden of $tlr2^{+/+}$, $tlr2^{+/-}$ and $tlr2^{-/-}$ were

also quantified at 3 dpi (d) and 4 dpi (e). Bacterial burdens were quantified by using bacterial

fluorescence pixels. Red stars in (e) indicate the data for the representative images shown in

(a-c). In (d, e), data (mean ± SEM) were combined from two independent experiments.

Statistical significance of differences was determined by one-way ANOVA with Tukey's

Multiple Comparison method as a post-hoc test for comparison between more than two

groups (d, e). ns, non-significant; ***, $P<0.001$; ****, $P<0.0001$.

Figure 4 Immune genes expression in $tlr2^{+/-}$ and $tlr2^{-/-}$ fish lines infected with Mm. The

expression levels of *il1b* (a), *tnfa* (b), *tnfb* (c), *irg1l* (d), *fosl1a* (e), *cebpb* (f), *cxcl11aa* (g) and

cxcl11ac (h) were determined at 4dpi by qPCR. Data (mean ± SEM) are derived from at least

three biological replicates (n = 10 embryos per group) and expressed relative to their

corresponding mock injection (PBS) control, which is set at 1. Statistical significance of

differences was determined by two-way ANOVA with Šidák Multiple Comparison test as a

post-hoc test. *$P<0.05$, **$P<0.01$, ***$P<0.001$

Figure 5 Overview of RNAseq results. (a, b) the number of DEGs of $tlr2^{+/-}$ and $tlr2^{-/-}$ strains

infected with Mm compared to the control at different *p*-value and fold change. (c), the work

flow of screening genes of which the regulation by infection is dependent on *tlr2*. (d), GO

analysis of the 97 upregulated genes. (e), GO analysis of the 92 down-regulated genes. FC,

fold-change.)

Figure 6 Overview of fold changes of representative genes selected from the gene

categories resulting from GO-term analysis. (a-e): *tlr2*-dependent genes with up regulation

corresponding to Fig. 5d. (f): *tlr2* specific genes with down regulation corresponding to Fig.

5e.

Supplementary figure 1 Pixel count analysis for double transgenic lines *Tg(mpeg1:mCherry-*

F);TgBAC(mpx:EGFP) of 2 dpf *tlr2*$^{+/+}$ and *tlr2*$^{-/-}$ embryos. a: mCherry reporter, b, EGFP

reporter.

Supplementary figure 2 RNAseq read counts of *tlr2* transcripts in heterozygotes (*tlr2*$^{+/-}$)

control versus *tlr2* mutant (*tlr2*$^{-/-}$) larvae. RNAseq data comparing reads mapped to *tlr2*

transcript (ENSDART0000012256). Heterozygotes and mutant reads are mapped to the entire

length of the mutant transcript indicating that the mutant transcript is not subjected to

nonsense mediated decay. Mutant data (MEAS) and heterozygotes (CTRL) data have been

submitted to the NCBI gene expression Omnibus database, accession number is GSE102766.

Supplementary figure 3 Analysis of differential expression of genes functioning in

glycolysis and gluconeogenesis between uninfected *tlr2*$^{+/-}$ and *tlr2*$^{-/-}$. The red boxes

represent up regulated genes (FC >2); blue boxes represent down regulated genes (FC<-2);

yellow boxes represent the genes that are differentially expressed with a *P* value lower than

0.05; green represent not-significantly differentially expressed genes.

Supplementary Figure 4 Volcano plots. Volcano plots showing the significance cutoff

applied to *tlr2*$^{+/-}$ infected with strain Mma20 versus control with PBS (a) and *tlr2*$^{-/-}$ infected

with strain Mma20 versus control with PBS (b). In these volcano plots, the transcripts were

considered significant (red) or non-significant (blue) by the conditions of |fold change| ≥

□1,45 and adjusted *P* value threshold□≤□0,05. Fold changes for each transcript was plotted

on the X-axis against -log10 transformed p-values on the Y-axis.

Supplementary figure 5 *tlr2*-dependent up regulation of genes with various GO terms.

Supplementary figure 6 *tlr2*-dependent down regulation of genes with various GO terms.

Supplementary figure 7 Sub-network enrichment analysis. Networks of common targets of

the 97 up regulated genes (Fig. 5d) in *tlr2*$^{+/-}$ (a) and *tlr2*$^{-/-}$ (b) with Mma20 infection. Red

represents up regulation, blue represents down regulation and grey represents genes for which

no expression was detected.

Supplementary figure 8 Sub-network enrichment analysis. The networks of common targets

of the 92 down regulated genes (Fig. 5e) in *tlr2*$^{+/-}$ (a) and *tlr2*$^{-/-}$ (b) with Mma20 infection.

Red represents up-regulation, blue represents down-regulation and grey represents genes for

which no expression was detected.

Supplementary figure 9 Sub-network enrichment analysis between zebrafish and human.

The Tlr8 pathway in zebrafish (a) with Mm infection and human macrophages (b) with Mtb

infection. Red represents up-regulation, blue represents down-regulation.

Supplementary   figure   10 Manual   counting   analysis   for   *Tg(mpeg1:mCherry-*

F);TgBAC(mpx:EGFP) of neutrophils (a) and macrophages (b) in 2 dpf *tlr2*$^{+/+}$ and *tlr2*$^{-/-}$

embryos.

Supplementary Table 1 Reads of *mpeg1.1* and *mpx* genes of the *tlr2*$^{+/-}$ control and the *tlr2*$^{-/-}$

mutant in RNAseq analysis.

Supplementary Table 2 GO analysis of genes that have a different basal expression level in

the absence of infection in the Tlr2 mutant versus the heterozygote control. Shown is the GO

analysis of the group of 878 genes that were different with a FDR *P* value of 10^{-10} indicated in

Fig. 2.

Supplementary Table 3 Two categories of genes including the c-Maf transcription factor

were significantly affected in the GO category

Supplementary Table 4 A-H Gene lists of GSEA and SNEA analysis

Supplementary Table 5 List of primers

a

Tlr2-201 ENSDART 00000122568.2

LRR domain — LRR domain — TIR domain

541   L   R   M   F   N   Q   S   S   L   R   S   F   K   Wt Tlr2

1621 TTA CGG ATG TTC AAC CAA AGC TCT TTG AGG AGT TTT AAA 1659

1621 TTA CGG ATG TTC AAC CAA AGC TCT TAG AGG AGT TTT AAA 1659

541   L   R   M   F   N   Q   S   S   STOP   Mut Tlr2

LRR domain — LRR domain

b *il1b*

RNA relative fold-change

****
****
****

$tlr2^{+/+}$ $tlr2^{+/-}$ $tlr2^{-/-}$

Pam3CSK4

c *fosl1a*

RNA relative fold-change

****
***
*

$tlr2^{+/+}$ $tlr2^{+/-}$ $tlr2^{-/-}$

Pam3CSK4

d *cebpb*

RNA relative fold-change

****
****
***

$tlr2^{+/+}$ $tlr2^{+/-}$ $tlr2^{-/-}$

Pam3CSK4

e *il1b*

RNA relative fold-change

ns
ns
ns

$tlr2^{+/+}$ $tlr2^{+/-}$ $tlr2^{-/-}$

Flagellin

f *fosl1a*

RNA relative fold-change

ns
ns
ns

$tlr2^{+/+}$ $tlr2^{+/-}$ $tlr2^{-/-}$

Flagellin

g *cebpb*

RNA relative fold-change

ns
ns
ns

$tlr2^{+/+}$ $tlr2^{+/-}$ $tlr2^{-/-}$

Flagellin

h

Macrophages $tlr2^{+/+}$ *Neutrophils* $tlr2^{+/+}$ *Merged* $tlr2^{+/+}$

100 μm

i

Macrophages $tlr2^{-/-}$ *Neutrophils* $tlr2^{-/-}$ *Merged* $tlr2^{-/-}$

j

Number of macrophages in tail region

ns

$tlr2^{+/+}$ $tlr2^{-/-}$

2 dpf

k

Number of neutrophils in tail region

ns

$tlr2^{+/+}$ $tlr2^{-/-}$

2 dpf