Inositol pyrophosphate kinase (VIH2) impart drought resistance by promoting

plant cell wall homeostasis

Author names and affiliations: Anuj Shukla[1], Mandeep Kaur[1,2], Swati Kanwar[1], Gazaldeep

Kaur[1], Shivani Sharma[1], Koushik Mazumder[1], Pratima Pandey[3], Vikas Rishi[1] and Ajay K

Pandey[1,*]

[1]National Agri-Food Biotechnology Institute (Department of Biotechnology), Sector 81,

Knowledge City, S.A.S. Nagar, Mohali-140306, Punjab, India.

[2]Department of Biotechnology, Panjab University, Punjab, India.

[3]Department of Biological Sciences, Indian Institute of Education and Research, Mohali

140306.

Corresponding author:

Dr. Ajay K Pandey, Scientist-E

Email: pandeyak@nabi.res.in; pandeyak1974@gmail.com

ORCID iD: https://orcid.org/0000-0003-1064-139X

Abstract

Inositol pyrophosphates (PPx-InsPs) are important signalling molecules, those participate in multiple physiological processes across wide range of species. However, limited knowledge is available for their role in plants. Here, we characterized two diphosphoinositol pentakisphosphate kinase (PPIP5K) wheat homologs, *TaVIH1* and *TaVIH2* for their spatio-temporal expression and physiological functions. We demonstrated the presence of functional VIH-kinase domains through biochemical assays where high energy pyrophosphate forms ($IP_{7/8}$) were generated. Our GUS-reporter assays in *Arabidopsis*, suggested the role of *TaVIH2* in drought stress. Yeast two-hybrid screen of TaVIH2 by utilizing wheat library yielded multiple cell-wall related interacting partners. *TaVIH2* overexpression in Arabidopsis provided growth advantage and drought tolerance. Further, transcriptomic studies of these overexpressing lines showed activation of genes encoding for abscisic acid metabolism, cell-wall biosynthesis and drought responsive element binding proteins. Biochemical analysis of their cell-wall components, confirmed enhanced accumulation of polysaccharides (arabinogalactan, cellulose and arabinoxylan) in transgenics. These results reveal novel function of VIH proteins in modulating cell wall homeostasis thereby providing drought tolerance.

Keywords: Inositol pyrophosphate kinase, wheat, drought stress, phytic acid, transcriptome, cell wall

Introduction

Inositol pyrophosphates (InsPPs) are an important member of the inositol phosphate family that have emerged as distinct molecules possessing array of phosphates around an inositol ring [1]. The role of high energy InsPPs including InsP7 and InsP8 has been studied in human cell lines and yeast. They were shown to participate in DNA recombination, vacuolar morphology, cell wall integrity, gene expression, pseudohyphal growth and phosphate homeostasis [2–7]. These InsPPs are predominantly synthesized by two different classes of enzymes that exhibit catalytic activity towards different positions on the six-C inositol ring. The first class of enzymes, Inositol hexakisphosphate kinases (IP6Ks) place a phosphate group at 5^{th} position of the fully phosphorylated ring of InsP6/IP6 to form 5PP-IP5 ie IP7 [8,9] and also generates two isomers of IP8 including; 1or 3,5PP-IP5 and 5PPP-IP5 [10]. The second class of enzymes referred as diphosphoinositol pentakisphosphate kinases (PP-IP5Ks) phosphorylate the 1^{st} position of IP6 synthesizing 1PP-IP5 [11–13]. These enzymes also catalyse the conversion of 5PP-IP5 to 1,5(PP)2IP4 i.e. IP8 which was first demonstrated in mammalian cells [1]. During the past two decades three isoforms of IP6K (IP6K1, IP6K2 and IP6K3) and two PP-IP5K (PP-IP5K1 and PP-IP5K2) were identified in mammals [1]. In yeast, a single IP6K (also referred as Kcs1) and PP-IP5K (also known as VIP1/IP7K) are present that are involved in synthesis of the respective forms of IP7 and IP8 [12,13]. VIPs are also referred as VIH in plants and are dual domain containing proteins; including a "rim-K" or ATP-grasp superfamily domain at the N-terminal and a C-terminal histidine acid-phosphatase domain [13].

In plant seeds, the most abundant inositol polyphosphate referred as, Inositol hexa*kis*phosphate (Phytic acid,IP 6) is the primary source of stored phosphorus (P) which is utilized by the plants to draw energy during the process of seed germination. Earlier, the presence of high anionic form of IP6 was speculated in plant species such as barley and potato [14,15]. The quest to identify the plant genes encoding for these inositol pyrophosphate kinases remained elusive till the identification of two plant genes referred as VIP/VIH from *Arabidopsis* [16]. These VIH proteins were characterized to bear PPIP5K activity and shown to be involved in plant defense response that is mediated through jasmonate levels [16,17]. These identified VIH show functional activity in yeast mutants with their ability to rescue invasive growth with hyphae formation.

Recent evidence also implicates that IP8 binds the SPX region of the SPX1 protein and control its interaction with a phosphate starvation response1, a central regulator of phosphate starvation [18]. Furthermore, in yeast, role of inositol pyrophosphate kinase was also implicated in vacuolar morphology and cell wall integrity [3]. Histone H3/H4 chaperone, Asf1p is known

to interact with the VIP1 (VIH) protein, thereby, suggesting this interaction important for
transcription elongation [19]. Therefore, the function of the protein could also be studied in
context to their interacting partners. Such comprehensive interaction studies for VIH protein is
also missing in plants that could provide new insight into their functional roles.

In the current study, we identified two functionally active wheat VIH genes capable
of utilizing IP_6 and IP_7 as substrates under In-vitro condition to generate higher InsPP.
Promoter fused GUS-reporter assays during different stress condition revealed specific
response of VIH2-3B promoter during drought condition. We also demonstrated that, at the
protein level wheat VIH2-3B interact with Fasciclin-like arabinogalactan protein (FLA6) and
other multiple cell-wall related proteins. Furthermore, we concluded that wheat homoeolog
VIH2-3B could impart tolerance to drought in transgenic *Arabidopsis* by altering the
composition of plant cell-wall and regulating distinct transcriptomic re-arrangements. Taken
together, our study provides novel insight for the possible function of plant VIH protein in
drought stress tolerance.

Results

Phylogeny and spatial-temporal expression characterization of VIH genes in wheat tissue
Our efforts to identify potential wheat VIH-like sequences resulted in the identification of six
genes with three homoeolog for each, *TaVIH1* and *TaVIH2* those were mapped to chromosome
3 and 4 respectively. The Kyte-Doolittle hydropathy plots indicated that wheat VIH proteins
were devoid of any transmembrane regions (Supplementary Figure S1A). The analysis
clustered plant VIH homologs together with TaVIH proteins close to their *Brachypodium*
counterparts in the monocot specific clade (Figure 1A). Amino acid sequence alignment of
wheat VIH protein sequences suggested the presence of conserved dual domain architecture
with two distinct domains consisting of amino-terminal rimK/ATP GRASP fold and a histidine
acid-phosphatase (HAP) of PPIP5K/VIP1 family (Supplementary Figure 1B). Wheat VIH1 and
VIH2 show 72% sequence identity at the protein level. Among the two wheat VIH proteins,
TaVIH1 show high identity of 78% with AtVIH proteins whereas, TaVIH2 show 70.0 and 70.6
% identity to both AtVIH1 and AtVIH2 respectively.

Transcript accumulation of *TaVIH* genes showed similar expression profiles for both
genes with highest expression in leaf tissue followed by flag leaf, root and with the least
expression in stem (Figure 1B). The transcript accumulation of *TaVIH1* was 1.5-fold higher
than *TaVIH2* in all the tissues investigated. These findings suggest that both VIH genes are

preferentially expressed in leaf. The highest expression of both VIH genes was observed at late

stages of grain filling with high transcript accumulation at 28 DAA stage (Figure 1C).

Interestingly, the pattern of gene expression remained same for both wheat VIH, wherein

TaVIH1 showed 3-fold higher expression in comparison to *TaVIH2* at all stages. The

expression profile in different grain tissues revealed a high expression of *TaVIH* genes in the

aleurone layer which is ~ 4-fold higher than in the endosperm. Similar levels of transcript

accumulation in the remaining grain tissues *viz.* embryo, glumes, and rachis; suggesting a

ubiquitous expression in these tissues (Figure 1C).

In general, expression levels of *TaVIH1* were more than *TaVIH2* during all stages of

wheat development (Figure 1D). Overall, our expression analysis using ExpVIP [20] database

showed high expression of *TaVIH1* transcripts from B and D genomes. *TaVIH2* did not show

any significant expression in majority of tissues, however a weak expression was found in the

spikes (Supplementary Figure S2A). Our analysis shows differential expression patterns of

VIH during abiotic and biotic stresses suggesting their role in different development processes

and stress response (Supplementary Figure S2B).

Wheat inositol pyrophosphate kinase demonstrate PPIP5K activity

The homology modelled structure of VIH proteins, based on human PPIP5K2 (hPPIP5K2; PDB

ID:3T9C) predicted two antiparallel β-sheets to co-ordinate the nucleotide analog AMP-PNP

(Supplementary Figure 3A). Likewise, human and *Arabidopsis* homologs, wheat VIH proteins

also appear to use only arginine and lysine residues in the inositol phosphate binding pocket,

except for a serine residue (Supplementary Figure 3B). Next, yeast complementation assay for

wheat VIH genes was carried out using growth assay on SD-Ura plates supplemented with 0,

2.5 and 5 mM 6-azuracil. Both the yeast strains transformed with wheat VIH genes were

confirmed for their protein expression by using Western analysis (Fig. The wild type strain

BY4741 showed an unrestricted growth phenotype, whereas *vip1Δ* transformed with empty

pYES2 vector showed growth sensitivity at 2.5 and 5mM concentrations (Supplementary

Figure 3C). To our surprise, the mutant strain transformed with pYES2-TaVIH1 could not

revive growth defect on selection plates whereas, the pYES2-TaVIH2-3B was able to rescue

the growth phenotype of *vip1Δ* strain. Under stress condition unlike wild type yeast, *vip1Δ*

mutant do not form pseudo-hyphae. The complemented *vip1Δ* strain with pYES2-TaVIH2-

3B rescued this phenotype during stress (Supplementary Figure 3D). Overall, our data

suggest that *TaVIH2* derived from the B genome is capable of complementing the growth

defects of *vip1Δ* strain.

In-vitro kinase assay was performed with the pure wheat VIH-kinase (KD) proteins
(Supplementary Figure S4) to check its activity. The relative luminescence units (RLU) were
recorded for TaVIH1-KD and TaVIH2-KD while using IP_6 as a substrate (Figure 2A). Our
assays show a very high increase in the RLU for both the proteins in the presence of IP_6.
Interestingly, the wheat VIH2 show ~15-fold higher luminescence response, suggesting its high
activity when compared to the VIH1 protein (Figure 3C). This kinase activity was diminished
in VIH (D-VIH) post denaturation, and the activity was not significantly different when
compared to either Enzyme control (Ec) or substrate control (IP_6) reactions.

To check the substrate utilizing ability of wheat VIH proteins, reactions were performed
using IP_6 and IP_7 that phosphorylated inositol molecule species which were visualized on PAGE
as reported earlier [21]. Our assays suggested that wheat VIH proteins could utilize IP_6 as a substrate
to generate IP_7 (Figure 2B; left panel; lane 3, 4), as was also found in the case of ScVIP protein
(Figure 2B; left panel; lane 1). Next, we tested if IP_7 (generated by ScVIP) could be utilized by
wheat VIH proteins. We observed that both the wheat VIH proteins (TaVIH1 and TaVIH2) could
utilize IP_7 to generate IP_8 (Figure 2B; right panel, lane 1, 3). Moreover, ScVIP was also able to
utilize IP7 generated by TaVIH1 and TaVIH2 (Figure 2B; right panel, lane 2,4). These
experiments conclude that TaVIH proteins are functionally active, capable of using both IP_6 and
IP_7 as a substrate under In-vitro conditions and may possess $PPIP_5K$ like activity. To further
confirm the nature of the phosphorylated inositol molecule during In-vitro reactions, MALDI-
Tof- mass spectroscopy (MS) was performed. MS analysis indicated a major signal at m/z of
739.009 that correspond to the mass of IP_7 for all the generated species (Figure 2C; left panel).
Similarly, the VIH1 generated IP_8 showed phosphorylated inositol molecule with m/z 820.1 and
additional expected acetonitrile adduct at 860.9 (Figure 2C; right panel). Whereas VIH2 derived
IP_8 showed m/z at 860.9 corresponding to the IP_8-acetonitrile adduct (Supplementary Figure S5).
This confirms formation of the IP_7 and IP_8 as phosphorylated forms of inositol molecules.
Overall, our analysis confirms the functionality of VIH proteins and their ability to generate both
IP_7 and IP_8.

Expression of *35S:TaVIH2* transgenic *Arabidopsis* display robust growth
To characterize the biological functions of TaVIH2, cDNA was overexpressed in *Arabidopsis*.
In total, seven transgenic lines were pre-selected based on TaVIH2 expression as shown by
western analysis and four transgenic lines (#Line2, #Line 4, #Line 5 and #Line 6) were selected
for further characterization (Figure 3B). We observed that, at the vegetative stage TaVIH2
transgenic *Arabidopsis* show robust growth. Plants (14 days old seedlings) showed enhanced

rosette area cover and increased number of leaves compared to controls (Col-0 and Col-0-Ev
(empty vector)) respectively (Figure 3B, C and D). These transgenic *Arabidopsis* also show
enhanced branching with an overall increase in the length of the main shoot axis and leaf size
compared to controls (Figure 4A). Primary and secondary shoots numbers were also enhanced
in the transgenic *Arabidopsis* (Figure 4B). In general, no significant differences during the
flowering stage were observed, yet the increased number of (20-24) of secondary shoots were
evident when compared to control plants (12-15 shoots). These data suggest that expression of
TaVIH2 in *Arabidopsis* impacts the overall growth of the plant.

VIH2-3B imparts resistance to drought stress

To investigate the promoter activities of *TaVIH1* and *TaVIH2*, 5' flanking regions (~ 1.5 Kb)
of these genes were cloned and the comparative analysis revealed the presence of various
hormones and abiotic stress responsive cis elements (Supplementary Figure S6A). The
presence of these elements suggested that wheat *VIH* could be regulated by stress. Interestingly,
the promoter of *VIH* contains multiple drought/dehydration responsive domains, a P1BS motif
(GNATATNC) and PHO4 binding sites in promoter regions of both *TaVIH* genes
(Supplementary Figure S6A). This motivated us to perform a preliminary screening
experiments using *TaVIH*-promoter fused to β-glucuronidase (GUS)-reporter gene
(pVIH1/2:GUS) in *Arabidopsis* (Col-0). Significant increase in GUS reporter activity of
pVIH2:GUS lines indicate the ability of this promoter to sense the given stress and drive GUS
reporter expression. To our surprise, *TaVIH2* promoter responded strongly to dehydration,
drought stress and Pi-starvation (Supplementary Figure S6B and Figure 5A). A strong GUS
expression was also visualized along the mid-vein and leaf base of the seedlings subjected to
Pi-stress. A weak expression of *VIH2* promoter was observed in the presence of ABA and GA$_3$.
Control (Empty-vector-EV) seedlings showed no visible GUS-staining. Based on our reporter
assays, we speculate that TaVIH2 participates during drought stress response that was further
investigated.

To further test the effect of drought like conditions, seedlings were exposed to drought-
like conditions using mannitol (125 mM) and glycerol (10 %)[21]. Our data indicate the
retardation of root growth in control *Arabidopsis* suggesting that the plants were sensitive to
the presence of these metabolites. In contrast, TaVIH2 overexpression in *Arabidopsis* was able
to escape the detrimental root growth (Figure 5B). This observation was highly significant
among the transgenic lines suggesting that TaVIH2 could contribute to drought stress (Figure

5C). Next, rate of water loss was studied to account for its possible role in withstanding drought

stress. The rate of water loss was very significant in the control plants as compared to the

transgenic (Figure 5D). The water loss percentage was lowest in the transgenic *Arabidopsis*

(40-46%) when compared to control plants (16-18%) after 8 hrs of incubation (Figure 5D). To

further, confirm its contribution towards drought tolerance, the transgenic *Arabidopsis* and

control plants were subjected to 14 days of water withholding (drought). This caused a dramatic

withering of both control and transgenic *Arabidopsis* plants. However, when the plants were

re-watered, high survival rates of ~50-65 % was observed in the transgenic plants, whereas no

or very low (3%) rates were observed in control (Figure 5E). This indicates that the transgenic

Arabidopsis overexpressing TaVIH2, escapes the effect of drought and improves survival rate

by imparting drought tolerance.

Wheat VIH2 interacts with Fasciclin-like arabinogalactan protein (FLA)

To investigate the mechanism of TaVIH2 proteins in stress tolerance, we used yeast two-hybrid

(Y2H) cDNA screening for its interacting partners. Protein expression of bait (VIH2) in the

yeast cells was confirmed by Western blot analysis (Supplementary Figure S7A). Two pooled

wheat cDNA libraries were prepared that resulted in the identification of 89 putative yeast

colonies with a mating efficiency of 3.8 % (Supplementary Figure S7B and C). Subsequent

stringent screening (α-Gal) of the colonies lead to the identification of ~52 putative interactors

(Supplementary Figure S7D). Upon sequencing of the ORFs, clones appearing more than twice

were considered for further studies. Careful analysis lead to the shortlisting of eleven strong

potential (3-6 clones) interactors with high occurrence of the clones those were related to

proteins involved in cell-wall related function (Table 1). Amongst these interactions most of

the genes encode for cell-wall related function, including Fasciclin-like arabinogalactan protein

(FLA), glycosyl-transferases and glycine-rich structural proteins. The most frequently (six

times) interacting clone was identified as FLA protein and its detailed analysis was done. To

further confirm this interaction, full-length cloning of *TaFLA6* cDNA (1.2 kb) was performed

and transformed (TaFLA6:AD+TaVIH2:BD) into yeast. Growth of yeast colonies on -His/-

Leu/-Trp (Auerobasidin) media (Figure 6A) and the In-vivo pull-down assay with co-expressed

VIH2 (cMYC-tagged) and FLA6 fusion protein (HA-tagged) in yeast confirmed the full-length

interaction of these two proteins (Figure 6B).

The FLA proteins are known to respond for different stress conditions and are to be

involved in plant growth and development [22,23]. These observations suggest that VIH2 and its

interacting partners may participate in similar pathway. Our localization studies in yeast suggest that FLA6 was present on the yeast plasma membrane (Figure 6C). Similarly, localization of VIH2 suggest their presence in the cytoplasm and the rim of the plasma membrane. Wheat FLA6 encodes a 367 aa protein containing FAS-like arabinogalactan protein with presence of typical trans-membrane domain (TMD) and glycophosphatidyl-inositol (GPI) domain [24,25]. The protein hydropathy plot identified a hydrophobic region near GPI region at the C-termini (Figure 6D). Transcript expression study of interacting partners (*TaFLA6, TaGT* and *TaXat1),* showed significant upregulation in the shoot tissue when subjected to drought stress (Supplementary Figure S8). Similar observation for *TaFLA6, TaGT* and *TaXat1* was also obtained through exVIP gene expression analysis (Supplementary Figure S8). Significant increase in transcript accumulation was observed during grain development and seed maturation (Supplementary Figure S8). These data indicated that *TaFLA6* and *TaGT/Xat* show transcriptional changes during desiccation, a prerequisite step in grain maturation. Important interacting clones, including *TaGT* and *TaXat* were also expressed in other tissue including developing roots and grains (Supplementary Figure S8). Taken together, our data suggest that wheat *VIH2* and *FLA6* are co-expressed under drought stress and they interact at the protein level.

Transcriptomes suggest that *VIH2-3B* stimulate genes related to drought stress

We next explored the reasoning for robust phenotype and resistance to drought in transgenic *Arabidopsis*. For this, transcriptomic changes in 25 days old seedlings of control and two transgenic plants (#Line4 and #Line6) were compared. PCA of normalized expression abundances revealed a high level of correlation among biological replicates (n=3) in each transgenic line. PCA also indicates distinct cluster for overexpressing transgenic lines and controls (Supplementary Figure S9A). Based on analysis involving respective three biological replicates, a total of 626 and 261 genes were significantly up- and down-regulated (-1>Log FC >1.0) in #Line4 while 797 and 273 genes were up- and downregulated in #Line6 transgenic *Arabidopsis* lines compared to control plants (Supplementary Table S3). Overall, 605 genes were commonly differentially altered in the two transgenic lines with respect to control plants (Col-0(Ev)).

Multiple genes were commonly up-regulated in transgenic *Arabidopsis* compared to control plants (Figure 7A and Supplementary Table S3). Interestingly, high number of genes are constitutively activated in the transgenic *Arabidopsis* belongs to the dehydration response element binding (DREB) protein including Integrase-type DNA-binding superfamily proteins

and glycine rich proteins. Upon analysis of the GO terms, the highest number of genes for "stress related" and "cell-wall related activities" were enriched in the biological process and cellular component categories (Figure 7B). Strikingly, multiple genes involved in cell-wall biosynthesis, modification and degradation were also up-regulated in the transgenic plants (Figure 8A). In addition to that, distinct cluster of genes involved in Abscisic acid (ABA) biosynthesis were also significantly up-regulated among the different lines of transgenic *Arabidopsis* (Figure 8B). Notably, genes encoding for 9-*cis*-epoxycarotenoid dioxygenase (*AtNCED6* and *AtNCED9*) involved in ABA biosynthesis were also up-regulated. Multiple DREB encoding genes and cytochrome P450 (CYPs) related family genes (*CYP71A23, CYP94B3, CYP71B12, CYP96A2, CYP702A1, CYP707A3, CYP82C2, CYP76G1, CYP705A4, CYP71B10, CYP706A2, CYP81D11*) were also differentially regulated in the transgenic *Arabidopsis* (Figure 8C and D) Overall, we conclude that distinct cluster of genes potentially involved in drought and ABA stress were significantly up-regulated in these transgenic plants.

VIH2 affects ABA levels and regulates plant cell-wall composition

Multiple genes related to ABA biosynthesis were differentially expressed in VIH2 overexpressing *Arabidopsis*. To verify if the de-novo gene expression response to the ABA related genes could be correlated with the In-vivo levels, ABA levels were quantified in the leaves. We observed that accumulation of ABA was significantly higher (~3-4 fold) in transgenic *Arabidopsis* when compared to the control plants (Figure 9A). This average increase of ABA in all the four transgenic lines was statistically significant ($p < 0.05$, Student's t test). Our data confirmed the involvement of ABA in the drought tolerance of transgenic lines.

To further draw the commonality between the gene expression pattern in VIH2 overexpressing *Arabidopsis* and due to drought, we analysed previously reported RNAseq data SRA:SRP075287 (under drought stress) for overlap of de-regulated genes. In total, 295 and 309 genes were commonly regulated in #Line4 and #Line6 (Figure 9B). Most of listed genes those were commonly regulated belong to the category of hormone metabolism, signalling, stress response, development and cell wall related functions (Figure 9C). Multiple genes related to CYPs and glycosyl transferases were highly enriched in the dataset (Table S5). These extended analysis supports the notion that VIH2-3B could impart activation of genes pertaining to drought. Overall, our molecular analysis helped in identifying sub-set molecular of components in transgenic plants that could impart basal drought resistance.

Multiple interacting partners of the VIH2 were identified with their possible role in the biosynthesis of cell-wall structural proteins or in membrane plasticity. Therefore, we

speculated that VIH2 protein could modulate the cell-wall composition. To address this, we measured the different cell-wall components of control and transgenic *Arabidopsis* using standard methods that resulted in comparable yields and without starch interference. Our analysis indicated a consistent increase in the accumulation of cellulose (from 1.3 to 2.5-fold) in the transgenic lines that was consistent among the biological replicates and multiple transgenic lines. Additionally, arabinoxylan (AX) and arabinogalactan (AG) was also increased (1.8- 2.2 and 1.47- 1.5-fold) in the transgenic lines as compared to the controls (Supplementary Table S4). Our extraction procedures for control plants show the ratio of 1::1.2 to 1.5 for arabinose/galactose and arabinose/xylans, this validates our extraction procedures. To further validate the role of VIH proteins, *Atvih2-3* mutant line was used for measuring the biochemical composition of the shoots cell wall. Our analysis showed a significant reduction of the AG, AX and cellulose content in this mutant line (Figure 9D). Altogether, our data demonstrate that overexpression of wheat VIH resulted in the compositional change in the cell-wall biosynthesis-related sugars and these changes could be linked to the enhanced drought response in leaves.

Discussion

Recently, studies investigating inositol pyrophosphates have gained much attention due to the presence of high energy pyrophosphate moieties speculated to regulate metabolic homeostasis in organisms [16,17,26–28]. This study was performed to characterize and identify the functional mechanism of VIH proteins involved in the biosynthesis of PP-InsPx. We have identified and characterized two wheat inositol pyrophosphate kinase (TaVIH1 and Ta VIH2) encoding genes and demonstrated that homoeolog *TaVIH2-3B* interacts with cell wall related proteins. Overexpression of TaVIH2 in *Arabidopsis* could enhance growth and provide tolerance to drought stress by modulating cell-wall and ABA related proteins through altered cell-wall polysaccharide composition (AG, AX and cellulose).

Hexaploid bread wheat has one of the most complex genomes comprising of three related sub-genomes that have originated from three separate diploid ancestors thus forming an allohexaploid genome [29,30]. Therefore, to consider the appropriate homoeolog-transcript for further studies, Wheat-Exp expression database was used to analyse VIH2 homoeolog expression in different tissues and also during the developmental time course (Figure S2). VIH2 is known to be involved in defense response via a jasmonate-dependent resistance in Arabidopsis [17]. Wheat VIH genes were also induced upon infection of plants with pathogens (Supplementary Figure S2). Thus, the role of plant VIH genes during plant-microbe interaction

was found to be conserved. TaVIH protein was authentic kinase protein since its kinase domain

could catalyse the phosphorylation and harbors yeast VIP1-like activity as demonstrated by its

utilization of both IP_6 and IP_7 as potential substrates. In the past, *AtVIH* genes have been shown

to be biochemically active for kinase activity that generates $InsP_7$ and $InsP_8$ [16,17]. Similarly,

yeast and human enzymes also show IP_6 and IP_7 kinase activity [13,31,32].We propose that

additional studies needs to be performed in future to confirm the chemical-forms of IP_7 and IP_8

generated by TaVIH2 using Nuclear Magnetic Resonance spectroscopy. TaVIH2-3B showed

the highest homology to AtVIH2 (70.6 %).

The presence of various *cis*-acting elements in the promoter region plays essential

roles in transcriptional regulation of genes in response to multiple environmental factors. The

transcriptional activity of *TaVIH2* promoter and differential expression analysis link TaVIH2

with Pi-starvation response (Figure S2). This function of inositol pyrophosphate kinases in the

regulation of Pi homeostasis seems to be evolutionarily conserved [28,32]. Very recently in

Arabidopsis, it was demonstrated that VIH derived $InsP_8$ is required to sense the cellular Pi

status and also binds to the intracellular Pi sensor SPX1 to control Pi homeostasis in plants [33].

We found that in addition to Pi homeostasis, *TaVIH2* promoter also responds to drought

conditions. Y2H study led to the identification of TaFLA6 and multiple cell-wall

reinforcement proteins as potential interactions of plant VIH proteins. Our qRT-PCR

specifically indicates that both *TaVIH2* and *FLA6* are co-expressed under drought condition,

suggesting that they are involved in the similar post-transcriptional response pathway

(Supplementary Figure S8). Previously, it was shown that FLA proteins were involved in cell-

wall reinforcement, plasticity, cell to cell adhesion and drought tolerance [22,34–37]. Cereal grains

such as wheat are also rich in arabinogalactans such as FLA[38] indicating that associated VIH

proteins might be responsible for these physiological responses including late stages of grain

maturation (Figure 1C).

The high numbers of reoccurring clones related to the cell wall biosynthesis suggest an

important role that VIH proteins may offer during developmental stages. Previously, multiple

Arabidopsis FLA proteins were reported to be perturbed under drought conditions [39,40]. FLA-

like protein was shown to be involved in molecular responses of Pi deficiency that is mediated

by the induced root hair elongation [41]. The domain analysis of FLA6 suggests it belonged to

the category-IV of FLAs and show presence of all the necessary domains, typical of this gene

family (Figure 6D). It is important to notice that additional VIH2 interacting clones encode for

glycosyltransferases, xylan-arabinosyl transferase and glycine-rich cell-wall structural like-

protein. Surprisingly, genes encoding for glycosyltransferases, xylan-arabinosyl transferases

are homoeologs (*TaGT* and *TaXat*) and belong to the family of transferases. The glycosyl and

xylan arabinosyl transferases are involved in the biosynthesis of polysaccharides for cell-wall

[42,43]. Similarly, glycine-rich cell-wall proteins are recognized for their role in cell-wall

reinforcement by callose deposition [44].

Protein-protein interaction studies could provide associated functional clues. Yeast

VIP1 was shown to interact with histone H3/H4 chaperone, ASF1. VIP1 and ASF1

counterparts in *S. pombe* functionally regulate actin-related protein-2/3 complexes and thereby

participate in the fate of cell morphology [19]. Protein-protein interaction studies using human

PPIP5K1 identified multiple proteins involved in vesicle-mediated trafficking, lipid

metabolism and cytoskeletal organization [45]. This protein interaction has been accorded to

the presence of long C-terminal intrinsic disorder region (IDR) of PPIP5K1. Here, in case of

wheat VIH2 18.59 % of the predicted disorder content was observed that reflect the presence

of IDR boundaries[46]. Presence of such IDR in VIH2 could support interaction with cell-wall

scaffolding proteins, akin to the interaction ability of human PPIP5K1 [45,47].

Earlier the double mutants of VIH genes in *Arabidopsis* show severe growth defects,

implicating their unexplored role in overall growth and development[18]. We hypothesize that

the molecular and biochemical changes in transgenic *Arabidopsis* provide the overall

mechanical strength to the plant cell and in turn tolerance to stress condition. These

observations were also supported by our transcriptome analysis of two independent TaVIH2

overexpressing *Arabidopsis* lines that show consistent high expression of cell-wall, ABA and

drought related genes (Figure 8 and Figure 9B). Multiple genes were differentially regulated

by full length TaVIH2 overexpression. This suggest that high protein levels of VIH2 could

cause changes in gene expression pattern. Classically, VIH proteins contains evolutionarily

conserved two distinct domains including a N terminal rimK/ATP GRASP kinase and

phosphatase domain. It remains to be dissected if the change in transcriptome response in these

transgenic *Arabidopsis* is due to the kinase or phosphatase domain. Earlier, multiple inositol-

1,3,4 tris*kis*phosphate 5/6-kinase (devoid of phosphatase domain) were also implicated for their

role in drought tolerance [48,49]. This may suggest that the tolerance for the drought could arise

by the presence of functional kinase domain.

Multiple studies have implicated that enhanced level of ABA leads to drought tolerance

[50]. The elevated levels of ABA in our transgenic plants could be the reason for the high

expression of genes for cell wall maintenance and biosynthesis. Cell wall related remodelling

and ABA regulated signalling are the primary response against abiotic stress including drought

[51–53]. ABA dependent increased expression of *CYP*s and *DREBP* have been reported earlier in

plants with their role implicated in drought stress [50,54,55]. Our study shows high basal expression
of genes encoding for DREBP and CYPs. The constitutive high expression of these gene
families in our transgenic *Arabidopsis* could account for their better adaptability for drought
stress. Earlier, changes in cellular levels of $InsP_7$ and $InsP_8$ have been attributed to guard cell
signalling, ABA sensitivity and resistance to drought in maize *mrp5* mutants [28,56]. This suggests
a molecular link between VIH, ABA levels and drought resistance.

Atvih2-3 mutant lines lacking mRNA expression also show alteration in the cell wall
composition despite its typical growth as wild type Col-0 (Figure 9D). Interestingly, *vih1* and
vih2 double mutants display severe growth defect that was rescued by the gene
complementation [18]. Our overexpression data showing enhanced branching and robust growth
collectively reinforce the notion that VIH are also involved in providing support for plant
growth. The *vih2* mutant in *Arabidopsis* are more susceptible to infestation by caterpillar
(*Pieris rapae*) and thrips [17]. The resistance against herbivore pathogens such as *P. rapae*, could
be gained by modulating the genes associated with cell-wall modification [57]. *Arabidopsis VIH2*
mutant lines showed compositional changes in the cell-wall extracted polysaccharides
especially in the AG level. The decreased resistance in *vih2* mutants against herbivores could
be accounted for the defect in signalling pathway via COI1-dependent gene regulation and
changes in the structural composition of the cell-wall.

Taken together we propose a working model, where wheat VIH participate in the
drought resistance in plants by modulating the changes in cell-wall gene expression, enhanced
ABA levels and change in biochemical composition to provide more mechanical strength
(Figure 10). In future, it will be interesting to quantitate the level of higher inositol
pyrophosphates in these plants. In summary, our work along with the previous functional
reports suggested an emerging novel role of plant VIH proteins in cell wall scaffolding
functions to provide resistance against drought stress.

Methods
Plant materials and growth conditions
The experimentation in this study was conducted using Bread wheat (*Triticum aestivum* L.)
variety C306, a rain-fed cultivar which is well known for its better processing quality. For
collection of the tissue materials, the spikes were tagged at the first day after anthesis (DAA).
Samples were collected in the form of spikes at 7, 14, 21 and 28 DAA stages and various
tissues, including root, stem, leaf and flag leaf at 14 DAA stage respectively. To further dissect

the expression levels in spikelet's, 14 DAA seed was used to separate different tissues,
including aleurone, endosperm, embryo, glumes and rachis as done previously [58].

All the experiments for the stress conditions were performed in three biological
replicates. Wheat seeds were surface sterilized as described earlier [58] and were allowed to
germinate on Whatman filter paper soaked with water for 3-5 days. The germinated seedlings
(8-10) with their residual endosperm excised, were transferred to Hoagland's nutrient media in
phytaboxes. For phosphate starvation (-Pi) experiment, seven days old seedlings were
subjected to Pi-sufficient (+Pi) or -Pi nutrient condition as described previously [58]. The root
and shoot samples were harvested at four different stages: 5, 10, 15 and 20 days of starvation
and snap frozen. For drought stress experiment, the seedlings were allowed to grow in
Hoagland's media containing 5% PEG-8000 [59] and samples collected 72 hrs after stress
treatment. The hydroponic culture was carried out in a growth chamber set at 22 ± 1 °C, 50–
70 % relative humidity and a photon rate of 300 µmol photons $m^{-2}\,s^{-1}$ with a 16 h light/8 h
dark cycle.

Isolation of total RNA, cDNA synthesis and quantitative real time PCR analysis
Total RNA from various tissues was extracted by manual method using TRIzol® Reagent
(Invitrogen™). The integrity of RNA and concentration was measured and contamination
of genomic DNA was removed by subjecting the RNA samples to DNase treatment using
TURBO™ DNase (Ambion, Life Technologies). 2µg of total RNA was used for cDNA
preparation using The Invitrogen SuperScript III First-Strand Synthesis System SuperMix
(Thermo Fisher Scientific) as per the manufacturer's guidelines.

In order to quantify the gene expression, qRT-PCR was performed using the QuantiTect
SYBR Green RT-PCR Kit (Qiagen, Germany). The gene specific primers capable of
amplifying 150-250 bp region from all the three homoeologous of both *TaVIH* genes were
carefully designed using Oligocalc software. Four technical replicates for each set of primers
and minimum of two to three experimental replicates were used to validate the experiment.
Gene specific primer (with similar primer efficiencies) used in the study are listed in
Supplementary Table S6. ADP-ribosylation factor gene (*TaARF*) was used as an internal
control in all the expression studies. The Ct values obtained after the run were normalized
against the internal control and relative expression was quantified using $2^{-\Delta\Delta C_T}$ method [60]. For
In-silico expression for *TaVIH* genes in different tissues and stresses, wheat VIH RefSeq IDs
were used to extract expression values as TPMs from expVIP database. For different tissues
and stages, the expression values were used to build a heatmap. In case of abiotic and biotic

stress conditions, the expression values from the control and stressed conditions were used to
get fold change values, which were then used to plot heatmaps using MeV software.

Identification and cloning of two wheat VIH genes

Two *Arabidopsis* (AT5G15070.2 and AT3G01310.2) and the previously reported yeast VIP1
sequences were used to perform Blastx analysis against the IWGSC (www.wheatgenome.org/)
and wheat EST databases. The identified EST sequences were checked for the presence of the
typical dual domain structure. Further, screening of these sequences resulted in the
identification of two different genomic locations (Table S1). Furthermore, the Pfam domain
identifiers of the signature ATP-Grasp Kinase (PF08443) and Histidine Acid Phosphatase
(PF00328) domains were used to identify VIH proteins in Ensembl database using BioMart
application. The corresponding predicted homoelogous transcripts were identified and
compared to the other VIH sequences. DNA STAR Lasergene 11 Core Suite was used to
perform the multiple sequence alignment and to calculate the sequence similarity. Gene
specific primers capable of amplifying the transcript from the specific genome was designed
after performing 5`and 3`-RACE to ascertain the completed open reading frame (ORF).
Subsequently, full length primers were designed to amplify the *VIH* genes. The generated full-
length sequence information was further used for qRT-PCR related studies.

Homology modelling, hydropathy plot and IDR prediction

Homology modelling was performed for VIH1-4D & VIH2-3B based on their ATP-grasp
domains (residues 7 to 332 for VIH1-4D and 12 to 339 for VIH2-3B), which share an identity
of ~57% with hPPIP5K2 (residues 41 to 366). In both cases, the align2d command in
MODELLER [61](V9.21) was used to align TaVIHs against hPPIP5K2 and the 3D models with
ANP, IHP and 4 Mg^{2+} ions fitted in were calculated using the automodel class. Best models
were selected based on the MODELLER objective function. The models were visualized using
UCSF Chimera [62]. The hydropathy profile for proteins was calculated according to Kyte and
Doolittle., 1982. The positive values indicate hydrophobic domains and negative values
represent hydrophilic regions of the amino acid residues. To identify the % similarity with IDR
boundaries, MFDp2 (http://biomine.cs.vcu.edu/servers/MFDp2 was used to predict the
disorder content in the input sequence [46].

Construct preparation for expression vector and yeast functional complementation

For complementation assays, pYES2, a galactose-inducible yeast expression vector was used.
The functional complementation of yeast by TaVIH proteins was studied using 6-azauracil based
assay. The wild type BY4741 (MATa; his3D1; leu2D0; met15D0; ura3D0) and *vip1Δ* (BY4741;
MATa; ura3Δ0; leu2Δ0; his3Δ1; met15Δ0; YLR410w::kanMX4) yeast strains were used for the
growth assays. The CDS corresponding to the catalytic domain of *ScVIP1* (1-535 amino acids)
cloned into pYES2 expression vector was used as a positive control. *TaVIH1/2* along with
ScVIP1 and empty vector were transformed individually into wild type and mutant strains by
lithium acetate method with slight modifications. For growth assay, the wild type and mutant *S.*
cerevisiae strains carrying different plasmids were allowed to grow overnight in minimal media
without uracil. The primary culture was used to re-inoculate fresh media to an OD_{600} of 0.1 and
allowed to grow till the culture attained an optical density of 0.6-0.8. The cell cultures were then
adjusted to O.D of 1 and further serially diluted to the final concentrations of 1:10, 1:100 and
1:1000. 10 μl each of these cell suspensions were used for spotting on SD(-Ura) plates containing
2% galactose, 1% raffinose and varying concentrations of 6-azauracil (0, 2.5 and 5 mM). The
colony plates were incubated at 30ºC and pictures were taken after 4 days.
555

Protein expression of wheat VIH1 and VIH2, In-Vitro Kinase assays, PAGE analysis and
MADLI-Tof analysis
The TaVIH1-KD and TaVIH2-KD kinase domain was cloned and expressed in *E. coli* BL21
cells using 0.5 mM IPTG and lysis buffer having pH 7.4 containing 50 mM sodium phosphate,
300 mM NaCl and protein inhibitor cocktail. Post sonication and centrifugation purification
was done on Cobalt resin affinity chromatography column (ThermoFisher Scientific, Waltham,
MA, USA). After column saturation overnight at 4°C it was washed with washing buffer
containing 7.5 mM imidazole and subsequently eluted with elution buffer containing 100 mM
EDTA. The eluate was pooled and concentrated using a concentrator having a molecular
weight cut-off of 10 kDa by spinning at conditions mentioned in the concentrator's manual.
The concentrated enzyme preparation was washed thrice with sodium phosphate buffer and
finally concentrated in Tris-HCl buffer, pH 7.4. To check expression western was done by
using Mouse anti-HIS primary antibody and Goat Anti-Mouse secondary antibody [HRP IgG
(H + L): 1:5000 dilutions; Invitrogen].
TaVIH1 and TaVIH2 kinase assays were performed in 20 mM HEPES (pH 7.5), 5 mM
$MgCl_2$, 20 mM ATP, 2 mM IP_6 and 1 mM DTT with 30 ng of respective protein. ScVIP1 was
taken as a control. The reaction was incubated at 28°C for 8-9 hrs, separated by PAGE and
visualized by toluidine blue. Inositol polyphosphates were resolved using 18 cm gel using 33.3

% polyacrylamide gel in Tis-Borate EDTA as mentioned earlier [21]. Gels were pre-run for 75

min at 300 volts and samples were loaded mixed with Dye (10 mM TrisHCl pH 7.0; 1 mM

EDTA; 30 % glycerol; 0.08 % Orange G). Gels were run at 5-6 mA for overnight at 4°C until

the Orange G dye front reached 6 cm from the bottom of the gel. Bands were visualized by

toluidine blue (0.1 % w/v) stain. TBE-PAGE gel purified products of TaVIH reaction was

used for MALDI-ToF-MS analysis. MALDI-ToF-MS was performed from gel extract

solutions which were pipetted on a α-Cyano-4-hydroxycinnamic acid (≥98%, Sigma) prepared

on a stainless-steel plate (0.5 μL of a 10 mg/mL ACN/H2O 1:1 solution). Negative ionization

mode was used for acquiring spectra on spectrometer (AB SCIEX TOF/TOF^{TM} 5800; equipped

with a 337 nm laser) operating in the linear mode.

Wheat cDNA Library construction, yeast two-hybrid screening and pull-down assays

The total RNA (~120 µg) samples were pooled from the vegetative tissues including shoots

and roots. From the isolated total RNA, mRNA purification was performed by using

(NucleoTrap mRNA mini kit, Macherey-Nagel, Germany). A total of 0.25 µg mRNA was used

to make the cDNA library (Make & Plate^{TM} Library System, Clontech, USA). The wheat

cDNA library was prepared and purified using CHROMA SPIN+TE-400 columns. A cDNA

fragment sizes of <2kb was used for the library screening using TaVIH2 as a bait. The library

shows the titre value of ~0.8x10^9 cfu/ml. Yeast two-hybrid assays and screening were

performed by using GAL4-based screening system (Matchmaker Gold Yeast Two-hybrid

System, Takara Inc., USA). Most of the steps were followed as per manufacturer's instructions

unless mentioned. Briefly, putative positive interacting clones were obtained when the

competent yeast strain Y187 was co-transformed with cDNA library+AD (pGADT7-Rec

vector) and Y2H-Gold containing BD vector (*TaVIH2-3B:pGBKT7* bait vector) respectively.

Following stringent screening procedures, putative clones were obtained and screened for their

reporter assays (Aureobasidin A). Full length ORF was cloned for the gene of interest and one-

on-one interaction was also done to confirm its interaction. Routine yeast transformation was

done by using Yeastmaker Yeast transformation System 2 (Clontech, USA).

Yeast cell lysate was prepared for performing pull down assay by using glass bead in

buffer (1% SDS, 100mM NaCl,100mM Tris-Cl, 1mM EDTA, 2% Triton and 1mM protease

inhibitor (100X Halt protease inhibitor, Thermofisher, USA). The protein concentration in

lysate was calculated at two dilutions and processed further using 2 ul of anti-c-Myc antibody.

The proteins were transfer to PVDF membrane and the blot was separated into two parts to

detect TaVIH2 and TaFLA6 respectively. Different primary antibodies were used for probing

(mouse Anti-c-myc and rabbit anti-HA with 1:2000 dilution). After washing the blots with

TBST, they were treated with the secondary antibody (Goat Anti-Mouse IgG (H + L); and Goat

Anti-Rabbit IgG (H + L) with 1:5000 dilution. Blot was developed by using BIO-RAD clarity

western ECL Substrate.

Protein localization

For the localization experiments *TaFLA6* was cloned in pGADT7 vector at *Eco*R1 and *Bam*H1

sites. *TaVIH2* cDNA was cloned in pGBKT7 vector using *Sma*1 and *Not*1 restriction sites. The

constructs were transformed in Y2H Gold yeast strain and selected on SD-Leu or SD-Trp plates

respectively. Yeast spheroplasts were prepared for localization study. Mouse monoclonal Anti-

HA (HA:FLA6-pGADT7): or rabbit Anti-c-Myc (cMYC:VIH2-pGBKT7) primary antibody

(Invitrogen, USA) was used for the respective preparations, at a ratio of 1:200 followed by 5

washing with blocking buffer. Yeast cells were incubated with Goat Anti- Mouse IgG (H+L)

Alexa Flour Plus 488 or Goat Anti- Rabbit IgG (H+L) Alexa Flour Plus 647 (Invitrogen, USA)

at a ratio of 1:500 for 4hr at room temperature. Cells were washed with blocking buffer and

mounted with Fluor mount (Sigma, USA). Representative fluorescent images were taken using

Zeiss fluorescence microscope Axio Imager Z2 with an Axiocam MRm camera at 63X of

magnification.

Cloning of VIH promoter, cDNA and *Arabidopsis* transformation

For promoter, ~2000 bp fragments upstream of the start codon were PCR amplified from

genomic DNA of cv. C306. The cloned DNA fragments (in pJET1.2) were sequenced

confirmed and inserted into pCAMBIA1391z, a promoter-less binary vector containing GUS

reporter gene, using forward and reverse primers with Bam*HI* and Nco*I* sites respectively to

form a *TaVIHpromoter:GUS* in pCAMBIA1391z. The promoter sequences of *TaVIH* genes

were analysed for the presence of cis-regulatory elements using PLANTCARE database

(http://bioinformatics.psb.ugent.be/webtools/plantcare/). For VIH2 cDNA (~3117 bp), blunt

ended cloning was done at *Spe*1 generated site in pCAMBIA1302 along with the C-terminal

His (pCAMBIA1302:*TaVIH*-His) tag. The generated transcription units were introduced into

Arabidopsis seedlings using *Agrobacterium* mediated transformation by floral dip method [63].

Three to four weeks old plants grown at 22 ± 1 °C, 16 h light/8 h dark cycle and a photon rate

of 100 µmol photons m^{-2} s^{-1} were used for transformation. The independent transformants

were screened on 0.5X MS media containing 30 mg/L hygromycin and 0.8% agar. The

transformed seedlings with long hypocotyls and green expanded leaves at 4-leaf stage were

separated out from the non-transformed seedlings and transferred to soil after about 3 weeks.

In a similar manner T_1 and T_2 generation seeds were also selected and allowed to grow till

maturity. The transgenic seedlings were confirmed for the presence of recombinant cassette

using PCR based approach. The transgenic lines harbouring empty pCAMBIA1391Z or

pCAMBIA1302 vector was used as respective negative control. The PCR positive lines were

further used for functional characterization.

GUS-reporter assays and characterization of transgenic lines *in Arabidopsis*

For promoter analysis, the seeds of PCR positive lines were surface sterilized and grown on

0.5X MS (Murashige and Skoog media) agar plates containing 30 mg/L Hygromycin B for 15

days before they were subjected to various abiotic stress and hormonal treatments. For

dehydration stress, the seedlings were air dried by placing them on Whatman filter paper for

1hr. Heat treatment was given by incubating the seedlings at 37˚C for 8hrs. Exposure to ABA

(100 µM), GA_3 (20 µM), NaCl (300 mM) and drought (20% and 30% PEG) were given by

placing the seedlings on filter paper impregnated with 0.5X MS solution containing the

respective chemical for 24 hrs. For Pi deficiency, seedlings were allowed to grow on 0.5X MS

agar plates without KH_2PO_4 for 96 hrs. Histochemical staining of seedlings after respective

treatments were performed by incubated overnight in GUS staining solution [64] with 2 mM X-

Gluc (5-bromo-4-chloro-3-indolyl-beta-D-glucuronic acid, HiMedia, India) at 37 ˚C in a 48-

well microplate containing about ten seedlings/well. Chlorophyll was removed from tissues by

dipping in 90% ethanol. The staining was visualized and photographed under Leica DFC295

stereomicroscope (Wetzlar, Germany) at magnification of 6.3X. MS solution without any

chemical served as a control.

For characterization of transgenic lines overexpressing VIH proteins parameters such

as rosette area, number of leaves, leaves size, length of main root axis and number of shoots

(primary and secondary). Four independent confirmed homozygous transgenic lines were used

for this study. Each parameter was calculated using three experimental replicates, each

consisting of twelve plants. For stress experiments, three days old seedlings of transgenic and

control pre-grown on 0.5X MS were transferred to 0.5X MS plates consisting of either 125

mM mannitol or 10 % glycerol for mimic drought condition. Ten seedlings were used and the

experiments were repeated for three times with similar phenotype. For control, seedlings

continued to grow on ½ MS plates. Root lengths were measured and graphs were plotted (using

three experimental replicate) and pictures were taken after nine days of growth. To calculate

the relative water loss %, twenty-five leaves per five plants with similar developmental stage

for each of the transgenic lines and control plants were subjected to incubation (27 ˚C) for the

period of 12 hrs. Fresh weight of the detached leaf was taken and continued for the

measurements after every 2 hrs. The experiment was repeated twice with similar observations.

For drought response minimum of fifty-five seedlings were pre-grown for the period of

fourteen days and were subjected to drought for additional fourteen days. The plants were then

re-watered for the period of seven days and % survival rates were calculated.

RNAseq profiling

Col-0(Ev) and overexpressing *TaVIH2-3B* Arabidopsis (#Line4 and 6) seedlings were grown

for the period of 25 days. For each genotype, total RNA was extracted from three independent

biological replicates by using RNeasy Plant Mini Kits (Qiagen, CA). Genomic DNA

contamination was removed by digestion with Turbo DNase (BioRad, CA). RNA quantity was

checked by Bioanalyzer for quality control (RIN>8). Library construction and sequencing were

performed by Eurofins, Bangalore, India; using pair end library preparation. About 9.5 to 13.8

million raw reads were obtained for each sample. Raw reads were processed to filter out the

adapter and low quality (QV<20) reads using trimmomatic v0.39 [65] which were then pseudo-

aligned against the reference transcriptome (ensembl release 48) using kallisto v0.46.2 [66]. The

obtained raw abundances were summarized to gene-level expression counts using tximport and

imported to DESeq2 [67,68] for differential expression (DE) analysis in R. The obtained log2 fold

change (LFC) values were further processed using apeglm package to reduce noise [18].Genes

with $1 > LFC < -1$ and padj < 0.05 were considered significantly DE. The expression correlation

across lines and within replicates was analyzed using Principal Component Analysis (PCA) in

ggplot2 [69] .

GC-MS analysis of Arabidopsis cell wall polysaccharides and ABA measurement

Extraction of cell wall components was performed as described earlier with minor modification

as depicted in the flowchart as Supplementary Figure S11 [70]. Since such chemical analysis

requires relatively large amounts of samples, pools from 3-5 independent plants (for each of

the three biological replicates) of the respective lines expressing wheat VIH2 were used for

chemical analysis. Briefly, five grams (fresh weight) of shoots from respective lines and control

at similar developmental stages (~25 days old) was crushed to a fine powder and processed

further. The derived pellet was used for extraction of Arabinoxylan (AX) and Cellulose;

whereas the supernatant was used for extracting arabinogalactan (AG). The extractions were

checked with Iodine solution to make sure that they are free of starch interference. The

compositional analysis the extracted AG, AX and Cellulose were determined by preparing their

alditol derivatives and processed for gas chromatography–mass spectrometry (GC-MS)

analysis as described [7172]. Two µl of samples were introduced in the split less injection mode

in DB-5 (60 m × 0.25 mm, 1 µm film thickness, Agilent, CA) using helium as a carrier gas.

The alditiol acetate derivative was separated using the following temperature gradient: 80 °C

for 2 min, 80-170 °C at 30°C/min, 170-240 °C at 4°C/min, 240 °C held for 30 min and the

samples were ionized by electrons impact at 70 eV. ABA was measured using Plant Hormone

Abscisic Acid (ABA) ELISA kit (Real Gene, Germany). Twenty-five days old plants leaves

were used for the measurement of the ABA content. One gram of fresh weight form eight plants

for each line was used for extractions. The experiments were repeated with at least three

independent extractions and concentration was calculated using standard graphs as per the

manual instructions. Standard graph was plotted using Log of concentration and colour

development for each line was measured at 430 nm (Supplementary Figure S12).

Acknowledgements

Authors thank Executive Director for facilities and support. This study was supported by

Department of Biotechnology, Basic Plant Biology Grant to AKP

[BT/PR12432/BPA/118/35/2014]. Part of this work was also supported by NABI-CORE grant

to AKP. Yeast strain *vip1Δ* and ScVIP1-cDNA was kindly gifted by Dr. Rashna Bhandari

(CDFD). MK thank UGC-CSIR for her research scholarship. Thanks to Dr. Gabriel Schaff for

sharing the *Arabidopsis vih2-3* mutant. AS thank DBT for SRF fellowship. DBT-eLibrary

Consortium (DeLCON) is acknowledged for providing timely support and access to e-

resources for this work.

Data availability

The resources including plasmids, constructs and transgenic *Arabidopsis* seeds will be available

upon request.

References:
1. Stephens, L. *et al.* The detection, purification, structural characterization, and
metabolism of diphosphoinositol pentakisphosphate(s) and bisdiphosphoinositol
tetrakisphosphate(s). *J. Biol. Chem.* **268**, 4009–4015 (1993).
2. Luo, H. R. *et al.* Inositol pyrophosphates are required for DNA hyperrecombination in
protein kinase c1 mutant yeast. *Biochemistry* **41**, 2509–15 (2002).
3. Dubois, E. *et al.* {InSaccharomyces} cerevisiae, the Inositol Polyphosphate Kinase
Activity of Kcs1p Is Required for Resistance to Salt Stress, Cell Wall Integrity, and

746   Vacuolar Morphogenesis. *J. Biol. Chem.* **277**, 23755–23763 (2002).

747 4.   Saiardi, A., Bhandari, R., Resnick, A. C., Snowman, A. M. & Snyder, S. H.
748   Phosphorylation of proteins by inositol pyrophosphates. *Science (80-.).* **306**, 2101–
749   2105 (2004).

750 5.   Norman, K. L. *et al.* Inositol polyphosphates regulate and predict yeast pseudohyphal
751   growth phenotypes. *PLoS Genet.* **14**, e1007493 (2018).

752 6.   Auesukaree, C., Tochio, H., Shirakawa, M., Kaneko, Y. & Harashima, S. Plc1p,
753   Arg82p, and Kcs1p, Enzymes Involved in Inositol Pyrophosphate Synthesis, Are
754   Essential for Phosphate Regulation and Polyphosphate Accumulation
755   {inSaccharomyces} cerevisiae. *J. Biol. Chem.* **280**, 25127–25133 (2005).

756 7.   Wilson, M. S., Jessen, H. J. & Saiardi, A. The inositol hexakisphosphate kinases
757   IP6K1 and -2 regulate human cellular phosphate homeostasis, including XPR1-
758   mediated phosphate export. *J. Biol. Chem.* **294**, 11597–11608 (2019).

759 8.   Voglmaier, S. M. *et al.* Purified inositol hexakisphosphate kinase is an ATP synthase:
760   Diphosphoinositol pentakisphosphate as a high-energy phosphate donor. *Proc. Natl.*
761   *Acad. Sci. U. S. A.* **93**, 4305–4310 (1996).

762 9.   Saiardi, A., Erdjument-Bromage, H., Snowman, A. M., Tempst, P. & Snyder, S. H.
763   Synthesis of diphosphoinositol pentakisphosphate by a newly identified family of
764   higher inositol polyphosphate kinases. *Curr. Biol.* **9**, 1323–1326 (1999).

765 10.   Draškovič, P. *et al.* Inositol Hexakisphosphate Kinase Products Contain Diphosphate
766   and Triphosphate Groups. *Chem. Biol.* **15**, 274–286 (2008).

767 11.   Choi, J. H., Williams, J., Cho, J., Falck, J. R. & Shears, S. B. Purification, Sequencing,
768   and Molecular Identification of a Mammalian {PP}-{InsP}5Kinase That Is Activated
769   When Cells Are Exposed to Hyperosmotic Stress. *J. Biol. Chem.* **282**, 30763–30775
770   (2007).

771 12.   Fridy, P. C., Otto, J. C., Dollins, D. E. & York, J. D. Cloning and characterization of
772   two human VIP1-like inositol hexakisphosphate and diphosphoinositol
773   pentakisphosphate kinases. *J. Biol. Chem.* **282**, 30754–62 (2007).

774 13.   Mulugu, S. *et al.* A conserved family of enzymes that phosphorylate inositol
775   hexakisphosphate. *Science (80-.).* **316**, 106–109 (2007).

776 14.   Dorsch, J. A. *et al.* Seed phosphorus and inositol phosphate phenotype of barley low
777   phytic acid genotypes. *Phytochemistry* **62**, 691–706 (2003).

778 15.   Lemtiri-Chlieh, F., MacRobbie, E. A. C. & Brearley, C. A. Inositol hexakisphosphate
779   is a physiological signal regulating the K$\mathplus$-inward rectifying conductance in
780   guard cells. *Proc. Natl. Acad. Sci.* **97**, 8687–8692 (2000).

781 16.   Desai, M. *et al.* Two inositol hexakisphosphate kinases drive inositol pyrophosphate
782   synthesis in plants. *Plant J.* **80**, 642–653 (2014).

783 17.   Laha, D. *et al.* VIH2 Regulates the Synthesis of Inositol Pyrophosphate InsP8 and
784   Jasmonate-Dependent Defenses in Arabidopsis. *Plant Cell* **27**, 1082–97 (2015).

785 18.   Zhu, J. *et al.* Two bifunctional inositol pyrophosphate kinases/phosphatases control
786   plant phosphate homeostasis. *Elife* **8**, (2019).

787 19.   Osada, S. *et al.* Inositol phosphate kinase Vip1p interacts with histone chaperone
788   Asf1p in Saccharomyces cerevisiae. *Mol. Biol. Rep.* **39**, 4989–4996 (2011).

789 20.   Borrill, P., Ramirez-Gonzalez, R. & Uauy, C. expVIP: A customizable RNA-seq data
790   analysis and visualization platform. *Plant Physiol.* **170**, 2172–2186 (2016).

791 21.   Losito, O., Szijgyarto, Z., Resnick, A. C. & Saiardi, A. Inositol pyrophosphates and
792   their unique metabolic complexity: analysis by gel electrophoresis. *PLoS One* **4**,
793   e5580–e5580 (2009).

794 22.   Johnson, K. L., Jones, B. J., Bacic, A. & Schultz, C. J. The Fasciclin-Like
795   Arabinogalactan Proteins of Arabidopsis. A Multigene Family of Putative Cell

Adhesion Molecules. *Plant Physiol.* **133**, 1911–1925 (2003).

23. Gillmor, C. S. *et al.* Glycosylphosphatidylinositol-Anchored Proteins Are Required for
Cell Wall Synthesis and Morphogenesis in Arabidopsis. *Plant Cell* **17**, 1128–1140
(2005).

24. Schultz, C., Gilson, P., Oxley, D., Youl, J. & Bacic, A. {GPI}-anchors on
arabinogalactan-proteins: implications for signalling in plants. *Trends Plant Sci.* **3**,
426–431 (1998).

25. Saha, S., Anilkumar, A. A. & Mayor, S. GPI-anchored protein organization and
dynamics at the cell surface. *Journal of Lipid Research* vol. 57 159–175 (2016).

26. Chakraborty, A., Kim, S. & Snyder, S. H. Inositol pyrophosphates as mammalian cell
signals. *Science Signaling* vol. 4 (2011).

27. Jadav, R. S., Chanduri, M. V. L., Sengupta, S. & Bhandari, R. Inositol Pyrophosphate
Synthesis by Inositol Hexakisphosphate Kinase 1 Is Required for Homologous
Recombination Repair. *J. Biol. Chem.* **288**, 3312–3321 (2012).

28. Williams, S. P., Gillaspy, G. E. & Perera, I. Y. Biosynthesis and possible functions of
inositol pyrophosphates in plants. *Front. Plant Sci.* **6**, (2015).

29. Huang, S. *et al.* Genes encoding plastid acetyl-CoA carboxylase and 3-
phosphoglycerate kinase of the Triticum/Aegilops complex and the evolutionary
history of polyploid wheat. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 8133–8 (2002).

30. Dvorak, J. & Akhunov, E. D. Tempos of Gene Locus Deletions and Duplications and
Their Relationship to Recombination Rate During Diploid and Polyploid Evolution in
the Aegilops-Triticum Alliance. *Genetics* **171**, 323–332 (2005).

31. Choi, K., Mollapour, E. & Shears, S. B. Signal transduction during environmental
stress: {InsP}8 operates within highly restricted contexts. *Cell. Signal.* **17**, 1533–1541
(2005).

32. Lee, Y. S., Mulugu, S., York, J. D. & O'Shea, E. K. Regulation of a cyclin-CDK-CDK
inhibitor complex by inositol pyrophosphates. *Science (80-.).* **316**, 109–112 (2007).

33. Dong, J. *et al.* Inositol Pyrophosphate InsP8 Acts as an Intracellular Phosphate Signal
in Arabidopsis. *Mol. Plant* **12**, 1463–1473 (2019).

34. Majewska-Sawka, A. & Nothnagel, E. A. The multiple roles of arabinogalactan
proteins in plant development. *Plant Physiology* vol. 122 3–9 (2000).

35. Shi, H., Kim, Y. S., Guo, Y., Stevenson, B. & Zhu, J. K. The Arabidopsis SOS5 locus
encodes a putative cell surface adhesion protein and is required for normal cell
expansion. *Plant Cell* **15**, 19–32 (2003).

36. MacMillan, C. P., Mansfield, S. D., Stachurski, Z. H., Evans, R. & Southerton, S. G.
Fasciclin-like arabinogalactan proteins: specialization for stem biomechanics and cell
wall architecture in Arabidopsis and Eucalyptus. *Plant J.* **62**, 689–703 (2010).

37. Huang, G.-Q. *et al.* Characterization of 19 novel cotton FLA genes and their
expression profiling in fiber development and in response to phytohormones and salt
stress. *Physiol. Plant.* **134**, 348–59 (2008).

38. Nirmal, R. C., Furtado, A., Rangan, P. & Henry, R. J. Fasciclin-like arabinogalactan
protein gene expression is associated with yield of flour in the milling of wheat. *Sci.*
Rep. **7**, (2017).

39. No, E.-G. & Loopstra, C. A. Hormonal and developmental regulation of two
arabinogalactan-proteins in xylem of loblolly pine (Pinus taeda). *Physiol. Plant.* **110**,
524–529 (2000).

40. Cagnola, J. I. *et al.* Reduced expression of selected {FASCICLIN}-{LIKE}
{ARABINOGALACTAN} {PROTEIN} genes associates with the abortion of kernels
in field crops of Zea mays (maize) and of Arabidopsis seeds. *Plant. Cell Environ.* **41**,
661–674 (2018).

41.   Kirchner, T. W. *et al.* Molecular Background of Pi Deficiency-Induced Root Hair
Growth in Brassica carinata {\textendash} A Fasciclin-Like Arabinogalactan Protein Is
Involved. *Front. Plant Sci.* **9**, (2018).

42.   Scheible, W.-R. & Pauly, M. Glycosyltransferases and cell wall biosynthesis: novel
players and insights. *Curr. Opin. Plant Biol.* **7**, 285–295 (2004).

43.   Whitehead, C. *et al.* A glycosyl transferase family 43 protein involved in xylan
biosynthesis is associated with straw digestibility in Brachypodium distachyon. *New
Phytol.* **218**, 974–985 (2018).

44.   Ueki, S. & Citovsky, V. Identification of an interactor of cadmium ion-induced
glycine-rich protein involved in regulation of callose levels in plant vasculature. *Proc.
Natl. Acad. Sci. U. S. A.* **102**, 12089–94 (2005).

45.   Machkalyan, G., Trieu, P., Pétrin, D., Hébert, T. & Miller, G. PPIP5K1 interacts with
the exocyst complex through a C-terminal intrinsically disordered domain and
regulates cell motility. *Cell. Signal.* **28**, (2016).

46.   Mizianty, M. J. *et al.* Improved sequence-based prediction of disordered regions with
multilayer fusion of multiple information sources. *Bioinformatics* **26**, i489–i496
(2010).

47.   Randall, T. A., Gu, C., Li, X., Wang, H. & Shears, S. B. A two-way switch for inositol
pyrophosphate signaling: Evolutionary history and biological significance of a unique,
bifunctional kinase/phosphatase. *Adv. Biol. Regul.* **75**, 100674 (2020).

48.   Marathe, A. *et al.* Exploring the role of Inositol 1,3,4-trisphosphate 5/6 kinase-2 (
GmITPK2) as a dehydration and salinity stress regulator in Glycine max (L.) Merr.
through heterologous expression in E. coli. *Plant Physiol. Biochem.* **123**, 331–341
(2018).

49.   Du, H. *et al.* Characterization of an inositol 1,3,4-trisphosphate 5/6-kinase gene that is
essential for drought and salt stress responses in rice. *Plant Mol. Biol.* **77**, 547–563
(2011).

50.   Daszkowska-Golec, A. & Szarejko, I. The Molecular Basis of ABA-Mediated Plant
Response to Drought. *Abiotic Stress - Plant Responses and Applications in Agriculture*
(2013) doi:10.5772/53128.

51.   Moore, J. P., Vicré-Gibouin, M., Farrant, J. M. & Driouich, A. Adaptations of higher
plant cell walls to water loss: drought vs desiccation. *Physiol. Plant.* **134**, 237–245
(2008).

52.   Tenhaken, R. Cell wall remodeling under abiotic stress. *Front. Plant Sci.* **5**, 771
(2015).

53.   Takahashi, F., Kuromori, T., Urano, K., Yamaguchi-Shinozaki, K. & Shinozaki, K.
Drought Stress Responses and Resistance in Plants: From Cellular Responses to Long-
Distance Intercellular Communication. *Front. Plant Sci.* **11**, 556972 (2020).

54.   Shinozaki, K. & Yamaguchi-Shinozaki, K. Gene networks involved in drought stress
response and tolerance. *J. Exp. Bot.* **58**, 221–227 (2006).

55.   Pandian, B. A., Sathishraj, R., Djanaguiraman, M., Prasad, P. V. V. & Jugulam, M.
Role of Cytochrome P450 Enzymes in Plant Stress Response. *Antioxidants (Basel,
Switzerland)* **9**, 454 (2020).

56.   Klein, M. *et al.* The plant multidrug resistance ABC transporter AtMRP5 is involved
in guard cell hormonal signalling and water use. *Plant Journal* vol. 33 119–129
(2003).

57.   De Vos, M. *et al.* The arabidopsis thaliana transcription factor AtMYB102 functions in
defense against the insect herbivore Pieris rapae. *Plant Signal. Behav.* **1**, 305–311
(2006).

58.   Shukla, V. *et al.* Tissue specific transcript profiling of wheat phosphate transporter

genes and its association with phosphate allocation in grains. *Sci. Rep.* **6**, (2016).

59.   Ji, H. *et al.* {PEG}-mediated osmotic stress induces premature differentiation of the
root apical meristem and outgrowth of lateral roots in wheat. *J. Exp. Bot.* **65**, 4863–
4872 (2014).

60.   Livak, K. J. & Schmittgen, T. D. Analysis of Relative Gene Expression Data Using
Real- Time Quantitative PCR and the $2^{-\Delta\Delta C_T}$ Method. *Methods* **408**, 402–408
(2001).

61.   Eswar, N. *et al.* Comparative Protein Structure Modeling Using Modeller. *Curr.*
Protoc. Bioinforma. **15**, 5.6.1--5.6.30 (2006).

62.   Pettersen, E. F. *et al.* {UCSF} Chimera?A visualization system for exploratory
research and analysis. *J. Comput. Chem.* **25**, 1605–1612 (2004).

63.   Zhang, X., Henriques, R., Lin, S. S., Niu, Q. W. & Chua, N. H. Agrobacterium-
mediated transformation of Arabidopsis thaliana using the floral dip method. *Nat.*
Protoc. **1**, 641–646 (2006).

64.   Jefferson, R. A. Assaying chimeric genes in plants: The GUS gene fusion system.
Plant Mol. Biol. Report. **5**, 387–405 (1987).

65.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

66.   Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Erratum: Near-optimal
probabilistic RNA-seq quantification. *Nat. Biotechnol.* **34**, 888 (2016).

67.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

68.   Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq:
transcript-level estimates improve gene-level inferences. *F1000Research* **4**, 1521
(2015).

69.   Gómez-Rubio, V. ggplot2 - Elegant Graphics for Data Analysis (2nd Edition). *J. Stat.*
Softw. **77**, (2017).

70.   Zablackis, E., Huang, J., Müller, B., Darvill, A. G. & Albersheim, P. Characterization
of the cell-wall polysaccharides of Arabidopsis thaliana leaves. *Plant Physiol.* **107**,
1129–1138 (1995).

71.   Blakeney, A. B., Harris, P. J., Henry, R. J. & Stone, B. A. A simple and rapid
preparation of alditol acetates for monosaccharide analysis. *Carbohydr. Res.* **113**, 291–
299 (1983).

72.   Bhagia, S., Nunez, A., Wyman, C. E. & Kumar, R. Robustness of two-step acid
hydrolysis procedure for composition analysis of poplar. *Bioresour. Technol.* **216**,
1077–1082 (2016).

List of Tables:

Table 1: List of genes identified as an interacting partners of wheat VIH2-3B. The table

summarises the information of the predicted gene function (based on the ensemble-BLAST) of

the sequenced clones. All the enlist genes resulted in blue colony formation when the screening

was performed on the SD-GALα (-AHLT) plates. Respective TRIAE IDs (RefSeq V1.0) are

mentioned. Most of these genes show more than once occurrence of the yeast colonies during
screening except for S.No 1, 2,3 clones that appeared more than thrice.

S.No	Predicted Biological Annotation	New TRIAE_ ID
1.	*Fasciclin-like arabinogalactan protein (FLA6)* *(protein ID:* ABI95396.1)	TraesCS2A02G165600
2.	*Glycosyl transferases (GT)*	TraesCS6B02G339100.1
3.	*Xylan arabinosyl transferase (Xat1)*	TraesCS6A02G309400.1
4.	*Glycine-rich cell-wall structural protein-like*	TraesCS2B02G541900
5.	*Alpha-amylase/trypsin inhibitor CM1*	TraesCS7B02G072000
6.	*NADH-ubiquinone reductase complex 1 MLRQ subunit;*	TraesCS7A02G238600
7.	*Hypothetical protein TRIUR3_12806*	TraesCS4A02G016700
8.	*ABC transporter B family member 25 A1-1*	TraesCS5A02G392600
9.	*Short chain dehydrogenase reductase*	TraesCS2B02G116700
10.	*WW domain binding protein (containing PPGPPP motif)*	TraesCS2A02G245500
11.	*Ethylene-responsive element binding protein 1 (EREB1) mRNA,*	TraesCS7B02G062200
12.	*Triticum aestivum mRNA, clone: tplb0058f16,*	TraesCS2A02G558900

942
943
Figure 1. Neighbourhood-Joining phylogenetic tree and expression analysis of wheat
genes encoding VIH. (A) Neighbourhood-Joining phylogenetic tree of VIP proteins. The full-
length amino acid sequences of VIH proteins from various taxonomic groups were used for the
construction of phylogeny using MEGA7.0. (B) *TaVIH1* and *TaVIH2* in different tissues of a
wheat plant. The cDNA prepared from 2µg of DNA free-RNA isolated from root, stem, leaf
and flag leaf tissues of a 14 DAA plant as template. (C) Quantitative expression analysis of
TaVIH genes at different seed maturation stages (7, 14, 21 and 28 days after anthesis and in 14

DAA seed (aleurone, Al; endosperm, En; embryo, Em; glumes, Gl and and rachis, Ra. For

qRT-PCR, cDNA was prepared from 2μg of DNA-free RNA isolated from respective tissues.

TaARF was used as an internal control for normalization of Ct values. Standard deviation was

calculated along with its level of significance at $p<0.05$ (*) with respect to the first tissue. (D)

Microarray based expression profiles of *TaVIH* genes.

Figure 2: Enzymatic activity and analysis of the substrate product on PAGE and MALDI-

Tof MS analysis. (A) The relative luminescence units for all reactions performed were

recorded using Spectramax optical reader. The kinase reactions were performed using 50 ng of

VIH1 and VIH2 purified proteins. (B) The products were resolved in the 33% PAGE and

stained with Toludene Blue using IP_6 (left panel) and IP_7 (right panel). Yeast VIP protein was

used as a positive control. These experiments were repeated 3 times using and similar assays

for product formation was obtained (C). MALDI-Tof MS analysis of synthesized products IP_7

and IP_8. MS analysis indicated a major signal at m/z of 739.009 that correspond to mass of IP_7

and IP_8 -adduct at 861.05 that matches with the expected generated species (details at

supplementary Figure X).

Figure 3: Generation of VIH2-3B transgenic Arabidopsis and its characterization. (A)

Western analysis and screening of Arabidopsis transgenic lines for VIH2-3B protein

overexpressing lines. Multiple transgenic lines were screened and Western was done using His-

Antibody using 20 ug of total protein. Coomassie Blue stain of the total protein (lower panel)

was used as a loading control. (B) Representative picture of the rosettte area of the Transgenic

Arabidopsis (#Line2, #Line4, #Line5 and #Line6) and controls. (C) Rosette area measurement

(in cm2) using Image-J for 4 different transgenic lines along with the controls. Measurement

was taken after 14 days of growth. (D) Number of Rosette leaves in transgenic Arabidopsis

and control lines. Three experimental replicates using 10 plants each were used to calculate the

parameters.

Figure 4: Morphological characterization of VIH2-3B transgenic Arabidopsis (A)

Representative pictures of transgenic Arabidopsis and controls post 25 days of growth

(flowering stage) and; length of main axis along with the leaves size (in mm). (B) Number of

total shoots (primary and secondary shoots) in transgenic Arabidopsis and control plants (right

panel). Left-panel show the morphological differences (as indicated by arrows) among the

lines. For each transgenic line, three experimental replicates were performed using 10 plants

each.

Figure 5: Drought response of VIH2-3B *Arabidopsis* transgenic lines. (A) Reporter assays

using promTaVIH2:GUS transgenic lines subjected to drought-mimic (30% PEG). Seedlings

with or without treatment (control) were stained overnight in GUS staining solution and
photographed using Leica stereomicroscope at 6.3X magnification. (B) Transgenic
Arabidopsis and control seedlings were subjected to drought mimic conditions with glycerol-
10 % and mannitol-125 mM. Ten seedlings were used for each transgenic line for each
treatment. These experiments were repeated three experimental replicates with similar
phenotype. (C) Root length of treated seedlings (in mm) for all the lines. Twenty seedlings
were used for the measurement of root length for each line. (D) Relative water loss in
Arabidopsis leaves post 12 hrs. Three experimental replicates each with ten leaves were used
to calculate the water loss %. (E) Drought treatment of soil grown plants. Fifty-five seedlings
pre-grown for the period of fourteen days were subjected to drought for additional fourteen
days. The plants were then re-watered for the period of seven days and % survival rates were
calculate. Representative pictures were taken post seven days of re-watering.

Figure 6: Interaction of TaVIH with TaFLA6 as identified during Y2H screening using
TaVIH2-3B as a bait. (A) Y2H assay for the FLA6 and VIH interaction as represented by
GAL plates containing -L-T and without GAL plates containing -HLT. (B) Pull down assay
and Western analysis of the wheat FLA6 in the yeast strains. (C) Localization of FLA6 using
HAtagged-TaFLA6 and TaVIH2 using cMYC tag. Fluorescence was measured using a Zeiss
fluorescence microscope (Axio Imager Z2) with an Axiocam MRm camera at 63X.
Representative images are shown and similar observations were noted for 3-4 independent
transformed colonies of yeast. (D) Protein domain arrangement and hydrophobic plot for FLA6
domains with negative values represent hydrophilic regions.

Figure 7: RNAseq analysis of Col-0 and #Line4 and 6. (A) Expression pattern (as Z-scores)
of top 56 genes commonly up-regulated among the transgenic lines w.r.t. Col-0(Ev) in 25 days
old seedlings. Heatmap depicts the comparative expression in Col-0(Ev) and over-expressing
lines of TaVIH2-3B (3 biological replicates; rep1-3). (B) Heatmap representing a graphical
summary of the Gene Ontology (GO) classification for DEGs in #Line4 and #Line6 w.r.t.
Control plants. Increasing intensities of brown and blue colors represent the comparative low
and high expression for each gene, as depicted by the color scale. Normalized expression
counts were used to plot the expression as Z-scores using heatmap.2 function from gplots
package in R. Significantly altered GO terms were identified using Classification SuperViewer
tool; x-axis represents the GO terms where bold terms represent significant alteration while y-
axis represents the normed frequency which when > 1 signifies over-representation while <1
signifies under-representation.

Figure 8: RNAseq analysis of Col-0(Ev) overexpressing *TaVIH2-3B* Arabidopsis (#Line4 and 6). Heatmaps for expression patterns (as Z-scores) for genes DE in both transgenic lines w.r.t. Col-0(Ev), encoding for (A) Heatmap representing the comparative expression response of genes involved in cell wall related homeostasis that were DE in both the transgenic lines w.r.t Col-0(Ev). (B) ABA biosynthesis related pathway genes, (C) DREB encoding genes; (D) cytochrome P450 (CYPs) genes. Increasing intensities of brown and blue colors represent the comparative low and high expression for each gene, as depicted by the color scale. Normalized expression counts were used to plot the expression as Z-scores using heatmap.2 function from gplots package (Warnes et al., 2005) in R. Genes encoding for respective pathways were extracted using MapMan (Thimm et al., 2004). R1, R2 and R3 represent the biological replicates for the RNAseq analysis of the respective lines.

Figure 9: ABA and polysaccharides composition of Arabidopsis shoots and speculative model for the functioning of wheat VIH2. (A) ABA measurement in the leaves of transgenic Arabidopsis overexpressing VIH2-3B and control plants. (B) Venn diagram representation for the genes differentially regulated by drought stress, and transgenic lines #Line4 and #Line6 w.r.t. respective Controls. Drought responsive genes were shortlisted using the Cufflinks pipeline after processing the datasets for 10 days drought stress and control. (C) Mapman pathway analysis using Classification SuperViewer for the genes that are commonly regulated by drought stress (SRA: SRP075287) as well as transgenic lines w.r.t. control plants (#Line4 and 6). Bold terms represent significant pathways, normed frequency > 1 signifies over-representation while < 1 signifies under-representation. (D) For cell wall composition analysis wildtype Col-O, *Arabidopsis* overexpressing TaVIH2-3B (Line#2, 4, 5 and 6) and *Arabidopsis vih2-3* representing Arabidopsis mutant defective for the expression of AtVIH2 were used. Total AG: arabinogalactan, AX: arabinoxylan and Cellulose (in µg/g) was measured as indicated in Methods. Analyses were made in triplicates with each experimental replicate representing at least five plants for each genotype. Vertical bars represent the standard deviation. * on the bar indicates that the mean is significantly different at $p < 0.001$ (#at p<0.05) with respect to their respective control plants.

Figure 10: Speculative model for the working of VIH2 to impart drought resistance to plants. VIH2 could be involved in cell wall related developments via FLA like proteins or by causing changes in the stomatal distribution. Bothe the effects might contribute for the drought response in transgenic Arabidopsis.

Legends for Supplementary Figures and Tables

Supplementary Figure S1: Kyte-Doolittle Hydropathy plots and conserved domains of wheat VIH proteins. (A) Kyte-Doolittle hydropathy plots with the positive values indicating the hydrophobic domains and negative values represent hydrophilic regions of the amino acid residues. The hydropathy profile for proteins was calculated according to Kyte and Doolittle., 1982. (B) Schematic representation of domain architecture of TaVIH proteins deduced from CDD database: light gray rectangles indicate ATP Grasp/RimK Kinase domain and dark gray colored heaxgon correspond to Histidine Phosphatase superfamily.

Supplementary Figure S2: Expression patterns of *TaVIH* gene homoeologous in different tissues and stress conditions. RNAseq datasets of (A) Tissues and developmental stages (B) Abiotic (phosphate starvation, heat and drought stress) and (C) Biotic stress conditions were used. The expression values were obtained from expVIP database in the form of TPM values and ratios of stressed to control condition were used to generate heatmaps using MeV software. Green and red colors represent down-regulation and up-regulation of the genes in the specific stresses, as shown by the color bar.

Supplementary Figure S3: Three-dimensional (3D) structure for TaVIH proteins based on homology modelling and yeast complementation assays. (A) TaVIH2-3B overall structure depicting the AMP-PNP ligand accommodated by two anti-paralled beta-sheets. (B) hPPIP5K2, TaVIH1-4D, and TaVIH2-3B depicting the conserved catalytic residues in the IP6/IHP binding pocket. The key conserved residues are depicted using sticks and the green spheres represent Mg^{2+} ions. (C) Yeast complementation assay for *TaVIH* genes. Representative image of spotting assay performed on SD-Ura plates containing 1% raffinose, 2% galactose and supplemented with 0, 2.5 and 5mM of 6-azauracil. The wild type BY4741 and *vip1Δ* strains were transformed with respective constructs using Li-acetate method. Representative images were taken 4 days after spotting assay was performed. Similar results were obtained with three independent repeats. (D) Filamentous growth assays were observed for wild type yeast (WT), yeast mutant- *vip1Δ* with empty pYES2 (*vip1Δ*) and *TaVIH2-3B* complementation in vip1Δ- (*TaVIH2-3B*+ Δvip1). Pictures were taken 20 days post incubation.

Supplementary Figure S4: Purification and western analysis of purified wheat VIH1 and VIH2 kinase domain (KD) proteins. Both the VIH proteins (VIH1 and VIH2) were expressed and purified as mentioned in the Methods section and the expression was confirmed by the Western analysis using His-antibody.

Supplementary Figure S5: MALDI-ToF-MS analysis of the IP_7 from gel extracts. Gel purified inositol pyrophosphates from (A) IP_6 at 660.25 (B) VIH1 derived IP_7 at 739.8 and (C) VIH2 derived IP_7 at 739.00 were subjected to mass spectrometry analysis and m/z spectrum of IP_7 is shown. The peaks in the spectra describing as IP_6 and IP_7 are in agreement with the theoretical values for molecular weight that are deduced to be 660 and 740 Da.

Supplementary Figure S5: MALDI-ToF-MS analysis of the IP_7 and IP_8 from PAGE gel extracts. Gel purified inositol pyrophosphates from (A) IP_7 at m/z of 739.82 (generated by VIH1) (B) IP_7 at m/z of 739.009 (generated by VIH2); (C) IP_8 at m/z of 820.00 or its ACN adduct at m/z of 861.05 (generated by VIH1); (D) IP_8 adduct (CAN) at m/z of 861.05 (generated by VIH12). The peaks in the spectra describing as IP_7 and IP_8 are in agreement with the theoretical values for molecular weight.

Supplementary Figure S6: Hormonal and abiotic stress response of *TaVIH* genes promoter. (A) Cis-element analysis of VIH1 and VIH2 promoters (~1.5kb) Multiple stress related domains are represented in a schematic form. (B) Representative images for histochemical GUS assay performed against different stresses for promTaVIH1:GUS and promTaVIH2:GUS transgenic lines raised in *Arabidopsis thaliana* Col-0 background. Two week old seedlings selected positive against hygromycin selection on 0.5XMS agar plates were subjected to respective treatments: 1hr air drying for dehydration, heat stress at 37 ˚C for 8hrs, (-)Pi condition: 0.5XMS medias without KH_2PO_4 for 96 hrs, ABA (100µM), GA_3 (20µM) and NaCl (300mM) and drought (20% PEG) for 24hrs. Seedlings with or without treatment (control) were stained overnight in GUS staining solution and photographed using Leica stereomicroscope at 6.3X magnification.

Supplementary Figure S7: Western blot analysis, cDNA preparation and screening of the VIH interacting proteins (representative image). (A) Western analysis of c-MYC fused TaVIH1 or TaVIH2 proteins in the yeast strain. (B) Double strand cDNA preparation of wheat seedling library used for yeast two hybrid interaction studies. The cDNA library was resolved on the 1.2 % agarose gel. Two different cDNA preparations (Rep1 and Rep2) was performed and pooled together for the library preparation. (C) Representative yeast colony map for calculating the mating efficiency ($-LT_{1/1000}$). (D) Representative picture of the yeast colonies with putative interacting clones obtained on the selection plates (-AHLT+αGal). Each of the independent streaked colonies represent single putative interacting clone. Only colonies showing strong-blue coloration were used further for further study.

Supplementary Figure S8: Relative quantification of *TaVIH2* and its interacting partner during drought and development stages. (A) *TaVIH2* transcripts in Roots and Shoots of wheat

seedlings after 72 hrs of treatment with 5% PEG-8000 (CR-control roots; RT-root treatment; CS- control shoot; ST-shoot treatment. cDNA templates were prepared from 2 µg RNA isolated from samples of CR-Control root, RT-Root treatment with PEG, CS-Control shoot and ST-Shoot treatment. The Ct values were normalized against the reference gene *TaARF*. Respective standard deviation was calculated with their level of significance at $p < 0.05$ (*) with respect to their respective control. (B Expression of *TaFLA6, TaGT* and *TaXat1 during different stages of development and* (C) during the drought condition for *(TaGT),* TraesCS6B02G339100 and *Xylan arabinosyl transferase (TaXat1),* TraesCS6A02G309400. Bar plot depicting fold change levels for *FLA6, TaGT* and *TaXat1* under drought condition post 1 and 6 hrs of stress.

Supplementary Figure S9: RNAseq analysis of transgenic Arabidopsis. (A) PCA analysis of the RNAseq for control (Col-0 (Ev)) and two transgenic Arabidopsis lines. (B) Map man analysis of the genes those are consistently represented in the two transgenic Arabidopsis lines with overexpressing TaVIH2-3B.

Supplementary Figure S10: Flow representation of the preparation and extraction of polysaccharides (Arabinogalactans, Arabinoxylans and Cellulose) form the shoots of *Arabidopsis*

Supplementary Figure S11: Standard graph for ABA measurement in plant leaves samples. (A) Y-axis indicates Log of concentration and X-axis indicates the optical density. Data was linearized by plotting the log of the target antigen concentrations versus the log of the OD and the best fit line was determined by regression analysis. (B) Panel showing the colour development for the quantitation of the ABA in different leaf samples, OD was taken at 420 nm.

Supplementary Table S1: List of *TaVIH* genes with computed physical and chemical parameters. The molecular weight and isoelectric point prediction were done using Expasy ProtParam tool (https://web.expasy.org/protparam/). The sub-cellular localization prediction was done using WoLF PSORT prediction tool (http://www.genscript.com/wolf-psort.html). RefSeq v1.1 for wheat Ensembl Plants was used for gene ID.

Supplementary Table S2: RPKM values of *TaVIH* genes' transcripts for normal (0 day) and phosphate starved (10 day) conditions in root and shoot tissues. The RNAseq data was used from Oono et al., 2014.

Supplementary Table S3: List of genes up- and down-regulated in #line4 (Sheets1,2) and line6 (Sheets3,4) w.r.t. Col-0(Ev) lines. DEGs were obtained using the Kallisto-DESeq2

pipeline; genes with LFC > 1 in either direction and padj < 0.05 were considered to be
differentially regulated.

Supplementary Table S4: GC-MS analysis in shoot of *Arabidopsis* overexpressing *TaVIH2-*
3B. Analysis was done for two independent extractions (independent extractions; Replicate1
and Replicate2; each using 4-5 pooled plants for respective lines). Inositol-derivative was used
as an internal control as mentioned in the Methods section. Four independent transgenic lines
(#Line2, 4, 5 and 6) along with control plants (Col-0 (Ev)) and mutant *vih2-3* line of
Arabidopsis was used for analysis of Arabinogalactans (AG), Arabinoxylans (AX) and
cellulose.

Supplementary Table S5: List of drought responsive genes that are differentially regulated in
#line4 (Sheet1), #line6 (Sheet2), and differentially regulated in both #line4 and line6 (Sheet3).
Drought responsive genes at 10days of drought stress w.r.t Control plants were extracted from
the SRA RNAseq dataset (SRP075287) using Cufflinks pipeline. Genes with 1 > LFC < -1
were considered to be drought responsive.

Figure 1

(A)

(B)

(C)

IP8 adduct
(IP8+Acetonitrile:861.05)

Figure 2

Figure 3

(A)

Col-0   Col-0(Ev)   #Line 2   #Line 4   #Line 5   #Line 6

25 Days

NS

*p <0.05

NS

Length of main
root axis (mm)

NS

Col-0   Col-0-Ev   #Line 2   #Line 4   #Line 5   #Line 6

* p <0.05

NS

Leaves Size (mm)

NS

Col-0   Col-0-Ev   #Line 2   #Line 4   #Line 5   #Line 6

(B)

*p<0.0001

NS

NS

Number of Shoots

Secondry shoot
Primary shoot

Col-0   Col-0-Ev   #Line 2   #Line 4   #Line 5   #Line 6

Col-0   Col-0-Ev   #Line 2   #Line 4   #Line 5   #Line 6

Figure 4

Figure 5

Figure 6

(A)

Col-0(Ev)	#Line4	#Line6

Integrase-type DNA-binding superfamily protein
Stress induced protein
Polyketide cyclase/dehydrase and lipid transport superfamily protein
mto 1 responding down 1
Integrase-type DNA-binding superfamily protein
nitrile specifier protein 2
histone acetyltransferase (DUF1264)
transmembrane protein
transmembrane protein
Integrase-type DNA-binding superfamily protein
Integrase-type DNA-binding superfamily protein
ARM repeat superfamily protein
hypothetical protein AT3G03280
Integrase-type DNA-binding superfamily protein
Glycine-rich protein family
nuclease
dicarboxylate carrier 2
Duplicated homeodomain-like superfamily protein
low-molecular-weight cysteine-rich 62
cotton fiber protein
ECA1 gametogenesis related family protein
SAUR-like auxin-responsive protein family
Polyketide cyclase/dehydrase and lipid transport superfamily protein
cytochrome P450, family 702, subfamily A, polypeptide 1
salt-inducible zinc finger 1
root meristem growth factor
ROTUNDIFOLIA like 12
cytochrome P450, family 707, subfamily A, polypeptide 3
glycine-rich protein
Calcium-binding EF-hand family protein
calmodulin-like 38
hypothetical protein AT4G27652
Polynucleotidyl transferase, ribonuclease H-like superfamily protein
Integrase-type DNA-binding superfamily protein
Copper amine oxidase family protein
Pollen Ole e 1 allergen and extensin family protein
transmembrane protein
redox responsive transcription factor 1
Pectin lyase-like superfamily protein
seed storage albumin 1
RmlC-like cupins superfamily protein
Plant thionin
RmlC-like cupins superfamily protein

-1.5 -1 -0.5 0 0.5 1 1.5
Row Z-Score

(B)

Figure 7

Figure 8

Figure 9

VIH
(Ins. pyrophosphate kinase)

FLA6, XAT

ABA/Drought
responsive genes

Cell wall
related developments

(Arabinoxylans/galactans,
cellulose)

Drought tolerance

Figure 10