Zygomorphic flowers have fewer potential pollinator species

Jeremy B. Yoder[1,3], Giancarlo Gomez[1], and Colin J. Carlson[2]

[1] Department of Biology, California State University Northridge, Los Angeles, CA, 91330, USA

[2] Department of Biology, Georgetown University, Washington, D.C. 20057, USA

[3] *To whom correspondence should be addressed: jeremy.yoder@csun.edu*

Keywords: Pollination, floral morphology, ecological specialization, interaction networks

2,705 words, 2 figures, no tables

Last saved by Jeremy Yoder, 6 August 2020

ABSTRACT

Botanists have long identified bilaterally symmetrical (zygomorphic) flowers with more

specialized pollination interactions than radially symmetrical (actinomorphic) flowers.

Zygomorphic flowers facilitate more precise contact with pollinators, guide pollinator behaviour,

and exclude less effective pollinators. However, whether zygomorphic flowers are actually

visited by a smaller subset of available pollinator species has not been broadly evaluated. We

compiled 53,609 floral visitation records in 159 communities and classified the plants' floral

symmetry. Globally and within individual communities, plants with zygomorphic flowers are

indeed visited by fewer species. At the same time, zygomorphic flowers share a somewhat larger

proportion of their visitor species with other co-occurring plants, and have particularly high

sharing with co-occurring plants that also have zygomorphic flowers. Visitation sub-networks for

zygomorphic species also show differences that may arise from reduced visitor diversity,

including greater connectance, greater web asymmetry, and lower coextinction robustness of

both plants and visitor species — but these changes do not necessarily translate to whole plant-

visitor communities. These results provide context for widely documented associations between

zygomorphy and diversification and imply that species with zygomorphic flowers may face

greater risk of extinction due to pollinator loss.

INTRODUCTION

An axiom of pollination ecology is that flowers with bilateral symmetry are more specialized than flowers with radial symmetry [1–4]. "Specialized," however, has multiple meanings in evolution and ecology, which are not mutually exclusive. Specialization may refer to a derived character state in a phylogenetic context [1,3], or the degree to which a flower manipulates pollinator behaviour [5], or it may refer to association with a particular set of pollinators (i.e., pollination syndromes; [6,7]) — or, finally, it may refer to association with fewer available pollinator species than seen for comparable plant species [8]. Zygomorphic flowers are derived within the angiosperms [9–11], and extensive research examines how floral structure attracts, guides, or excludes pollinators [5,6,12–15]. However, data addressing the fourth sense in which zygomorphic flowers may be specialized — association with fewer pollinators than otherwise expected — are surprisingly sparse.

Floral symmetry has been recognized as an important feature of angiosperm diversity since at least the 18th Century [4]. Modern treatments identified zygomorphy as derived, and hypothesized that zygomorphic forms facilitate more effective pollination [1,3,16]. Zygomorphy is associated with greater diversification rates [17–19], consistent with the hypothesis that using fewer or more constant pollinators creates more opportunities for reproductive isolation [20]. Greater pollination specialization might also interact with global patterns of diversity, such as latitudinal gradients [21,22]: recent syntheses find evidence that biotic interactions are stronger in the tropics [23,24], though assessments of latitudinal effects on pollination specifically have mixed results [25,26].

Floral symmetry has been considered as an element of pollination syndromes [6,7], but to our

knowledge, documentation that zygomorphic flowers associate with fewer pollinator species is

restricted to anecdotal observations (e.g. [1,3,5]). Broad confirmation of this understanding

would illuminate the research linking pollination to diversification [1,16,18,20,27]. Ecologically,

the use of fewer pollinators by zygomorphic flowers may have implications for factors ranging

from species' geographic extents to risks of extinction due to pollinator loss.

If floral symmetry creates systematic differences in pollinator associations, these differences

should manifest in floral visitation networks. Plant-pollinator associations have been prominent

case studies in investigations of ecological networks' structure and assembly [28–30], geographic

variation [25,26], and evolutionary stability [31]. Databases of floral visitation networks have

global coverage, recording visitor diversity and sharing among co-occurring angiosperm species

across a variety of contexts. Here, we compile a global dataset of floral visitation records to test

the hypothesis that zygomorphic flowers have fewer visitor species, and examine how this effect

may shape plant-pollinator networks.

METHODS

We compiled floral visitation networks from the Web of Life (www.web-of-life.es) and

Interaction Web DataBase (www.nceas.ucsb.edu/interactionweb). Networks varied widely in

size, attributable to heterogeneity in the time periods over which they were studied, their

observation protocols, and the geographic ranges of communities represented (Figure 1). We

accounted for this in subsequent analyses by simplifying visitation into a binary state (the most

common recording mode), and by examining contrasts within individual networks or treating

network identity as a random effect (see below).

We identified unique plant taxa across all networks (hereafter 'species'; identification of plants

and visitors varied in resolution, but 95% of records had plants identified to species), and

classified their floral symmetry based on taxonomic knowledge, formal descriptions of species or

higher taxa, and, when necessary, inspecting images of herbarium sheets or reliably identified

fresh flowers. We determined symmetry based on the perianth, but in ambiguous cases also

considered symmetry of the androecium or gynoecium; for example, a flower with a very nearly

radially symmetric corolla, but stamens arrayed in a bilaterally symmetric manner, would be

coded as zygomorphic. In some cases, we classified symmetry not based on individual flowers

but on flowering heads (e.g., we considered species in the Asteraceae actinomorphic). We

removed species from the working dataset if we were unable to find authoritative descriptions or

images, or if they were wind-pollinated ("n. det.", Figure 1B). Data and scripts are posted to

Dryad, at https://datadryad.org/10.5061/dryad.gxd2547j3

We conducted analysis in R v. 4.0 [32]. For each plant species in the dataset, we totalled the

visitor species recorded, and calculated an index of visitor species sharing, the proportion of

visitor species to each plant species that also visit each other co-occurring plant species,

averaged across the co-occurring plant species. We calculated sharing with all plants in the same

network, and sharing with plants in the same network having each type of floral symmetry. We

examined the structure of complete networks, and of the sub-networks of visitors to plants with

each type of floral symmetry in each community, calculating connectance (the realized

proportion of possible plant-visitor links [33]), web asymmetry (the degree to which visitor

species outnumber plant species or vice-versa), and coextinction curves (the relationship between

species losses in one trophic level and species losses in another [34]) using the

`networklevel()` and `second.extinct()` functions in the `bipartite` package [35].

To test for phylogenetic signal in floral symmetry, visitor species count, and visitor species

sharing, and to control for phylogeny in subsequent analysis, we mapped taxa in our dataset to a

recently published time-calibrated supertree of the seed plants (the "ALLMB" supertree of [36]),

using the `congeneric.merge()` function from the package `pez` [37] to add species to the

tree if they were not already included. We tested for phylogenetic signal using the

`phylosignal` package [38], estimating Blomberg's K and K* [39], Abouheif's C_{mean} [40],

Moran's I [41], and Pagel's λ [42] statistics. We performed principal component analysis of the

phylogenetic distance matrix for all plant taxa and used phylogenetic distance principal

components as covariates in models fitted to explain variation in visitor count and sharing.

We tested the hypotheses that visitor species count and visitor species sharing differed with

respect to floral symmetry by fitting Bayesian multilevel regression models using the `brms`

package [43,44]. Competing models explained visitor count and sharing with a group effect

(analogous to a random effect in a ML framework) of source network identity and possible

population (i.e., fixed) effects of floral symmetry, latitude, and the first two principal components

of phylogenetic distance (which jointly explained 61% of variation). We compared model fit in

terms of expected log pointwise predictive density (ELPD) using leave-one-out cross-validation,

implemented in `brms` with the `LOO()` function [45].

124
Figure 1. The global plant visitation dataset. (A) Global distribution of the plant-visitor
networks, with points scaled to indicate numbers of plant taxa. (B) Bar plots giving the number
of plant taxa for each network, coloured according to floral symmetry (left, full dataset, right, all
networks except for the largest, to provide better visibility. (C) Visitor species per plant, grouped
by floral symmetry. (D) Sharing of visitor species with all co-occurring plant species, and co-
occurring species with each type of floral symmetry. (E) Mean visitor species per plant in sub-
networks for plants with differing symmetries, with grey lines linking points for zygomorphic
and actinomorphic taxa from the same network, plotted against latitude of the network location.
In (C) and (D), asterisks indicate significant differences in one-tailed Wilcoxon tests: * $p < 0.01$,
** $p < 10^{-4}$, *** $p < 10^{-5}$.

RESULTS

We compiled 159 networks, recording 53,609 observed visits to 2,700 angiosperm species

(Figure 1A; Supplementary Table 1). We were able to classify floral symmetry for 2,685 plant

species, and were able to place 2,582 of these in the time-calibrated supertree [36]. Globally, and

in individual networks, zygomorphic flowers were a minority: 498 species (18%) were

zygomorphic; only five networks had more zygomorphic than actinomorphic species, while 70

lacked any (Figure 1B). Globally, the number of visitor species to zygomorphic flowers was

significantly smaller than that for actinomorphic flowers (median 4 pollinators per zygomorphic

species with zygomorphic flowers versus 5 per actinomorphic species; $p = 0.003$, one-tailed

Wilcoxon test; Figure 1C). We found significant phylogenetic signal for floral symmetry ($C_{mean} =$

0.85, Moran's $I = 0.10$, $K = 0.20$, $K^* = 0.22$, and $\lambda = 0.89$; $p < 0.001$ for all); visitor species

count deviated from the null models for C_{mean}, I, and λ ($p < 0.001$ for each) but not for K and K^*.

At the same time, zygomorphic flowers shared a higher proportion of their visitor species with

co-occurring plants (median 0.20 for zygomorphic versus 0.18 for actinomorphic, $p = 0.003$;

Figure 1D). Plants also had greater sharing with co-occurring plants of the same floral symmetry

(actinomorphic species, median sharing of 0.24 with actinomorphic species versus 0.19 with

zygomorphic species, $p = 0.003$; zygomorphic species, median sharing of 0.30 with zygomorphic

species versus 0.27 with actinomorphic species; $p < 10^{-5}$; Figure 1D).

Thirty-nine networks comprising 46,026 visitation records included enough plants of each

symmetry type (at least five) to compare sub-networks based on symmetry — that is, to compare

networks for plants with different symmetry having access to the same pool of possible visitor

Figure 2. Floral symmetry and plant-visitor network structure. (A) Network descriptive statistics for sub-networks based on floral symmetry. Grey lines link points representing zygomorphic and actinomorphic sub-networks from the same source network, horizontal black bars mark the median value within each floral symmetry type, and asterisks mark significant differences in one-tailed paired Wilcoxon tests. (B) Scatterplots of relationships between the proportion of zygomorphic flowers in a network and the set of network structure metrics in A; linear regression lines (white with grey 95% CI) are given when the correlation is significant with $p \le 0.05$.

species. Zygomorphic sub-networks had significantly fewer mean visitor species per plant (one-tailed paired Wilcoxon test, $p < 0.001$; Figure 2A), but only marginally greater visitor sharing ($p = 0.06$). Zygomorphic sub-networks had significantly greater connectance ($p < 10^{-4}$), greater asymmetry ($p < 0.001$), and lower coextinction robustness, as measured by the exponent of the coextinction curve, for both plants ($p < 10^{-5}$) and visitors ($p < 10^{-5}$). Among 89 complete

networks that included at least one zygomorphic plant, a higher proportion of zygomorphic

plants was associated with greater visitor sharing (Figure 2B; product-moment correlation on

log-transformed values = 0.22, p = 0.04) and greater connectance (cor = 0.21, p = 0.04),

paralleling the sub-network patterns. However, networks with more zygomorphic flowers also

had lower asymmetry (cor = -0.26, p = 0.01), and greater visitor coextinction robustness (cor =

0.27, p = 0.01), the opposite of patterns in symmetry-based sub-networks. The proportion of

zygomorphic flowers was not correlated with the mean number of visitors per plant or plant

coextinction robustness.

The best-fit model explaining visitor count included floral symmetry, latitude, an interaction

between symmetry and latitude, and phylogeny in addition to the group effect of network identity

(R^2 = 0.28; ΔELPD $\geq$ 3.8 versus all other models; Supplementary Tables 2, 3). For visitor

sharing, the best-fit model also included all terms; but all simpler models that included an effect

of symmetry had ΔELPD $\leq$ 0.6 (R^2 = 0.66 for all such models; Supplementary Tables 4, 5).

DISCUSSION

Zygomorphic flowers have long been considered to be more specialized, which could mean that

they are visited by fewer pollinator species. We find that, globally and at the level of individual

communities, plants with zygomorphic flowers do indeed have fewer visitor species, and that the

visitation networks of plants with zygomorphic symmetry may be structured by this difference

(Figure 1C,D; Figure 2A). Sub-networks of plants with zygomorphic flowers show greater

connectance, greater asymmetry, and lower coextinction robustness for both plants and visitors

(Figure 2A); however, these patterns do not necessarily translate to whole networks (Figure 2B).

Visitor species count is correlated with latitude north or south (Figure 1E), and both floral

symmetry and visitor count show significant phylogenetic structure. Bayesian multilevel

regressions accounting for these confounding effects nevertheless find significant effects of floral

symmetry on visitor species count and sharing.

The visitation records we examine have limitations for assessing pollination specialization. Many

of the original studies do not evaluate visitors' pollen transfer, and often record visits as a binary,

while in reality floral visitors vary considerably in visitation frequency and effectiveness as

pollinators. Thus, the effective pollinators of the plants in our dataset are likely a subset of

recorded visitors. However, we think it unlikely that data restricted to effective pollinators would

reverse the qualitative patterns we see, because a reversal would require systematic bias based on

floral symmetry in recording visits.

Our findings that zygomorphic flowers share more visitor species make sense given the other

aspects of pollinator specialization associated with zygomorphy. Arithmetically, a plant with

fewer visitors can more easily share a high proportion of them even if the absolute number of

visitors shared is low, but higher sharing by zygomorphic species probably also reflects network

structure. Plant-visitation networks are generally nested [30], so the fact that zygomorphic

species tend to have fewer visitor species means that their visitors are more likely to interact with

many other plant taxa. Zygomorphic flowers, which often manipulate pollinator behaviour, apply

pollen to specific parts of pollinators' bodies, or attract pollinators that show greater constancy

over a single foraging trip, may be better able to tolerate elevated risk of receiving heterospecific

pollen due to higher visitor sharing [1,20]. The fact that the highest rate of sharing we see in our

data is between zygomorphic flowers and other co-occurring zygomorphic species is consistent

with this hypothesis.

In our compiled dataset, plants with both types of floral symmetry had more visitor species in

communities farther from the equator (Figure 1E). This could reflect bias in the resolution of

species in more tropical communities, as there are more undescribed species in the tropics, and in

the Global South [46]. Networks farther from the equator do have more plant and visitor species

recorded (cor = 0.54 for plants, cor = 0.65 for visitors, $p < 10^{-5}$ for both; see Figure 1A).

However, the correlation weakens or disappears if the comparison is with latitude rather than

distance from the equator (i.e., treating distance south of the equator as different from distance

north: cor = 0.3, p = 0.69 for plants; cor = 0.16, p = 0.05 for visitors), suggesting this is not

simply an issue of better taxonomy in the Global North. Moreover, if plant and visitor species

numbers show similar correlations with distance from the equator, we might expect that the

number of visitors *per* plant would be constant across latitudes. An alternative explanation is that

floral visitation is more specialized in the tropics, independent of floral symmetry. Testing this

hypothesis with greater rigor is beyond the scope of our data, but it would be consistent with

recent syntheses finding stronger effects of biological interactions in the tropics [23,24].

An ecological association between floral symmetry and pollinator diversity may explain

evolutionary associations, across the angiosperms, between floral zygomorphy and

diversification [17,18,47]. The direction of the causal relationship, however, remains ambiguous.

It may be that association with fewer pollinators creates more opportunities to evolve

reproductive isolation [1,20,47] or increases the number of plant species that can be supported by

a given community of pollinators [18]; or it may be that more specialized pollination evolves in

response to greater diversity as co-occurring species must subdivide available pollinators more

finely [27].

Finally, our results have important implications for conservation. We find lower coextinction

robustness for sub-networks of plants with zygomorphic flowers, which may be explained by

higher connectance and asymmetry of these sub-networks (Figure 2A) and higher visitor sharing

among zygomorphic flowers (Figure 1D). These patterns are not necessarily borne out at the

level of whole communities (Figure 2B), suggesting that larger communities may be robust to

disturbances that endanger their most specialized members. Nevertheless, our analysis does

imply that plant taxa with zygomorphic flowers are at greater risk of extinction due to pollinator

loss. Despite significant uncertainty in the magnitude of losses, pollinators are widely known to

be in rapid decline due to pesticide use, habitat degradation, and emerging infections [48,49].

The patterns we find are coarse, but simple rubrics for triage are critical for conserving the more

than 300,000 species of angiosperms, most of which will never benefit from individualized

conservation assessment. Perhaps more importantly, our results support the idea that

"compartments" of the global plant-pollinator network must be targets for holistic conservation,

focused on preserving interactions and functionality where the network is most fragile [50].

ACKNOWLEDGEMENTS

JBY was supported by start-up funds from California State University Northridge, and CJC by a

postdoctoral fellowship from the Georgetown Environment Initiative. We thank Shweta Bansal

for helpful conversations and Stephen Sondheim for teaching us that a faboid legume can begin

an adventure.

DATA, CODE, AND MATERIALS

Data including floral symmetry annotations, visitor counts and sharing, and network structure

statistics are available with scripts on Dryad, https://datadryad.org/10.5061/dryad.gxd2547j3

COMPETING INTERESTS

The authors declare no competing interests.

AUTHORS' CONTRIBUTIONS

JBY and CJC conceived and designed the study; GG annotated floral symmetry with supervision

from JBY, and JBY and GG conducted analysis with code provided by CJC; JBY drafted the

paper in consultation with CJC and GG. All gave final approval for publication and agree to be

held accountable for the work performed.

REFERENCES

1.   Takhtajan A. 1969 *Flowering Plants: Origin and Dispersal*. Edinburgh: Oliver & Boyd.

2.   Darwin C. 1877 *The Various Contrivances by Which Orchids are Fertilised by Insects*.

2nd. New York: D. Appleton and Company.

3.   Stebbins GL. 1974 *Flowering Plants: Evolution Above the Species Level*. Belknap Press,

Harvard.

4.   Endress PK. 2012 The Immense Diversity of Floral Monosymmetry and Asymmetry

Across Angiosperms. *Bot. Rev.* **78**, 345–397. (doi:10.1007/s12229-012-9106-3)

5.   Fenster CB, Armbruster WS, Dudash MR, Wilson P, Fenster CCB, Thomson JD. 2004

Pollination syndromes and floral specialization. *Annu. Rev. Ecol. Evol. Syst.* **35**, 375–403.

(doi:10.1146/annurev.ecolsys.34.011802.132347)

6.   Rosas-Guerrero V, Aguilar R, Martén-Rodríguez S, Ashworth L, Lopezaraiza-Mikel M,

Bastida JM, Quesada M. 2014 A quantitative review of pollination syndromes: do floral

traits predict effective pollinators? *Ecol. Lett.* **17**, 388–400. (doi:10.1111/ele.12224)

7.   Ashworth L, Aguilar R, Martén-Rodríguez S, Lopezaraiza-Mikel M, Avila-Sakar G,

Rosas-Guerrero V, Quesada M. 2015 Pollination syndromes: A global pattern of

convergent evolution driven by the most effective pollinator. In *Evolutionary Biology:*

Biodiversification from Genotype to Phenotype, pp. 203–224. Cham: Springer

International Publishing. (doi:10.1007/978-3-319-19932-0_11)

8.   Poisot T, Bever JD, Nemri A, Thrall PH, Hochberg ME. 2011 A conceptual framework for

the evolution of ecological specialisation. *Ecol. Lett.* **14**, 841–851. (doi:10.1111/j.1461-

0248.2011.01645.x)

9.   Citerne H, Jabbour F, Nadot S, Damerval C. 2010 The evolution of floral symmetry. *Adv.*

Bot. Res. **54**, 85–137. (doi:10.1016/S0065-2296(10)54003-5)

10.   Sauquet H *et al.* 2017 The ancestral flower of angiosperms and its early diversification.

Nat. Commun. **8**, 16047. (doi:10.1038/ncomms16047)

11.   Reyes E, Sauquet H, Nadot S. 2016 Perianth symmetry changed at least 199 times in

angiosperm evolution. *Taxon* **65**, 945–964. (doi:10.12705/655.1)

12.   Fenster CB, Armbruster WS, Dudash MR. 2009 Specialization of flowers: is floral

orientation an overlooked first step? *New Phytol.* **183**, 502–506.

13. Kampny CM. 1995 Pollination and flower diversity in the Scrophulariaceae. *Bot. Rev.* **61**,
350–366.
14. Ushimaru A, Dohzono I, Takami Y, Hyodo F. 2009 Flower orientation enhances pollen
transfer in bilaterally symmetrical flowers. *Oecologia* **160**, 667–674.
(doi:10.1007/s00442-009-1334-9)
15. Macior LW. 2006 Behavioral aspects of coadaptations between flowers and insect
pollinators. *Ann. Missouri Bot. Gard.* **61**, 760. (doi:10.2307/2395027)
16. Takhtajan A. 1991 *Evolutionary Trends in Flowering Plants*. New York: Columbia
University Press.
17. Sargent RD. 2004 Floral symmetry affects speciation rates in angiosperms. *Proc. R. Soc. B*
271, 603–608.
18. Vamosi JC, Vamosi SM. 2010 Key innovations within a geographical context in flowering
plants: Towards resolving Darwin's abominable mystery. *Ecol. Lett.* **13**, 1270–1279.
(doi:10.1111/j.1461-0248.2010.01521.x)
19. O'Meara BC *et al.* 2016 Non-equilibrium dynamics and floral trait interactions shape
extant angiosperm diversity. *Proc. R. Soc. B Biol. Sci.* **283**. (doi:10.1098/rspb.2015.2304)
20. Grant V. 1949 Pollination systems as isolating mechanisms in angiosperms. *Evolution (N.*
Y). **3**, 82–97.
21. Mittelbach GG *et al.* 2007 Evolution and the latitudinal diversity gradient: speciation,
extinction and biogeography. *Ecol. Lett.* **10**, 315–331. (doi:doi:10.1111/j.1461-
0248.2007.01020.x)
22. Jansson R, Davies TJ. 2008 Global variation in diversification rates of flowering plants:
Energy vs. climate change. *Ecol. Lett.* **11**, 173–183. (doi:10.1111/j.1461-

333   0248.2007.01138.x)

334 23. Briscoe Runquist R *et al.* 2020 The relative role of abiotic and biotic environments in

335   local adaptation: A quantitative and qualitative synthesis. *Am. Nat.* (doi:10.1086/707322)

336 24. Hargreaves AL, Germain RM, Bontrager M, Persi J, Amy L. 2020 Local adaptation to

337   biotic interactions: A meta-analysis across latitudes. *Am. Nat.* (doi:10.1086/707323)

338 25. Olesen JM, Jordano P. 2002 Geographic patterns in plant-pollinator mutualistic networks.

339   *Ecology* **83**, 2416–2424.

340 26. Ollerton J, Cranmer L. 2002 Latitudinal trends in plant-pollinator interactions: are tropical

341   plants more specialised? *Oikos* **98**, 340–350.

342 27. Armbruster WS, Muchhala N. 2009 Associations between floral specialization and species

343   diversity: Cause, effect, or correlation? *Evol. Ecol.* **23**, 159–179. (doi:10.1007/s10682-

344   008-9259-z)

345 28. Olesen JM, Bascompte J, Dupont YL, Jordano P. 2007 The modularity of pollination

346   networks. *Proc. Natl. Acad. Sci.* **104**, 19891–19896.

347 29. Krishna A, Guimarães Jr PR, Jordano P, Bascompte J. 2008 A neutral-niche theory of

348   nestedness in mutualistic networks. *OIKOS*

349 30. Bascompte J, Jordano P, Olesen JM, Melián CJ, Olesen JM. 2003 The nested assembly of

350   plant–animal mutualistic networks. *Proc. Natl. Acad. Sci.* **100**, 9383–9387.

351   (doi:10.1073/pnas.1633576100)

352 31. Nuismer SL, Week B, Aizen MA. 2018 Coevolution Slows the Disassembly of Mutualistic

353   Networks. *Am. Nat.* **192**, 490–502. (doi:10.1086/699218)

354 32. R Core Team. 2020 R: A language and environment for statistical computing.

355 33. Dunne JA, Williams RJ, Martinez ND. 2002 Food-web structure and network theory: The

role of connectance and size. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 12917–12922.

(doi:10.1073/pnas.192407699)

34.   Memmott J, Waser N, Price M. 2004 Tolerance of pollination networks to species

extinctions. *Proc. R. Soc. B Biol. Sci.* **271**, 2605–2611.

35.   Dormann CF, Gruber B, Fründ J. 2008 Introducing the bipartite Package : Analysing

Ecological Networks. **8**, 8–11.

36.   Smith SA, Brown JW. 2018 Constructing a broadly inclusive seed plant phylogeny. *Am. J.*

Bot. **105**, 302–314. (doi:10.1002/ajb2.1019)

37.   Pearse WD, Cadotte MW, Cavender-Bares J, Ives AR, Tucker CM, Walker SC, Helmus

MR. 2015 pez: phylogenetics for the environmental sciences. *Bioinformatics* **31**, 2888–

2890. (doi:10.1093/bioinformatics/btv277)

38.   Keck F, Rimet F, Bouchez A, Franc A. 2016 Phylosignal: An R package to measure, test,

and explore the phylogenetic signal. *Ecol. Evol.* **6**, 2774–2780. (doi:10.1002/ece3.2051)

39.   Blomberg SP, Garland T, Ives AR. 2003 Testing for phylogenetic signal in comparative

data: Behavioral traits are more labile. *Evolution (N. Y).* **57**, 717–745. (doi:10.1111/j.0014-

3820.2003.tb00285.x)

40.   Abouheif E. 1999 A method for testing the assumption of phylogenetic independence in

comparative data. *Evol. Ecol. Res.* **1**, 895–909.

41.   Gittleman JL, Kot M. 1990 Adaptation: Statistics and a null model for estimating

phylogenetic effects. *Syst. Zool.* **39**, 227. (doi:10.2307/2992183)

42.   Pagel M. 1999 Inferring the historical patterns of biological evolution. *Nature* **401**, 877–

884. (doi:10.1038/44766)

43.   Bürkner P-C. 2017 brms: An R package for Bayesian multilevel models using Stan. *J.*

379   *Stat. Softw.* **80**, 1–28. (doi:10.18637/jss.v080.i01)

380 44. Bürkner P-C. 2018 Advanced Bayesian multilevel modeling with the R package brms. *R J.*

381   **10**, 395. (doi:10.32614/RJ-2018-017)

382 45. Vehtari A, Gelman A, Gabry J. 2017 Practical Bayesian model evaluation using leave-one-

383   out cross-validation and WAIC. *Stat. Comput.* **27**, 1413–1432. (doi:10.1007/s11222-016-

384   9696-4)

385 46. Scheffers BR, Joppa LN, Pimm SL, Laurance WF. 2012 What we know and don't know

386   about Earth's missing biodiversity. *Trends Ecol. Evol.* **27**, 501–510.

387   (doi:10.1016/j.tree.2012.05.008)

388 47. Kay KM, Sargent RD. 2009 The role of animal pollination in plant speciation: Integrating

389   ecology, geography, and genetics. *Annu. Rev. Ecol. Evol. Syst.* **40**, 637–656.

390   (doi:10.1146/annurev.ecolsys.110308.120310)

391 48. Lister BC, Garcia A. 2018 Climate-driven declines in arthropod abundance restructure a

392   rainforest food web. *Proc. Natl. Acad. Sci.* **115**, E10397–E10406.

393   (doi:10.1073/pnas.1722477115)

394 49. Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. 2010 Global

395   pollinator declines: Trends, impacts and drivers. *Trends Ecol. Evol.* **25**, 345–353.

396   (doi:10.1016/j.tree.2010.01.007)

397 50. Corbet SA. 2000 Conserving compartments in pollination webs. *Conserv. Biol.* **14**, 1229–

398   1231. (doi:10.1046/j.1523-1739.2000.00014.x)

399