Title Page

The maize *Hairy Sheath Frayed1* (*Hsf1*) mutant alters leaf patterning through
increased cytokinin signaling

Author names and affiliations

Michael G. Muszynski, [a,1] Lindsay Moss-Taylor, [b, 2] Sivanandan Chudalayandi, [b, 3]
James Cahill, [b, 4] Angel R. Del Valle-Echevarria, [a] Ignacio Alvarez-Castro, [c, 5] Abby
Petefish, [b] Hitoshi Sakakibara, [d] Dmitry M. Krivosheev, [e, 6], Sergey N. Lomin, [e] Georgy
A. Romanov, [e] Subbiah Thamotharan, [f] Bailin Li, [g] and Norbert Brugière [g]

[a] Department of Tropical Plant and Soil Sciences, University of Hawai□i at Mānoa,
Honolulu, HI 96822, USA.

[b] Department of Genetics, Development and Cell Biology, Iowa State University, Ames,
Iowa, 50011

[c] Department of Statistics, Iowa State University, Ames IA, 50011

[d] RIKEN Center for Sustainable Resource Science, Tsurumi, Yokohama 230-0045,
Japan.

[e] Timiryazev Institute of Plant Physiology, Russian Academy of Sciences, Moscow,
127276, Russia

[f] School of Chemical and Biotechnology, SASTRA University; Thanjavur, 613401, India

[g] Corteva Agriscience, Johnston, IA 50131

ORCID IDs: 0000-0002-0817-7594 (M.G.M.);

[1] Address correspondence to mgmuszyn@hawaii.edu.

[2] Present address: Department of Laboratory Medicine and Pathology, University of
Minnesota, Minneapolis, MN 55455, USA

[3] Genome Informatics Facility, Iowa State University, Ames, Iowa, 50011

[4] Present address: Corteva Agriscience, Johnston, IA 50131

[5] Present address: Instituto de Estadística, Universidad de la República, Montevideo,
Uruguay

[6] Present address: Vologda State University, Vologda 160000, Russia

Short Title: Cytokinin influences leaf development

39   Summary: Increased cytokinin signaling in the maize *Hairy Sheath Frayed1* mutant
40   modifies leaf development leading to changes in pattering, growth and cell identity.

ABSTRACT

Leaf morphogenesis requires growth polarized along three axes - proximal-distal, medial-lateral and abaxial-adaxial. Grass leaves display a prominent proximal-distal (P-D) polarity consisting of a proximal sheath separated from the distal blade by the auricle and ligule. Although proper specification of the four segments is essential for normal morphology, our knowledge is incomplete regarding the mechanisms which influence P-D specification in monocots like maize (*Zea mays*). Here we report the identification of the gene underlying the semi-dominant, leaf patterning, maize mutant *Hairy Sheath Frayed1* (*Hsf1*). *Hsf1* plants produce leaves with outgrowths consisting of proximal segments – sheath, auricle and ligule – emanating from the distal blade margin. Analysis of three independent *Hsf1* alleles revealed gain-of-function missense mutations in the ligand binding domain of the maize cytokinin (CK) receptor *Zea mays Histidine Kinase1* (*ZmHK1)* gene. Biochemical analysis and structural modeling suggest the mutated residues near the CK binding pocket affect CK binding affinity. Treatment of wild type seedlings with exogenous CK phenocopied the *Hsf1* leaf phenotypes. Results from expression and epistatic analyses indicated the *Hsf1* mutant receptor appears to be hypersignaling. Our results demonstrate that hypersignaling of CK in incipient leaf primordia can reprogram developmental patterns in maize.

INTRODUCTION

Proper leaf morphogenesis in higher plants requires defined patterns of growth polarized along three axes: adaxial-abaxial, medial-lateral and proximal-distal (McConnell and Barton, 1998; Tsukaya, 1998; Bowman et al., 2002; Byrne et al., 2002). Growth along the proximal-distal (P-D) axis is particularly evident in grass leaves, like maize, which are composed of four distinct segments; the sheath is proximal, the blade

is distal and the auricle and ligule form the boundary between the two (Figure 1A)

(Sylvester et al., 1996). A number of genes have been identified that influence P-D

patterning, with *BLADE-ON-PETIOLE* (*BOP*) genes affecting proximal identity in

eudicots and monocots (Ha et al., 2003, 2004; Norberg et al., 2005; Toriba et al., 2019;

Moon et al., 2013; Tavakol et al., 2015). In grasses, ectopic expression of class I

knotted1like homeobox (*knox*) transcription factor genes in developing leaf primordia

alters P-D patterning, primarily disrupting the formation of a defined sheath-blade

boundary (Freeling and Hake, 1985; Hake et al., 1989, 1991; Smith et al., 1992;

Schneeberger et al., 1995; Muehlbauer et al., 1997; Foster et al., 1999a; Tsiantis et al.,

1999; Byrne et al., 2001). Class I *knox* genes typically function in meristem formation

and maintenance, and their down-regulation is required for normal development of

determinant organs like leaves (Endrizzi et al., 1996; Long et al., 1996; Kerstetter et al.,

1994). In meristems, KNOX proteins function to increase cytokinin (CK) accumulation

by positive regulation of CK synthesis genes and simultaneously decrease gibberellic

acid (GA) accumulation by suppression of GA biosynthesis genes or activation of GA

catabolic genes (Ori et al., 2000; Sakamoto et al., 2001; Hay et al., 2002; Jasinski et al.,

2005; Yanai et al., 2005; Sakamoto et al., 2006; Bolduc and Hake, 2009). In addition, a

rice KNOX transcription factor was shown to also affect brassinosteroid (BR)

accumulation by upregulating BR catabolism in the shoot apical meristem (Tsuda et al.,

2014). Determinate leaf primordia form when *knox* expression is down-regulated by the

action of ROUGH SHEATH2 (RS2) and related proteins resulting in a decrease in CK

and increase in GA accumulation (Hay et al., 2006; Guo et al., 2008). In addition to the

action of CK and GA, auxin is required for proper leaf initiation and positioning. The

polar transport of auxin by PINFORMED1 (PIN1) auxin efflux carriers guides the

formation of auxin maxima, localized regions of high auxin accumulation, that is

required for initiation of leaf primordia (Pozzi et al., 2001; Scarpella et al., 2006;

Benjamins and Scheres, 2008; Zhao, 2008). The emerging model predicts that spatial

differences in cytokinin/auxin ratios control final cell fate (Shani et al., 2006; Muller and

Sheen, 2008). Ectopic *knox* expression presumably shifts critical phytohormone ratios

in developing leaf primordia but the exact molecular mechanisms by which

phytohormone ratios determine leaf patterning remain incomplete.

As phytohormones play pivotal roles in many developmental programs, the pathways that signal their perception and response have been well characterized. For example, the perception and response to the CK phytohormones relies on a two-component signal transduction system (Hwang and Sheen, 2001; Yonekura-Sakakibara et al., 2004; Hwang and Sakakibara, 2006; Du et al., 2007; To and Kieber, 2008). The perception of CK is mediated via a partially redundant signaling system of histidine kinases (HKs), histidine phosphotransfer proteins (HPTs) and response regulators (RRs). CK signaling begins with the perception of CK by binding to HK receptors at the ER, and probably also plasma membrane, which triggers receptor phosphorylation (Lomin et al., 2011). The activated receptors initiate phosphorelay by transferring phosphoryl groups to HPTs, which shuttle between the cytoplasm and nucleus. Once in the nucleus, phosphorylated HPTs transfer their phosphoryl groups to type-B RRs, which in turn activate expression of type-A RRs and other CK responsive genes (Rashotte et al., 2006). The type-A RRs and other CK-responsive genes mediate several CK-regulated processes including shoot and root growth, de-etiolation, leaf expansion, root vascular development, senescence, and cytokinin homeostasis (To and Kieber, 2008). In maize, multiple members of each of the CK signaling components have been identified (Yonekura-Sakakibara et al., 2004). Maize has seven HKs (ZmHKs), of which, three have been shown to bind and signal various types of CKs in heterologous assays (Lomin et al., 2011; Steklov et al., 2013). Three HPTs (ZmHPs), three type-B RRs and seven type-A RRs (ZmRRs) have also been identified in maize (Asakura et al., 2003). Of these signal transduction components, the function of only *ZmRR3*, a type-A RR, has been defined by null mutations and shown to underlie the *aberrant phyllotaxy1* (*abph1*) mutation (Jackson and Hake, 1999; Giulini et al., 2004). Our understanding of the functions of other components of the CK signal transduction pathway remains incomplete for cereal species like maize.

To gain a better understanding of the signaling mechanisms which mediate leaf pattern specification, we initiated a study of the semi-dominant *Hairy Sheath Frayed1* (*Hsf1*) mutation which alters P-D leaf development in maize (Bertrand-Garcia and Freeling, 1991a). Although *Hsf1* disrupts the P-D leaf pattern similar to dominant class I

knox mutations, *Hsf1* is not itself a *knox* gene, since it does not map to the location of

any maize *knox* genes (Bertrand-Garcia and Freeling, 1991b). In this report, we show

that the *Hsf1* phenotype results from specific missense mutations in the maize CK

receptor *Zea mays Histidine Kinase1* (*ZmHK1*) gene (Yonekura-Sakakibara et al.,

2004). Our analyses of mutant receptor function, the effects of exogenous CK

treatment on leaf development, and epistatic interaction suggest that the ZmHK1

receptor is hypersignaling in *Hsf1* mutants. Overall, our results indicate CK

hypersignaling can influence the specification of P-D leaf patterning in maize and

underscores the capacity of CK to alter developmental programs.

RESULTS

The *Hsf1* mutation induces specific alterations to maize leaf patterning

The original *Hsf1* mutation arose via ethyl methanesulfonate (EMS) mutagenesis of the

inbred line Mo17 and was designated *Hsf1-N1595* (also called *Hsf1-O*) (Bird and

Neuffer, 1985). A second mutation, *Hsf1-N1603* (hereafter called *Hsf1-1603*)*,* was

shown to be allelic or very closely linked (Bertrand-Garcia and Freeling, 1991a). We

isolated three additional alleles in independent EMS mutagenesis screens in different

inbred backgrounds: *Hsf1-AEWL* in A619, *Hsf1-2559* in Mo17, and *Hsf1-7322* in A632.

All *Hsf1* alleles have very similar phenotypes compared to the *Hsf1-N1595* (hereafter

called *Hsf1-1595*) reference mutation. As was shown previously for *Hsf1-1595*, plants

heterozygous for any of the *Hsf1* alleles display a highly penetrant mutant leaf

patterning phenotype with outgrowths consisting of sheath, auricle and ligule emanating

from the distal blade margin (Figures 1A to 1C) (Bertrand-Garcia and Freeling, 1991a).

The outgrowths have proximal identity and were termed "prongs", which we adopted to

describe this structure (Figure 1B). Although *Hsf1* mutant plants have proximal tissue

growing on the distal blade, they have a normal blade-sheath boundary (Figure 1A)

(Bertrand-Garcia and Freeling, 1991a). All the pleiotropic phenotypes described for

Hsf1-1595 in Bertrand-Garcia and Freeling (1991a) are shared by all the other *Hsf1*

alleles, including an increase in macrohair size and density on the abaxial sheath,

adaxial blade, and blade margin, an increase in leaf number, shorter stature, short and

narrow leaves, and reduced root growth (Supplemental Figures 1A to 1B; Supplemental

Table 1). Bertrand-Garcia and Freeling (1991a) also showed homozygous *Hsf1* plants

have a stronger mutant phenotype being extremely stunted, with multiple shoots arising

from the coleoptile node at germination, and having adventitious needle- or club-shaped

leaves (Supplemental Figures 1A to 1B).

Since plants heterozygous for ether *Hsf1-1595*, *Hsf1-1603*, or *Hsf1-AEWL* were

phenotypically very similar (Figure 1C), we chose the *Hsf1-1603* allele to characterize

the temporal and spatial patterns of prong formation to better understand how the *Hsf1*

mutation affected leaf patterning. In *Hsf1-1603* heterozygotes, prongs first appeared on

leaf 5 in a few plants, and most commonly appeared on leaf 6 or leaf 7 but never on

earlier arising leaves (Supplemental Figure 1C). The earliest sign of P-D leaf polarity

specification is the formation of the preligule band (PLB) which will differentiate into the

auricle and ligule (Sylvester et al., 1990; Johnston et al., 2014). Formation of the PLB

typically is first observed in plastochron 5 or 6 stage leaf primordia (P5 - P6) with the

initiating ligule becoming visible about plastochron 7 or 8 (P7 - P8) (Johnston et al.,

2014). Plastochron describes the stage of leaf primordia development and refers to the

position of the primordia relative to the shoot apical meristem (SAM) (Sylvester et al.,

1990). Thus, a P5 primordium has four younger primordia between it and the SAM. To

determine if the *Hsf1-1603* mutation influenced the timing of the acquisition of P-D

polarity, we examined leaf primordia in *Hsf1-1603/+* and wild type sib plants for signs of

early ligule development (see Methods). The initiating ligule was most commonly first

visible on P7 primordia in both wild type and *Hsf1-1603* heterozygotes indicating no

influence on P-D polarity acquisition (Supplemental Figure 1E). To determine if the

appearance of prong primordia on the blade margin coincided with the acquisition of P-

D polarity, developing leaf primordia from *Hsf1-1603* heterozygotes were dissected and

examined for the presence of initiating prongs. Prong initials were most commonly

observed on the blade margins of P5 or P6 leaf primordia but some were noted as early

as P4 (Supplemental Figure 1D, 1F to 1G), consistent with prong formation in *Hsf1-*

1595 heterozygotes (Bertrand-Garcia and Freeling, 1991b). Thus prongs typically

initiated from blade margins about the same plastochron stage as formation of the PLB.

Prongs were observed to occur in different sizes and at different positions along

the leaf blade margin (Figures 1B and 1F, and Supplemental Figure 2A). To determine

if prong formation was random or patterned, we measured the size and positon of each

prong from both margins of mature leaves collected from different positions on the shoot

of *Hsf1-1595*, *Hsf1-1603* and *Hsf1-AEWL* heterozygous plants. Results showed that

prong formation was more frequent on leaves higher on the shoot (Supplemental Table

2) with prongs occupying more of the blade margin in these upper leaves compared to

lower leaves (Supplemental Figures 2B and 2C). Next we determined where prongs

formed along the P-D axis of the blade. Analysis indicated prongs only formed in the

proximal 70% and never in the distal 30% of the blade, with the majority of prongs

forming within a region encompassing the proximal 15% to 40% of the blade

(Supplemental Figure 2D). Next we examined the range of prong sizes for each *Hsf1*

allele within this prong-forming region. For all three alleles, the majority of prongs were

about a centimeter in size but a few were larger, ranging from 3 to 6 centimeters

(Supplemental Figure 2E). With relative position and size known, we next asked

whether prong position was related to its size. In general, the largest prongs often

formed in the basal 20% of the blade and smaller prongs formed at any position within

the prong forming region (Supplemental Figure 2F). Thus, our analysis indicated prong

formation was not random but occurred in particular regions of the blade and initiated at

specific developmental stages.

Gain-of-function mutations in the maize cytokinin receptor gene *ZmHK1* underlie

the *Hsf1* mutation

Previous studies mapped *Hsf1-1595* to the long arm of chromosome 5 (Bertrand-Garcia

and Freeling, 1991b). To isolate the gene underlying the *Hsf1* locus, we screened a

backcross mapping population of over 3,000 plants with linked molecular markers

derived from the maize reference genome (B73 RefGen_v1). The *Hsf1* locus was localized to a 21-kb interval with a single gene model (GRMZM2G151223, B73 RefGen_v2). This gene model was well supported with abundant EST evidence and was annotated as encoding *Zea mays Histidine Kinase1* (*ZmHK1*), one of seven maize histidine kinase cytokinin receptors (Yonekura-Sakakibara et al., 2004; Steklov et al., 2013). To confirm *ZmHK1* was the correct gene and to identify the causative lesions, the *ZmHK1* gene was sequenced from all five *Hsf1* alleles. The entire *ZmHK1* genomic region, including ca. 2-kb upstream and downstream of the transcription start and stop, was sequenced from *Hsf1-1595*, *Hsf1-1603, Hsf1-2559, Hsf1-7322* and *Hsf1-AEWL* homozygotes and their progenitor inbred lines. As expected for EMS-generated mutations, single nucleotide transitions were identified in the five *Hsf1* alleles compared to their progenitor sequences. Although each allele arose independently, *Hsf1-1595* and *Hsf1-1603* had the exact same transition mutations as *Hsf1-7322* and *Hsf1-2559*, respectively. Thus, hereafter, we refer to the three different *Hsf1* alleles: *Hsf1-1595*, *Hsf1-1603* and *Hsf1-AEWL*. Each transition mutation produced a missense mutation in a highly conserved amino acid located in the CHASE (<u>c</u>yclases/<u>h</u>istidine-kinase-<u>a</u>ssociated <u>se</u>nsory) domain of the ZmHK1 protein, where CK binding occurs (Figure 1D) (Hothorn et al., 2011; Steklov et al., 2013). The *Hsf1-1595* mutation changed proline 190 to leucine (CCA>CTA), the *Hsf1-1603* mutation changed glutamate 236 to lysine (GAG>AAG), and the *Hsf1-AEWL* mutation changed leucine 238 to phenylalanine (CTT>TTT). The missense mutation in *Hsf1-AEWL* is particularly significant because this is the same type of amino acid substitution, although at a slightly different position in the CHASE domain, which was found in another gain-of-function mutation in a CK receptor, the *spontaneous nodule formation2* (*snf2)* mutation in the lotus *Lhk1* receptor, (Figure 1D) (Tirichine et al., 2007). The *snf2* mutation was shown to cause mutant LHK1 to signal independent of the CK ligand in a heterologous signaling assay suggesting the *snf2* mutation locked LHK1 in an active signaling state (Tirichine et al., 2007). Based on the location and nature of the amino acid substitutions in the three *Hsf1* mutations and the presumed mode of action of the *snf2* mutation in *Lhk1*, we hypothesized that the *Hsf1* mutations might also lock the ZmHK1 receptor in an active CK signaling state and signal independent of CK.

The *Hsf1* mutant CK receptors have altered histidine kinase signaling and ligand binding activities

To determine if the *Hsf1* mutant receptors are signaling independent of CK, we utilized a heterologous histidine kinase signaling assay system developed in the yeast *Saccharomyces cerevisae* (Suzuki et al., 2001). In the yeast assay, the cognate his-kinase of an endogenous two-component phosphorelay signal transduction system was deleted. Functional replacement of the endogenous his-kinase with the assayed CK receptor, in this case ZmHK1, allowed the activity of the receptor to be determined as the output of the endogenous yeast transduction system, which is the ability to grow on glucose media (Suzuki et al., 2001). We engineered the exact point mutation found in each *Hsf1* mutation into the *ZmHK1* cDNA in the p415CYC-ZmHK1 plasmid for expression in yeast (Suzuki et al., 2002; Higuchi et al., 2009). We next tested receptor activity in the *sln1* deletion yeast strain TM182 carrying each of the *Hsf1* missense mutations, the wild type *ZmHK1* cDNA, and the empty p415CYC vector grown on glucose media with and without the CK ligand (Figure 1E). As expected, the wild type ZmHK1 strain only grew well on glucose media supplemented with higher concentrations of the three CKs tested (Figure 1E) and, at lower CK concentrations, only grew robustly on glucose with the preferred ligand N^6-(Δ^2-isopentenyl) adenine (iP) (Supplemental Figures 3A to 3C). In the absence of added CK, strains carrying either ZmHK1-AEWL or ZmHK1-1603 grew robustly on glucose media (Figure 1E). This result indicated that the ZmHK1-AEWL and ZmHK1-1603 receptors signaled independent of added CK in this assay. To determine if the mutant receptors were still CK responsive, they were also grown on glucose media supplemented with the three tested CKs (Figure 1E). Growth on glucose supplemented with different CKs did not reveal any receptor activity differences between ZmHK1-AEWL and ZmHK1-1603. Surprisingly, growth of the ZmHK1-1595 strain was different than the other two mutant receptors and wild type. The ZmHK1-1595 strain did not grow on glucose media without added CK, similar to wild type ZmHK1 (Figure 1E). Instead, the ZmHK1-1595 strain showed strong growth on glucose media with 10 µM of the preferred CK iP and

weaker growth on glucose with 10 µM of two other bioactive CKs, *trans*-zeatin (tZ) and

cis-zeatin (cZ), suggesting ZmHK1-1595 had weak receptor activity in this assay (Figure

1E

To investigate ligand specificity differences, CK ligand binding affinities were

determined for the mutant and wild type receptors (Romanov et al., 2005; Lomin et al.,

2011). Affinities were determined for 6 different CKs and adenine (Ade) using two

binding assays with receptors expressed in bacterial spheroplasts (Romanov et al.,

2005) or residing in tobacco membranes after transient expression *in planta* (Lomin et

al., 2015, 2011). The ligand preferences for the wild type ZmHK1 receptor were

comparable to those determined previously (Table 1) (Lomin et al., 2015, 2011). The

mutant receptors, on the other hand, all showed increased affinities for most of the CKs

tested (Table 1). The preference ranking of the mutant receptors for different CKs was

mostly similar to wild type (Supplemental Figure 4) but the affinities were increased

between 2- to 8-fold (Table 2). The only exception was the affinity for the synthetic CK

thidiazuron, which was reduced for all the mutant receptors compared to wild type

ZmHK1. Thus, the missense mutations in the *Hsf1* alleles increased the relative binding

affinity of the receptor for all the natural CKs tested, suggesting the mutant receptors

might be hypersignaling.

The *Hsf1* missense mutations localize near the CK binding pocket in ZmHK1

To gain better insight into how each *Hsf1* missense mutation might impact CK binding,

we determined the effect these mutations had on the structure of the CHASE domain,

which was facilitated by the publication of the crystal structure of the *Arabidopsis*

thaliana histidine kinase4 (*AHK4*) gene CHASE domain (Hothorn et al., 2011). *AHK4* is

co-orthologous to *ZmHK1* (69% identical and 83% similar within 245 residues of the

CHASE domain) and three other paralogous histidine kinases in the maize genome

(Steklov et al., 2013). To explore the effects of the *Hsf1* mutations on receptor

structure, homology modeling was used first to model the 3D structure of the CHASE

domain of ZmHK1 using the structure of AHK4. This was done with and without CK

occupying the binding pocket, which did not change the results. Given the high degree

of amino acid identity between ZmHK1 and AHK4, the ZmHK1 CHASE domain

structure was resolved with high confidence. Next, each mutant receptor was modeled

based on the derived ZmHK1 structure. The models were subjected to dynamics

simulation with appropriate solvation (see Methods). The results of homology modeling

showed that the amino acids mutated in each *Hsf1* allele do not occur within the CK

binding pocket (Figure 1F) and thus do not contribute to direct polar contacts with the

ligand. Instead, each altered residue is located near a loop domain that forms one face

of the binding cavity. An indication of how the mutated residues at these positions

might affect ligand binding was provided by the structure model of the ZmHK1-1603

receptor. The residue altered in ZmHK1-1603 is E236, which is predicted to form an

ion-pair interaction with R192 located in the loop domain. This polar interaction may

help to stabilize the position of the loop domain (Figure 1F). The *Hsf1-1603* mutation

converts E236 to K, a negative to positive residue change, which is expected to break

the polar interaction with R192 and possibly destabilize the position of the loop due to

the nearness of the two positively charged residues. Altering the position of the loop

may change the overall conformation of the ligand binding pocket and, thus, account for

differences in ligand binding affinities. The missense residues in the other two mutant

receptors could potentially alter the conformation of the CK binding pocket via a

different mechanism, although our modeling results did not reveal an obvious one.

Exogenous CK treatment recapitulated the *Hsf1* phenotype

The biochemical and structural analyses suggested the *Hsf1* mutant receptor might be

hypersignaling the perception of CK which altered leaf development. To test the idea

that increased CK signaling could produce *Hsf1*-like phenotypes, wild type, B73 inbred

seeds were transiently treated with the CK 6-benzylaminopurine (6-BAP). The embryo

in a mature maize seed possesses about 5 leaf primordia and it is these primordia

which experienced the hormone treatment (Kerstetter and Poethig, 1998). Imbibed

seeds were treated for 6 days with 10 µM 6-BAP, rinsed and transplanted to soil (see

Methods). After growth for 3-weeks, the first 4 seedling leaves were examined for developmental changes (Figures 2A to 2E). Similar to *Hsf1*, 100% of the CK treated B73 seeds produced smaller seedling leaves covered with abundant macrohairs (Figures 2A to 2E). Leaf sheath length, blade length and blade width were reduced by 10% to 20% for leaf 3, similar to leaf size reductions in the *Hsf1* seedlings (Figure 2C). In addition, macrohair density increased on the abaxial sheath, near the auricle, on the adaxial blade, and blade margins in 100% of the CK-treated B73 seedlings (Figures 2D to 2E). This pattern of ectopic macrohair formation was similar to that seen in *Hsf1* seedlings (Bertrand-Garcia and Freeling, 1991a). In addition to alterations in leaf size and pubescence, nearly 20% of the CK treated B73 seeds produced seedlings with prongs on leaf 4 (Figures 2F). This was in contrast to *Hsf1* seedlings where prongs rarely, if ever, developed on leaf 4 (Supplemental Figure 1C). Increasing the concentration of exogenous 6-BAP to 100 µM increased the number of B73 seedlings with prongs on leaf 4 to nearly 90% (Figure 2F) Thus, transient, exogenous CK treatment recapitulated three prominent aspects of the *Hsf1* phenotype: reduced leaf size, increased macrohair abundance, and formation of prongs on blade margins, confirming these developmental changes can be induced by CK.

If CK hypersignaling in *Hsf1* was due to increased ligand affinity, then we would expect *Hsf1* to be hypersensitive to CK treatment. To test this idea, we performed six-day treatments on segregating *Hsf1-1603/+* seeds using 0.1 µM CK, a concentration that did not elicit leaf size changes in B73 inbred seed (Supplemental Figure 5A). To distinguish segregating heterozygous *Hsf1* plants from wild type sib plants, PCR genotyping was used to detect a size polymorphism in the *Hsf1-1603* allele (Supplemental Table 3). After CK treatment, seedlings were grown for 3 weeks, after which, leaf phenotypes were measured. While 0.1 µM CK treatment had no effect on wild type sibling leaf size (Supplemental Figure 5A), it did reduce the leaf size of *Hsf1-1603/+* plants 10% to 30% (Supplemental Figure 5B). Thus, *Hsf1-1603/+* seedlings were responsive to a lower concentration of CK that did not elicit a response in wild type sib or B73 inbred seedlings. Treatment with 10 µM 6-BAP was also used to assess effects on prong and macrohair formation in *Hsf1-1603/+* plants. Similar to earlier

results (Supplemental Figure 1C), seedlings from control water-treated *Hsf1-1603/+*

seeds first formed prongs on leaf 5 (ca. 5%) or leaf 6 (ca. 25%) but never on earlier

arising leaves (Figure 2G to 2I). In fact, about 60% of *Hsf1-1603/+* seedlings normally

first formed prongs on leaves arising on or after leaf 7 (Figure 2H). By contrast, of the

10 μM 6-BAP treated *Hsf1-1603/+* seeds, nearly 60% produced seedlings where prongs

first formed on leaf 4 and only about 30% formed prongs on leaves arising on or after

leaf 7 (Figures 2G to 2I). In addition, macrohair abundance appeared increased for CK-

treated *Hsf1-1603/+* compared to control *Hsf1-1603/+* or 6-BAP treated wild type sib

seedlings but this was not measured (Figure 2J). Thus, CK treatment of *Hsf1* resulted

in earlier arising and enhanced mutant phenotypes, indicating the mutation was

hypersensitive to the CK hormone, consistent with the biochemical analysis of the

mutant receptor.

CK responsive genes are up-regulated in *Hsf1* leaf primordia

Based on the *Hsf1* mutant plant phenotypes, we presumed that hypersignaling in

developing leaf primordia gave rise to the alterations in P-D leaf patterning and other

phenotypes. To test this idea, we determined the expression of *ZmHK1* and several CK

responsive genes in *Hsf1-1603/+* and wild type sibling plants. Published qPCR and *in*

silico expression analyses

(https://www.maizegdb.org/gene_center/gene/Zm00001d017977#rnaseq) indicated

ZmHK1 was expressed broadly across several tissues including leaves, roots, stem,

and tassel (Yonekura-Sakakibara et al., 2004). We reverse transcribed cDNA from

three tissues, shoot apices (shoot apical meristem plus 3 youngest leaf primordia),

immature leaf, and mature green leaf from two-week old seedlings. Using quantitative

PCR (qPCR) we assessed expression in plants heterozygous for the three *Hsf1* alleles

compared to their wild type sibs (Figure 3A). We did not detect an increase in *ZmHK1*

transcript accumulation in the *Hsf1/+* mutants compared to their wild type controls.

Next, we examined expression of CK-responsive genes; two type-A response

regulators, *ZmRR3* and *ZmRR6,* and a cytokinin oxidase gene, *ZmCKO4b* (Asakura et

al., 2003; Giulini et al., 2004). We found increased transcript accumulation for all three CK-responsive genes in the *Hsf1/+* mutants, although there was some inconsistencies across genotypes and tissues (Figure 3A).

Using *in situ* hybridization, we assessed transcript localization of *ZmHK1* and *ZmRR3* in wild type and *Hsf1-1603/+* shoot apices (Figure 3B). The *ZmHK1* transcript was found to be distributed broadly within developing leaf primordia and shoot apices in both genotypes (Figure 3B). As was demonstrated previously, *ZmRR3* was expressed in a specific wedge-shaped domain in the apical meristem in both longitudinal and transverse sections of wild type apices but no signal was detected in leaf primordia (Figure 3B) (Giulini et al., 2004). However, the spatial expression of *ZmRR3* was expanded in *Hsf1-1603/+* apices. Strong *ZmRR3* expression was visible in its normal meristem domain but signal was also detected in leaf primordia and was particularly evident at the margins (Figure 3B). Given the expanded pattern of *ZmRR3* expression in *Hsf1-1603/+* leaf primordia margins and that *ZmRR3* expression is CK responsive, we interpreted this to indicate increased CK signaling in the tissue where prongs will form.

Mutation of *ZmRR3*, a negative regulator of CK signaling, enhances the *Hsf1* phenotype

To test if the increased transcript accumulation of the CK responsive genes was biologically relevant, we made use of a null allele of *ZmRR3*, also known as *aberrant phyllotaxy1* (*abph1*). Plants homozygous for the recessive *abph1* reference allele have an altered phyllotactic pattern and develop leaves paired 180° at each node instead of having the normal alternating pattern (Figures 4A to 4B) but have no P-D patterning defects (Jackson and Hake, 1999). Backcross families were produced which segregated four phenotypes – wild type, heterozygous *Hsf1-1603*, homozygous *abph1*, and heterozygous *Hsf1-1603* plus homozygous *abph1* – in equal frequencies (Figures 4A to 4B). Double mutant plants, heterozygous for *Hsf1* and homozygous for *abph1*,

had paired leaf phyllotaxy and a strongly enhanced *Hsf1* phenotype, including very

stunted stature, increased shoot branching, very slow growth, extremely short and

narrow leaves, and severe leaf patterning defects including abundant prongs and bi- or

trifurcation of leaf blades (Figure 4B). The synergistic interaction of *Hsf1* and *abph1*

was consistent with *ZmRR3* functioning as a negative regulator of CK signaling and

indicated the loss of *abph1* function enhanced the *Hsf1* phenotype.

DISCUSSION

CK influences specific developmental programs in maize leaves

In this study we showed that the *Hsf1* mutation conditions a CK hypersignaling

phenotype that has multiple effects on plant growth and development, including specific

effects on (i) leaf patterning, (ii) leaf size and (iii) leaf epidermal cell fate (Bertrand-

Garcia and Freeling, 1991a). Supporting this idea, we also show exogenous CK

treatment of wild type maize seeds produced similar changes in these developmental

programs. Prominent among the developmental changes was a specific alteration in P-

D leaf patterning where ectopic outgrowths with proximal identity (prongs) formed in the

distal blade (Figures 1A to 1C and Supplemental Figure 2A). Although growth along the

P-D axis is fundamental to normal leaf development and morphology, its molecular

control has not been fully characterized. In eudicots, the activities of several

transcription factor genes, such as, *BLADE ON PETIOLE1* (*BOP1*), *LEAFY PETIOLE*

(*LEP*), and *JAGGED* (*JAG*), have been linked to the control of P-D leaf development

(van der Graaff et al., 2000, 2003; Ha et al., 2004; Ohno et al., 2004; Norberg et al.,

2005). *BOP* genes have also been shown to influence P-D leaf patterning in monocots

like barley and recently, the activity of three, redundant *OsBOP* genes was shown to be

required for sheath identity in rice (Tavakol et al., 2015; Toriba et al., 2019). In several

monocots, the misexpression of several class I *knox* genes perturb P-D patterning by

potentially altering phytohormone ratios in developing leaf primordia (Reiser et al., 2000;

Schneeberger et al., 1995; Foster et al., 1999b; Ramirez et al., 2009). Our analysis of

Hsf1, the second characterized mutation of a maize CK signaling gene, has uncovered

a connection between CK and the specification of P-D leaf patterning that is consistent

with this hypothesis. How CK drives prong formation is not clear, although the interplay

of CK and GA are known to control the degree of leaf complexity in eudicots like

Arabidopsis and tomato, through the specification of marginal lobes or leaflets (Jasinski

et al., 2005; Bar and Ori, 2015). Whether there is any overlap between the

mechanism(s) of prong formation in *Hsf1* and leaflet formation in species like tomato will

require further analysis. Prong formation itself appears developmentally regulated as

prong initiation seems to be coordinated with formation of the ligule suggesting the

signals establishing the P-D axis might be transmitted across the entire leaf primordium

(Supplemental Figures 1D to 1E). Moreover, prong formation is not random as prongs

form only within a certain domain of the blade, with the largest prongs forming more

basally (Supplemental Figures 2D to 2F). Intriguingly, this prong-formation region has

some overlap with the domain of the leaf blade deleted by mutation of the duplicate

wuschel-related homeobox (*wox*) genes *narrow sheath1* and *narrow sheath2*

(Nardmann et al., 2004). This implies that the marginal domain specified by these

duplicate *wox* transcription factors may provide a permissive context for prongs to form.

This hypothesis can be tested by analysis of prong formation in the triple mutant.

Leaf sheath and blade length, and blade width were reduced in *Hsf1*

heterozygotes compared to wild type sib plants at seedling and mature growth stages,

consistent with previous reports, and CK treatment recapitulated this phenotype in wild

type inbred seedlings (Figures 2A to 2C) (Bertrand-Garcia and Freeling, 1991b). Since

CK activity typically promotes cellular proliferation, how CK hypersignaling reduces

growth in the shoot is not known, although increased CK signaling is known to reduce

root growth (Werner et al., 2001, 2003). Typically, reducing CK accumulation or

signaling results in smaller leaves and other above ground organs, suggesting

increased CK activity might be expected to enhance growth (Werner et al., 2001;

Nishimura et al., 2004). Growth of the maize leaf is organized linearly along its

longitudinal axis into distinct zones of cell division, cell expansion and differentiation

(Freeling and Lane, 1992). Recent transcriptome, proteome and hormone profiling

studies have enumerated multiple regulatory pathways controlling the size of and
transitions between the different growth zones, with GA playing a prominent role (Li et
al., 2010; Nelissen et al., 2012; Facette et al., 2013). How increased CK signaling
impacts these growth zones to determine final leaf size will require further analysis
building upon these previous studies.

In addition to a change in P-D patterning and reduction in leaf size, the *Hsf1*
mutation and CK treatment of wild type seed promoted increased macrohair formation
in the leaf epidermis (Figures 2D to 2E and 2J). Macrohairs are normally found on adult
leaves on the abaxial sheath, at high density near the ligule but declining basipetally, on
the adaxial blade and along the blade margin. *Hsf1* increased macrohair production not
only on the abaxial sheath, adaxial blade, auricle and blade margins of adult leaves but
also on juvenile and transitional leaves which are typically glabrous. CK treatment
phenocopied the increased pubescence phenotype of *Hsf1* (Figures 2D to 2E). The
epidermis of the maize leaf has three types of pubescence – bicellular microhairs,
macrohairs and prickle hairs – with macrohairs being the most prominent (Freeling and
Lane, 1992). Macrohairs form by differentiation of specialized epidermal cells organized
in patterned files beginning in the fifth or sixth leaf (Moose et al., 2004). Little is known
regarding the signals specifying macrohair formation, although a recessive mutation
affecting macrohair initiation, *macrohairless1*, has been reported (Moose et al., 2004).
By contrast, trichome differentiation in the leaves of eudicots, like Arabidopsis, is known
to be controlled by a core network of positive and negative transcriptional regulators
(Ishida et al., 2008; Grebe, 2012; Pattanaik et al., 2014). And trichome initiation on the
inflorescence organs in Arabidopsis is jointly stimulated by the activity of CK and GA,
and downstream transcription factors (Gan et al., 2007; Zhou et al., 2013). The
increase in macrohair formation mediated by CK treatment or the *Hsf1* mutant suggests
CK can reprogram epidermal cell fate in maize leaves as well.

Missense Mutations in the Maize CK Receptor ZmHK1 underlie the *Hsf1*
phenotype

Our data indicate gain-of-function mutations of the maize CK receptor *ZmHK1* underlie

the semi-dominant *Hsf1* mutations. CK signaling, which is well described (To and

Kieber, 2008; Hwang et al., 2012), regulates several developmental and physiological

processes, although influences on leaf patterning are not among them. For example,

combinations of loss of function mutations of the three Arabidopsis CK receptors

demonstrate this gene family has partially overlapping and redundant functions in the

control of shoot and root growth, seed size, germination and leaf senescence (Higuchi

et al., 2004; Nishimura et al., 2004; Riefler et al., 2006). CK receptors were shown to

also possess phosphatase activity by analysis of a specific mutation of *AHK4/CRE1*, the

recessive *wooden leg* (*wol*) allele (*CRE1 T278I*) (Mahonen et al., 2006). Plants

homozygous for the *wol* allele have abnormal root vascular development due to the

dose-dependent constitutive phosphatase activity of this allele. A gain-of-function

mutation in the CHASE domain of AHK3 (*ore12-1*) revealed this receptor plays a major

role in CK-mediated leaf senescence; although how this mutation affected receptor

activity was not explored (Kim et al., 2006). The study of gain-of-function mutations has

revealed additional information on CK receptor function. Novel, dominant, missense

mutations in *AHK2* and *AHK3*, the *repressor of cytokinin deficiency* alleles (*rock2* and

rock3) enhanced CK signaling, increased CK hypersensitivity, and increased transcript

accumulation of CK-responsive genes, similar to the *Hsf1* mutations (Figure 3) (Bartrina

et al., 2017). In contrast, the *rock* mutations had the opposite effect on phenotype

compared to *Hsf1*, producing early flowering, enlarged rosette leaves and shoots, and

longer roots. The contrasting phenotypic effects might be due to differences in signaling

strength between the *rock* and *Hsf1* mutations or reflect differences in the downstream

circuitry between the two species.

Mutations near the CK binding pocket alter ligand affinity and receptor signaling

To clarify how the function of ZmHK1 was altered in the *Hsf1* mutants, we

analyzed their activity in heterologous his-kinase signaling and ligand binding assays.

Our results indicate two of the *Hsf1* mutant receptors signal independent of added CK in

yeast and all three have increased binding affinities for the natural CKs tested (Figure

1E and Table 1). The mutant receptors may be in a "locked on" state, similar to what

was hypothesized for the *snf2* mutation or the increased ligand affinities of the *Hsf1*

receptors may explain their ability to signal independent of CK action. We favor the

second idea and think the increased CK affinity explains the ability of the mutant

receptors to signal in heterologous hosts. Many microbes, including *E. coli* and yeast,

contain low concentrations of iP as a normal constituent of tRNA which can become

free due to tRNA decay (Skoog and Armstrong, 1970; Hall, 1973; Romanov, 1990; Mok

and Mok, 2001). The three mutant receptors all have increased affinity for iP (Table 2).

This stronger affinity may be due to stronger complex formation, or longer receptor

occupancy and, as a consequence, stronger signaling even in the presence of low iP

concentration. Thus, the ability of the ZmHK1-AEWL and ZmHK1-1603 receptors to

signal in yeast without added CKs may be due to their increased affinity for iP already

present at low concentration in yeast cells (Figure 1E). In fact, it has been shown that

expressing other HK receptors in the *sln1* deletion yeast strain TM182 permits this

strain to grow on glucose without added CKs, albeit at a much slower rate than with

CKs present, and recombinant HKs synthesized in *E. coli* cannot be crystalized without

iP complexed in the binding pocket (Higuchi et al., 2009; Hothorn et al., 2011). Since all

three mutant receptors have increased ligand affinities (Table 1), have nearly identical

mutant plant phenotypes in several different genetic backgrounds (Figure 1 and

Supplemental Table 1), and show similar misexpression patterns of CK responsive

genes (Figures 3A) we conclude all three *Hsf1* mutant receptors function similarly *in*

planta.

Our structural analysis localized each residue mutated in *Hsf1* to the ligand-

binding Per-Arnt-Sim-like (PAS) subdomain of the CHASE domain in ZmHK1 (Figure

1F) (Steklov et al., 2013; Hothorn et al., 2011). Notably, none are within the CK binding

pocket or predicted to make contact with the ligand. Rather all are located near a loop

domain comprising one face of the pocket suggesting interactions with this loop may

affect the binding pocket resulting in increased ligand affinity. Interestingly, amino acid

substitutions that rendered *AHK4* constitutively active in a heterologous *E. coli* his-

kinase assay were located downstream of the CHASE domain in the second

transmembrane domain and near the kinase domain (Miwa et al., 2007). In addition,

none of the *rock* mutations are located in the ligand-binding PAS domain (Bartrina et al.,

2011). Rather two are in the N-terminal α–helices and one is in the C-terminal

transmembrane domain. Therefore, further structure-function studies will be needed to

define which residues are crucial for activity and to resolve the precise mechanism(s) by

which individual missense mutations alter ligand binding and receptor signaling.

Hsf1 affects downstream components of CK signaling

More ZmHK1 signaling in developing *Hsf1* leaf primordia resulted in increased

transcript accumulation of several early CK response genes in all three *Hsf1* mutant

alleles (Figure 3A). Although not all CK reporters responded the same within an allele

or tissue, overall our data are consistent with *Hsf1* upregulating CK responsive genes.

The most consistent effect was upregulation of *ZmRR3* where its normally meristem-

confined expression was expanded in *Hsf1-1603* to include expression near newly

arising leaf primordia and in primordia margins (Figure 3B). Notably, the increased CK

signaling reported by *ZmRR3* marks the margins of early stage leaf primordia (Figure

3B) which is where prongs will form later in development (Supplemental Figures 1F to

1G). Although we found ectopic *ZmRR3* signal along the entire margin, outgrowths do

not emanate from the entire blade margin but, rather, occur sporadically, with

outgrowths interspersed with regions of normal blade margin (Figures 1B to 1C and

Supplemental Figure 2A). This observation suggests even though CK hypersignaling

can promote proximalization of blade margin cells, not all cells at the margin are

competent to respond to this signal. Double mutants heterozygous for *Hsf1-1603* and

homozygous for *abph1*, a null allele of *ZmRR3*, show a synergistic interaction (Figures

4A to 4B). Several type-A RRs function to negatively regulate CK signal transduction,

as well as, regulate circadian rhythms, phytochrome function and meristem size (To et

al., 2004). The increased severity of growth defects in *Hsf1* heterozygotes which lack

abph1 activity suggests upregulation of *ZmRR3* (*abph1*) partially ameliorates CK

hypersignaling. This also suggests that *ZmRR3* normally functions to attenuate CK

signal transduction in maize shoot apices, in addition to specifying leaf phyllotaxy.

The identification of the CK receptor *ZmHK1* as the gene underlying the leaf

patterning *Hsf1* mutation adds to our understanding of the role CK can play in basic

developmental programs. Future studies to determine the molecular determinants

functioning downstream of CK signaling that promote prong formation should illuminate

mechanisms important for developmental reprogramming and cell fate acquisition.

METHODS

Plant Material, Genetics, Phenotypic Measurements, and Analysis.

The *Hsf1-1595*, *Hsf1-1603* and *Hsf1-2559* mutants arose via EMS mutagenesis

of the inbred Mo17 and seed was obtained from the Maize Genetic Cooperation Stock

Center (http://maizecoop.cropsci.uiuc.edu/). *Hsf1-AEWL* arose via EMS mutagenesis of

the inbred A619 and *Hsf1-7322* via EMS mutagenesis of the inbred A632 in

independent screens. Homozygous *Hsf1* mutants of all five alleles were identified for

sequence analysis from progeny of self-pollinated heterozygous B73 introgressed

plants by phenotype and also by PCR screening of linked sequence polymorphisms

unique to the progenitor inbred lines and the backcross inbred B73. Since *Hsf1-1595*

and *Hsf1-1603* were the same transition mutations as *Hsf1-7322* and *Hsf1-2559*,

respectively, further analysis was only performed on three mutants: *Hsf1-1595*, *Hsf1-*

1603 and *Hsf1-AEWL*. All phenotypic, molecular and epistatic analyses were

performed on the three alleles that had been backcrossed a minimum of six times to the

inbred B73. The *Hsf1* phenotype of the three alleles was fully penetrant as a

heterozygote in all backcross generations. Progeny from self- or sib-pollinated *Hsf1*

heterozygotes of the three alleles segregated 25% severely stunted, very slow growing,

multi-shoot plants that only survived when grown in the greenhouse but were sterile.

The *abph1* mutant seed was backcrossed a minimum of three times to the inbred B73

before making the double mutant family segregating with *Hsf1-1603*. *Hsf1-1603*

heterozygotes were crossed by *abph1* homozygotes and double heterozygous progeny

plants were backcrossed by *abph1* homozygotes creating double mutants families

segregating 25% +/+, *abph1*/+ (WT); 25% +/+, *abph1*/*abph1* (single *abph1* mutant);

25% *Hsf1*/+, +/*abph1* (single *Hsf1* mutant); and 25% *Hsf1*/+, *abph1*/*abph1* (double *Hsf1*

abph1 mutant). Allele specific PCR genotyping was done to confirm phenotypes of

Hsf1 heterozygotes and *abph1* heterozygotes and homozygotes (Supplemental Table

3).

Measurement of adult plant traits of the three *Hsf1* mutant alleles was performed

on field grown families segregating 50% wild type: 50% *Hsf1* heterozygotes.

Measurements were taken on 7-11 plants of each genotype in 1-row plots with two

biological replicates. For analysis of prong position, prong size and percent prong

margin, the third leaf above the ear of adult *Hsf1* heterozygous plants was collected

from 1-row plots of field grown plants in three replicates in summer 2013.

Approximately, 6 to 10 leaves were collected per plot for each allele. For each leaf,

measurements were made for (1) total blade length, (2) prong position by measuring the

distance from the base of the blade to the mid-point of each prong on each blade

margin, and (3) prong size by measuring from the basal to the distal position along the

margin where proximal tissue emerged from the blade for each prong (Figure 1B).

Percent prong margin was defined as the proportion of leaf blade margin that is

occupied by tissue having proximal (sheath, auricle and/or ligule) identity and was

calculated by summing the size of all prongs from both sides of the leaf blade divided by

twice the length of the leaf blade.

Analysis of prong position, prong size and the relationship between prong

position and size was estimated with kernel smoothing methods (Silverman, 1986;

Wand and Jones, 1995). For all cases a Gaussian kernel was used and the data

reflection method was applied for boundary correction since both position and size are

positive variables. The bandwidth were selected using least squares cross validation

(Bowman, 1984). All computations were performed using R software, kernel density

estimation was performed using the ks package (Duong, 2007) and figures were

created with the ggplot2 package (Wickham, 2009).

Map-based cloning of *Hsf1*.

Hsf1-1595 was introgressed into B73 and crossed to PRE84 to generate a BC1

mapping population. Genetic mapping with 96 BC1 individuals defined *Hsf1* between

two SNP markers on chr5: PHA12918-F (204590502 bp, B73 RefGen_v2) and

PHA5244-F (206614542 bp, B73 RefGen_v2). The two flanking markers were used to

screen a BC1 population of 1500 individuals from B73_*Hsf1* x A632 and 1600 individual

from B73_*Hsf1* x PRE84. 224 recombinants were identified, and these individuals were

used for further fine mapping. Additional markers derived from the *Hsf1* interval were

developed and used to fine map the *Hsf1* mutation with the recombinants, as described

in Jiang et al., 2012 (Jiang et al., 2012). The gene underlying the *Hsf1* mutation was

finally delimited to a 21 kb interval, between Indel marker 410984 (205538463 bp, B73

RefGen_v2, with one recombinant between this marker and *Hsf1*) and SNP marker

391087 (205559234 bp, B73 RefGen_v2, with three recombinants between this marker

and *Hsf1*). There is only one annotated gene model (B73 RefGen_v3

GRMZM2G151223, B73 RefGen_v4 Zm00001d017977) in this interval that was also

annotated in NCBI as LOC541634 *histidine kinase1*, a putative cytokinin receptor.

Heterologous histidine kinase assays. Signaling of the wild type and *Hsf1* mutant

ZmHK1 receptors in yeast was performed as described previously (Inoue et al., 2001).

The exact point mutations for each of the three *Hsf1* missense mutations were

engineered into the cDNA of *ZmHK1* in the plasmid P415-CYC1-ZmHK1 plasmid with

the QuikChange II Site-Directed Mutagenesis kit (Agilent Technologies) using the

manufacturer's specifications.

Cytokinin binding affinity determination. Cytokinin binding assays were performed

with recombinant maize cytokinin receptors expressed in *E. coli* cells. Spheroplasts

were prepared from cell lines expressing the wild type ZmHK1, and mutant ZmHK1-

AEWL and ZmHK1-1603 receptors. Competitive cytokinin binding assays were

performed as previously described (Lomin et al., 2011). Transient expression of

receptors for the homologous binding assay was done by transformation of tobacco

Nicotiana benthamiana as previously described (Sparkes et al., 2006). Agrobacteria *A.*

tumefaciens carrying cytokinin receptor genes fused to GFP were grown in parallel with

a helper agrobacterial strain p19 (Voinnet et al., 2003). Five to six week old tobacco

plants were infiltrated with the mixture of two agrobacterial strains and the expression

level of receptor genes was checked after 4 days using a confocal microscope. For

those cases with sufficient expression, leaves were processed further for plant

membrane isolation. For plant membrane isolation, all manipulations were done at 4

°C. Tobacco leaves were homogenized in buffer containing 300 mM sucrose, 100 mM

Tris-HCl (pH 8.0), 10 mM Na_2-EDTA, 0.6% polyvinylpyrrolidone K30, 5 mM $K_2S_2O_5$, 5

mM DTT and 1 mM PMSF. The homogenate was filtered through Miracloth

(Calbiochem), and the filtrate was first centrifuged for 10 min at 10000 g, and then for

30 min at 100000 g. The microsome pellet was resuspended in PBS (pH 7.4), frozen

and stored at -70 °C before using.

ZmHK1 structure modeling.

The amino acid sequence of the ZmHK1 CHASE domain (86-270) was obtained from

the protein sequence database of NCBI (accession id: NP_001104859). It shares 69%

sequence identity with the *Arabidopsis HK4* sensor domain. The homology model for

ZmHK1 was generated using Swiss model server (http://swissmodel.expasy.org) with

the published crystal structure of AHK4 (pdb code: 3T4J) as a template. Subsequently

the model was solvated and subjected to energy minimization using the steepest

descent followed by conjugate gradient algorithm to remove clashes. The

stereochemical quality of the ZmHK1 model was assessed using the PROCHECK

program. None of the residues were in the disallowed regions of the Ramachandran

map.

Exogenous CK treatment.

Exogenous CK treatments were performed with 6-benzylaminopurine (6-BAP) (Sigma

Aldrich) dissolved in 10 drops 1N NaOH and brought to 1mM concentration with distilled

water. All water control treatments were done using a similar stock of 10 drops 1N

NaOH and diluted in parallel to the CK stock. Further dilutions to the desired CK

concentration were done with distilled water. Maize kernels were surface sterilized with

two 5 minute washes of 80% ethanol followed by two 15 minute washes of 50% bleach

and rinsed five times in sterile distilled water. Kernels were imbibed overnight with

sterile distilled water prior to the start of the hormone treatment. For hormone

treatments, 20 imbibed kernels per replicate were placed embryo-side down on two

paper towels in a petri dish, covered with two more layers of paper towel and filled with

15 mL of CK treatment or the water control solution. Petri dishes were sealed with

parafilm and placed in a lab drawer in the dark at room temperature for 6 days. After

treatment, germinating kernels were rinsed with sterile, distilled water and planted in 4

cm square pots in soilless potting medium (Metro-Mix 900, SunGro Horticulture) and

grown in the greenhouse (day: 16 hr./28°C, night: 8 hr./21°C) with supplemental lighting

(high pressure sodium and metal halide lights) and standard light intensity (230 μE m^{-2}

s^{-1} at height of 3.5 feet). Growth was monitored and leaf measurements were taken

after the fourth leaf collar (auricle and ligule) had fully emerged from the whorl after 3 to

4 weeks. For measurements, individual leaves were removed from the plant and each

component measured. Leaf sheath length was defined as the site of insertion of the

leaf base to the culm (stem) to the farthest point of sheath adjoining the ligule. Leaf

blade length was defined as the most proximal point of blade adjoining the ligule to the

distal blade tip. Leaf blade width was measured margin to margin at half of the leaf

blade length. All leaf measurements were analyzed using JMP PRO 12 software using

a student's t-test to determine significance with two comparisons, and Tukey's HSD test

to determine significance with more than two comparisons. To examine macrohair

abundance, epidermal impressions were made using Krazy Glue Maximum Bond®

cyanoacrylate glue applied to a Fisherbrand Superfrost Plus® microscope slide. The

adaxial blade of leaf one was pressed firmly into the glue for about 30 seconds, followed

by immediate removal of the leaf. Slides were imaged on an Olympus BX60 light

microscope.

Expression analysis.

In situ **hybridization***:*

For *in situ* hybridization, we slightly modified an online protocol from Jeff Long. For
complete details refer to http://pbio.salk.edu/pbiol/in_situ_protocol.html. *In situ* probes
were made using T7/SP6 promoter based *in vitro* transcription in the cloning vector
pGEMT (Promega). FAA (Formaldehyde Acetic Acid) fixed and paraffin embedded
maize shoot apices were sectioned at 10 µm thickness and laid on Probe-On-Plus
slides (Fisher) and placed on a warmer at 42°C. After overnight incubation, the slides
were deparrafinized using Histo- Clear (National Diagnostics), treated with proteinase K
and dehydrated. Probes were applied on the slides and pairs of slides were
sandwiched carefully and incubated at 55°C overnight. The following day the slides
were rinsed and washed. Diluted (1:1250) anti-DIG-antibody (Roche) was applied to
the slides and incubated for 2 hours. After thoroughly washing the slides, sandwiched
slides were placed in NBT-BCIP (Roche) solution (200 µl in 10ml buffer C; 100mM Tris
pH9.5/100mM NaCl/50mM $MgCl_2$) in dark for 2 to 3 days for color development. Color
development reaction was stopped using 1x Tris EDTA. The slides were mounted using
Immu-Mount (Thermo Scientific) and observed and imaged under a bright field
microscope.

RT qPCR:

Seedling tissue was collected from two-week old, stage V3 to V4 *Hsf1/+* and wild type
sib seedlings for each allele and included (1) ca. 2 cm of mature green leaf blade from
the distal half of leaf #4, (2) ca. a 2 cm cylinder of immature leaf tissue, cut ca. 1 cm
above the insertion point of leaf #5 after removing leaf #4, and (3) the remaining 1 cm
cylinder of tissue above the insertion point of leaf #5, consisting of the SAM, young leaf
primordia and the apical part of the stem. Tissue was bulked from three different plants

for each biological replicate and three replicates were collected. Total RNA was

extracted from these tissues using Trizol reagent, adhering to the manufacturer's

protocol (http://tools.lifetechnologies.com/content/sfs/manuals/trizol_reagent.pdf).

cDNA was synthesized from total RNA using the SuperScript® III First-Strand

(Invitrogen) synthesis system for reverse transcriptase PCR (RT-PCR) and oligo-d(T)

primers. Quantitative real-time PCR was performed on the cDNA using an LC480

(Roche) and the SYBR green assay. The primers were designed near the 3' end of the

gene with an amplicon size of between 120 bp to 250 bp. Folylpolyglutamate synthase

(FPGS) was used as an endogenous control as it was shown to have very stable

expression across a variety of maize tissues and range of experimental conditions

(Manoli et al., 2012). Two technical replicates were included for each gene.

Comparative ΔΔCt method was used to calculate fold change compared to the

endogenous control. ΔCt of mutant (*Hsf1*) and ΔCt of wild type (WT) was expressed as

the difference in Ct value between target gene and the endogenous control. ΔΔCt was

then calculated as the difference of ΔCt (*Hsf1*) and ΔCt (WT). Finally, fold change in

target gene expression between *Hsf1* and WT was determined as $2^{-\Delta\Delta Ct}$.

Acknowledgements

We thank Dave Jackson (Cold Spring Harbor Labs) for the *abph1* mutant seed, Erica

Unger-Wallace (Iowa State University) for technical assistance and Erik Vollbrecht

(Iowa State University) for support during the early phase of this research. D.M.K.,

S.N.L. and G.A.R. were partly supported by the Molecular and Cell Biology Program of

the Presidium of RAS. . This work was supported by the National Science Foundation

under Grant Number 1022452.

Author contributions

REFERENCES

Asakura, Y., Hagino, T., Ohta, Y., Aoki, K., Yonekura-Sakakibara, K., Deji, A.,
Yamaya, T., Sugiyama, T., and Sakakibara, H. (2003). Molecular characterization
of His-Asp phosphorelay signaling factors in maize leaves: Implications of the
signal divergence by cytokinin-inducible response regulators in the cytosol and the
nuclei. Plant Mol. Biol. **52**: 331–341.

Bar, M. and Ori, N. (2015). Compound leaf development in model plant species. Curr.
Opin. Plant Biol. **23**: 61–69.

Bartrina, I., Jensen, H., Novák, O. rej, Strnad, M., Werner, T., and Schmülling, T.
(2017). Gain-of-Function Mutants of the Cytokinin Receptors AHK2 and AHK3
Regulate Plant Organ Size, Flowering Time and Plant Longevity. Plant Physiol.
173: 1783–1797.

Bartrina, I., Otto, E., Strnad, M., Werner, T., and Schmülling, T. (2011). Cytokinin
Regulates the Activity of Reproductive Meristems, Flower Organ Size, Ovule
Formation, and Thus Seed Yield in Arabidopsis thaliana. Plant Cell Online **23**: 69–
80.

Benjamins, R. and Scheres, B. (2008). Auxin: The Looping Star in Plant Development.
Annu. Rev. Plant Biol. **59**: 443–465.

Bertrand-Garcia, R. and Freeling, M. (1991a). Hairy-Sheath Frayed #1-O: A Systemic,
Heterochronic Mutant of Maize That Specifies Slow Developmental Stage
Transitions. Am. J. Bot. **78**: 747–765.

Bertrand-Garcia, R. and Freeling, M. (1991b). Hsf1-O (Hairy sheath frayed): 5L
linkage data. Maize Genet. Coop. News Lett.: 30.

Bird, R.M. and Neuffer, M.G. (1985). Developmentally Interesting New Mutants in
Plants Odd New Dominant Mutations Affecting the Development of the Maize Leaf.
Free. M. (Ed.). Ucla (University Calif. Los Angeles) Symp. Mol. Cell. Biol. New Ser.
Vol. 35. Plant Genet. Third Annu. Arco Plant Cell Res. Institute-Ucla Symp. Plant

Biol. Keystone, Colo., USA, Apr. 13-: 818–822.

Bolduc, N. and Hake, S. (2009). The Maize Transcription Factor KNOTTED1 Directly

Regulates the Gibberellin Catabolism Gene ga2ox1. Plant Cell **21**: 1647–1658.

Bowman, A.W. (1984). An alternative method of cross-validation for the smoothing of

density estimates. Biometrika **71**: 353–360.

Bowman, J.L., Eshed, Y., and Baum, S.F. (2002). Establishment of polarity in

angiosperm lateral organs. Trends Genet. **18**: 134–141.

Byrne, M., Timmermans, M., Kidner, C., and Martienssen, R. (2001). Development of

leaf shape. Curr. Opin. Plant Biol. **4**: 38–43.

Byrne, M.E., Simorowski, J., and Martienssen, R.A. (2002). ASYMMETRIC

LEAVES1 reveals knox gene redundancy in Arabidopsis. Development **129**: 1957–

1965.

Du, L., Jiao, F., Chu, J., Jin, G., Chen, M., and Wu, P. (2007). The two-component

signal system in rice (Oryza sativa L.): A genome-wide study of cytokinin signal

perception and transduction. Genomics **89**: 697–707.

Duong, T. (2007). ks: Kernel density estimation and kernel discriminant analysis for

multivariate data in R. J. Stat. Softw. **21**: 1–16.

Edgar, R.C. (2004). MUSCLE: multiple sequence alignment with high accuracy and

high throughput. Nucleic Acids Res. **32**: 1792–1797.

Endrizzi, K., Moussian, B., Haecker, A., Levin, J.Z., and Laux, T. (1996). The

SHOOT MERISTEMLESS gene is required for maintenance of undifferentiated

cells in Arabidopsis shoot and floral meristems and acts at a different regulatory

level than the meristem genes WUSCHEL and ZWILLE. Plant J. **10**: 967–979.

Facette, M.R., Shen, Z., Björnsdóttir, F.R., Briggs, S.P., and Smith, L.G. (2013).

Parallel Proteomic and Phosphoproteomic Analyses of Successive Stages of Maize

Leaf Development. Plant Cell **25**: 2798–2812.

Foster, T., Veit, B., and Hake, S. (1999a). Mosaic analysis of the dominant mutant,

Gnarley1-R, reveals distinct lateral and transverse signaling pathways during maize

leaf development. Development **126**: 305–313.

Foster, T., Yamaguchi, J., Wong, B.C., Veit, B., and Hake, S. (1999b). Gnarley1 Is a

Dominant Mutation in the knox4 Homeobox Gene Affecting Cell Shape and Identity.

Plant Cell **11**: 1239–1252.

Freeling, M. and Hake, S. (1985). Developmental genetics of mutants that specify

Knotted leaves in maize. Genetics **111**: 617–634.

Freeling, M. and Lane, B. (1992). The maize leaf. In The Maize Handbook, M. Freeling

and V. Walbot, eds (Springer-Verlag: New York LB - PBS Record: 390), p. in

press.

Gan, Y., Liu, C., Yu, H., and Broun, P. (2007). Integration of cytokinin and gibberellin

signalling by Arabidopsis transcription factors GIS, ZFP8 and GIS2 in the regulation

of epidermal cell fate. Development **134**: 2073–2081.

Giulini, A., Wang, J., and Jackson, D. (2004). Control of phyllotaxy by the cytokinin-

inducible response regulator homologue ABPHYL1. Nature **430**: 1031–1034.

van der Graaff, E., Dulk-Ras, A.D., Hooykaas, P.J., and Keller, B. (2000). Activation

tagging of the LEAFY PETIOLE gene affects leaf petiole development in

Arabidopsis thaliana. Development **127**: 4971–4980.

van der Graaff, E., Nussbaumer, C., and Keller, B. (2003). The Arabidopsis thaliana

rlp mutations revert the ectopic leaf blade formation conferred by activation tagging

of the LEP gene. Mol. Genet. Genomics **270**: 243–252.

Grebe, M. (2012). The patterning of epidermal hairs in Arabidopsis — updated. Curr.

Opin. Plant Biol. **15**: 31–37.

Guo, M., Thomas, J., Collins, G., and Timmermans, M.C.P. (2008). Direct

Repression of KNOX Loci by the ASYMMETRIC LEAVES1 Complex of

Arabidopsis. Plant Cell **20**: 48–58.

Ha, C.M., Jun, J.H., Nam, H.G., and Fletcher, J.C. (2004). BLADE-ON-PETIOLE1

encodes a BTB/POZ domain protein required for leaf morphogenesis in

Arabidopsis thaliana. Plant Cell Physiol **45**: 1361–1370.

Ha, C.M., Kim, G.T., Kim, B.C., Jun, J.H., Soh, M.S., Ueno, Y., Machida, Y.,
Tsukaya, H., and Nam, H.G. (2003). The BLADE-ON-PETIOLE 1 gene controls
leaf pattern formation through the modulation of meristematic activity in
Arabidopsis. Development **130**: 161–172.

Hake, S., Sinha, N., Veit, B., Vollbrecht, E., and Walko, R. (1991). Mutations of
Knotted alter cell interactions in the developing maize leaf. In Plant Molecular
Biology, R.G. Herrmann and B. Larkins, eds (Plenum Press: New York), pp. 555–
562.

Hake, S., Vollbrecht, E., and Freeling, M. (1989). Cloning Knotted, the dominant
morphological mutant in maize using Ds2 as a transposon tag. EMBO J. **8**: 15–22.

Hall, R.H. (1973). Cytokinins as a probe of developemental processes. Annu. Rev.
Plant Physiol. **24**: 415–444.

Hay, A., Barkoulas, M., and Tsiantis, M. (2006). ASYMMETRIC LEAVES1 and auxin
activities converge to repress BREVIPEDICELLUS expression and promote leaf
development in Arabidopsis. Development **133**: 3955–3961.

Hay, A., Kaur, H., Phillips, A., Hedden, P., Hake, S., and Tsiantis, M. (2002). The
gibberellin pathway mediates KNOTTED1-type homeobox function in plants with
different body plans. Curr Biol **12**: 1557–1565.

Higuchi, M. et al. (2004). In planta functions of the Arabidopsis cytokinin receptor
family. PNAS **101**: 8821–8826.

Higuchi, M., Kakimoto, T., and Mizuno, T. (2009). Cytokinin Sensing Systems Using
Microorganisms Plant Hormones S. Cutler and D. Bonetta, eds (Humana Press).

Hothorn, M., Dabi, T., and Chory, J. (2011). Structural basis for cytokinin recognition
by Arabidopsis thaliana histidine kinase 4. Nat Chem Biol **7**: 766–768.

Hwang, I. and Sakakibara, H. (2006). Cytokinin biosynthesis and perception. Physiol.
Plant. **126**: 528–538.

Hwang, I. and Sheen, J. (2001). Two-component circuitry in Arabidopsis cytokinin
signal transduction. Nature **413**: 383–389.

Hwang, I., Sheen, J., and Müller, B. (2012). Cytokinin Signaling Networks. Annu. Rev.
Plant Biol. **63**: 353–380.

Inoue, T., Higuchi, M., Hashimoto, Y., Seki, M., Kobayashi, M., Kato, T., Tabata, S.,
Shinozaki, K., and Kakimoto, T. (2001). Identification of CRE1 as a cytokinin
receptor from Arabidopsis. Nature **409**: 1060–1063.

Ishida, T., Kurata, T., Okada, K., and Wada, T. (2008). A Genetic Regulatory Network
in the Development of Trichomes and Root Hairs. Annu. Rev. Plant Biol. **59**: 365–
386.

Jackson, D. and Hake, S. (1999). Control of phyllotaxy in maize by the abphyl1 gene.
Development **126**: 315–323.

Jasinski, S., Piazza, P., Craft, J., Hay, A., Woolley, L., Rieu, I., Phillips, A., Hedden,
P., and Tsiantis, M. (2005). KNOX action in Arabidopsis is mediated by coordinate
regulation of cytokinin and gibberellin activities. Curr Biol **15**: 1560–1565.

Jiang, F., Guo, M., Yang, F., Duncan, K., Jackson, D., Rafalski, A., Wang, S., and
Li, B. (2012). Mutations in an AP2 Transcription Factor-Like Gene Affect Internode
Length and Leaf Shape in Maize. PLoS One **7**: e37040.

Johnston, R., Wang, M., Sun, Q., Sylvester, A.W., Hake, S., and Scanlon, M.J.
(2014). Transcriptomic Analyses Indicate That Maize Ligule Development
Recapitulates Gene Expression Patterns That Occur during Lateral Organ Initiation.
Plant Cell **26**: 4718–4732.

Kerstetter, R., Vollbrecht, E., Lowe, B., Veit, B., Yamaguchi, J., and Hake, S.
(1994). Sequence analysis and expression patterns divide the maize knotted1-like
homeobox genes into two classes. Plant Cell **6**: 1877–1887.

Kerstetter, R.A. and Poethig, R.S. (1998). The specification of leaf identity during
shoot development. Annu. Rev. Cell Dev. Biol. **14**: 373–398.

Kim, H.J., Ryu, H., Hong, S.H., Woo, H.R., Lim, P.O., Lee, I.C., Sheen, J., Nam,
H.G., and Hwang, I. (2006). Cytokinin-mediated control of leaf longevity by AHK3
through phosphorylation of ARR2 in Arabidopsis. PNAS **103**: 814–819.

Li, P. et al. (2010). The developmental dynamics of the maize leaf transcriptome. Nat.

Genet. **42**: 1060–1067.

Lomin, S.N., Krivosheev, D.M., Steklov, M.Y., Arkhipov, D.V., Osolodkin, D.I., Schmülling, T., and Romanov, G.A. (2015). Plant membrane assays with cytokinin receptors underpin the unique role of free cytokinin bases as biologically active ligands. J. Exp. Bot. **66**: 1851–1863.

Lomin, S.N., Yonekura-Sakakibara, K., Romanov, G.A., and Sakakibara, H. (2011). Ligand-binding properties and subcellular localization of maize cytokinin receptors. J. Exp. Bot. **62**: 5149–5159.

Long, J.A., Moan, E.I., Medford, J.I., and Barton, M.K. (1996). A member of the KNOTTED class of homeodomain proteins encoded by the STM gene of Arabidopsis. Nature **379**: 66–69.

Mahonen, A.P., Higuchi, M., Tormakangas, K., Miyawaki, K., Pischke, M.S., Sussman, M.R., Helariutta, Y., and Kakimoto, T. (2006). Cytokinins Regulate a Bidirectional Phosphorelay Network in Arabidopsis. Curr. Biol. **16**: 1116–1122.

Manoli, A., Sturaro, A., Trevisan, S., Quaggiotti, S., and Nonis, A. (2012). Evaluation of candidate reference genes for qPCR in maize. J. Plant Physiol. **169**: 807–815.

McConnell, J.R. and Barton, M.K. (1998). Leaf polarity and meristem formation in Arabidopsis. Development **125**: 2935–2942.

Miwa, K., Ishikawa, K., Terada, K., Yamada, H., Suzuki, T., Yamashino, T., and Mizuno, T. (2007). Identification of Amino Acid Substitutions that Render the Arabidopsis Cytokinin Receptor Histidine Kinase AHK4 Constitutively Active. Plant Cell Physiol **48**: 1809–1814.

Mok, D.W. and Mok, M.C. (2001). CYTOKININ METABOLISM AND ACTION. Annu. Rev. Plant Physiol. Plant Mol. Biol. **52**: 89–118.

Moon, J., Candela, H., and Hake, S. (2013). The Liguleless narrow mutation affects proximal-distal signaling and leaf growth. Development **140**: 405–412.

Moose, S.P., Lauter, N., and Carlson, S.R. (2004). The Maize macrohairless1 Locus Specifically Promotes Leaf Blade Macrohair Initiation and Responds to Factors Regulating Leaf Identity. Genetics **166**: 1451–1461.

Muehlbauer, G.J., Fowler, J.E., and Freeling, M. (1997). Sectors expressing the
homeobox gene liguleless3 implicate a time-dependent mechanism for cell fate
acquisition along the proximal-distal axis of the maize leaf. Development **124**:
5097–5106.

Muller, B. and Sheen, J. (2008). Cytokinin and auxin interaction in root stem-cell
specification during early embryogenesis. Nature **453**: 1094–1097.

Nardmann, J., Ji, J., Werr, W., and Scanlon, M.J. (2004). The maize duplicate genes
narrow sheath1 and narrow sheath2 encode a conserved homeobox gene function
in a lateral domain of shoot apical meristems. Development **131**: 2827–2839.

Nelissen, H., Rymen, B., Jikumaru, Y., Demuynck, K., Van Lijsebettens, M.,
Kamiya, Y., Inzé, D., and Beemster, G.T.S. (2012). A Local Maximum in
Gibberellin Levels Regulates Maize Leaf Growth by Spatial Control of Cell Division.
Curr. Biol. **22**: 1183–1187.

Nishimura, C., Ohashi, Y., Sato, S., Kato, T., Tabata, S., and Ueguchi, C. (2004).
Histidine Kinase Homologs That Act as Cytokinin Receptors Possess Overlapping
Functions in the Regulation of Shoot and Root Growth in Arabidopsis. Plant Cell
16: 1365–1377.

Norberg, M., Holmlund, M., and Nilsson, O. (2005). The BLADE ON PETIOLE genes
act redundantly to control the growth and development of lateral organs.
Development **132**: 2203–2213.

Ohno, C.K., Reddy, G.V., Heisler, M.G.B., and Meyerowitz, E.M. (2004). The
Arabidopsis JAGGED gene encodes a zinc finger protein that promotes leaf tissue
development. Development **131**: 1111–1122.

Ori, N., Eshed, Y., Chuck, G., Bowman, J.L., and Hake, S. (2000). Mechanisms that
control knox gene expression in the Arabidopsis shoot. Development **127**: 5523–
5532.

Pattanaik, S., Patra, B., Singh, S., and Yuan, L. (2014). An overview of the gene
regulatory network controlling trichome development in the model plant,
Arabidopsis. Front. Plant Sci. **5**: 259.

Pozzi, C., Rossini, L., and Agosti, F. (2001). Patterns and symmetries in leaf
development. Semin Cell Dev Biol **12**: 363–372.

Ramirez, J., Bolduc, N., Lisch, D., and Hake, S. (2009). Distal Expression of knotted1
in Maize Leaves Leads to Reestablishment of Proximal/Distal Patterning and Leaf
Dissection. Plant Physiol. **151**: 1878–1888.

Rashotte, A.M., Mason, M.G., Hutchison, C.E., Ferreira, F.J., Schaller, G.E., and
Kieber, J.J. (2006). A subset of Arabidopsis AP2 transcription factors mediates
cytokinin responses in concert with a two-component pathway. PNAS **103**: 11081–
11085.

Reiser, L., Sanchez-Baracaldo, P., and Hake, S. (2000). Knots in the family tree:
evolutionary relationships and functions of knox homeobox genes. Plant Mol Biol
42: 151–166.

Riefler, M., Novak, O., Strnad, M., and Schmulling, T. (2006). Arabidopsis Cytokinin
Receptor Mutants Reveal Functions in Shoot Growth, Leaf Senescence, Seed
Size, Germination, Root Development, and Cytokinin Metabolism. Plant Cell **18**:
40–54.

Romanov, G.A. (1990). Cytokinins and tRNAs: a hypothesis on their competitive
interaction via specific receptor proteins. Plant. Cell Environ. **13**: 751–754.

Romanov, G.A., Spíchal, L., Lomin, S.N., Strnad, M., and Schmülling, T. (2005). A
live cell hormone-binding assay on transgenic bacteria expressing a eukaryotic
receptor protein. Anal. Biochem. **347**: 129–134.

Ronquist, F. and Huelsenbeck, J.P. (2003). MrBayes 3: Bayesian phylogenetic
inference under mixed models. Bioinformatics **19**: 1572–1574.

Sakamoto, T., Kamiya, N., Ueguchi-Tanaka, M., Iwahori, S., and Matsuoka, M.
(2001). KNOX homeodomain protein directly suppresses the expression of a
gibberellin biosynthetic gene in the tobacco shoot apical meristem. Genes Dev **15**:
581–590.

Sakamoto, T., Sakakibara, H., Kojima, M., Yamamoto, Y., Nagasaki, H., Inukai, Y.,
Sato, Y., and Matsuoka, M. (2006). Ectopic Expression of KNOTTED1-Like

Homeobox Protein Induces Expression of Cytokinin Biosynthesis Genes in Rice. Plant Physiol. **142**: 54–62.

Scarpella, E., Marcos, D., Friml, J., and Berleth, T. (2006). Control of leaf vascular patterning by polar auxin transport. Genes Dev. **20**: 1015–1027.

Schneeberger, R.G., Becraft, P.W., Hake, S., and Freeling, M. (1995). Ectopic expression of the knox homeo box gene rough sheath1 alters cell fate in the maize leaf. Genes Dev. **9**: 2292–2304.

Shani, E., Yanai, O., and Ori, N. (2006). The role of hormones in shoot apical meristem function. Curr. Opin. Plant Biol. **9**: 484–489.

Silverman, B.W. (1986). Density estimation for statistics and data analysis (Chapman and Hall).

Skoog, F. and Armstrong, D. (1970). Cytokinins. Annu. Rev. Plant Physiol. **21**: 359–384.

Smith, L.G., Greene, B., Veit, B., and Hake, S.L.B.-P.B.S.R. 3380 (1992). A dominant mutation in the maize homeobox gene, Knotted-1, causes its ectopic expression in leaf cells with altered fates. Development **116**: 21–30.

Sparkes, I.A., Runions, J., Kearns, A., and Hawes, C. (2006). Rapid, transient expression of fluorescent fusion proteins in tobacco plants and generation of stably transformed plants. Nat. Protoc. **1**: 2019–2025.

Steklov, M., Lomin, S.N., Osolodkin, D.I., and Romanov, G.A. (2013). Structural basis for cytokinin receptor signaling: an evolutionary approach. Plant Cell Rep. **32**: 781–793.

Suzuki, T., Ishikawa, K., Yamashino, T., and Mizuno, T. (2002). An Arabidopsis Histidine-Containing Phosphotransfer (HPt) Factor Implicated in Phosphorelay Signal Transduction: Overexpression of AHP2 in Plants Results in Hypersensitiveness to Cytokinin. Plant Cell Physiol **43**: 123–129.

Suzuki, T., Miwa, K., Ishikawa, K., Yamada, H., Aiba, H., and Mizuno, T. (2001). The Arabidopsis Sensor His-kinase, AHK4, Can Respond to Cytokinins. Plant Cell Physiol. **42**: 107–113.

Sylvester, A.W., Cande, W.Z., and Freeling, M. (1990). Division and differentiation
during normal and liguleless-1 maize leaf development. Development **110**: 985–
1000.

Sylvester, A.W., Smith, L., and Freeling, M. (1996). Acquisition of identity in the
developing leaf. Annu. Rev. Cell Dev. Biol. **12**: 257–304.

Tavakol, E. et al. (2015). The Barley Uniculme4 Gene Encodes a BLADE-ON-
PETIOLE-Like Protein That Controls Tillering and Leaf Patterning. Plant Physiol.
168: 164 LP – 174.

Tirichine, L., Sandal, N., Madsen, L.H., Radutoiu, S., Albrektsen, A.S., Sato, S.,
Asamizu, E., Tabata, S., and Stougaard, J. (2007). A Gain-of-Function Mutation
in a Cytokinin Receptor Triggers Spontaneous Root Nodule Organogenesis.
Science (80-.). **315**: 104–107.

To, J.P.C., Haberer, G., Ferreira, F.J., Deruere, J., Mason, M.G., Schaller, G.E.,
Alonso, J.M., Ecker, J.R., and Kieber, J.J. (2004). Type-A Arabidopsis Response
Regulators Are Partially Redundant Negative Regulators of Cytokinin Signaling.
Plant Cell **16**: 658–671.

To, J.P.C. and Kieber, J.J. (2008). Cytokinin signaling: two-components and more.
Trends Plant Sci. **13**: 85–92.

Toriba, T., Tokunaga, H., Shiga, T., Nie, F., Naramoto, S., Honda, E., Tanaka, K.,
Taji, T., Itoh, J.-I., and Kyozuka, J. (2019). BLADE-ON-PETIOLE genes
temporally and developmentally regulate the sheath to blade ratio of rice leaves.
Nat. Commun. **10**: 619.

Tsiantis, M., Schneeberger, R., Golz, J.F., Freeling, M., and Langdale, J.A. (1999).
The maize rough sheath2 gene and leaf development programs in monocot and
dicot plants. Science (80-.). **284**: 154–156.

Tsuda, K., Kurata, N., Ohyanagi, H., and Hake, S. (2014). Genome-Wide Study of
KNOX Regulatory Network Reveals Brassinosteroid Catabolic Genes Important for
Shoot Meristem Function in Rice. Plant Cell **26**: 3488–3500.

Tsukaya, H. (1998). Genetic evidence for polarities that regulate leaf morphogenesis. J.

Plant Res. **111**: 113–119.

Voinnet, O., Rivas, S., Mestre, P., and Baulcombe, D. (2003). An enhanced transient
expression system in plants based on suppression of gene silencing by the p19
protein of tomato bushy stunt virus. Plant J. **33**: 949–956.

Wand, M.P. and Jones, M.C. (1995). Kernel Smoothing Vol. 60 of Monographs on
statistics and applied probability. (Chapman and Hall, London).

Werner, T., Motyka, V., Laucou, V., Smets, R., Van Onckelen, H., and Schmülling,
T. (2003). Cytokinin-Deficient Transgenic Arabidopsis Plants Show Multiple
Developmental Alterations Indicating Opposite Functions of Cytokinins in the
Regulation of Shoot and Root Meristem Activity. Plant Cell **15**: 2532–2550.

Werner, T., Motyka, V., Strnad, M., and Schmülling, T. (2001). Regulation of plant
growth by cytokinin. PNAS **98**: 10487–10492.

Wickham, H. (2009). ggplot2 (Springer New York: New York).

Yamada, H., Suzuki, T., Terada, K., Takei, K., Ishikawa, K., Miwa, K., Yamashino,
T., and Mizuno, T. (2001). The Arabidopsis AHK4 Histidine Kinase is a Cytokinin-
Binding Receptor that Transduces Cytokinin Signals Across the Membrane. Plant
Cell Physiol **42**: 1017–1023.

Yanai, O., Shani, E., Dolezal, K., Tarkowski, P., Sablowski, R., Sandberg, G.,
Samach, A., and Ori, N. (2005). Arabidopsis KNOXI Proteins Activate Cytokinin
Biosynthesis. Curr. Biol. **15**: 1566–1571.

Yonekura-Sakakibara, K., Kojima, M., Yamaya, T., and Sakakibara, H. (2004).
Molecular Characterization of Cytokinin-Responsive Histidine Kinases in Maize.
Differential Ligand Preferences and Response to cis-Zeatin. Plant Physiol. **134**:
1654–1661.

Zhao, Y. (2008). The role of local biosynthesis of auxin and cytokinin in plant
development. Curr. Opin. Plant Biol. **11**: 16–22.

Zhou, Z., Sun, L., Zhao, Y., An, L., Yan, A., Meng, X., and Gan, Y. (2013). Zinc Finger
Protein 6 (ZFP6) regulates trichome initiation by integrating gibberellin and
cytokinin signaling in Arabidopsis thaliana. New Phytol. **198**: 699–708.

1116

 FIGURE LEGENDS

1118

Figure 1. *Hsf1* mutants alter leaf patterning and are caused by missense mutations

in the *ZmHK1* cytokinin receptor. **(A)** Adaxial view of half-leaves from WT and *Hsf1-*

1603/+ sibs showing the proximal-distal organization of the sheath (s), ligule (l), auricle

(a) and blade (b) and a prong outgrowth (red triangle). Bar = 5 cm. **(B)** Close-up of a

blade margin (b) from WT and *Hsf1-1603/+* showing a prong consisting of proximal leaf

segments – sheath (s), ligule (l) and auricle (a) juxtaposed to the blade (b). Bar = 1 cm.

(C) Comparison of leaf phenotypes between the three *Hsf1* alleles. L4 (top), 4th leaf

below tassel; L5 (bottom, 5th leaf below tassel. Bar = 10 cm. **(D)** Amino acid alignment

of a portion of the CHASE domain from different plant his-kinase cytokinin receptors

and the three *Hsf1* mutant alleles. Missense residues are marked by black triangles for

the *Hsf1* alleles and by a white triangle for the *Lotus snf2* allele. Amino acid sequences

derived from AT2G01830 (*AHK4*), AM287033 (*LHK1* and *LHK1-snf2*), XM_003570636

(*BdHK1*), XM_002454271 (*SbHK1*), NM_001111389 (*ZmHK1-NCBI*),

GRMZM2G151223 (*ZmHK1-MaizeGDB*), *ZmHK1* from the A619 inbred (*ZmHK1-*

AEWL) and the Mo17 inbred (*ZmHK1-1603* and *ZmHK1-1595*). **(E)** ZmHK1 receptors

with *Hsf1* mutations show CK independent growth in a yeast his-kinase signaling assay.

Growth of *S. cerevisiae sln*Δ mutant transformed with an empty vector, the ZmHK1

vector or one of the *Hsf1* mutant *ZmHK1* vectors on glucose media with no CK (DMSO)

or supplemented with different cytokinins - iP, tZ, or cZ. Growth on galactose media of

the *sln*Δ mutant transformed with each of the assayed vectors. DMSO, dimethyl

sulfoxide; iP, N^6-(Δ^2-isopentenyl)adenine; tZ, *trans*-zeatin; cZ, *cis*-zeatin. Dilutions of

yeast cultures (O.D.$_{600}$ = 1.0) for each yeast strain are noted on the left of each image.

(**F**) Ribbon diagram of the ZmHK1 CHASE domain with the *Hsf1* mutations (magenta)

noted and one molecule of N^6-(Δ^2-isopentenyl)adenine (blue and aqua) complexed in

the binding pocket. Arginine 192 (blue), in the loop domain (red) forming one face of

the binding cavity, is predicted to form a salt bridge with E236, the residue altered in
Hsf1-1603. Hsf1-1595 is P190L, *Hsf1-1603* is E236K and *Hsf1-AEWL* is L238F.

Figure 2.   Exogenous CK treatment phenocopies the *Hsf1* leaf development defects
and enhances the *Hsf1* mutation. (**A**) Phenotype of 3-week old wild type and
heterozygous *Hsf1*-1603/+ seedlings. Bar = 2 cm. (**B**) Phenotypes of 3-week old B73
water (- CK) and 10 µM 6-BAP treated (+ CK) seedlings. Bar = 2 cm. (**C**) Boxplots of
leaf sizes comparing wild type (WT) to *Hsf1-1603/+* sib seedlings, and B73 water (- CK)
and 10 µM 6-BAP treated (+ CK) seedlings. Horizontal bars represent the maximum,
third quantile, median, first quantile, and minimum values respectively, dots outside of
the plot are outliers, and the * indicates a *P*-value ≤ 0.0001 calculated from a two-tailed
Student's t-test. (**D**) Macrohair production on the abaxial sheath and auricle (white
triangles) of 2-week old B73 water (- CK) and 10 µM 6-BAP treated (+ CK) seedlings.
Insets show an adaxial view of the sheath-blade boundary of leaf 1. (**E**) Glue
impressions of adaxial leaf 1 blade from 2-week old B73 water (- CK) and 10 µM 6-BAP
treated (+ CK) seedlings showing increased macrohair presence in the medial blade
and at the margin. (**F**) CK-induced prong formation in B73 seedlings (n ≥ 12 for each
treatment). (**G**) Effect of CK treatment on prong formation in 2-week old *Hsf1-1603*/+
seedlings (yellow arrows mark prongs). Bar = 2 cm. (**H**) Frequency and leaf number
where the first prong formed in *Hsf1-1603/+* with (red) and without (blue) 10 µM 6-BAP
treatment (n ≥ 12 for each treatment). (**I**) Close-up of prongs formed on leaf 4 from CK-
treated and control *Hsf1-1603*/+ seedlings (in [G]). (**J**) Macrohair production on 2-week
old seedlings due to CK treatment or *Hsf1-1603*/+ mutation or both.

Figure 3.   Expression of CK signaling and responsive genes. (**A**) Relative mRNA
accumulation of CK genes in different tissues of 2-week old seedlings of the three *Hsf1*
alleles and WT sibs measured by qPCR. For each genotype, values are the means
(±SE) of three biological replicates consisting of tissue pooled from at least 3 plants. .

Asterisks indicate significant differences between WT and *Hsf1*/+ sib (Student's *t* test, P

≤ 0.05). GL – Green leaf, IL – immature leaf, SA –shot apex. (**B**) Pattern of *ZmHK1* and

ZmRR3 transcript accumulation in WT and *Hsf1-1603*/+ shoot apex. Longitudinal and

transverse sections were hybridized with *ZmHK1* or *ZmRR3* specific antisense probes.

The longitudinal section of *ZmRR3* hybridized to WT is not medial and so *ZmRR3*

expression appears to be apically localized, but it is not. Initiating leaf primordia (yellow

arrows) and leaf primordia margins (red triangles) are marked in the *Hsf1/*+ sections

probed with *ZmRR3*. Bar = 30 µm.

1180

Figure 4.   The *Hsf1* phenotype is enhanced by loss of *ZmRR3* function. (**A**)

Phenotypes of 30-day old (left to right) WT, *abph1/abph1*, *Hsf1-1603/*+, and *Hsf1-*

1603/+, abph1/abph1 mutants. This family segregated 9 wild type, 12 *abph1/abph1*, 10

Hsf1-1603/+, and 15 double *Hsf1-1603/*+, *abph1/abph1*, which fits a 1:1:1:1 expected

ratio. Inset shows a close-up of a double *Hsf1*, *abph1* mutant. Bar = 15 cm. (**B**)

Phenotypes of 60-day old plants segregating the same four genotypes in (**A**). Bar = 10

cm. Insets in the double mutant images show close-ups of prongs from that genotype.

Yellow and red arrowheads mark paired leaves on the *abph1* mutant and prongs on the

Hsf1/+ mutant, respectively.

1190

1191

1192

Supplemental Data

1193

1194

Supplemental Figure 1. *Hsf1* phenotypes.

Supplemental Figure 2. Prong formation is patterned in *Hsf1* leaves.

Supplemental Figure 3. ZmHK1 activity in heterologous yeast his-kinase assay.

Supplemental Figure 4. Comparison of ligand binding affinity constants of wild type
and mutant ZmHK1 receptors.

Supplemental Figure 5. Effects of CK treatment on leaf growth.

Supplemental Table 1. Mature plant phenotypes of the three *Hsf1* alleles.

Supplemental Table 2.   Frequency of prongs for the three *Hsf1* alleles by leaf position in the
upper shoot..

Supplemental Table 3: Primers used for positional cloning or genotyping

Supplemental Table 4. Primers used for expression analysis

1206

1207   **Tables**

1208

1209   **Table 1. Apparent affinity constants K$_D$* for wild type and mutant ZmHK1**
1210   **receptors with different cytokinins**

| Assay | Receptor | \multicolumn{8}{c}{K$_D$* for cytokinins (nM)} | | | | | | | |

Assay	Receptor	iP	BA	tZ	cZ	Kin	TD	DZ	Ade
Bacterial spheroplasts	ZmHK1	2.90	3.69	31.8	37.5	33.0	37.6	312.0	>10000
	AEWL	0.36	0.56	6.38	5.56	7.62	93.7	61.6	>10000
	1603	0.59	0.91	7.27	6.74	7.50	111.0	88.0	>10000
Tobacco membrane	ZmHK1	0.52	1.42	7.16	8.31	-	49.2	114.0	>10000
	1595	0.23	0.31	1.65	2.14	-	71.9	14.1	>10000

1211   iP, N^6-(Δ^2-isopentenyl)adenine; BA, 6-benzylaminopurine; tZ, *trans*-zeatin; cZ, *cis*-
1212   zeatin; Kin, kinetin; TD, thidiazuron; DHZ, dihydrozeatin, Ade, adenine.

1213

1214

1215

1216

1217

1218

1219

1220

1221 **Table 2. Fold increase of affinity to various cytokinins of mutant receptors**

1222 **compared to ZmHK1**

Cytokinin	Receptor		
	ZmHK1-AEWL	ZmHK1-1603	ZmHK1-1595
iP	8.06	4.92	2.26
BA	6.59	4.05	4.58
tZ	4.98	4.37	4.34
cZ	6.74	5.56	3.88
Kin	4.33	4.40	-
TD	0.4	0.39	0.68
DZ	5.06	3.55	8.09
Assay	Bacterial spheroplasts		Tobacco membrane

1223 iP, N^6-(Δ^2-isopentenyl)adenine; BA, 6-benzylaminopurine; tZ, *trans*-zeatin; cZ, *cis*-

1224 zeatin; Kin, kinetin; TD, thidiazuron; DHZ, dihydrozeatin, Ade, adenine.

1225

1226

1227
1228
Figure 1. *Hsf1* mutants alter leaf patterning and are caused by missense mutations
in the *ZmHK1* cytokinin receptor. **(A)** Adaxial view of half-leaves from WT and *Hsf1-*
1603/+ sibs showing the proximal-distal organization of the sheath (s), ligule (l), auricle
(a) and blade (b) and a prong outgrowth (red triangle). Bar = 5 cm. **(B)** Close-up of a
blade margin (b) from WT and *Hsf1-1603/+* showing a prong consisting of proximal leaf
segments – sheath (s), ligule (l) and auricle (a) juxtaposed to the blade (b). Bar = 1 cm.
(C) Comparison of leaf phenotypes between the three *Hsf1* alleles. L4 (top), 4th leaf
below tassel; L5 (bottom, 5th leaf below tassel. Bar = 10 cm. **(D)** Amino acid alignment
of a portion of the CHASE domain from different plant his-kinase cytokinin receptors
and the three *Hsf1* mutant alleles. Missense residues are marked by black triangles for

the *Hsf1* alleles and by a white triangle for the *Lotus snf2* allele. Amino acid sequences

derived from AT2G01830 (*AHK4*), AM287033 (*LHK1* and *LHK1-snf2*), XM_003570636

(*BdHK1*), XM_002454271 (*SbHK1*), NM_001111389 (*ZmHK1-NCBI*),

GRMZM2G151223 (*ZmHK1-MaizeGDB*), *ZmHK1* from the A619 inbred (*ZmHK1-*

AEWL) and the Mo17 inbred (*ZmHK1-1603* and *ZmHK1-1595*). **(E)** ZmHK1 receptors

with *Hsf1* mutations show CK independent growth in a yeast his-kinase signaling assay.

Growth of *S. cerevisiae sln*Δ mutant transformed with an empty vector, the ZmHK1

vector or one of the *Hsf1* mutant *ZmHK1* vectors on glucose media with no CK (DMSO)

or supplemented with different cytokinins - iP, tZ, or cZ. Growth on galactose media of

the *sln*Δ mutant transformed with each of the assayed vectors. DMSO, dimethyl

sulfoxide; iP, N^6-(Δ^2-isopentenyl)adenine; tZ, *trans*-zeatin; cZ, *cis*-zeatin. Dilutions of

yeast cultures (O.D.$_{600}$ = 1.0) for each yeast strain are noted on the left of each image.

(**F**) Ribbon diagram of the ZmHK1 CHASE domain with the *Hsf1* mutations (magenta)

noted and one molecule of N^6-(Δ^2-isopentenyl)adenine (blue and aqua) complexed in

the binding pocket. Arginine 192 (blue), in the loop domain (red) forming one face of

the binding cavity, is predicted to form a salt bridge with E236, the residue altered in

Hsf1-1603. *Hsf1-1595* is P190L, *Hsf1-1603* is E236K and *Hsf1-AEWL* is L238F.

1256

1257

1258

1259

1260

1261

1262

1263

1264
1265

1266 **Figure 2.** Exogenous CK treatment phenocopies the *Hsf1* leaf development defects
1267 and enhances the *Hsf1* mutation. (**A**) Phenotype of 3-week old wild type and
1268 heterozygous *Hsf1*-1603/+ seedlings. Bar = 2 cm. (**B**) Phenotypes of 3-week old B73
1269 water (- CK) and 10 μM 6-BAP treated (+ CK) seedlings. Bar = 2 cm. (**C**) Boxplots of

leaf sizes comparing wild type (WT) to *Hsf1-1603/+* sib seedlings, and B73 water (- CK) and 10 µM 6-BAP treated (+ CK) seedlings. Horizontal bars represent the maximum, third quantile, median, first quantile, and minimum values respectively, dots outside of the plot are outliers, and the * indicates a *P*-value ≤ 0.0001 calculated from a two-tailed Student's t-test. (**D**) Macrohair production on the abaxial sheath and auricle (white triangles) of 2-week old B73 water (- CK) and 10 µM 6-BAP treated (+ CK) seedlings. Insets show an adaxial view of the sheath-blade boundary of leaf 1. (**E**) Glue impressions of adaxial leaf 1 blade from 2-week old B73 water (- CK) and 10 µM 6-BAP treated (+ CK) seedlings showing increased macrohair presence in the medial blade and at the margin. (**F**) CK-induced prong formation in B73 seedlings (n ≥ 12 for each treatment). (**G**) Effect of CK treatment on prong formation in 2-week old *Hsf1-1603*/+ seedlings (yellow arrows mark prongs). Bar = 2 cm. (**H**) Frequency and leaf number where the first prong formed in *Hsf1-1603/+* with (red) and without (blue) 10 µM 6-BAP treatment (n ≥ 12 for each treatment). (**I**) Close-up of prongs formed on leaf 4 from CK-treated and control *Hsf1-1603*/+ seedlings (in [G]). (**J**) Macrohair production on 2-week old seedlings due to CK treatment or *Hsf1-1603*/+ mutation or both.

1294

1295
1296
Figure 3.   Expression of CK signaling and responsive genes. (**A**) Relative mRNA
accumulation of CK genes in different tissues of 2-week old seedlings of the three *Hsf1*
alleles and WT sibs measured by qPCR. For each genotype, values are the means
(±SE) of three biological replicates consisting of tissue pooled from at least 3 plants. .
Asterisks indicate significant differences between WT and *Hsf1/+* sib (Student's *t* test, P
≤ 0.05). GL – Green leaf, IL – immature leaf, SA –shot apex. (**B**) Pattern of *ZmHK1* and
ZmRR3 transcript accumulation in WT and *Hsf1-1603/+* shoot apex. Longitudinal and
transverse sections were hybridized with *ZmHK1* or *ZmRR3* specific antisense probes.
The longitudinal section of *ZmRR3* hybridized to WT is not medial and so *ZmRR3*
expression appears to be apically localized, but it is not. Initiating leaf primordia (yellow
arrows) and leaf primordia margins (red triangles) are marked in the *Hsf1/+* sections
probed with *ZmRR3*. Bar = 30 μm.

1314

1315
1316
Figure 4. The *Hsf1* phenotype is enhanced by loss of *ZmRR3* function. (**A**)
Phenotypes of 30-day old (left to right) WT, *abph1/abph1*, *Hsf1-1603/+*, and *Hsf1-*
1603/+, abph1/abph1 mutants. This family segregated 9 wild type, 12 *abph1/abph1*, 10
Hsf1-1603/+, and 15 double *Hsf1-1603/+*, *abph1/abph1*, which fits a 1:1:1:1 expected
ratio. Inset shows a close-up of a double *Hsf1*, *abph1* mutant. Bar = 15 cm. (**B**)
Phenotypes of 60-day old plants segregating the same four genotypes in [A]. Bar = 10
cm. Insets in the double mutant images show close-ups of prongs from that genotype.
Yellow and red arrowheads mark paired leaves on the *abph1* mutant and prongs on the
Hsf1/+ mutant, respectively.

1326
1327
1328