Title: Refinement and reactivation of a taste-responsive hippocampal network

Authors: Linnea E. Herzog[1], Donald B. Katz[1,2,3], and Shantanu P. Jadhav[1,2,3,*]

Affiliations:

[1]Graduate Program in Neuroscience, Brandeis University, Waltham, MA, 02453, USA

[2]Neuroscience Program, Department of Psychology, and Volen National Center for Complex

Systems, Brandeis University, Waltham, MA, 02453, USA

[3]Lead Contact

*Correspondence to: shantanu@brandeis.edu (S.P.J.)

Summary

Animals need to remember the locations of nourishing and toxic food sources for survival, a fact

that necessitates a mechanism for associating taste experiences with particular places. We

have previously identified such responses within hippocampal place cells [1], the activity of

which is thought to aid memory-guided behavior by forming a mental map of an animal's

environment that can be reshaped through experience [2-7]. It remains unknown, however,

whether taste-responsiveness is intrinsic to a subset of place cells, or emerges as a result of

experience that reorganizes spatial maps. Here, we recorded from neurons in the dorsal CA1

region of rats running for palatable tastes delivered *via* intra-oral cannulae at specific locations

on a linear track. We identified a subset of taste-responsive cells that, even prior to taste

exposure, had larger place fields than non-taste-responsive cells overlapping with stimulus

delivery zones. Taste-responsive cells' place fields then contracted, as a result of taste

experience, leading to a stronger representation of stimulus delivery zones on the track. Taste-

responsive units exhibited increased sharp-wave ripple co-activation during the taste delivery

session and subsequent rest periods, which correlated with the degree of place field

contraction. Our results reveal that novel taste experience evokes responses within a

preconfigured network of taste-responsive hippocampal place cells with large fields, whose

spatial representations are refined by sensory experience to signal areas of behavioral salience.

This represents a possible mechanism by which animals identify and remember locations where

ecologically relevant stimuli are found within their environment.

Keywords

Hippocampus, Taste, Chemosensation, Place cells, Sharp-wave ripple, Co-activation,

Reactivation

Results and Discussion

To investigate how hippocampal place cells responded to tastes during running, we recorded

the activity of CA1 neurons (n = 143, mean ± SEM: 28.6 ± 4.49 neurons/session; **Figure S1A**)

as rats (n = 5) ran back and forth on a linear track during a novel taste experience (**Figure 1A**).

Each experimental day consisted of three track-running sessions ('Pre,' 'Tastes,' and 'Post')

interleaved with rest sessions in a sleep box. During the 'Pre' and 'Post' probe sessions, rats

explored the track in the absence of tastes. During the 'Tastes' session, rats received aliquots of

either sweet or salty tastes, delivered directly to the tongue *via* intra-oral cannula when the rats

reached either of two designated locations along the track. Only experimental days that marked

rats' first exposure to this paradigm (in which animals were naïve to both the tastants and

delivery locations) were considered for further analysis. Isolated units were classified as putative

pyramidal cells (82.5%, 118/143) or interneurons (17.5%, 25/143) based on spike width and

firing rate (**Figure S1B**). Most pyramidal cells (99.2%, 117/118) could be categorized as place

cells (see STAR Methods, [1, 8]) and exhibited spatially reliable firing over the course of a

session (**Figure S1C**); a subset of these units (32.5%, n = 38/117) responded to tastes (**Figure**

1B, **Table S1**), consistent with previous findings [1]. Taste responses were observed only in

cells whose place fields overlapped with delivery zones during the 'Tastes' session (**Figure 1C**),

supporting previous work demonstrating that hippocampal sensory responses are gated by

place cells' spatial firing properties [1, 8, 9].

We previously established, using a passive taste delivery paradigm: 1) that a subset of

hippocampal neurons encode taste palatability; 2) that these taste responses are gated by (i.e.,

only appear when the rat is in its) place fields; and 3) that taste-responsive neurons tend to

have larger place fields than non-taste-responsive cells [1]. However, it remains unclear

whether or how these differences in spatial selectivity emerge as a consequence of taste

experience, and how they evolve with experience. We therefore compared the place field sizes

of taste-responsive (n = 38) and non-taste-responsive (n = 79) place cells in the track-running

session immediately preceding a novel taste experience (**Figure 2A**). Taste-responsive cells

had larger place fields than non-taste-responsive cells before the rats' first exposures to

saccharin and NaCl solutions (**Figure 2B**, **S2A**), revealing a relationship between initial place

field size and subsequent responsiveness to tastes (**Figure 2C, S2B**).

Cells identified as initially having large place fields were far more likely to later be

activated by taste stimuli. However, we found no direct relationship between taste response

magnitude and place field size (n = 42 cells, Pearson's correlation: r = -0.23, permutation test, p

= 0.14), arguing against the hypothesis that larger place fields simply connote higher sensitivity

to responses—while the likelihood of taste responsiveness is much higher in cells with larger

place fields, the magnitude of taste sensitivity is similar for taste-responsive cells with different

place field sizes. In addition, we have previously shown that hippocampal gustatory responses

specifically code for taste palatability (rather than responding to any and all tastes) [1], rendering

the possibility unlikely that spatially sparse neurons are more indiscriminately sensitive to

sensory modulation, regardless of modality.

These results suggest that gustatory (or perhaps reward) receptivity is to some degree a

basic "feature" of the hippocampal system – perhaps reflecting differential input from the medial

(MEC) and lateral (LEC) entorhinal cortices, which represent the two major cortical inputs to the

hippocampus, encoding spatial and non-spatial information, respectively [10-13] — with initial

differences in place field size potentially arising from preexisting synaptically connected

ensembles [14-16].

Further, following taste exposure, taste-responsive cells' place fields shrink, becoming

comparable to those of non-taste-responsive place cells; non-taste-responsive cells' place

fields, meanwhile, did not change in size from the Pre to the Post session (**Figure 2B**).

Analyses of local information content [17] yielded similar results. Since place fields are known to

accumulate in rewarded locations [18-19], it is reasonable to hypothesize that this

reorganization might function to localize place fields to taste delivery zones. However, we

observed comparable degrees of location remapping in taste- and non-taste-responsive cells

(**Figure 3A**) — no overall differences in Pre-Post place field location or place field-taste zone

distance were found between groups (**Figure 3B**). Instead, taste-responsive cells exhibited high

spatial overlap with taste delivery zones during the Pre session. Contraction of their place fields

with taste experience decreased overall overlap (**Figure 3C**), but increased place field density in

the delivery zones, such that the place field peaks of taste-responsive cells tended to more

reliably overlap with taste zones in the Post session (**Figure 3D**). This refinement was observed

in taste-responsive cells whose place fields initially overlapped with taste delivery zones (**Figure**

3E); lower proportions of non-taste-responsive cells exhibited overlap (29/38 vs. 44/79, χ^2 test, χ

$= 4.65$, $p = 0.031$). These changes in spatial selectivity could not be attributed to interneuron

firing in the taste zone, which tended to decrease during the Tastes session, then increase

again during Post, becoming comparable to the Pre-session rate (1-way ANOVA, $p = 3e-04$,

multiple comparisons tests, Pre vs. Taste: $p = 2e-04$, Tastes vs. Post: $p = 0.032$).

In summary: taste-responsive cells initially have large place fields that overlap with

future locations of taste delivery as taste responses are gated by place fields; these place fields

then contract following a novel taste experience, leading to a more precise representation of

taste delivery locations in animals' mental maps of the track. Non-taste-responsive cells do not

exhibit this effect, instead representing the track continuously throughout taste experience.

One possible mechanism underlying this representational reorganization is the co-firing

of goal-related assembly patterns during sharp-wave ripples (SWRs; [20-21]), a high-frequency

hippocampal oscillation that is necessary for memory consolidation during wake [22] and sleep

[23]. During ripples, the activity of sequences of place cells is typically "replayed" in a time-

compressed manner [24-25]; sensory cues are known to influence this SWR reactivation [26]

and replay content [27]. We therefore hypothesized that taste-responsive place cells would

exhibit particularly strong SWR co-activation during waking behavior and sleep, as

reorganization occurred.

To test this hypothesis, we identified SWR co-activation events (**Figure S3A**) in taste-

and non-taste-responsive cell pairs (**Figure S3B**) whose place fields exhibited high and low

spatial overlap. As expected [24, 28-30], highly overlapping cell pairs co-fired more during wake

and sleep SWRs than pairs with low overlap (**Figure 4A**). This effect persisted when we

calculated the mean co-activation differences for all possible combinations of cell pairs

(Taste/Taste or Non/Non – Taste/Non) across sleep and wake (**Figure 4B**). Highly overlapping

taste-responsive cell pairs exhibited higher co-activation differences than similarly overlapping

non-taste-responsive cell pairs during taste experience and post-experience sleep (**Figure 4B**,

S4C). This effect parallels the changes observed in place field size and spatial coverage on the

track—we saw that taste-responsive cells' place fields contract over areas of taste delivery

following the Tastes session, while non-taste-responsive cells remain stable across taste

experience. Furthermore, we observed a significant correlation between place field contraction

(from Pre to Post) and SWR reactivation probability during the Tastes session (**Figure 4C**,

S4D). These data support the hypothesis that reactivation during SWRs can help stabilize

animals' mental maps as they encounter tastes within their environment, with taste-responsive

units coming to more precisely signify areas where food can be found.

Overall, our data show that taste responsivity is intrinsic to a subset of hippocampal

place cells, which initially possess larger place fields than non-taste-responsive cells prior to

taste exposure. The introduction of novel tastes in a familiar environment led to the contraction

of taste-responsive cells' place fields over delivery zones—a refinement of animals' internal

representations of the track to emphasize locations of taste consumption. This sharpening of

place fields only took place during animals' first exposure to sweet and salty tastes (place field

size, second experimental day: 2-way ANOVA, cell type: $p = 1$, epoch: $p = 0.57$, interaction: $p = $

0.96), revealing that the introduction of novel tastes in distinct spatial locations is what drives

response plasticity, not unlike how place fields themselves classically emerge and stabilize over

animals' first exposure to a new environment [31].

The idea that rewarding or aversive experiences can reorganize hippocampal

representations of an environment to encode points of behavioral salience is not new [18-19,

32-33], but we have shown that these shifts arise from changes in the spatial properties of a

preexisting network of sensory-responding cells, which refine their spatial representations. One

possible mechanism underlying this shift is the increased SWR co-activation of taste-responsive

units with high spatial overlap, which correlated with the degree of place field contraction that

we observed over taste experience. Our findings are consistent with previous observations that

introducing novel stimuli, whether rewarding [18-19] or aversive [32-33], leads to the increased

reactivation of behaviorally relevant trajectories during SWRs and reorganizes animals'

hippocampal representations to map out areas of high salience within an environment. These

results provide further evidence for a flexible memory framework where hippocampal neurons

can multiplex contextual information and sensory cues to update animals' mental maps of an

environment [4, 34-37].

Acknowledgments

This work was supported by the Alfred P. Sloan Foundation (Sloan Research Fellowship in

Neuroscience to S.P.J.), the Whitehall Foundation (S.P.J.), and the National Institutes in Health

(NIH Grants R01 DC006666 and R01 DC007703 to D.B.K. and Training Grant T90 DA032435

to L.E.H.).

Author Contributions

All authors contributed to all aspects of this report.

Declaration of Interests

The authors declare no competing interests.

Figure Legends

Figure 1. Experimental design and example taste responses.

A, Experimental schematic. Each experiment day consisted of seven sessions: three track-

running sessions (Pre, Tastes and Post), interleaved with four sleep sessions (S1-4). For the

Pre and Post probe sessions, rats ran on the track in the absence of tastes. During the Tastes

session, rats received a randomized delivery of either sweet (green = S, 4 mM saccharin) or

salty (yellow = N, 100 mM sodium chloride) taste solution upon entering either of the two

designated regions on the track (Zone 1 and Zone 2). Neurons from the dorsal CA1

hippocampal region were recorded continuously throughout all sessions (see also **Figure S1**

and **Table S1**). *B,* Example in- and out-of-field taste responses. Top panels depict the firing rate

(FR) maps (with peak FR denoted in Hz) and line plots for an example taste- and non-taste-

responsive cell during the Tastes session, calculated with taste delivery periods either included

or excluded from the analysis. Below, taste trial locations for the two example sessions are

shown, with in-field trials (shown here using the taste-times-excluded place field) outlined in

pink. Middle panels depict raster plots of each cell's in- and out-of-field responses to each taste.

Black dots indicate spike times during each individual trial aligned to the time of taste delivery.

Bottom panels show the mean FR in response to saccharin or NaCl deliveries occurring in- and

out-of-field for each cell. Note that taste responses are only found within the taste-responsive

cell's place field, as previously reported [1]. *C*, Magnitude of taste responsiveness/specificity (η^2)

for the in- and out-of-field regions of taste- (*n* = 20) and non-taste-responsive (*n* = 22) cells that

met our criteria ($\geq$ 10 trials in- and out-of-field). η^2 was highest (2-way ANOVA, cell type: **p =

0.0038, epoch: **p = 0.0019, interaction: *p* = 0.13) for taste-responsive cells in-field (multiple

comparisons tests, Taste/Out-of-field: *p = 0.012; Non/In-field: **p = 0.0094; Non/Out-of-field:

***p = 2.1e-04), indicating that taste responses were observed only in cells with place fields

overlapping taste delivery zones. Similar results were obtained when taste delivery times were

included in place map generation.

Figure 2. Taste-responsive place cells become spatially selective after taste experience.

A, Example taste- and non-taste-responsive place cell firing during the Pre and Post sessions.

Top panels depict trial raster plots, with black dots indicating spike locations along each run of

the track. Middle panels show the average linearized firing rate (FR) maps for each cell, with the

peak FR denoted in Hz. Bottom panels depict these FR maps as a line plot, with place field (PF)

boundaries indicated by pink shading. ***B***, Mean PF size for taste- ($n = 38$) and non-taste-

responsive ($n = 79$) cells, normalized as a fraction of track coverage. Taste-responsive cells

initially had larger PF (2-way ANOVA, cell type: ***$p = 9.8$e-06, epoch: **$p = 0.0037$, interaction:

$p = 0.052$), which contracted after taste experience (multiple comparisons tests, Pre/Non: ***$p =$

4.6e-05; Post/Tastes: *$p = 0.020$; Post/Non: ***$p = 1.9$e-06; see also **Figure S2**). ***C***, Likelihood

of taste-responsiveness as a function of Pre PF size. Each bar represents the percentage of

taste-responsive (TR) cells in each size bin, which increases with increasing place field size.

Figure 3. Taste-responsive cells' place fields contract around areas of taste delivery.

A, Spatial firing maps of example taste- and non-taste-responsive cells during the Pre and Post

track-running sessions. Numbers on each plot denote peak spatial firing rate (FR) in Hz. Place

field (PF) boundaries during each session are indicated by pink shading, while the peak

distributions of trials (taste deliveries) during the Tastes session (taste zone, or TZ) are

indicated by gray boxes. Note how taste-responsive cells' place fields overlap with TZ across

taste experience, while non-taste-responsive cells' do not. ***B***, Change in PF peak location and

PF-TZ distance from Pre to Post. No differences in the mean Δ PF peak location (unpaired t-

test, $p = 0.54$) or Δ PF-TZ distance (unpaired t-test, $p = 0.55$) were observed between groups or

shuffled controls (paired t-tests, Δ PF peak location: Taste vs. Shuffle: $p = 0.25$, Non vs. Shuffle:

$p = 0.17$; Δ PF-TZ distance, Taste vs. Shuffle: $p = 0.89$, Non vs. Shuffle: $p = 0.31$), indicating

that taste- and non-taste-responsive cells exhibited comparable location remapping from the

213 Pre to Post session. *C*, Mean PF-TZ overlap for taste- and non-taste-responsive cells. The

214 place fields of taste-responsive cells exhibited high overlap with taste delivery zones during the

215 Pre session (2-way ANOVA, cell type: ***p = 9.6e-08, epoch: ***p = 6.3e-08, interaction: *p =

216 0.020; multiple comparisons tests, Pre/Taste vs. Pre/Non: ***p = 4.5e-07), which decreased with

217 experience for both taste- (***p = 1.5e-05) and non-taste-responsive units (*p = 0.029). *D*,

218 Proportion of taste- and non-taste-responsive cells with PF peaks in TZ. Bars show the

219 percentage of cells with PF peaks located within the TZ, compared to chance level (calculated

220 based on average TZ coverage of the track). A greater-than-chance proportion of taste-

221 responsive cells had PF peaks in the TZ during Post (binomial test, ***p = 3.9e-05),

222 underscoring increased density of taste-responsive cell field peaks in TZ after experience. *E*,

223 Mean PF size as a function of PF-TZ overlap during Pre. TZ-overlapping taste-responsive cells

224 had larger PF (3-way ANOVA, cell type: *p = 0.015, epoch: p = 0.10, overlap: ***p = 3.7e-06,

225 cell type x epoch: p = 0.42, cell type x overlap: *p = 0.013, epoch x overlap: *p = 0.018, cell type

226 x epoch x overlap: p = 0.097) than any other group (multiple comparisons tests, vs.

227 Pre/Non/Overlap: ***p = 5.8e-05; Pre/Taste/No overlap: ***p = 1.6e-04; Pre/Non/No overlap:

228 ***p = 1.5e-08; Post/Taste/Overlap: **p = 0.0039; Post/Non/Overlap: ***p = 3.9e-07;

229 Post/Taste/No overlap: **p = 0.0026; Post/Non/No overlap: ***p = 1.0e-08), revealing that the

230 contraction effect shown in **Figure 2B** is confined to taste-responsive cells with PF initially

231 overlapping the TZ.

233 **Figure 4.** Taste-responsive cells exhibit increased SWR co-activation during taste experience.

234 *A*, Spatial firing plots and awake and sleep SWR co-activation for three example cell pairs

235 (taste-responsive pairs with high and low overlap, non-taste-responsive pair with high overlap)

236 during track running sessions (see also **Figure S3**). Numbers on each place map plot denote

237 peak spatial firing rate (FR) in Hz. Place field (PF) boundaries during each session are indicated

238 by pink lines. The bar plots on the right show the Z-scored co-activation for each cell pair during

wake and sleep SWRs. Note how the high overlap pairs exhibit more co-activation than the low

overlap pairs. **B**, SWR co-activation differences between taste- and non-taste-responsive cell

pairs across taste experience. Bars display the mean difference in SWR co-activation between

same (Taste/Taste or Non/Non) and different (Taste/Non) taste pairs for all significant co-

activation events. Significant differences were found for both wake (3-way ANOVA, cell type:

***$p = 3.7e-04$, overlap: ***$p = 9.1e-74$, cell type x epoch: **$p = 0.0067$, epoch x overlap: ***$p =$

$1.1e-33$) and sleep (cell type: ***$p = 5.6e-09$, epoch: ***$p = 4.6e-08$, overlap: ***$p = 2.9e-39$, cell

type x epoch: ***$p = 2.3e-14$). Post hoc comparisons revealed that taste-responsive cell pairs

with high overlap had higher co-activation differences than non-taste-responsive cell pairs

during taste experience (*$p = 0.046$) and post-experience sleep (***$p = 1.1e-15$). **C,** Each cells'

reactivation probability during the Tastes session is plotted against the decrease in PF size from

the Pre to Post session. The significant correlation ($n = 117$ cells, Pearson's correlation: $r =$

0.29; permutation test: **$p = 0.002$) suggests that the observed increase in reactivation plays a

role in reorganizing animal's mental maps to better represent locations of taste delivery after

experience.

STAR Methods

Contact for Reagent and Resource Sharing

Further information and requests for resources should be directed to the Lead Contact, Dr.

Shantanu P. Jadhav (shantanu{at}brandeis.edu).

Experimental Model and Subject Details

Five adult male Long-Evans rats (450-550 g, 4-6 months) served as subjects in this study. Rats

were individually housed and kept on a 12 h light/dark cycle, with all sessions taking place

around the same time during the light period. All surgical and experimental procedures were

conducted in accordance with the National Institutes of Health guidelines and approved by the

Brandeis University Institutional Animal Care and Use Committee.

Method Details

Experimental design

After several weeks of habituation to daily handling, animals were pre-trained to seek water

rewards delivered via reward well at either end of an elevated linear track, and habituated to an

elevated, opaque sleep box [22]. After animals learned to run back and forth on the track, they

were chronically implanted with a multi-tetrode drive and an intra-oral cannula (see *Surgical*

implantation and electrophysiology).

Following recovery from the implantation surgery (~7-8 days), rats were water-deprived

to 85-90% of their *ad libitum* weight and re-trained using the same paradigm as described

above, but with one change – water rewards delivered directly via intra-oral cannula instead of

at reward wells. At ~14 d after implantation, we performed daily recording sessions in which rats

ran back and forth on the linear track during a novel taste experience (see **Figure 1A** and

Zoned taste paradigm), where animals received sweet and salty tastes at two locations, located

~1/3 and 2/3 of the way down the track. Palatable taste solutions, which naïve rats prefer over

water [38-39], were used in order to motivate the animals' behavior. Animals were naïve to

these tastes prior to experiment day, as well as taste exposure in these locations, which were

distinct from those experienced during pre-training. Only data from animals' first day of

exposure to this paradigm were used for the analyses described in the paper, unless otherwise

specified. Following the conclusion of these zoned taste experiments, additional recordings

were conducted while rats received a battery of basic tastes; data from these experiments has

been presented previously to demonstrate that hippocampal neurons respond to tastes [1].

Following the conclusion of all experiments, we made electrolytic lesions through each

electrode tip to mark recording locations. After 12-24 hours, animals were euthanized and

intracardially perfused with 4% formaldehyde using approved procedures. Brains were fixed for

24-48 hours, cryoprotected (30% sucrose in 4% formaldehyde), and stored at 4°C. Brains were

sectioned into 50 μm slices and stained with cresyl violet to confirm electrode placement in the

hippocampal cell layer (**Figure S1A**).

Zoned taste paradigm

Each recording session typically lasted between 2 and 3 hours, and consisted of three running

sessions on a 150 cm elevated linear track interleaved with four 15-20 minute sleep sessions in

a ~30 x 30 x 40 cm black box (rest box). The first (Pre) and last (Post) sessions on the linear

track consisted of 15-20 minute periods in which animals ran back and forth on the track in the

absence of tastes (mean number of track runs per session: 17.2 ± 3.7 during Pre, 14.0 ± 3.2

during Post). During the 30-90 minute Tastes session, rats received a pseudo-randomized

delivery of either a sweet [4 mM saccharin (S)] or salty [100 mM sodium chloride (N)] taste

solution at designated areas ("taste zones," or TZ) located approximately 1/3 and 2/3 of the way

down the track (mean number of taste trials per session: 91.2 ± 17.7). Rats consistently slowed

down after taste delivery (average speed in the 2.5 s before taste delivery: 11.8 +/- 0.32 cm/s,

average speed in the 2.5 s after taste delivery: 3.11 +/- 0.10 cm/s; unpaired t-test, $p = 1.1e\text{-}$

122). The rat's position was detected in real-time with an overhead monochrome CCD camera

(30 fps) and LEDs affixed to the recording headstage, and used to trigger taste deliveries when

animals entered into one of the two taste delivery zones. Taste solutions were delivered directly

onto the tongue in ~40 μL aliquots via two polyamide tubes inserted into the intra-oral cannula,

with a separate tube for each solution to prevent the mixing of tastes. Animals had access to an

additional 15-20 mL of water in their home cage after the end of each experiment day.

Surgical implantation and electrophysiology

Following pre-training on the linear track and habituation to the sleep box, animals were

implanted with a microdrive array consisting of 25-30 independently moveable tetrodes in the

right dorsal hippocampal region CA1 (-3.6 mm AP, 2.2 mm ML), and an intra-oral cannula. Each

intra-oral cannula consisted of a polyethylene tube inserted beneath the left temporalis muscle

and terminating anterolateral to the first maxillary molar, allowing for the precise delivery of taste

solutions onto the rat's tongue [38-40].

316   Electrophysiological recordings were conducted using a SpikeGadgets system [41].

Spikes were sampled at 30 kHz and bandpass filtered between 600 Hz and 6 kHz. Local field

potentials (LFPs) were sampled at 1.5 kHz and bandpass filtered between 0.5 and 400 Hz. Over

~14 d following surgery, tetrodes were gradually advanced to the CA1 hippocampal cell layer,

as identified by characteristic EEG patterns (SWRs; theta rhythm) as previously described [22,

41-42]. Tetrodes were readjusted after each day's recordings. Each animal had one

hippocampal reference tetrode in corpus callosum, which was also referenced to a ground

screw installed in the skull overlying cerebellum.

Unit inclusion

Spikes were sorted as described previously [1, 41-42]. In brief, single units were isolated offline

based on peak amplitude and principal components (Matclust, M.P. Karlsson). Only well-

isolated units with stable waveforms that fired at least 100 spikes per session were included in

our analysis. Putative interneurons (*Int*) were identified on the basis of firing rate (> 8.5 Hz) and

spike width (< 0.35 ms) parameters (**Figure S1B**). All other isolated units were classified as

putative pyramidal cells (*Pyr*). We isolated a total of 143 neurons from five rats, conducted

across five experiments, and recorded throughout the experiments. **Table S1** shows the

distribution of cells across all five animals.

Quantification and Statistical Analysis

Sharp-wave ripple detection

SWRs were detected as previously described [41-42] using the ripple-band (150-250 Hz)

filtering of LFPs from multiple tetrodes. A Hilbert transform was used to determine the envelope

of band-passed LFPs, and events that exceeded a threshold (mean + 3 SD) were detected.

SWR events were defined as the times around initially detected events when the envelope

exceeded the mean. SWR periods were excluded from place field analysis, similar to previous

studies [1, 41-42].

Taste response properties

Hippocampal responses to tastes were characterized as described previously ([1]; for

characterization of response properties in other parts of the taste system, see [40, 43-46]. Since

we have already extensively described the coding content of taste responsiveness in these

neurons [1], and given our use of only two tastes in the current context, here we perform (except

when noted, see *Taste selectivity* below) a simpler test of taste responsiveness. In brief,

neurons were classified as taste-responsive if they exhibited responses to taste presence

(divergence from baseline) and/or identity (taste-specific differences in evoked firing) in the

2500 ms of spiking activity following each taste delivery; all other units were classified as non-

taste-responsive (see **Figure 1B** for examples).

Taste selectivity

The magnitude of taste responsiveness and specificity for each cell was quantified using eta-

squared (η^2), a standard measure of ANOVA effect sizes that describes the proportion of

variance in a dependent variable explained by each factor:

355              $\eta^2 = SS_{(factor)}/SS_{(total)}$,

where SS is the sum of squares [1, 47]. In our analysis, we used the summed SS of the two

main factors (time + taste) to calculate η^2. As described below (see *In-field vs. out-of-field*

analysis), a 2-way ANOVA was used to assess differences in η^2 for the in-field and out-of-field

regions of taste-responsive and non-taste-responsive place cells that fit our analysis criteria (> 1

Hz place field peak, $\geq$ 10 in- and out-of-field trials; **Figure 1C**). To determine if there was a

general relationship between taste response magnitude and place field size, we used

permutation tests to compare the Pearson's correlation between in-field η^2 and PF size during

the Tastes session (*n* = 42 cells that fit our analysis criteria with taste delivery times excluded in

place map generation, see *In-field vs. out-of-field analysis*) with 1000 cell ID-shuffled controls.

Spatial maps

To visualize the spatial firing properties of neurons, 2D position data were converted to linear

positions, and one-dimensional occupancy-normalized firing rate maps (**Figure 1B**, **2A**, **3A**, **4A**)

were constructed using 2 cm bins and smoothed with a 1D Gaussian (σ = 5 cm) as previously

described [22, 41]. Place fields from each session ('Pre,' 'Tastes,' and 'Post') were detected

individually. Data from SWR periods (see *Sharp-wave ripple detection*) and periods of

immobility (< 3 cm/s) were excluded from spatial map analysis.

Peak rates for each cell were defined as the maximum firing rate across all spatial bins

in the spatial map. Each cell's place field was defined as the largest group of neighboring spatial

bins which had firing rates $\geq$ 20% of the peak rate (in some cases, multiple regions on the track

met this criteria, in which case only the largest is depicted as the place field); sizes were then

calculated by multiplying the number of bins by the bin size [48]. Only place cells (*n* = 117,

defined as pyramidal cells whose unsmoothed peak firing rate exceeded 1 Hz [1, 8]) were

included in analyses of spatial firing.

Within-session spatial reliability (**Figure S1C**) was measured by dividing each of the

'Pre', 'Tastes' and 'Post' sessions into three segments. The correlation coefficients of firing

maps generated in each segment (first vs. middle, middle vs. last, first vs. last) were compared

separately for taste-responsive ($n = 38$) and non-taste-responsive ($n = 79$) cells using 1-way

ANOVAs [49].

To compare place field sizes across sessions, place field size was normalized as a

fraction of the track. A 2-way ANOVA was used to determine whether the mean place field size

differed significantly between taste-responsive and non-taste-responsive place cells before and

after animals' first exposure to the zoned tastes paradigm (**Figure 2B**; this analysis was also

conducted separately for repeated exposures; $n = 6$ taste-responsive and $n = 49$ non-taste-

responsive cells across 4 sessions). Positional information at peak firing locations [17] was

similarly analyzed across sessions and cell types. Importantly, these comparisons of place field

size and local information content were between the Pre and Post sessions, which were

identical in nature (neither involves taste delivery) and speeds maintained by rats traversing the

track. The mean PF size for individual animals is depicted in **Figure S2A**. The percentages of

taste- and non-taste-responsive cells in each spatial bin (0.2*n as a fraction of track coverage)

were compared using χ^2 tests (**Figure S2B**). A plot of the percentage of taste-responsive cells

in each spatial bin was also constructed (**Figure 2C**).

To determine how taste experience affects place field position (**Figure 3B**), the mean

change in place field peak location (Δ PF peak location) and place field-taste zone distance (Δ

PF-TZ distance) from Pre to Post were compared for taste- and non-taste-responsive cells

using unpaired t-tests. The taste zones were defined separately for each experiment day using

a Gaussian fit to determine the peak distribution of trial locations during the Tastes session

(defined as μ ± σ for the normal distribution of all trial locations, calculated separately for each

delivery zone).The closest taste zone to a place field's peak location was used to calculate PF-

TZ distance. Paired t-tests were used to compare both the mean Δ PF peak location and Δ PF-

TZ distance for each group with controls that were calculated by randomly shuffling values from

the Post session.

To compare the PF-TZ overlap across sessions (**Figure 3C**), the track was divided into

100 spatial bins, and we calculated which bins each PF and TZ spanned along the normalized

track. PF-TZ overlap was defined as the number of taste zone locations that overlapped with a

cell's place field, divided by the total number of taste zone locations. 2-way ANOVAs were used

to compare the PF-TZ overlap between taste- and non-taste-responsive cells before and after

taste experience.

To determine if greater-than-chance numbers of cells in any group had place field peaks

inside the taste delivery zone (**Figure 3D**), binomial tests were used to compare the observed

vs. expected proportions of taste-responsive and non-taste-responsive cells with PF peaks in

the TZ during Pre and Post. The expected proportion (25.4%) was calculated based on the

average TZ coverage of the track, operating under the assumption that place fields exhibit equal

coverage of the track.

To determine how initial overlap with taste delivery zones affects place fields (**Figure**

3E), taste- and non-taste responsive cells were grouped according to their TZ overlap during the

Pre session ($n = 29$ overlapping taste-responsive cells, $n = 9$ non-overlapping taste-responsive

cells, $n = 44$ overlapping non-taste-responsive cells, $n = 35$ non-overlapping non-taste-

responsive cells); place field sizes were then compared between all groups using a 3-way

ANOVA. The proportions of TZ-overlapping units were also compared using a χ^2 test.

In-field vs. out-of-field analysis

To analyze how cells responded to tastes delivered inside or outside of their place fields (**Figure**

1B), only place cells that contained at least ten in- and out-of-field deliveries of each taste were

considered. This analysis was conducted separately using spatial maps calculated with taste

delivery times (from 500 ms before to 2500 ms after taste delivery) included ($n = 22$ taste-

responsive and $n = 21$ non-taste-responsive cells) or excluded ($n = 20$ taste-responsive and $n =$

22 non-taste-responsive cells). A 2-way ANOVA was used to assess differences between the

mean in-field and out-of-field taste response magnitude (calculated using η^2, see *Taste*

selectivity) of taste-responsive and non-taste-responsive cells (**Figure 1C**).

Analysis of interneuron firing rates

To determine whether changes in place cell firing were influenced by interneuron activity in the

taste zone, a 1-way ANOVA was used to compare the average normalized firing rates of

interneurons in the TZ across sessions [50].

SWR co-firing

SWRs co-activation events were defined as ripples during which both cells in a given cell pair

fired spikes (example in **Figure S3A**; locations of all awake ripples and co-firing events are

plotted in **Figure S3B**). Co-activation of cell pairs during awake and sleep SWRs was calculated

as previously described [30]:

If n_A and n_B represent instances when neurons A and B were independently active during N

events, then the expected number of events during which both neurons were active (n_{AB})

follows a hypergeometric distribution [51] with mean

$E[n_{AB}|N, n_A, n_B] = n_A n_B / N$

and variance

$\sigma^2[n_{AB}|N, n_A, n_B] = \dfrac{n_A n_B (N - n_A)(N - n_B)}{N^2(N - 1)}$

We use a Z-score to normalize across cell pairs with different activity levels (see right panels of

Figure 4A):

$Z = \dfrac{n_{AB} - E[n_{AB}|N, n_A, n_B]}{\sigma^2[n_{AB}|N, n_A, n_B]}$

Only cell pairs that fired at least 100 spikes per session, had a peak firing rate exceeding 3 Hz

during run sessions, and co-fired at least 5 times during SWRs were considered for analysis.

Place field overlap for pairs of cells was defined as twice the sum of the overlapping areas of

the linear rate curves divided by the sum of the areas of each curve [22]; all cell pairs with

overlap $\geq$ 0.5 were classified as high overlap (see examples in top and bottom panels of **Figure**

4A), while all cell pairs with overlap < 0.5 were classified as low overlap (see example in middle

panel of **Figure 4A**). For sleep SWRs, the closest running session was used to calculate spatial

overlap.

460   The mean co-activation difference was defined as the difference in Z-scored co-

activation between all combinations of same (Taste/Taste or Non/Non) – different (Taste/Non)

cell pairs during awake running sessions (Pre, Tastes and Post) as well as the Pre rest session

before the run sessions and the Post rest sessions after taste experience (averaged across the

two rest sessions, S3 and S4). For these comparisons, same cell pairs (Taste or Non) with a

greater spatial overlap than their corresponding different cell pair were classified as high overlap

(High), while same cell pairs with a smaller spatial overlap than their corresponding different cell

pair were classified as low overlap (Low). A 3-way ANOVA was used to compare the cell type,

overlap and epoch effects on the mean co-activation difference during awake and sleep SWRs

(**Figure 4B**; number of same-different cell pair combinations; wake: $n = 96$ Pre/Taste/High, $n =$

332 Pre/Non/High, $n = 96$ Pre/Taste/Low, $n = 310$ Pre/Non/Low, $n = 74$ Tastes/Taste/High, $n =$

320 Tastes/Non/High, $n = 118$ Tastes/Taste/Low, $n = 322$ Tastes/Non/Low, $n = 107$

Post/Taste/High, $n = 305$ Post/Non/High, $n = 85$ Post/Taste/Low, $n = 337$ Post/Non/Low; sleep:

$n = 261$ Pre/Taste/High, $n = 2221$ Pre/Non/High, $n = 182$ Pre/Taste/Low, $n = 4475$ Pre/Non/Low,

$n = 248$ Post/Taste/High, $n = 3990$ Post/Non/High, $n = 195$ Post/Taste/Low, $n = 2706$

Post/Non/Low). Means for individual animals are shown in **Figure S3C**.

476   In order to assess the relationship between SWR activation and place field refinement

(**Figure 4C**), the reactivation probability was computed for all units as previously described [19];

a Pearson's correlation between reactivation probability and the difference in PF size from the

Pre to Post sessions was then calculated. Permutation tests using 1000 cell ID-shuffled controls

were used to assess the significance of this correlation. Means for individual animals are

depicted in **Figure S3D**.

Statistical analysis

All statistical tests were performed using custom routines in MATLAB (MathWorks, Natick, MA,

RRID: SCR_001622) and evaluated at a level of $\alpha = 0.05$, with a Bonferroni correction applied

for multiple comparisons. Unless otherwise noted, values and error bars in the text denote

means ± SEM.

Data and Software Availability

The data and code used in this study are available upon reasonable request by contacting the

Lead Contact, Dr. Shantanu P. Jadhav (shantanu{at}brandeis.edu).

Supplemental Information titles and legends

Figure S1. Histology, unit classification and spatial reliability, related to Figure 1.

A, Histological verification of tetrode locations in intermediate dorsal CA1. *B*, Classification of

putative interneurons (Int, gray crosses) from pyramidal cells (Pyr, black circles) based on spike

width (> 8.5 Hz) and firing rate (< 0.35 ms) parameters. *C*, Spatial reliability of place cells across

sessions. There were no significant changes in the mean correlation coefficient of taste-

responsive (*n* = 38) and non-taste-responsive (*n* = 79) cells' place field maps across each third

of the session, indicating that spatial firing was reliable within each session (1-way ANOVA,

Taste responsive, Pre: *p* = 0.99, Tastes: *p* = 0.21, Post: *p* = 0.55; Non-taste-responsive, Pre: *p*

= 0.27, Tastes: *p* = 0.72, Post: *p* = 0.98).

Table S1. Cell distribution across animals, related to Figure 1.

Summary of the number of taste-responsive and total CA1 cells recorded from each animal.

Only cells meeting the inclusion criteria (see STAR Methods) are reported. Putative pyramidal

cells (Pyr) and interneurons (Int) were identified on the basis of firing rate and spike width

parameters. Neurons were classified as "taste-responsive" if they exhibited responses to taste

presence or identity.

Figure S2. Taste responsiveness is related to place field size during the Pre session,

related to Figure 2.

A, Place field size in **Figure 2B** shown for individual animals. *B*, Histogram of taste-responsive

and non-taste-responsive place field sizes. Stacked bars represent the percentage of cells

within each cell type that fell within a given size bin (n*0.2 as a fraction of the track). Cells with

smaller place fields were more likely to be non-taste-responsive (PF size < 0.2, χ^2 test, χ = 6.5,

p = 0.011), while cells with larger place fields were more likely to be taste-responsive (PF size

= 0.4-0.6, χ = 7.5, **p* = 0.0063; PF size > 0.6, χ = 6.4, *p* = 0.011).

Figure S3. SWR co-activation and distribution of significant events along the track,

related to Figure 4.

A, Example of one co-activation event following a taste trial, with saccharin delivery denoted by

the vertical green line. Spikes from two taste-responsive cells are shown in black, with spikes

that occurred during the same SWR inside the purple bar. SWRs were detected using the

simultaneously recorded EEG filtered at 150-250 Hz (black traces above raster plot). The top

trace shows the SWR in higher magnification. *B*, Probability of SWRs and significant co-

activation events for taste- and non-taste-responsive cell pairs at each location on the track

during awake running sessions. For visualization, taste zones (TZ, gray shaded areas) are

depicted as an average across sessions. *C*, Mean co-activation differences from **Figure 4B**

shown for individual animals (each column corresponds to a bar on the main plot). For each

epoch, only animals with sufficient numbers of cell pairs are shown. *D*, Mean reactivation

probability vs. decrease in place field (PF) size from **Figure 4C** shown for individual animals.

References

1.   Herzog, L.E., Pascual, L.M., Scott, S.J., Mathieson, E.R., Katz, D.B., and Jadhav, S.P.

(2019). Interaction of taste and place coding in the hippocampus. J Neurosci *39*, 3057-

3069.

2.   Muller, R. (1996). A quarter of a century of place cells. Neuron *17*, 813-822.

3.   Martin, S.J., and Clark, R.E. (2007). The rodent hippocampus and spatial memory: from

synapses to systems. Cell Mol Life Sci *64*, 401-431.

4.   Colgin, L.L., Moser, E.I., and Moser, M.B. (2008). Understanding memory through

hippocampal remapping. Trends Neurosci *31*, 469-477.

5.   Eichenbaum, H. (2017). The role of the hippocampus in navigation is memory. J

Neurophysiol *117*, 1785-1796.

6.   Schiller, D., Eichenbaum, H., Buffalo, E.A., Davachi, L., Foster, D.J., Leutgeb, S., and

Ranganath, C. (2015). Memory and space: towards an understanding of the cognitive

map. J Neurosci *35*, 13904-13911.

7.   Ekstrom, A.D., and Ranganath, C. (2018). Space, time, and episodic memory: the

hippocampus is all over the cognitive map. Hippocampus *28*, 680-687.

8.   Moita, M.A., Rosis, S., Zhou, Y., LeDoux, J.E., and Blair, H.T. (2003). Hippocampal

place cells acquire location-specific responses to the conditioned stimulus during

auditory fear conditioning. Neuron *37*, 485-497.

9.   Shan, K.Q., Lubenov, E.V., Papadopoulou, M., and Siapas, A.G. (2016). Spatial tuning

and brain state account for dorsal hippocampal CA1 activity in a non-spatial learning

task. eLife *5*.

10.   Li, Y., Xu, J., Liu, Y., Zhu, J., Liu, N., Zeng, W., Huang, N., Rasch, M.J., Jiang, H., Gu,

X., et al. (2017). A distinct entorhinal cortex to hippocampal CA1 direct circuit for

olfactory associative learning. Nat Neurosci *20*, 559-570.

11.   Van Cauter, T., Camon, J., Alvernhe, A., Elduayen, C., Sargolini, F., and Save, E.

(2013). Distinct roles of medial and lateral entorhinal cortex in spatial cognition. Cereb

Cortex *23*, 451-459.

12.   Wang, C., Chen, X., Lee, H., Deshmukh, S.S., Yoganarasimha, D., Savelli, F., and

Knierim, J.J. (2018). Egocentric coding of external items in the lateral entorhinal cortex.

Science *362*, 945-949.

13.   Hargreaves, E.L., Rao, G., Lee, I., and Knierim, J.J. (2005). Major dissociation between

medial and lateral entorhinal input to dorsal hippocampus. Science *308*, 1792-1794.

14.   Dragoi, G., and Tonegawa, S. (2013). Selection of preconfigured cell assembles for

representation of novel spatial experiences. Philos Trans R Soc Lond B Biol Sci *369*,

20120522.

15.   Villette, V., Malvache, A., Tressard, T., Dupuy, N., and Cossart, R. (2015). Internally

recurring hippocampal sequences as a population template of spatiotemporal

information. Neuron *88*, 357-366.

16.   Liu, K., Sibille, J., and Dragoi, G. (2018). Generative predictive codes by multiplexed

hippocampal neuronal triplets. Neuron *99*, 1329-1341.e6.

17.   Olypher, A.V., Lansky, P., Muller, R.U., and Fenton, A.A. (2003). Quantifying location-

specific information in the discharge of rat hippocampal place cells. J Neurosci Methods

127: 123-135.

18.   Hollup, S.A., Molden, S., Donnett, J.G., Moser, M.B., and Moser, E.I. (2001).

Accumulation of hippocampal place fields at the goal location in an annular watermaze

task. J Neurosci *21*, 1635-1644.

19.   Singer, A.C., and Frank, L.M. (2009). Rewarded outcomes enhance reactivation of

experience in the hippocampus. Neuron *64*, 910-921.

20.   Dupret, D., O'Neill, J., Pleydell-Bouverie, B., and Csicsvari, J. (2010). The reorganization

and reactivation of hippocampal maps predict spatial memory performance. Nat

Neurosci *13*, 995-1002.

21.   Girardeau, G., and Zugaro, M. (2011). Hippocampal ripples and memory consolidation.

Curr Opin Neurobiol *21*, 452-459.

22.   Jadhav, S.P., Kemere, C., German, P.W., and Frank, L.M. (2012). Awake hippocampal

sharp-wave ripples support spatial memory. Science *336*, 1454-1458.

23.   Girardeau, G., Benchenane, K., Wiener, S.I., Buzsaki, G., and Zugaro, M.B. (2009).

Selective suppression of hippocampal ripples impairs spatial memory. Nat Neurosci *12*,

1222-1223.

24.   Lee, A.K., and Wilson, M.A. (2002). Memory of sequential experience in the

hippocampus during slow wave sleep. Neuron *36*, 1183-1194.

25.   Carr, M.F., Jadhav, S.P., and Frank, L.M. (2011). Hippocampal replay in the awake

state: a potential substrate for memory consolidation and retrieval. Nat Neurosci *14*, 147-

153.

26.   Eschenko, O., Ramadan, W., Molle, M., Born, J., and Sara, S.J. (2008). Sustained

increase in hippocampal sharp-wave ripple activity during slow-wave sleep after

learning. Learn Mem *15*, 222-228.

27.   Ji, D., and Wilson, M.A. (2007). Coordinated memory replay in the visual cortex and

hippocampus during sleep. Nat Neurosci *10*, 100-107.

28.   Wilson, M.A., and McNaughton, B.L. (1994). Reactivation of hippocampal ensemble

memories during sleep. Science *265*, 676-679.

29.   Foster, D.J., and Wilson, M.A. (2006). Reverse replay of behavioural sequences in

hippocampal place cells during the awake state. Nature *440*, 680-683.

30.   Cheng, S., and Frank, L.M. (2008). New experiences enhance coordinated neural

activity in the hippocampus. Neuron *57*, 303-313.

31.   Frank, L.M., Stanley, G.B., and Brown, E.N. (2004). Hippocampal plasticity across

606   multiple days of exposure to novel environments. J Neurosci *24*, 7681-7689.

607   32.   Okada, S., Igata, H., Sasaki, T., and Ikegaya, Y. (2017). Spatial representation of

608   hippocampal place cells in a T-Maze with an aversive stimulation. Front Neur Circ *11*,

609   101.

610   33.   Mamad, O., Agayby, B., Stumpp, L., Reilly, R.B., and Tsanov, M. (2019). Extrafield

611   activity shifts the place field center of mass to encode aversive experience. eNeuro *6*.

612   34.   Breese, C.R., Hampson, R.E., and Deadwyler, S.A. (1989). Hippocampal place cells:

613   stereotypy and plasticity. J Neurosci *9*, 1097-1111.

614   35.   Bostock, E., Muller, R.U., and Kubie, J.L. (1991). Experience-dependent modifications of

615   hippocampal place cell firing. Hippocampus *1*, 193-205.

616   36.   Anderson, M.I., and Jeffery, K.J. (2003). Heterogeneous modulation of place cell firing

617   by changes in context. J Neurosci *23*, 8827-8835.

618   37.   Gauthier, J.L., and Tank, D.W. (2018). A dedicated population for reward coding in the

619   hippocampus. Neuron *99*, 179-193.e177.

620   38.   Grill, H.J., and Norgren, R. (1978). The taste reactivity test. I. Mimetic responses to

621   gustatory stimuli in neurologically normal rats. Brain Res *143*, 263-279.

622   39.   Travers, J.B., and Norgren, R. (1986). Electromyographic analysis of the ingestion and

623   rejection of sapid stimuli in the rat. Behav Neurosci *100*, 544-555.

624   40.   Katz, D.B., Simon, S.A., and Nicolelis, M.A. (2001). Dynamic and multimodal responses

625   of gustatory cortical neurons in awake rats. J Neurosci *21*, 4478-4489.

626   41.   Tang, W., Shin, J.D., Frank, L.M., and Jadhav, S.P. (2017). Hippocampal-prefrontal

627   reactivation during learning is stronger in awake compared with sleep states. J Neurosci

628   *37*, 11789-11805.

629   42.   Jadhav, S.P., Rothschild, G., Roumis, D.K., and Frank, L.M. (2016). Coordinated

630   excitation and inhibition of prefrontal ensembles during awake hippocampal sharp-wave

631   ripple events. Neuron *90*, 113-127.

632   43.   Fontanini, A., Grossman, S.E., Figueroa, J.A., and Katz, D.B. (2009). Distinct subtypes

633   of basolateral amygdala taste neurons reflect palatability and reward. J Neurosci *29*,

634   2486-2495.

635   44.   Li, J.X., Yoshida, T., Monk, K.J., and Katz, D.B. (2013). Lateral hypothalamus contains

636   two types of palatability-related taste responses with distinct dynamics. J Neurosci *33*,

637   9462-9473.

638   45.   Piette, C.E., Baez-Santiago, M.A., Reid, E.E., Katz, D.B., and Moran, A. (2012).

639   Inactivation of basolateral amygdala specifically eliminates palatability-related

640   information in cortical sensory responses. J Neurosci *32*, 9981-9991.

641   46.   Sadacca, B.F., Rothwax, J.T., and Katz, D.B. (2012). Sodium concentration coding gives

642   way to evaluative coding in cortex and amygdala. J Neurosci *32*, 9999-10011.

643   47.   Maier, J.X., Blankenship, M.L., Li, J.X., and Katz, D.B. (2015). A multisensory network

644   for olfactory processing. Curr Biol *25*, 2642-2650.

645   48.   Brun, V.H., Otnass, M.K., Molden, S., Steffenach, H.A., Witter, M.P., Moser, M.B., and

646   Moser, E.I. (2002). Place cells and place recognition maintained by direct entorhinal-

647   hippocampal circuitry. Science *296*, 2243-2246.

648   49.   Zhang, S., Schönfeld, F., Wiskott, L., and Manahan-Vaughan, D. (2014). Spatial

649   representations of place cells in darkness are supported by path integration and border

650   information. Front Behav Neurosci *8*, 222.

651   50.   Schoenenberger, P., O'Neill, J., and Csicsvari, J. (2016). Activity-dependent plasticity of

652   hippocampal place maps. Nat Commun *7*, 11824.

653   51.   Sheskin, D.J. (2004). Handbook of Parametric and Nonparametric Statistical Procedures

654   (New York: Chapman & Hall/CRC).

Figure 1

A

Taste responsive | Non taste responsive

Pre | Post | Pre | Post

Cell 1: 6.1 | 6 | 6.4 | 11.8

Cell 2: 5.8 | 6.2 | 10.1 | 3.3

150 cm

FR (Hz)
Max
Min
PF

B

Norm. PF size

*** * ***

Taste
Non

Pre   Post

C

% TR/total

<0.2   0.2-0.4   0.4->0.6   >0.6

Pre PF size

Figure 2

Figure 3

Figure 4