Regional differences in the abiotic environment contribute to genomic divergence within a

wild tomato species

Running title: Landscape genomics of a wild tomato

Matthew J.S. Gibson[1] & Leonie C. Moyle[1]

[1]Department of Biology, Indiana University, Bloomington, Indiana, USA

Corresponding author: Matthew J.S. Gibson (gibsomat@indiana.edu)

ABSTRACT

The wild currant tomato *Solanum pimpinellifolium* inhabits a wide range of abiotic habitats across

its native range of Ecuador and Peru. Although it has served as a key genetic resource for the

improvement of domestic cultivars, little is known about the genetic basis of traits underlying local

adaptation in this species, nor what abiotic variables are most important for driving differentiation.

Here we use redundancy analysis (RDA) and other multivariate statistical methods (structural

equation modeling (SEM) and generalized dissimilarity modeling (GDM)) to quantify the

relationship of genomic variation (44,064 single nucleotide polymorphisms) with climate and

geography, among 140 wild accessions. RDA, SEM, and GDM each identified environment as

explaining more genomic variation than geography, suggesting that local adaptation to

heterogeneous abiotic habitats may be an important source of genetic diversity in this species.

Environmental factors describing temporal variation in precipitation and evaporative demand

explained the most SNP variation among accessions, indicating that these forces may represent

key selective agents. Lastly, by studying how SNP-environment associations vary throughout the

genome, we mapped the location and investigated the functions of loci putatively contributing to

climatic adaptations. Together our findings indicate an important role for selection imposed by the

abiotic environment in driving genomic differentiation between populations.

KEYWORDS: adaptation, eigenanalysis, redundancy analysis, population structure, landscape

genetics

INTRODUCTION

Local adaptation is thought to underlie a large proportion of phenotypic variation segregating within species. Despite this, the primary environmental agents driving selection between natural populations are still unknown in most systems. Across landscapes, genetic divergence can result from selection imposed by environmental forces, but also from the effects of restricted gene flow and genetic drift when subpopulations are partially isolated (Wright, 1943). Disentangling the concurrent effects of spatial isolation and natural selection on generating genetic diversity is critical for understanding the relative importance of adaptation in driving diversification, as well as for identifying the environmental factors underlying its occurrence. This remains a major technical challenge, mainly because collinearity is common among spatial and environmental variables (Kissoudis et al., 2016; Lasky et al., 2012; Lee & Mitchell-Olds, 2011; Wiens, 1989). New statistical methods and the increasing availability of fine-scale environmental and genomic data provide one approach to estimate the independent contributions of environment and space to explaining patterns of genetic variation. These strategies can complement traditional tests for local adaptation (i.e., reciprocal transplants) and begin to characterize the nature of selection acting in the wild, as well as its genetic targets (Fournier-Level et al., 2011; Hancock et al., 2011; Lasky et al., 2012, 2015).

Solanum pimpinellifolium L. (currant tomato) is one of the closest wild relatives of domesticated tomato (*Solanum lycopersicum* L.) and has historically served as a key resource for the identification and breeding of agronomically-important traits (Pedley & Martin, 2003; Pitblado & Kerr, 1979; Tanksley, 2004; Tanksley et al., 1996). Several accessions are known to be tolerant to acute abiotic stressors such as salinity or drought (Bolarin et al., 1991; Cuartero et al., 1992; Kissoudis et al., 2016; Foolad & Yin, 1997; Razali et al., 2018), however there is

limited data on the nature of genetic and phenotypic variation segregating within this species

(Foolad, 2007), especially as it relates to potential climatic adaptations. *S. pimpinellifolium*

occupies a large geographic range from Ecuador to southern Peru **(Figure 1**); populations

experience cool mid-elevation habitats in the Andes as well as dry, hot habitats along the pacific

coast. Both the abiotic environment (climate and soil characteristics; **Figure 1**) and ecologically

relevant traits vary considerably throughout the range (Chitwood et al., 2012a, 2012b; Nakazato

et al., 2008; Nakazato et al., 2010; Rick et al., 1977; Zuriaga et al., 2009), and previous studies

have suggested that a large fraction of this phenotypic variation could be attributable to spatially

varying selection imposed by the abiotic environment (Zuriaga et al., 2009; Blanca et al., 2012,

2015). For example, using common garden experiments Nakazato et al. (2008) identified

significant intraspecific variation for drought tolerance between accessions from coastal and

highland regions. Using Mantel tests, Blanca et al. (2015) showed that major patterns of genetic

structure among *S. pimpinellifolium* accessions correspond strongly with broad patterns of

bioclimatic variation (also see Zuriaga et al., 2009 & Blanca et al., 2012). In addition to these

intraspecific patterns, between species, distribution models have further suggested that local

climatic variation (especially in temperature and water availability), rather than geographic

isolation, is the main factor explaining contemporary distributions (Nakazato et al., 2010).

Whether this pattern holds for variation within species, as well as the identity of the key

environmental/selective gradients responsible, has not yet been thoroughly assessed in a genomic

context. Nonetheless, describing the factors shaping intraspecific genetic variation is essential for

uncovering the processes that initiate phenotypic divergence, including the relative role of local

adaptation in driving diversification.

The main goals of this study were to characterize the environmental forces that might

impose selection on ecologically relevant traits across the range of *S. pimpinellifolium,* and to

determine the degree to which these environmental factors explain genetic variation independent

of demographic history. In the absence of confounding population structure, associations

between allele frequencies and environmental variables are considered evidence for local

adaptation (Endler, 1986). These associations can be used to identify both the forces responsible

for imposing selection, as well as the specific loci involved in adaptive evolutionary responses.

However, since both adaptation and spatially structured gene flow can result in correlations

between allele frequencies and the environment (Endler, 1977), demographic history can cloud

such inferences of local adaptation. Moreover, abiotic selection is often the result of a

combination of many partially colinear variables, making simple gene-environment correlations

incomplete and potentially misleading. To address these issues, multivariate statistical methods

commonly used in community ecology—in particular the multivariate regression technique

"redundancy analysis" (RDA; Legendre & Legendre, 1998) and other constrained ordination

procedures—have recently been applied to landscape genomics to study how climatic variation

affects spatial patterns of genomic variation (Lasky et al., 2012, 2016; Forester et al., 2018;

Wang et al., 2013). These methods have been used to identify both the sources of selection

imposed by the multivariate environment and their genetic targets (individual loci) (Forrester et

al., 2018; Lasky et al., 2012, 2016; Lee & Mitchell-Olds, 2011; Manel et al., 2010; Salathe &

Schmid-Hempel, 2011; Sork et al., 2010). In doing so, they have begun to address the challenges

of disentangling geographical structure from environmental variation in order to quantify the

independent contribution of environmental forces to shaping genomic variation across space.

Here we employ RDA to study the relationship between environmental and SNP data

matrices while modeling population structure as a covariate, in order to estimate the relative

influence of abiotic adaptation on genomic variation across the native range of *S.*

pimpinellifolium. Specifically, we use RDA to 1) estimate the independent quantitative

contributions of climate, space, and their colinear portion to explaining patterns of genome-wide

SNP variation; 2) characterize the specific abiotic variables with the largest influence on genetic

variation; and 3) identify genomic regions showing strong associations with major axes of

multivariate climate and investigate their potential functional roles in mediating local adaptation.

We complement these RDA analyses with other methods designed to quantify the roles of

selection and spatial isolation. Our analyses reveal that models including several key abiotic

variables explain more genomic variation among genotypes than models considering only

geographic isolation. They also allow us to identify loci that have significant associations with

environmental variation, after incorporating population structure via partial regression. These

results suggest a prominent role for local adaptation in driving genetic divergence among

populations of this agronomically important crop wild relative.

METHODS

Plant material and genotyping. Seeds from 140 wild georeferenced *S. pimpinellifolium*

accessions (**Figure 1**; **Table S1**) spanning the entire species range were obtained from the

Charles M. Rick Tomato Genetic Resource Center (TGRC; Davis, Calif.). Seeds were soaked in

a 50% bleach solution (as per the TGRC protocol), germinated, and transplanted into 1-gallon

pots at the Indiana University greenhouses. Fresh leaf material was collected from one plant per

accession at 30 days post-germination. DNA was extracted using DNeasy Plant Mini Kits

(Qiagen, Valencia, Calif, USA) and two double-digest restriction site associated sequencing

(ddRAD) libraries were prepared using *PstI* and *EcoRI* enzymes by the Indiana University

Center for Genomics and Bioinformatics. These enzymes were selected to enrich for cut sites in

coding regions based on an *in silico* digestion of the *S. lycopersicum* reference genome ver.

SL3.0 (Tomato Genome Consortium, 2012). Libraries were sequenced across two Illumina

NextSeq flowcells (150 bp, paired-end, high-output). Raw reads were filtered for quality,

trimmed of Illumina adapter sequences using *fastp* (Chen et al., 2018), and then demultiplexed

by individual using the process_radtags program in *Stacks* (ver. 2; Catchen et al., 2013). After

demultiplexing, reads were mapped to the *S. lycopersicum* reference genome version SL3.0

using BWA (Li & Durbin, 2009). Variants were called using the *Stacks* ref_map pipeline. Only

bi-allelic SNPs typed in at least 30 percent of individuals were retained. Further filtering was

applied prior to running specific analyses (see below).

Environmental data. For each accession, data were compiled for 54 abiotic environmental

parameters available from public databases. 19 bioclimatic variables as well as monthly solar

radiation, vapor pressure, and wind speed data were obtained directly from the WorldClim2

database (Fick & Hijmans, 2017). Additional transformations of the WorldClim data were

obtained from the Climate South America (ClimateSA v1.0; Hamann et al., 2013) database.

Global aridity measures were obtained from the Consultative Group on International Agricultural

Research (CGIAR; Zomer et al., 2008). Lastly, soil pH, type, density (fine earth fraction), and

water capacity at a depth of 5cm were obtained from the SoilGrids initiative

(http://soilgrids.org). All variables were centered, scaled, and tested for normality using Shapiro-

Wilks tests. If appropriate, variables were log-transformed. One-hot encoding was used for all

factor variables prior to running RDA. The full list of all 54 climate and soil variables can be

found in **Table S2**.

Spatial data. We represented spatial (non-genetic) structure in RDA using distance-based Moran

Eigenvector Maps (dbMEMs; formally referred to as principle components of neighborhood

matrices or PCNMs). Briefly, dbMEMs—much like principle components in a PCA—represent

structure in a dataset with orthogonal axes. dbMEMs describe complex patterns of spatial

structure in rectangular form suitable for use as a co-factor in downstream statistical analyses

such as RDA (Borcard et al., 2018). Importantly, they are able to model both broad- and fine-

scale patterns (Borcard et al., 2018; see **Figure S1**), both of which may affect the partitioning of

genetic variation across large and highly heterogenous landscapes. dbMEMs were calculated

using the quickMEM function included in Borcard et al. (2018). This function uses forward

selection and permutation tests to identify the appropriate number of axes needed to explain

patterns of spatial genetic variation, without overfitting.

Population genetic structure. Both model-based (*fastStructure*; Raj et al., 2014) and non-model

based (PCA) methods were used to quantify genetic structure in the data. Markers in high

linkage disequilibrium ($r^2 > 0.9$) were pruned from the dataset prior to running PCA and

fastStructure, using *bcftools* (Li et al., 2011). PCA was implemented in the R (R Core Team,

2019) package *adegenet* (Jombart & Ahmed, 2011). Prior to running PCA, missing calls were

imputed as the mean numeric value for each marker, to speed up computation. Model complexity

in *fastStructure* was determined by minimizing the marginal likelihood of the data over values of

K from one to ten. Clusters in the data were also identified using the K-means algorithm for all

PCs by minimizing the Bayesian information criterion (BIC) across values of K from one to 100.

Partitioning total SNP variation. To evaluate the relative contribution of demographic history

and the abiotic environment to explaining patterns of spatial genetic variation, the amount of

genome-wide SNP diversity attributable to environment and geography was estimated using

three methods: variance partitioning by redundancy analysis (RDA), generalized dissimilarity

modeling (GDM), and structural equation modeling (SEM). RDA is a form of constrained

ordination first applied in community ecology that allows one to assess the explanatory power of

multivariate predictors (in this case, environmental and geographic variables) for multivariate

responses (genotypes; van den Wollenberg, 1977; Legendre & Legendre, 1998). RDA identifies

linear combinations of explanatory variables (canonical axes) that best explain linear

combinations of response variables. As with other forms of eigenanalysis, these canonical axes

are orthogonal to each other. Here, these axes represent correlated SNPs that are also correlated

with environment. Using multiple partially constrained models, it is possible to partition total

genetic variation into that explained independently by climate, space, and their colinear portion

(Borcard et al., 2018). All RDA analyses were performed using the R package *vegan* (Oksanen,

2018) and significance was assessed using 1,000 permutations of the genotype matrix. Since

RDA requires no missing data, only markers containing no missing calls were included (6,830

SNPs, compared to 44,064 in the total dataset). Distances between collection sites were

represented using dbMEMs (described above). Because collinearity among environmental

predictors can lead to inflation of variance components in RDA (Borcard et al., 2018), we

performed forward variable selection using the forward.sel function in the R package *adespatial*

to identify predictive and non-redundant environmental variables for partitioning.

In addition to RDA, structural equation modeling (SEM) was performed using the R

package *lavaan* (Rosseel, 2012). SEM is a statistical framework similar to path analysis that has

been applied to estimating the relative contributions of isolation-by-environment (IBE) and

isolation-by-distance (IBD) to explaining spatial genetic variation (Wang et al., 2013). While

powerful, this method is not able to estimate the independent contribution of the colinear portion

of environment and space on explaining SNP variation, unlike RDA. For SEM, geographic

distance was represented using Euclidean distance between accessions. All 54 environmental

variables were used to infer the latent environmental distance term (SEM accommodates

correlated inputs; Wang et al., 2013). SEM was performed using the same set of markers as used

for variance partitioning in RDA.

Lastly, generalized dissimilarity modeling (GDM; Ferrier et al., 2007) was applied to the

SNP data. GDM is similar in principle to RDA and SEM but relies on different statistical

methods. GDM uses nonlinear matrix regression to accommodate nonlinear statistical

relationships that are often observed in ecological datasets; in contrast, RDA and SEM—as

implemented here—do not model nonlinear relationships. GDM analyses were performed in R

using the *gdm* package (Manion et al., 2018).

Importance of specific climate variables. Forward variable selection with RDA was used to infer

which specific environmental variables may be most important for generating selective

differences across the range of *S. pimpinellifolium*. We performed this procedure with the

forward.sel function in *adespatial*, using 10,000 permutations to assess significance at each

iteration and with RDA models both including and not including spatial structure as a covariate.

The relative contribution of each individual variable to the constrained model was calculated as a

weighted sum of correlation coefficients across the first four RDA axes. The contribution of each

variable to the full RDA model was then calculated as the ratio of this value and total genetic

variance (Lasky et al., 2012).

Identifying outlier loci. RDA was also used to identify outlier loci showing especially strong

relationships with multivariate environmental axes. The ability of RDA to identify loci

responding to combinations of environmental predictors, instead of testing and treating each

variable independently, is an advantage of RDA outlier analysis over classical mixed-model

association methods. Moreover, because it simultaneously evaluates combinations of SNPs,

RDA may also be better able to identify more subtle patterns environmental association

involving multiple minimally diverged loci (Forester et al., 2018)—suggestive of polygenic

adaptation—that can be obscured by traditional approaches. Here, outliers were defined as the

SNPs having loadings along the first RDA axis +/- 4 SDs from the mean for that axis (Forester et

al., 2018; Lasky et al., 2012). To reduce the potential for spurious associations due to shared

ancestry, a partial RDA conditioned on dbMEMs was implemented. Instead of using the reduced

set of markers with no missing calls for this analysis, we imputed SNPs at the markers with

missing data in the full dataset based on the mean allelic value at each site, and used this full

dataset for outlier identification. Top candidate SNPs were investigated for their proximity to

known gene models, using the ITAG 3.2 tomato genome annotation (Tomato Genome

Consortium, 2012).

RESULTS

Genotype coverage and distribution

Our ddRAD pipeline identified 359,796 SNPs segregating within *S. pimpinellifolium*. After

stringent filtering for depth and missing data, this set of SNPs was reduced to 44,064. One

variant occurred roughly every 18 kb. SNPs were enriched (75.6%) for sites in or nearby (within

5 kb) annotated gene models (9.4% within exons), suggesting that our enrichment based on *in*

silico digestion was successful (determined by SnpEff; Cingolani et al., 2012). Of annotated

SNPs in coding regions, 60.9% resulted in changes to amino acid composition (i.e., were non-

synonymous).

Population structure follows a latitudinal gradient and defines regional genetic clusters

The first two axes of the multi-locus PCA based on 17,358 SNPs (LD pruned) explained 15.8%

and 6.24% of genetic variation, respectively. Minimizing BIC in K-means clustering identified

five clusters in the data (**Figure 2**), which followed a latitudinal gradient and roughly

corresponded to variation along the first principal component axis. The model-based analysis

conducted in *fastStructure* identified four groups in the data; these groups largely corresponded

to the clusters identified using K-means, except that clusters four and five were not separated and

instead were identified as a single group (**Figure 2**). These patterns agree with other recent

genomic analyses of population structure in this species (Lin et al., 2019). Varying degrees of

admixture were inferred. The most heterogeneous individuals were found at the center of the

species range (cluster 1, light blue in **Figure 2**), an observation which agrees with previous

marker analyses in this species (Lin et al., 2019; Rick et al., 1977; Zuriaga et al., 2009) and

which matches general expectations for patterns of genetic diversity across species ranges

(Vucetich & Waite, 2003). Similarly, heterozygosity increased towards the equator (**Figure S2**),

also in general agreement with previous findings (Rick et al., 1977; Zuriaga et al., 2009; Lin et

al., 2019).

RDA identifies climate and spatial variables that predict SNP variation

Spatial variables. Forward selection of the dbMEM variables identified 10 axes as sufficient to

explain geographic structure among accessions. Four of these axes described broad patterns of

latitudinal structure (dbMEM1, 2, 3, and 4) and the remaining six described fine-scale patterns

(dbMEM6, 7, 8, 9, 11 and 17; **Figure S1**). The axes describing fine-scale spatial structure

cumulatively explained the most SNP variation among accessions (7.01%; **Table 1**). Broad-scale

spatial variables explained a total of 5.42% of SNP variation (**Table 1**). dbMEM2 explained the

most variation of any single variable (1.74%).

Climate variables. When analyzed without accounting for spatial structure using partially

constrained ordination, we identified 14 abiotic variables as significantly predictive of genetic

variation among accessions (**Table 2**). The most predictive variable was degree days above 18C,

however several variables contributed nearly equally to the RDA model. These include annual

maximum solar radiation, average annual wind speed, isothermality (a measure of temperature

variability), annual maximum wind speed, and variation in vapor pressure. In contrast, forward

selection constrained on dbMEM spatial axes identified 6 environmental variables as predictive

of genetic variation (**Table 2**); these variables represent gradients predictive of genetic variation

after accounting for the proportion explained by geographic distance. Importantly, these will not

include environmental gradients that are entirely colinear with space—including some that may

contribute to clinal adaptation—and thus represent a conservative subset of the variables

potentially contributing to geographical variation in selection. The contribution of these 6

gradients in RDA space is shown in **Figure 3**. The strongest predictor was the coefficient of

monthly variation in vapor pressure, followed by precipitation seasonality, soil texture, annual

maximum solar radiation, maximum potential evapotranspiration, and minimum potential

evapotranspiration (**Table 2**).

The correlations of these six variables with the first two RDA axes (loadings in **Figure 3**)

suggests that the contribution of each variable to SNP variation differs geographically across the

species range. To test for the presence of regional climatic differences that may contribute to

divergent selection, we studied how each environmental predictor varied by genetic cluster

(**Figure S3**). Significant differences among genetic clusters were identified for all 6 predictors

based on ANOVA tests ($P < 1 \times 10^{-8}$ for all tests), suggesting that each regional cluster occupies a

more or less unique abiotic habitat. This finding is recapitulated when breaking down all 58

environmental variables into orthogonal components (**Figure S4**); the first two components of

climatic variation separate accessions by region.

Regional climatic differences explain more SNP variation than geography

Variance partitioning by RDA. Using RDA, the abiotic environment (14 variables identified

through forward selection) and spatial structure (10 dbMEMs identified through forward

selection) explained 22% of SNP variation among accessions. The majority of this (17%) was

attributable to the colinear portion of environment + space (**Figure 4**). This fraction could

represent either the contribution of clinal environmental factors or latent spatial population

structure that is not captured by the dbMEMs. Considering the independent effects of

environment and space, the environment alone explained more SNP variation (3%) than space

alone (2%; **Figure 4**). The observed proportions of variation explained by climate and space

were greater than the proportions explained in all of 1000 permuted data sets (P < 0.001).

Structural equation modeling (SEM). Regression components of environmental distance and

geographic distance on genetic distance identified using SEM were 0.788 (SE = 0.044) and 0.174

(SE = 0.012), respectively (**Table S4**). This corresponds to 7% (6.4% - 7.7%) and 3% (2.6% -

3.5%) of total genetic variance being attributable to environmental distance and geographic

distance between accessions, respectively, which is consistent with our findings from RDA

(**Figure 4**). Only the environmental regression component was higher than those obtained from

all of 1000 permuted data sets (P < 0.001). As with RDA, this suggests that environmental

differences between accessions are shaping the distribution of genetic variation to a larger degree

than spatial isolation.

Generalized dissimilarity modeling (GDM). The percent deviance in genetic variation among

accessions explained by the full (environment + space) GDM model was 30.2% (**Figure 4**; see

Table S5 for full model fit). 29.2% and 23.9% of deviance was explained by models considering

only environmental and geographic effects on genetic variation between accessions, respectively

(**Figure 4**). All of these models were significant (P < 0.001) based on permutation tests. These

results agree with RDA and SEM; a larger fraction of SNP variation is attributable to

environmental variation than to spatial isolation.

RDA identifies outlier loci associated with multivariate climate

We observed significant variation among SNPs in the degree to which they were associated with

the abiotic environment, as measured by their loadings along the first RDA axis (**Figure 5**). We

identified candidate loci for local adaptation by examining SNPs displaying loadings along the

first RDA axis +/- 4 standard deviations from the mean (Forester et al., 2018; Lasky et al., 2012).

This identified 252 SNPs showing particularly strong associations with multivariate climate. In

the model containing spatial structure as a covariate, the strongest signal of association spanned

23.5 Mbp on chromosome four (**Figure 5**). This region contained 333 annotated genes. The top

associated SNP in this region was 30 bp upstream of a 60S ribosomal subunit gene (**Table 3**).

Other significant associations in this region include intergenic substitutions proximal to DNA

glycosylase, exocyst complex component 7, and cellulose synthase; a non-synonymous

substitution in MYB transcription factor 73; and intronic substitutions in sterol

glusoyltransferase 4 (SGT4), nuclease S1, BY-2 kinesin protein 5, RNA helicase DEAH-Box 13,

and translation initiation factor 3, among others (**Table 3**; **Table S8**).

DISCUSSION

Local adaptation to the abiotic environment is often thought to play a major role in diversification both within and between species, however its contribution relative to other evolutionary forces is rarely quantified. In this study, we investigated the relationship of 44,064 SNPs with fine-scale climatic and spatial data among 140 accessions of the ecologically diverse crop wild relative *S. pimpinellifolium*. We found that the abiotic environment—as described by climate and soil characteristics—explained more SNP variation than spatial structure (**Figure 4**), demonstrating that our evaluated environmental factors are more influential in shaping the geographic distributions of SNPs in this species, compared to spatial isolation. Both GDM and SEM—two methods which have also been applied to quantifying the contributions of space and environment to patterns of genetic variation—agree with our RDA results. From this we conclude that ecological differences among populations are closely associated with the maintenance of contemporary genetic variation across space and that local adaptation is likely a major driver of these patterns.

Our RDA analysis identified that a large (17%) portion of SNP variation among accessions was explainable by the colinear fraction of environment and space (**Figure 4**). This fraction could represent the contribution of clinal environmental forces that are highly correlated with space (e.g., latitudinal gradients in temperature). Clinal gradients are considered to be important drivers of adaptive evolution in many species (Adrion et al., 2015; Kooyers et al., 2015), and across the range of *S. pimpinellifolium*, many environmental variables show strong geographic trends and are correlated with both genetic (i.e., PCs) and spatial (i.e., dbMEMs) structure (**Table S8**). Previous studies in *S. pimpinellifolium* have demonstrated strong correlations between environmental and genetic distances among accessions of this species,

largely associated with latitudinal gradients (Blanca et al., 2015; Zuriaga et al., 2009).

Accordingly, when we did not correct for spatial structure with constrained ordination, a highly

clinal variable—degree days above 18C—explained the most total SNP variation (**Table 2**).

Although geographic clines have historically been considered important for understanding the

genetics of adaptation (Adrion et al., 2015; Endler, 1977, 1986), disentangling geographically

structured variation from adaptive variation remains a major challenge—as evidenced here—and

will likely require novel statistical methods to address. In *S. pimpinellifolium*, further population

sampling across clinal gradients coupled with *in situ* common garden experiments could be one

way of disentangling the independent effects of space and climate in these cases.

Despite the substantial collinearity we observed between space and environment, our

analysis discerned several axes of spatial variation that uniquely explained SNP variation. The

second spatial dbMEM variable explained the largest portion of SNP variation among accessions

(**Table 1**). This variable describes broad patterns of spatial structure and separates accessions

from the center of the range (northern Peru) from those at the northern and southern extremes

(**Figure S1**). This finding is consistent with our genetic analysis of population structure—as well

as previous spatial genetic data (Rick et al., 1977; Zuriaga et al., 2009; Lin et al., 2019)—which

demonstrate a strong latitudinal pattern of genetic relatedness. Interestingly, while the second

dbMEM axis explained the most variance of any single spatial variable, cumulatively more

variance was explained by fine-scale dbMEM axes (dbMEM5, 6, 7, 8, 9, & 17; cumulative R^2 =

0.0701). Such a pattern may be driven by the geographic landscape of the Andes; large and

patchy altitudinal clines (sea level to > 6500 m) may limit local dispersal between adjacent

populations, amplifying genetic differentiation over small spatial scales.

395   Apart from spatial variation associated with SNPs, our analysis also identified specific

396 environmental factors that uniquely explained additional SNP variation. Temporal fluctuations in

397 precipitation (seasonality) and vapor pressure (the coefficient of monthly variation) explained the

398 most genome-wide SNP variation when we simultaneously considered the effects of spatial

399 structure in our RDA analysis (**Table 2**). This reinforces some previous empirical findings—

400 especially the role of precipitation seasonality as a key factor—but also points to additional

401 environmental forces as potential abiotic selective agents in this species. Precipitation was

402 previously implicated in driving ecological adaptation in both this and other tomato species

403 (Nakazato et al., 2008, 2010). Specifically, among accessions of *S. pimpinellifolium*, variation in

404 days to wilting (a measure of drought tolerance) is strongly associated with differences in annual

405 precipitation between coastal and highland regions (Nakazato et al., 2008), consistent with

406 selection being imposed by this environmental axis. Interestingly, our findings at the

407 intraspecific scale are also similar to previous inferences of important ecological drivers among

408 wild tomato species. Distribution models have identified precipitation and temperature as the

409 major determinants of differences in contemporary species ranges across the wild tomato clade

410 (Nakazato et al., 2010). Moreover, these analyses indicate that precipitation seasonality in

411 particular is important for determining the allopatric distributions of *S. pimpinellifolium* and its

412 sister taxon *S. lycopersicum* var. *cerasiforme*.

413   In contrast to precipitation, the importance of vapor pressure as a selective agent within

414 or among wild tomato species has not been evaluated, although vapor pressure's role in

415 modulating plant-water relations has been well studied in domesticated tomato (Guichard et al.,

416 2005; Zhang et al., 2017). Deficits in vapor pressure between the mesophyll and external leaf

417 surface promotes transpiration (Zhang et al., 2017), and if not met with additional water uptake

from the soil, large differentials in vapor pressure will lead to reduced growth and desiccation.

Our findings from RDA suggest that regional differences in vapor pressure are a strong selective

force in *S. pimpinellifolium* (**Table 2**). The importance of *variation* in vapor pressure—and not a

measure of annual or monthly mean values—may point to a role for specific physiological

mechanisms which maximize fitness during acute fluctuations in evaporative demand (e.g., rapid

stomatal closure responses, induction of root growth, or other strategies for increasing water use

efficiency under temporary stress). This hypothesis is testable in the future with direct

manipulative experiments across accessions that are predicted to differ at specific SNPs

associated with this axis of the environment.

While variation in vapor pressure and precipitation were the most predictive of genomic

variation across the range as a whole (**Table 2**), more specifically, regional environmental

differences and our RDA analysis suggest that different factors are most influential in different

regions of the species range (**Figure 3**; Figure **S3**). Genetic clusters we identify within *S.*

pimpinellifolium are, in general, significantly diverged from each other for their environments

(**Figure S3**; **Figure S4**). For example, coastal accessions (genetic clusters 1, 4, and partially 3)

are exposed to higher levels of solar radiation compared to the rest of the range (**Figure S3**,

panel 3); northern Peruvian accessions (cluster 2) experience highly seasonal precipitation

(**Figure S3**; panel 2); and Ecuadorian accessions (cluster 5) experience high soil type diversity

(i.e., all four soil types that found throughout the species range are also present in this region;

Figure S3, panel 6). Together with our RDA results—which identified these and other variables

as significantly predictive of SNP variation independent of spatial population structure—these

patterns of environmental variation are consistent with regional differences in the primary agents

of ecological selection. Unlike precipitation (see above), we are not aware of any studies that

have examined genetic variation for tolerance to these environmental factors (i.e., solar radiation,

soil texture, and potential evapotranspiration) that would further support them as local selective

agents. Further common garden manipulation experiments would be helpful to understand the

exact role these forces have in shaping local climatic adaptation in *S. pimpinellifolium*.

Nonetheless, these observations broadly mirror comparative findings from among wild

tomatoes—that detect clear niche differentiation between species (Nakazato et al., 2010)—by

uncovering similar niche variation intraspecifically at the regional level.

In addition to these findings, and just as with other genetic (Lasky et al., 2012, 2016) and

community ecological studies using similar methods (Borcard et al., 2018), a large portion of

variance remained unexplained. This is likely due to the influence of several factors that cannot

be fully addressed in this type of experiment. First, although we included a large set of

environmental variables in our analyses, many other unmeasured ecological forces could also be

acting; most importantly, this includes biotic interactions that are likely only partially reflected in

the abiotic factors considered here. Second, axes of population structure may not be fully

explained by spatial dbMEM variables, potentially due to unmodeled geographic heterogeneity

present in the Andes. Third, other evolutionary forces that maintain local allelic diversity (e.g.,

balancing selection) may weaken signals of SNP-environment associations. Lastly, as

implemented here, both RDA and SEM only model linear associations between climate/space

and SNPs (Borcard et al., 2018), and thus will not capture nonlinear statistical relationships if

they exist. Interestingly, our GDM analysis—which does model nonlinear associations—

explained significantly more total SNP variation than the other two methods (**Figure 4**),

suggesting the presence of nonlinear SNP-environment relationships across the range of *S.*

pimpinellifolium. Explicitly modeling and estimating the contribution of nonlinear terms to

patterns of SNP-environment associations may be crucial to understanding local adaptation in the

context of multivariate environments, but such methods are currently limited (Rellstab et al.,

2015).

Our SNP outlier analysis allowed us to map the location of genomic regions showing

strong associations with major axes of the environment and identify candidate loci potentially

involved in local adaptation (**Figure 5**). In contrast to traditional scans that treat each

environmental variable independently (e.g., Fournier-Level et al., 2011; Hancock et al., 2011),

our analysis considered how loci are associated with linear combinations of environmental

gradients. This framework may be more appropriate for identifying ecologically relevant loci;

rarely can a single environmental factor be considered the only contributor to selection. We

identified 252 signals of putatively adaptive loci in *S. pimpinellifolium* using genomic scans

based on RDA loadings. Of foremost interest are two regions containing known regulators of

abiotic stress tolerance: *Solyc04g051150*—sterol glucosyl transferase 4 (SGT4)—and

Solyc04g050080—MYB transcription factor 73 (**Table 3**). SGT genes have been shown to have

functional roles in abiotic stress tolerance in several species (Griebel & Zeier, 2010; Hugly et al.,

1990; Kumar et al., 2015; Senthil-Kumar et al., 2013). Expression of SGT4 and its paralog SGT2

is rapidly induced upon exposure to cold and salt stress in domesticated tomato (Ramirez-Estrada

et al., 2017). In *Arabidopsis thaliana* SGT knock-out mutants, plants show reduced tolerance to

several abiotic stressors, including cold and salinity (Mishra et al., 2015). The other major

candidate identified here, MYB 73, acts as a salt stress-specific transcription factor and

modulates whole-plant responses to salinity in *A. thaliana* (Kim et al., 2013). Expression profiles

of domesticated tomato under hormone-induced stress suggest a general role for MYB

transcription factors in responding to abiotic cues (Li et al., 2016), although the specific function

of MYB 73 has not been evaluated in *Solanum*. While genotypic variance in salt-stress tolerance

has been documented in *S. pimpinellifolium* (Rao et al., 2013)—suggesting that functional

variation in salt-response loci may be prevalent—our analyses were not able to evaluate the

contribution or association off this factor to genome-wide SNP variation, since fine-scale data on

soil salinity in South America is not yet available. Nonetheless, these scans provide new

information on the location of loci relevant for adaptive responses to abiotic stress, generating

prime candidates for future functional investigation, as well as for breeding into domesticated

lines.

In conclusion, all three multivariate methods employed here (RDA, SEM, and GDM)

identified that climatic variation among the 140 wild *S. pimpinellifolium* accessions explained

more genomic variation than geographic distance between accessions. Environmental variables

describing temporal variation in precipitation and evaporative demand explained the most SNP

variation overall, however we also observe significant geographical differences in other key

climatic factors suggestive of intraspecific niche differentiation between geographic regions. Our

outlier analysis has uncovered potential genomic targets of ecological selection, although further

studies will be required to understand their exact genetic and phenotypic contributions to this

pattern. To do so, for example, SNP-environment associations can be incorporated into breeding

value prediction frameworks to test the functional roles of genomic regions showing strong

environmental associations (Lasky et al., 2015; Yoder et al., 2012). Together, our results

complement comparative findings from the wild tomato clade (Nakazato et al., 2010), indicating

that spatially varying selection imposed by divergent abiotic habitats may be an important source

of genetic variability in this species.

ACKNOWLEDGEMENTS

The authors thank the UC Davis Tomato Genetic Resource Center for providing access to the *S.*

pimpinellifolium germplasm used in this study, the Indiana University Center for Genomics and

Bioinformatics for library preparation and sequencing, and the Indiana University greenhouse

staff. This research was supported by National Science Foundation award DEB-1136707 to

LCM.

REFERENCES

Adrion, J.R., Hahn, M.W., and Cooper, B.S. (2015). Revisiting classic clines in Drosophila melanogaster in the age of genomics. *Trends in Genetics* 31(8): 434-444.

Blanca, J., Cañizares, J., Cordero, L., Pascual, L., Diez, M.J., & Nuez, F. (2012). Variation Revealed by SNP Genotyping and Morphology Provides Insight into the Origin of the Tomato. *PLoS One* 7(10): e48198

Blanca, J., Montero-Pau, J., Sauvage, C., Bauchet, G., Illa, E., Diez, M.J., Francis, D., Causse, M., van der Knaap, E., Cañizares, J. (2015). Genomic variation in tomato, from wild ancestors to contemporary breeding accessions. *BMC Genomics* 16: 257

Bolarin, M., Fernandez, F., Cruz, V., and Cuartero, J. (1991). Salinity tolerance in four wild tomato species using vegetative yield-salinity response curves. *Journal of the American Society for Horticultural Science*, 116: 286–290.

Borcard, D., Gillet, F., and Legendre, P. (2018). *Numerical Ecology with R*. Springer International, 2nd edition.

Catchen, J., Hohenlohe, P., Bassham, S., Amores, A., and Cresko, W. (2013). Stacks: an analysis tool set for population genomics. *Molecular Ecology*, 22: 3124–3140.

Chen, S., Zhou, Y., and Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*, 34(17): i884–i890.

Chitwood, D., Headland, L., Filiault, D., Kumar, R., Jimenez-Gomeez, J., Schrager, A., Park, D., Peng, J., Sinha, N., and Maloof, J. (2012a). Native environment modulates leaf size and response to simulated foliar shade across wild tomato species. *PLoS One*, 7(1): e29570.

Chitwood, D., Headland, L., Kumar, R., Peng, J., Maloof, J., and Sinha, N. (2012b). The developmental trajectory of leaflet morphology in wild tomato species. *Plant Physiology*, 158(3): 1230–1240.

Cingolani, P., Platts, A., Wang, L., Coon, M., Nguyem, T., Wang, L., LandmS.J, Lu, X., and Ruden, D. (2012). A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118. *Fly*, 6(2): 1–13.

Cuartero, J., Yeo, A., and Flowers, T. (1992). Selection of donors for salt-tolerance in tomato using physiological traits. *New Phytologist*, 121: 63–69.

Des Marais, D. and Juenger, T. (2010). Pleiotropy, plasticity, and the evolution of plant abiotic stress tolerance. *Annals of the New York Academy of Sciences*, 1206: 56–79.

Des Marais, D., McKay, J., Richards, J., Sen, S., Wayne, T., and Juenger, T. (2012). Physiological genomics of response to soil drying in diverse Arabidopsis accessions. The *Plant cell*, 24:893–914.

Endler, J. (1977). Geographic Variation, Speciation, and Clines. Princeton University Press, Princeton, NJ.

Endler, J. (1986). Natural Selection in the Wild. Princeton University Press, Princeton, NJ.

Ferrier, S., Manion, G., Elith, J., and Richardson, K. (2007). Using generalized dissimilarity modelling to analyse and predict patterns of beta diversity in regional biodiversity assessment. *Diversity and Distributions*, 13(3): 252–264.

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, 37(12), 4302-4315.

Foolad, M. and Yin, G. (1997). Genetic potential for salt tolerance during germination in Lycopersicon species. *HortScience*, 32(2): 296–300.

Foolad, M. R. (2007). Current status of breeding tomatoes for salt and drought tolerance. In *Advances in molecular breeding toward drought and salt tolerant crops* (pp. 669-700). Springer, Dordrecht.

Forester, B., Lasky, J., Wagner, H., and Urban, D. (2018). Comparing methods for detecting multilocus adaptation with multivariate genotype-environment associations. *Molecular Ecology*, 27: 2215–2233.

Fournier-Level, A., Korte, A., Cooper, M., Nordborg, M., Schmitt, J., and Wilczek, A. (2011). A map of local adaptation in Arabidopsis thaliana. *Science*, 334(6052): 86–89.

Griebel, T. & Zeier, J. (2010). A role for beta-sitosterol to stigmasterol conversion in plant-pathogen interactions. *Plant Journal* 63(2): 254-268.

Guichard, S., Gary, C., Leonardi, C., and Bertin, N. (2005). Analysis of Growth and Water Relations of Tomato Fruits in Relation to Air Vapor Pressure Deficit and Plant Fruit Load. *Journal of Plant Growth Regulation*, 24(3): 201.

Hamann, A., Wang, T., Spittlehouse, D., and Murdock, T. (2013). A comprehensive, high-resolution database of historical and projected climate surfaces for western North America. *Bulletin of the American Meteorological Society*, 94: 1307–1309.

Hancock, A., Brachi, B., Faure, N., Horton, M., Jarymowycz, L., Sperone, G., Tooma- jian, C., Roux, F., and Bergelson, J. (2011). Adaptation to Climate Across the Arabidopsis thaliana Genome. *Science*, 334(6052): 83–86.

Hugly, S., McCourt, P., Browse, J., Patterson, G.W., Somrville, C. (1990). A Chilling Sensitive Mutant of Arabidopsis with Altered Steryl-Ester Metabolism. *Plant Physiology* 93(3): 1053-1062.

Jombart, T. and Ahmed, I. (2011). Adegenet 1.3-1: new tools for the analysis of genome-wide SNP data. *Bioinformatics*, 27(21): 3070–3071.

Kim, J., Nguyen, N., Jeong, C., Nguyen, N., Hong, S., and Lee, H. (2013). Loss of the R2R3 MYB, AtMyb73, causes hyper-induction of the SOS1 and SOS3 genes in response to high salinity in Arabidopsis. *Journal of Plant Physiology*, 170(16): 1461–1465.

Kissoudis, C., Sunarti, S., Van De Wiel, C., Visser, R.G.F., Van Der Linden, C.G., & Bai, Y. (2016). Responses to combined abiotic and biotic stress in tomato are governed by stress intensity and resistance mechanism. *Journal of Experimental Botany*, 67(17): 5119–5132.

Kooyers, N., Greenlee, A., Colicchio, J., Oh, M., and Blackman, B. (2015). Replicate altitudinal clines reveal that evolutionary flexibility underlies adaptation to drought stress in annual Mimulus guttatus. *New Phytologist*, 206(1): 152–165.

Kumar, M.S.S., Ali, K., Dahuja, A., and Tyagi, A. (2015). Role of phytosterols in drought stress tolerance in rice. *Plant Physiology and Biochemsitry* 96: 83-89.

Lasky, J., Des Marais, D., McKay, J., Richards, J., Juenger, T., and Keitt, T. (2012). Characterizing genomic variation of Arabidopsis thaliana: the roles of geography and climate. *Molecular Ecology*, 21: 5512–5529.

Lasky, J., Upadhyaya, H., Ramu, P., Deshhhpande, S., Hash, C., Bonnette, J., Juenger, T., Hyma, K., Acharya, C., Mitchell, S., Buckler, E., Brenton, Z., Kresovich, S., and Morris, G. (2015). Genome-environment associations in sorghum landraces predict adaptive traits. *Science Advances*, 1(6): e1400218.

Lee, C. and Mitchell-Olds, T. (2011). Quantifying effects of environmental and geographical factors on patterns of genetic differentiation. *Molecular Ecology*, 20: 4631–4642.

Legendre, P. and Legendre, L. (1998). Numerical Ecology. Elsevier, *Amsterdam, New York*, 2nd edition.

Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association mapping
and population genetical parameter estimation from sequencing data. *Bioinformatics*,
27(21): 2987–2993.
Li, H. and Durbin, R. (2009). Fast and accurate short read alignment with Burrows- Wheeler
Transform. *Bioinformatics*, 25: 1754–1760.
Li, Z., Peng, R., Tian, Y., Han, H., Xu, J., and Yao, Q. (2016). Genome-Wide Identification and
Analysis of the MYB Transcription Factor Superfamily in Solanum lycopersicum. *Plant*
Cell and Physiology 57(8): 1657-1677.
Lin, Y., Liu, C., and Chen, K. (2019). Assessment of Genetic Differentiation and Linkage
Disequilibrium in Solanum pimpinellifolium Using Genome-Wide High-Density SNP
Markers. *G3* 9(5): 1497-1505.
Manel, S., Poncet, B., Legendre, P., Gugerli, F., and Holderegger, R. (2010). Common factors
drive adaptive genetic variation at different spatial scales in Arabis alpina. *Molecular*
Ecology, 19: 3824–3835.
Manion, G., Lisk, M., Ferrier, S., Nieto-Lugilde, D., Mokany, K., and Fitzpatrick, M. (2018). gdm:
Generalized Dissimilarity Modeling. R package version 1.3.11.
Mishra, M., Singh, G., Tiwari, S., Singh, R., Kumari, N., and Misrra, P. (2015). Characterization
of Arabidopsis sterol glycosyltransferase TTG15/UGT80B1 role during freeze and heat
stress. *Plant Signaling & Behavior*, 10(12): e1075682.
Nakazato, T., Bogonovich, M., and Moyle, L. (2008). Environmental factors predict adaptive
phenotypic differentiation within and between two wild Andean tomatoes. *Evolution*,
62(4): 774– 792.
Nakazato, T., Warren, D., and Moyle, L. (2010). Ecological and geographic modes of species
divergence in wild tomatoes. *American Journal of Botany*, 97(4): 680–693.
Oksanen, J., Blanchet, F., Kindt, R., Legendre, P., Minchin, P., O'Hara, R., Simpson, G., Solymos,
P., Stevens, M., Wagnerr, H., and Oksanen, M. (2013). Package 'vegan'. Community
ecology package, version 2.9.
Pedley, K. and Martin, G. (2003). Molecular basis of Pto-mediated resistance to bacterial speck
disease in tomato. *Annual Review of Phytopathology*, 41: 215–243.
Pitblado, R. and Kerr, E. (1979). A source of resistance to bactereial speck– Pseudomonas tomato.
Tomato Genetics Cooperative, 29: 30.
Raj, A., Stephen, M., and Pritchard, J. (2014). fastSTRUCTURE: Variational Inference of
Population Structure in Large SNP Data Sets. *Genetics*, 197: 573–589.
Rao, E.S., Kadirvel, P., Symonds, R.C., and Ebert, A.W. (2013). Relationship between survival
and yield related traits in Solanum pimpinellifolium under salt stress. *Euphytica* 190(2):
215-228.
Ramirez-Estrada, K., Castillo, N., Lara, J., Arro, M., Boronat, A., Ferrer, A., and Altabella, T.
(2017). Tomato UDP-Glucose Sterol Glycosyltransferases: A Family of Developmental
and Stress Regulated Genes that Encode Cytosolic and Membrane-Associated Forms of
the Enzyme. *Frontiers in Plant Science*, 8, 984.
Razali, R., Bougouffa, S., Morton, M., Lightfoot, D., Alam, I., Essack, M., Arold, S., Kamau, A.,
Schmockel, S., Pailles, Y., Shahid, M., Michell, C., Salim, A., Ho, Y., Tester, M., Bajic,
V., and Negrao, S. (2018). The Genome Sequence of the Wild Tomato Solanum
pimpinellifolium Provides Insights into Salinity Tolerance. *Frontiers in Plant Science*, 9,
1402.

Rellstab, C., Gugerli, F., Eckert, A.J., Hancock, A.M., and Holderegger, R. (2015). A practical
guide to environmental association analysis in landscape genomics. *Molecular Ecology*
24(17): 4348-4370.
Rick, C., Fobes, J., and Holle, M. (1977). Genetic variation in Lycopersicon pimpinelli- folium:
evidence of evolutionary change in mating systems. *Plant Systematics and Evolution*, 127:
139–170.
Rosseel, Y. (2012). lavaan: An R Package for Structural Equation Modeling. *Statistical Software*,
48(2): 1–36.
Salathe, R. and Schmid-Hempel, P. (2011). The genotypic structure of a multi-host bumblebee
parasite suggests a role for ecological niche overlap. *PLoS One*, 6: e22054.
Senthil-Kumar, M., Wang, K., and Mysore, K.S. (2013). AtCYP710A1 gene-mediated
stigmasterol production plays a role in imparting temperature stress tolerance in
Arabidopsis thaliana. *Plant Signaling and Behavior* 8: e23142.
Sork, V., Davis, F., Westfall, R., Flint, A., Ikegami, M., Wang, H., and Grivet, D. (2010). Gene
movement and genetic association with regional climate gradients in California valley oak
(Quercus lobata Nee) in the face of climate change. *Molecular Ecology*, 19: 3806–3823.
Tanksley, S. (2004). The genetic, developmental, and molecular bases of fruit size and shape
variation in tomato. *The Plant cell*, 16: 181–189.
Tanksley, S., Grandillo, S., Fulton, T., Zamir, D., Eshed, Y., Petiard, V., Lopez, J., and Beck-
Bunn, T. (1996). Advanced backcross QTL analysis in a cross between an elite processing
line of tomato and its wild relative L. pimpinellifolium. *Theoretical and Applied Genetics*,
92: 213–224.
Tomato Genome Consortium (2012). The tomato genome sequence provides insights into fleshy
fruit evolution. *Nature*, 485: 635–641.
van den Wollenberg, A. (1977). Redundancy analysis: an alternative for canonical correlation
analysis. *Psychometrika*, 42: 207–219.
Vucetich, J.A. and Walte, T.A. (2003). Spatial patterns of demography and genetic processes
across the species' range: Null hypotheses for landscape conservation genetics.
Conservation Genetics 4(5): 639-645.
Wang, I., Glor, R., and Losos, J. (2013). Quantifying the roles of ecology and geography in spatial
genetic divergence. *Ecology Letters*, 16: 175–182.
Wiens, J. (1989). Spatial Scaling in Ecology. *Functional Ecology*, 3(4): 385–397.
Wright, S. (1943). Isolation by distance. *Genetics*, 28(2): 114–138.
Yoder, J., Stanton-Geddes, J., Zhou, P., Briskine, R., Young, N., and Tiffin, P. (2012). Genomic
signature of adaptation to climate in Medicago truncatula. *Genetics*, 196: 1263–1275.
Zhang, D., Du, Q., Zhang, Z., Jiao, X., Song, X., and Li, J. (2017). Vapour pressure deficit control
in relation to water transport and water productivity in greenhouse tomato production
during summer. *Scientific Reports*, 7: 43461.
Zomer, R., Trabucco, A., Bossio, D., van Straaten, O., and Verchot, L. (2008). Climate Change
Mitigation: A Spatial Analysis of Global Land Suitability for Clean Development
Mechanism Afforestation and Reforestation. Agriculture, Ecosystems, and Environment,
126: 67–80.
Zuriaga, E., Blanca, J.M., Cordero, L., Sifres, A., Blas-Cerdan, W.G., Morales, R., and Nuez, F.
(2009). Genetic and bioclimatic variation in Solanum pimpinellifolium. *Genetic Resources*
and Crop Evolution 56: 39-51.
698

DATA ACCESSIBILITY

Demultiplexed raw read (FASTQ) files from all 140 accessions are deposited in the NCBI SRA

(BioProject: PRJNA561198). All NGS pipeline and data analysis scripts as well as aggregated

environmental data are available on GitHub at https://github.com/gibsonmatt/pimpGEA.

AUTHOR CONTRIBUTIONS

MJSG and LCM designed the experiments; MJSG performed the greenhouse and molecular

work and conducted the bioinformatic analyses; MJSG and LCM wrote the manuscript.

Figure 1: 140 South American accessions included in our analyses. Ecuador (North) and Peru
(South) are outlined in black. (A) Accession collection sites overlain on average vapor pressure
(in units of kPa). Variation in vapor pressure contributed the most to explaining patterns of spatial
genetic variation using RDA. (B) Accessions overlain on precipitation seasonality (coefficient of
monthly variation).
714

Figure 2: Genetic population structure in *S. pimpinellifolium* as revealed by principle components (PC) analysis (top panels) and fastStructure (bottom panel). (A) First two PC axes. Points are colored by membership in the five K-means clusters. (B) Accession collection sites colored by K-mean cluster membership. (B) Admixture proportions of all accessions identified with *fastStructure*, ordered by clusters identified in K-means to show correspondence between the methods.

Table 1: List of dbMEM variables and their contribution to SNP variation as determined by forward variable selection with RDA. Cumulatively more variation was explained by variables describing fine-scale geographic structure (0.070) than large-scale structure (0.054). Significance in forward selection was determined by permutation tests.

Variable	Class	R^2	F	P-val
MEM2	Broad	0.017	2.439	0.001
MEM8	Fine	0.015	2.122	0.001
MEM4	Broad	0.015	2.104	0.001
MEM6	Fine	0.014	1.980	0.002
MEM1	Broad	0.013	1.888	0.001
MEM5	Fine	0.013	1.822	0.003
MEM17	Fine	0.010	1.464	0.02
MEM9	Fine	0.009	1.386	0.038
MEM7	Fine	0.009	1.373	0.037
MEM3	Broad	0.009	1.328	0.041

742 **Table 2:** Top environmental variables and their contribution to spatial SNP variation as determined
743 by forward variable selection with RDA. Results from both the full RDA model selection (not
744 conditioned on space) and partial RDA model selection (conditioned on space) are reported.

RDA (not constrained on space)		RDA (constrained on space)	
Variable	**Contribution to RDA model**	**Variable**	**Contribution to RDA model**
Degree days above 18C	13.60	CV vapor pressure	2.76
Annual max solar radiation	13.45	Prec. seasonality	2.43
Annual average wind speed	13.45	Soil texture	2.25
Isothermality	13.44	Annual max solar radiation	2.23
Annual max wind speed	13.33	Max potential evapotransp.	2.16
CV vapor pressure	13.31	Min potential evapotransp.	1.64
Temperature seasonality	13.17		
Extreme min temperature (30 yr)	13.10		
Max soil water stress	11.55		
Climatic moisture deficit	10.63		
Annual temperature range	9.44		
Mean diurnal range	8.02		
Max potential evapotransp.	6.05		
CV wind speed	3.21		

745
746

Figure 3: Biplot of site (accession) scores for the first two RDA axes, constraining on space with dbMEMs. Points are colored to correspond with genetic clusters identified in PCA (shown in inset, from Figure 2). Top and right axes (red) show the correlation of each environmental predictor with RDA axes 1 and 2, respectively.

Figure 4: Summary of the three variance partitioning methods. Note that SEM and GDM are unable to quantify the independent contribution of environment + space.

Figure 5: Values of squared RDA1 SNP loadings throughout the genome. Top panel: SNP loadings on the first RDA axis of raw SNP variation. Bottom panel: SNP loadings on the first RDA axis accounting for spatial structure among accessions. Note the change in y-axis scale between the two panels.

761 **Table 3:** Annotated gene models located within 10kb of the top 15 outlier SNPs along the first
762 RDA axis, after accounting for spatial structure. We also do not report SNPs within 10kb of SNPs
763 with higher loadings. The full list of outlier SNPs and their annotations can be found in table S6.

Chr.	SNP position	Locus	SNP Category	Distance from Locus (bp)	RDA 1 Loading	Locus Description
4	46907724	Solyc04g050390	Intergenic	30	0.015	60S ribosomal subunit
4	37812488	Solyc04g047830	Intergenic	4389	0.013	DNA glycosylase
4	44823446	Solyc04g049930	Missense	0	0.013	Unknown protein
4	49678371	Solyc04g051150	Intron	0	0.013	Sterol glucosyl transferase 4 (SGT4)
3	66381751	Solyc03g115060	Intergenic	66	0.012	Exocyst complex component 7
5	3609439	Solyc05g009440	Intron	0	0.012	Nuclease S1
1	88554548	Solyc01g098080	Intron	0	0.011	BY-2 kinesin-like protein 5
4	45372222	Solyc04g050080	Missense	0	0.011	MYB transcription factor 73
8	26166364	Solyc08g041710	Intron	0	0.011	Transmembrane protein
6	39445574	Solyc06g062360	Intron	0	0.011	Syntaxin-like protein
11	417966	Solyc11g005560	Intergenic	658	0.011	Cellulose synthase
8	27570643	Solyc08g023500	Intron	0	0.011	Metallohydrolase/oxioreductase
4	5709089	Solyc04g015490	3` UTR	0	0.011	Magnesium chelatase subunit D
4	45599110	Solyc04g050150	Intron	0	0.011	RNA helicase DEAH-Box 13
8	23509033	Solyc08g042140	Intron	0	0.011	Translation initiation factor 3 subunit

765