Evaluation of eight protocols for genomic DNA extraction of *Hypostomus*

commersoni (Valenciennes, 1836), Siluriformes: Loricariidae.

Priscila Mezzomo[1,#a,*], Albanin A. Mielniczki-Pereira[1,¶], Tanise L. Sausen[1,¶], Jorge R. Marinho[1,¶],

Rogério L. Cansian[1,¶]

[1]Department of Biological Sciences, URI, Erechim, RS, Brazil

[#a]Current Adress: Department of Biological Sciences-Structural Genomics Consortium,

UNICAMP, Campinas, SP, Brazil

*Corresponding author

E-mail: priscilamez@gmail.com

¶These authors contributed equally to this work.

Word count: 2704.

Abstract word count: 198.

Number of figures: 1.

Number of tables: 4.

Conflicts of Interest: The authors have declared that no competing interests exist.

Abstract

The principle and techniques applied in DNA extraction play a pivotal role in the obtention of a considerable and purified yield of genetic material. In this work, we evaluate the efficiency of seven protocols chosen from literature for the DNA extraction of *Hypostomus commersoni* (Valenciennes, 1836), an important component of South American freshwater ichthyofauna. We also proposed and evaluate an improved protocol, based on the combination of variables from the selected methodologies that demonstrated higher potential for extracting *H. commersoni* DNA. The quality of samples was assessed through spectrophotometry, gel electrophoresis and PCR-RAPD amplification. The efficiency of DNA extraction was influenced both by the method applied and target-tissue of choice. Higher concentrations and yield of DNA were obtained from ocular tissue, with a positive spectrum of incubation in lysis buffer for up to 36 hours after sample collection, without maceration, using fresh tissues and in the presence of high concentration of Proteinase K. In these conditions, samples were successfully amplified using a set of primers from Operon Technologies. To date, there is no record of description for the parameters analyzed in this work, neither the description of RAPD markers for *H. commersoni*, evidencing the importance of our findings.

Key-words: DNA extraction; PCR-RAPD; eye tissue; fresh fish tissue; proteinase K.

Introduction

A crucial step for high quality genetic analysis is considered to be a satisfactory DNA extraction. Nevertheless, this purpose is not always easy to achieve, since different biological compounds, such as lipids and proteins, can act as contaminants and interfere in the final quality of the product [1]. The principles and techniques applied in DNA extraction play a pivotal role in obtaining a considerable and purified yield of this molecule [2].

A variety of methods has been established to isolate DNA from biological materials [3,6]. Regardless the method of choice, the procedure usually consists of three steps: lysis, purification and DNA recovery, being the first the most critical stage, since is at this point that the chances of causing DNA damages are higher [7].

The quality and amount of available DNA strand template have a major influence on genetic analyzes based on PCR (Polymerase Chain Reaction), for example [8,9]. Therefore, an ideal extraction technique should optimize DNA yield, minimize DNA degradation, and be efficient in terms of cost, time, labor, and supplies [10].

With the availability of a plethora of molecular analysis machinery, there is an exponential demand for high quality samples, capable to generate great results with minimum of errors. To meet these requirements, commercial kits were developed, aiming for DNA extractions of excellence [11-13].

However, the market price of this products it is not always easily affordable to every laboratory [10, 14]. In the other hand, the extraction of genetic material with non-commercial reagents is less costly due to the use of homemade reagents and, therefore, such methodologies can be applied in laboratories with less financial resources.

Stablish a well succeed protocol for DNA extraction is a very important step in genetic analysis, considering that the same method may present variations between

different organisms [15]. Furthermore, the quality of isolated DNA through methods described for other taxa, may present unsatisfactory results when applied to fish species [2].

H. commersoni (Valenciennes, 1836), commonly known as suckermouth catfish, is an important component of freshwater ichthyofauna, inhabiting the Paraná and Uruguay basins, in South America [16, 17]. This taxon is marked by its ecological and commercial relevance [18-22], and, despite that, there is no record of description of well-succeed DNA extraction protocol nor PCR-RAPD markers to date.

In this work, we evaluated seven protocols chosen from literature in the extraction of genomic DNA of *H. commersoni* and, also, we propose an improved method, aiming to obtain high quality and purified DNA samples of this specie.

Materials and methods

Ethics Statement

This study was approved by the CEUA (Ethics Committee for Animal Use) of URI (Regional Integrated University). Written informed consent was obtained from the regulatory organ.

Sample collection

Specimens of *H. commersoni* were subjected to euthanasia with 0.1% clove oil in aquarium water, and the tissues selected for DNA extraction were removed and placed in a sterile 2 mL eppendorf tube for subsequent analysis according to each protocol particular necessity.

DNA Extraction

101 All DNA was extracted and quantified at the biochemistry and molecular biology

102 laboratory of URI (Regional Integrated University, Erechim-RS/Brazil).

Protocols and variables analyzed in the DNA extraction

Seven protocols from literature were selected for DNA extraction of *H. commersoni*,

whose methodologies had previously been described for fish species or whose parameters

were similar to those intended to be analyzed in this work, such as the same extraction

conditions and target tissues (Box 1). An eighth protocol, elaborated by the combination of

the variables that presented better potential to extract *H. commersonii* DNA in each of the

seven previously tested protocols, was proposed and evaluated too.

Box 1 – Protocols evaluated for DNA extraction.

Protocol	Reference[1]
1	[26] Doyle and Doyle (1987) *CTAB.
2	[27] Medrano et al. (1990) *STE/EDTA.
3	[29] Whitmore et al. (1992) *PHENOL.
4	[23] Aljanabi and Martinez (1997) *SALTS.
5	[28] Faleiro et al. (2003) *CTAB.
6	[24] Barrero et al. (2008) *SALTS.
7	[25] Coppini (2009) *STE/EDTA.
8	Developed in this work[2] *STE/EDTA.

113 [1] The full description of protocols 1 to 7 is available as supplementary material (S1).
114 [2] Described in full in the results section.
115 *Principle of the method.
116

117 In combination to the protocols, we also evaluate eight different variables that, based on

118 literature findings and our previous analysis, was tough to have an influence on the purity

119 and amount of DNA obtained, as follows:

1. **Preservation of specimens**: fresh tissues (1. collection, 2. euthanasia, 3. tissue extraction and immediate use) or stored in 70% and 95% ethanol at temperatures of 25 ° C, -10 ° C and -20 ° C;

2. **Maceration**: mechanic lysis (vortex), liquid nitrogen or manual lysis (with sterile glass stick);

3. **Target-tissue**: dorsal fin, caudal fin, muscle, gills, blood and ocular tissue;

4. **Time of incubation in extraction buffer**: 1 hour (55 ºC), 12 hours (27 ºC), 34 hours (27 ºC) and 36 hours (27 ºC);

5. **RNAse**: presence and absence, with gradients of concentration;

6. **Proteinase K**: presence and absence, with gradients of concentration;

7. **Final storage**: at 4 ºC, -10 ºC and -20 ºC;

8. **Integrity and quantification**: on the day of extraction, after 10, 20 and 30 days of storage.

Several tests aiming for experimental optimization were performed and for each variable, the factors capable to generate better concentration and purity of DNA were selected. The results of extractions for each variable were categorized into a qualitative matrix including three classes, according to the effect on the final extracted product:

(i) **Efficient**: satisfactory electrophoresis and spectrophotometry;

(ii) **Indifferent**: did not influence the parameters evaluated;

(iii) **Inefficient**: low concentration of genetic material, low purity or degradation of samples.

Analysis of DNA integrity

The integrity of the obtained samples was verified through agarose gel (0,8%) electrophoresis, stained with ethidium bromide (1 mg/ml). The reaction run for 150 minutes at 90 volts and the visualization and image caption of the gel were obtained under ultraviolet (UV) light.

Evaluation of concentration and purity of DNA

The samples whose extraction was successful and did not result in DNA degradation according to the electrophoresis, where submitted to spectrophotometry and analyzed for protein contamination (purity). Samples were quantified in a spectrophotometer (model NI 1600UV - Nova Instruments), at A_{260} and A_{280} nanometers (nm). The purity of the DNA was confirmed by the ratio of absorbance values at the two wavelengths measured [33]. The DNA concentration of each sample was determined by the equation: [DNA] = 50 µg/ml x D (dilution factor) x A_{260} (read value obtained in the wavelength of 260 nm). Satisfactory purity values were considered to be ranging between 1.7-2.0.

DNA amplification by PCR-RAPD (Polymerase Chain Reaction - Random Amplified Polymorphic DNA)

Samples that reached the aforementioned parameters of DNA concentration and yield were submitted to amplification with different primers of PCR-RAPD. In total, 32 primers from Operon Technologies kits OPA (01, 02, 03, 04, 05, 06, 07, 08, 09), OPB (03, 05, 07, 10, 11, 15), OPF (10, 11, 12 e 13), OPW (01, 02, 03, 04, 05, 06, 07, 08, 09, 10) and OPY (01,02,03) were tested. Ten nanograms of genomic DNA was amplified in PCR buffer (20 mM Tris-Hcl pH8.4, 50 mM KCL), 4 mM MgCl2, 0.7 mM dNTPs, 0.3 µM primers e 1

unity of Taq DNA polymerase enzyme (Invitrogen). The reaction was conducted in a final

volume of 25 µL, in the following conditions: denaturation at 95ºC for 30 s, annealing at

50ºC for 1 min e extension at 72ºC for 5 min, totaling 40 cycles. The amplification

products were applied in agarose gel (1.5%), stained with ethidium bromide (1 mg/ml).

The reaction run for 150 minutes at 150V. Visualization and image caption of band

patterns were obtained under ultraviolet (UV) light.

Statistical analysis

177 Statistical analysis was performed using Python (version 3.6.0). Differences between DNA

178 extraction methods in total DNA concentration were determined using ANOVA with

179 Tukey HSD post-hoc test.

Results

DNA extraction method impacts DNA yield

Our work corroborated that even a method that uses in-house produced reagents instead of

commercial kits is capable to extract DNA with sufficient quality to be used in molecular

analysis, as PCR-RAPD. The amount of DNA extracted was dependent on the DNA

extraction technique and the target-tissue of choice. For all variables examined, the

improved extraction method developed and proposed at this work, produced the greatest

total DNA mean yields, for all tissues, resulting in significantly higher absorbance ratios

when compared to the seven other extraction methods. Different variables also affect in

different ways the extraction of DNA, as showed in Table 1.

Table 1. Effect of different variables in the DNA extraction of *H. commersoni*.

Variables	Efficient[1]	Indifferent[2]	Inefficient[3]
Specimens conservation	Fresh (27 ºC)	-	Fresh (-10 ºC and -20 ºC) 70% and 95% ethanol (4 ºC, -10 ºC and -20 ºC)
Maceration	Lysis with buffer	Liquid nitrogen	In vortex Manual lysis
Target-Tissue	Ocular tissue	–	Caudal fin Dorsal fin Muscle Gills Blood
Time of incubation	1h (55 ºC) 12h, 24h and 36h (27 ºC)	–	–
RNAse	–	Presence	–
Proteinase K	Presence in high concentration ($\geq$ 20 mg.ml^{-1})	–	–
Storage	-10 ºC -20 ºC	–	4 ºC
Quantification	In the extraction day After 10, 20 and 30 days of storage	–	–

194 [1]Efficient: satisfactory electrophoresis (integrity) and spectrophotometry (quantification);

195 [2]Indifferent: did not influenced the final amount and the purity of DNA extracted;

196 [3]Inefficient: sample degradation and low purity observed.

Improved protocol for DNA extraction of *H. commersoni*

The eighth protocol evaluated, improved and proposed in this work, in general, presented

the highest efficiency in the extraction of *H. commersoni* DNA. The samples submitted to

extraction through this protocol were the only ones that amplified with PCR-RAPD

primers, since they had the best DNA concentration and yield of purify DNA in the

samples (according to the absorbance ratio of 260/280 nanometers).

The complete methodology of protocol 8, including reagents and steps of

extraction, is describe in Box 2.

206
207

208 **Box 2.** Description of reagents and steps for DNA extraction of *H. commersoni* through the
209 method proposed in this work.

Reagents	
STE Buffer	0,1 M NaCl + 10 mM Tris + 1 mM EDTA (pH 8,0)
Lysis Buffer	2,5 mM Tris-HCl pH 8,0 + 0,1 mM EDTA pH 8,0 + 0,1% SDS + 8,0 mM NaCl
Proteinase K	20 mg.mL^{-1}
CIA	Chloroform + isoamylic alcohol (1:24 v/v)
TE Buffer	10 mM Tris pH 8,0 + 1 mM EDTA pH 8,0
TBE Buffer	500 mM Tris-HCl + 60 mM boric acid + 83 mM EDTA
Steps of extraction	

1. Collect 20 mg of fresh tissue.
2. Wash in 500 µl of STE (cold) and discard.
3. Add 500 µl of lysis buffer, 2 µL of β-Mercaptoetanol and 13 µl of proteinase K, in this order, and incube at 55 ºC for 1 hora.
4. Add 700 µl of CIA and homogenize by inversion.
5. Centrifuge for 10 minutes at 9,000 rpm and transfer the supernatant to a new tube.
6. Repeat the steps 4 and 5 (3x).
7. Transfer the aqueous phase to a new tube containing 900 µl of absolute isopropanol (cold), homogenize by inversion.
8. Incubate for 2 hours at –20 ºC.
9. Centrifuge for 15 minutes at 9000 rpm and discard the supernatant.
10. Wash the resulting *pellet* with ethanol analytical grade (cold).
11. Discard all the volume and incube at 37 ºC for 15 minutes.
12. Add 100 µL of TE buffer and incube for 1 hour at ≈ 25 ºC.
13. Storage at – 10 ºC until required.

210 *This protocol described the sample incubation in lysis buffer for 1 hour (step 3) but can also be
211 applied for the spectrum of 6, 12 or 36 hours previous to the continuity of extraction (steps 4-13).
212

Analysis of DNA integrity in electrophoresis gel

All samples from the six tissues and the eight applied conditions were submitted to gel

electrophoresis in triplicate, resulting in a total count of 144 samples. The ocular tissue

presented the highest integrity of extracted DNA, in all combinations of variables and

protocols. For all tissues, the main bands of DNA were around 46-48 kb in size.

Comparing protocols, the EDTA methods (P2, P7 and P8) showed relatively lighter smear

tails, indicating less DNA degradation.

Quantification and assessment of purity

The resulting values of quantification for each DNA sample extracted under the different

conditions and through the different protocols evaluated, are presented in Table 2 and 3

and Fig 1A–1F. The amount of DNA obtained, and final yield mean values varied between

different tissue samples, as well as between samples of the same tissue but extracted using

different protocols.

Table 2. Mean of total DNA amount (ng/µl) extracted using eight different methods for six
different tissues.

Method (Protocol)	CF	DF	MC	GL	BD	OT	Methods Mean (ng/µl)
	MEAN OF TOTAL DNA AMOUNT (ng/µl)						
	Mean ± SD						
1. Doyle & Doyle (1987)	120 ±21.2	139 ±26.9	198 ±10.6	110 ±22	365 ±60.4	400 ±26.8	222
2. Medrano et al. (1990)	175 ±8.6	114 ±23.7	155 ±32.8	137 ±27.8	375 ±115.2	520 ±62.8	246
3. Whitmore et al. (1992)	116 ±18.0	74 ±19	173 ±35.3	125 ±6.5	335 ±70.3	455 ±48.8	216
4. Aljanabi & Martinez (1997)	181 ±17.3	90 ±12.3	120 ±33.2	110 ±12.6	455 ±60.1	465 ±50.9	237
5. Faleiro et al. (2003)	168 ±39.9	150 ±24.5	205 ±28.1	155 ±10.6	425 ±20.4	505 ±63.9	268
6. Barreto et al. (2008)	157 ±29.0	97 ±9.6	207 ±32.6	175 ±18.7	460 ±51.9	555 ±47.3	275
7. Coppini (2009)	186 ±6.6	160 ±44.4	170 ±19.1	135 ±1.6	330 ±17.1	516 ±22.2	250
8. Improved protocol (2019)	209 ±15.7	220 ±33.5	216 ±0.8	210 ±4.1	463 ±9.7	712 ±18.8	438
Tissues Mean	**164**	**131**	**180**	**145**	**404**	**516**	-

229 ***Tissues:** CF: caudal fin. DF: dorsal fin. MC: muscle. GL: gills. BD: blood. OT: ocular tissue.
230 **Sample values from read in the wavelength of 260 nm where applied to the equation [DNA] = 50
231 µg/mL x D (dilution factor) x A_{260} to obtain its concentration in ng per µl.
232

233 **Table 3.** Mean of DNA amount from ratio 260/280 nm, extracted using eight different
234 methods for six different tissues.

Method (Protocol)	CF	DF	MC	GL	BD	OT	Methods Mean
	ABSORBANCE RATIO (OD 280/260 nm)						
	Mean ± SD						
1. Doyle & Doyle (1987)	1.5 ±0.016	0.8 ±0.08	1.4 ±0.08	1 ±0	1.9 ±0.08	1.8 ±0.12	1.4
2. Medrano et al. (1990)	1.8 ±0.08	0.6 ±0.08	1.6 ±0.08	1 ±0.08	1.4 ±0.08	1.6 ±0.08	1.3
3. Whitmore et al. (1992)	1 ±0.14	0.7 ±0.08	1.6 ±0.08	1.2 ±0	1.6 ±0.08	1.8 ±0.08	1.3
4. Aljanabi & Martinez (1997)	1.1 ±0.16	0.8 ±0.08	1 ±0.08	1 ±0	1.7 ±0.08	1.6 ±0.08	1.2
5. Faleiro et al. (2003)	1.1 ±0.08	0.9 ±0.08	1.6 ±0.08	1.1 ±0.14	1.6 ±0	1.9 ±0.08	1.4
6. Barreto et al. (2008)	1.1 ±0.08	0.9 ±0.08	1.3 ±0.08	1.2 ±0	1.8 ±0.08	1.7 ±0.08	1.3
7. Coppini (2009)	1.5 ±0.08	1.1 ±0.12	0.8 ±0.08	1.1 ±0.08	1.6 ±0	1.8 ±0.08	1.3
8. Improved protocol (2019)	2 ±0	1.5 ±0.08	1.7 ±0.08	1.5 ±0.08	1.9 ±0.08	2 ±0.08	1.8
Tissues Mean	**1.4**	**0.9**	**1.3**	**1.1**	**1.7**	**1.8**	-

235 ***Tissues:** CF: caudal fin. DF: dorsal fin. MC: muscle. GL: gills. BD: blood. OT: ocular tissue.
236 **Sample values from ratio of the reads in the wavelength of 260 and 280 nanometers, to quantify
237 the purity of DNA in the samples.
238

Fig 1. Mean of total DNA amount (± standard deviation) extracted from six different tissues using eight different protocols. Samples were extracted and total DNA was measured by spectrophotometry. Statistical significance was determined using one way ANOVA with Tukey-HSD post-hoc test. Significance defined by $p \leq 0.05$ and denoted by lower case letters (samples marked with the same letter are significantly equal). Whiskers denoting lower standard deviation are cropped at zero.

When comparing samples, the ocular tissue had the highest mean yield of extracted DNA (516 ng/µl), while the lowest concentration was for the dorsal fin (131ng/µl). In relation to the protocols, means between different tissues indicated that the lowest value was for

extractions performed using protocol 3 (216 ng/µl), while the highest mean concentration

of DNA was obtained with protocol 8 (438 ng/µl). Statistically, as indicated by the lower

case letters, there is no significant difference between most of the samples whose DNA

was extracted from the tissues dorsal fin, caudal fin, gills and muscle. The major

exceptions were blood and ocular tissue, which had similar p value for most of the

evaluated protocols.

Although the total amount of DNA present high values and several samples were

statistically similar, the mean values of the absorbance ratio are the important data, once

concerns the purity of samples and DNA amount. The absorbance ratio values were 1.2

(protocol 4), 1.3 (protocols 2, 3, 6 and 7), 1.4 (protocols 1 and 5) and 1.8 (protocol 8),

confirming its higher efficiency in comparison to the others, indicated also by the lowest

observed standard deviation.

DNA extraction method influences PCR-RAPD amplification

All samples were subject to PCR-RAPD analysis, but only the ones extracted with protocol

8 amplified with the selected primers. Of the 34 primers tested, 9 amplified the DNA of the

evaluated specie in the concentrations of 40 and 20 ng/µl and were described in this work.

These markers present potential to be used in studies evaluating the genetic structure of

ocular tissue samples of *H. commersoni* through PCR-RAPD markers.

The score of bands varied between the amplifications using different concentrations

of DNA (Table 4). The concentration of 20 ng/µl generated the major count of fragments,

followed by the concentration of 40 ng/µl. The concentration of 10 ng/µl of DNA did not

amplify with the primers OPW05, OPW06, OPW07, OPW08 and OPW09, and, therefore,

this concentration is the less indicated for use in the PCR-RAPD reaction with this set of

primers and specie, according to our findings.

Table 4. Score of DNA bands amplified for each primer.

Primer	Sequence 5' - 3'	Score of Amplified DNA bands		
		40 ng/µl	20 ng/µl	10 ng/µl
OPW-01	CTCAGTGTCC	6	6	6
OPW-02	ACCCCGCCAA	6	8	8
OPW-03	GTCCGGAGTG	4	8	7
OPW-04	CAGAAGCGGA	7	9	7
OPW-05	GGCGGATAAG	4	4	-
OPW-06	AGGCCCGATG	7	8	-
OPW-07	CTGGACGTCA	7	8	-
OPW-08	GACTGCCTCT	5	5	-
OPW-09	GTGACCGAGT	6	6	-

278 **Discussion**

The relevance in evaluating different protocols for DNA extraction and the description of

an efficient method for the specie *H. commersoni* is justified by the fact that obtaining high

quality samples is crucial for studying organisms at genetic level. Specifically, in the case

of *H. commersoni*, there are few molecular, genetic and/or phylogenetic data available in

the literature. The information compilated at this work may allow future analyzes of

molecular characterization of the group and, nevertheless, the techniques described here

have potential to be used to study other organisms or fish species

In this work, we proposed a methodology based in the DNA extraction from ocular

tissue and, to our knowledge, there is no record of this parameter for any fish species to

date. The use of this tissue may present some limitations, such its availability in number

and size, but our results indicate that this organ have a great potential as source for extract

good quality genomic DNA in *H. commersoni* and possibly in other fish species.

This technique can be useful in cases where DNA extraction is performed using

samples belonging to scientific collections, where the specimens have already bein

euthanized, or when an alternative model of extraction source is required, once the

classical ones have been exhausted without present a positive result.

During the test period, all the steps that presented promising results regarding the

primary objective of our study were selected, aiming to group them into a single protocol,

considered the "ideal" for the *H. commersoni* species. In this way, it is possible to discuss

some of the characteristics that presented better efficiency in terms of quantity and quality

of isolated DNA.

The use of liquid nitrogen for tissue maceration in DNA isolation, especially in

solid tissues, is a method described as efficient according to several findings [35].

However, in this work, it has not been shown to be a factor influencing the quality or

quantity of product obtained. This data is important because it results in a protocol of lower

cost and greater ease of execution, facilitating the application of this methodology in a

larger number of laboratories and research groups.

The addition of RNAse is suggested as important in protocols for extracting fish

DNA, being used in the steps of breaking tissues and flakes, in order to eliminate the

contamination of the samples by RNA and to increase DNA yield [30]. However, in this

study the use of RNAse had no effect on DNA extraction. For all protocols and variables

tested, RNAse did not involve more pure or concentrated samples, even when evaluated in

several gradients and at different stages of each protocol. The possibility of eliminating the

use of RNAse during DNA extraction also results in the development of a lower cost

protocol.

Other variables, however, corroborated with information found in the literature,

such as the use of proteinase K to increase extraction efficiency, especially at temperatures

of sample incubation up to 55 ºC. Proteinase K digestion is characterized by the

purification of quality DNA for genetic analyzes such as PCR [31, 32].

One of the most promising results was the possibility of exploring different

incubation times of the samples in the extraction buffer, with a proven spectrum of

efficiency in 1, 12, 24 and 36 hours, using the ocular tissue of *H. commersoni*. The

advantages related to this discovery are the possibility of extracting the DNA in a single

day (1 hour at 55 ºC), incubating the samples overnight, or, furthermore, being able to

postpone extraction after its incubation in lysis buffer, until 36 hours at room temperature

of approximately 25 ° C. This result is also important because it facilitates the transport of

material where the specimens can be euthanized in the field and the ocular tissue

maintained in lysis buffer for up to 36 hours without the need for refrigeration or any other

form of preservation until the DNA can be extracted in adequate conditions.

In general, all 7 protocols chosen from the literature and evaluated in this work

presented relative efficiency in the extraction of *H. commersoni* DNA, despite some

limitations, when evaluating the concentration indices obtained from each sample.

However, when the values were submitted to the absorbance ratio equation, it is evident

that the protocol proposed in this paper was the one that obtained the highest DNA purity

values in the samples.

The protocol that presented higher efficiency is a simple and economic method to

obtain DNA from *H. commersoni*, with the potential to be applied in other similar species.

Following the methodology described in our improved protocol, it was possible to submit

the samples to the amplification reaction, using primers for PCR-RAPD. The samples from

the other protocols were submitted to the same analysis, but without success.

Under optimal conditions, a single cell contains broadly DNA to serve as a template

for PCR reactions. The usefulness of this protocol in PCR-based genetic studies was

confirmed through the amplification of samples using PCR-RAPD markers. Our findings

brought unprecedented data for *H. commersoni*, and this information can be applied in

further studies of genetic characterization of this taxon.

Acknowledgements

The authors thanks CAPES, CNPq, FAPERGS and the Post-Graduation Program in Ecology from URI (Integrated Regional University-Erechim/RS, Brazil), for the financial support and scholarships provided.

References

1. Bauer M, Patzelt D (2003) A method for simultaneous RNA and DNA isolation from dried blood and semen stains. Forensic Science International 136:76-80.

2. Muhammad H, Iqbal Z, Iqbal MU, Younas T, Bashir Q (2016) An efficient method for DNA isolation from fish fin. Pak. Journal of Agricultural Science 53:843-850.

3. Milligan BG (1998) Total DNA isolation. In: Hoelzel AR, ed. Molecular Genetic Analysis of Population: A Practical Approach, 2nd Edition. Oxford, New York, Tokyo: Oxford University Press. pp 29–64.

4. Reischl U, Linde HJ, Metz M, Leppmeier B, Lehn N (2000) Rapid identification of methicillin-resistant Staphylococcus aureus and simultaneous species confirmation using real-time fluorescence PCR. Journal of Clinical Microbiology 38:2429–2433.

5. Rantakokko-Jalava K, Jalava J (2002) Optimal DNA isolation method for detection of bacteria in clinical specimens by broad-range PCR. Journal of Clinical Microbiology 40:4211–4217.

6. Tongeren SP, Degener JE, Harmsen HJM (2011) Comparison of three rapid and easy bacterial DNA extraction methods for use with quantitative real-time PCR. European Journal of Clinical Microbiology & Infectious Diseases 30:1053–1061.

7. Barbosa C, Nogueira S, Gadanho M, Chaves S (2015) DNA extraction: Finding the most suitable method. In Molecular microbial diagnostic methods: Pathways to implementation for the food and water industries. Elsevier Inc., pp. 135–154.

8. Ward RD, Hanner R, Hebert PDN (2009) The campaign to DNA barcode all fishes, FISH-BOL. Journal of Fish Biology 74:329-356.

9. Wang TY, Wang L, Zhang JH, Dong WH (2011) A simplified universal genomic DNA extraction protocol suitable for PCR. Genetic Molecular Research 10:519-525.

10. Chen H, Rangasamy M, Tan SY, Wang H, Siegfried BD (2010) Evaluation of Five Methods for Total DNA Extraction from Western Corn Rootworm Beetles. PLoS ONE 5:11963.

11. Coombs NJ, Gough AC, Primrose JN (1999) Optimization of DNA and RNA extraction from archival formalin-fixed tissue. Nucleic Acids Res 27:16-12.

12. Barea JA, Pardini MIMC, Gushiken T (2004) Extração de DNA de materiais de arquivo e fontes escassas para utilização em reação de polimerização em cadeia (PCR). Rev Bras Hematol Hemoter 26:274-81.

13. Bueno V (2004) DNA e aperfeiçoamento das técnicas de extração. Rev Bras Hematol Hemoter 26:233-4.

14. Mega NO, Revers LF (2011) Developing a rapid, efficient and low cost method for rapid DNA extraction from arthropods. Ciência Rural 41:1563-1570.

15. Lucentini L, Caporali S, Palomba A, Lancioni H, Panara F (2006) A comparison of conservative DNA extraction methods from fins and scales of freshwater fish: A useful tool for conservation genetics. Conservation Genetics 7:1009-2012.

16. Burgess WE (1989). An atlas of freshwater and marine catfishes. A preliminary survey of the Siluriformes. TFH Publications, Inc., Neptune City, New Jersey (USA). pp 784.

17. Reis RE, Kullander SO, Ferrari JR CJ (2003) Check list of the freshwater fishes of South and Central. América. Porto Alegre: EDIPUCRS, 742.

18. Buck S, Sazima I (1995) An assemblage of mailed catfishes (Loricariidae) in southeastern Brazil: distribution, activity, and feeding. Ichthyological Exploration of Freshwaters 6:325–332.

19. Garavello JC, Garavello JP (2004) Spatial distribution and interaction of four species of the catfish genus Hypostomus Lacépède with bottom of Rio São Francisco, Canindé do São Francisco, Sergipe, Brazil (Pisces, Loricariidae, Hypostominae). Brazilian Journal of Biology 64:103-141.

20. Baldisserotto B (2009) Piscicultura continental no Rio Grande do Sul: situação atual, problemas e perspectivas para o futuro. Ciência Rural 39:291-299.

21. Gonçalves AP (2011) Ecologia e etnoecologia de *Hypancistrus zebra* (Siluriformes: Loricariidae) no Rio Xingu, Amazônia Brasileira. Dissertação (Mestrado) - Programa de Pós-graduação em Ecologia Aquática e Pesca, Instituto de Ciências Biológicas, Universidade Federal do Pará. Belém, 137-pp.

22. Lujan NK, Winemiller KO, Armbruster JW (2012) Trophic diversity in the evolution and community assembly of loricariid catfishes. BMC Evolutionary Biology 12:124.

430 23. Aljanabi SM, Martinez I (1997) Universal and rapid salt-extraction of high quality
431 genomic DNA for PCR-based techniques. Nucleic Acid Research 25:4692-3.
432
433 24. Barrero NML, Povh JA, Ribeiro RP, Gomes PC, Jacometo CB, Lopes TS (2008)
434 Comparison of DNA extraction protocols of fish fin and larvae samples: modified salt
435 (NaCl) extraction. Ciencia e Investigacion Agraria 35:65-74.
436
437 25. Coppini VJ (2009) Variabilidade genética em *Oligoryzomys nigripes* através de
438 marcadores RAPD e cariotipagem, tendo o rio Uruguai como barreira geográfica. Master
439 Thesis in Ecology - URI (Universidade Regional Integrada do Alto Uruguai e das
440 Missões), Erechim/RS - Brazil.
441
442 26. Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of
443 fresh leaf tissue. Phytochemical Bulletin 19:11-15.
444
445 27. Medrano JF, Aasen E, Sharrow L (1990) DNA extraction from nucleated red blood
446 cells. Biotechniques 8:43.
447
448 28. Faleiro FG, Faleiro ASG, Cordeiro MCR, Karia CT (2003) Metodologia para
449 operacionalizar a extração de DNA de espécies nativas do cerrado. Planaltina: Embrapa
450 Cerrados 1:1-6.
451
452 29. Whitmore DH, Thai TH, Craft M (1992) Gene Amplification Permits Minimally
453 Invasive Analysis of Fish Mitochondrial DNA. Transactions of the American Fisheries
454 Society 12:170-177.
455
456 30. Wasko AP, Martin C, Oliveira C, Forresti F (2003) Non-destructive genetic sampling
457 in fish. An improved method for DNA extraction from fish fins and scales. Hereditas
458 138:161-165.
459
460 31. Sweeney PJ, Walker JM (1993) Enzymes of molecular biology. In: *Methods in*
461 *Molecular Biology*, M.M. Burrell, ed., Humana Press 16:305.
462
463 32. Goldenberger D, Perschil I, Ritzler M, Altwegg M (1995) A simple "universal" DNA
464 extraction procedure using SDS and proteinase K is compatible with direct PCR
465 amplification. PCR Methods and Applications 4:368-70.
466
467 33. Barbosa MM (1998) Quantificação e controle da qualidade do DNA genômico. In:
468 Milach SCK, ed. Marcadores moleculares em plantas. Porto Alegre: Sandra Milach. pp 99-
469 106.
470
471 34. Mullis KB, Faloona FA (1987) Specific synthesis of DNA in vitro via a polymerase-
472 catalyzed chain reaction. Methods in Enzymology 155:335-350.
473

474   35. Chen H, Leibenguth F (1995) Studies on multilocus fingerprints, RAPD markers, and
475   mitochondrial DNA of a gynogenetic fish *Carrasius auratusfibelio*. Biochemical genetics
476   3:297-306.

477
478
479