The inhibition of LSD1 via sequestration contributes to tau-mediated neurodegeneration

Amanda K. Engstrom[1], Alicia C. Walker[1], Rohitha A. Moudgal[1], Dexter A. Myrick[1], Stephanie M. Kyle[1], Yu Bai,[1] M Jordan Rowley[2], David J. Katz[1*]

[1] Department of Cell Biology, Emory University School of Medicine, Atlanta GA 30322, USA.
[2] Department of Genetics, Cell Biology, and Anatomy, University of Nebraska Medical Center, Omaha, NE 68198

*Correspondence to: David J

Email: djkatz@Emory.edu.

Classification

Biological Sciences: Neuroscience

Keywords

LSD1, Neurodegeneration, Tauopathy, Alzheimer's disease

Author Contributions

A.K.E and D.J.K. conceived and designed the study and wrote the manuscript. A.K.E. performed experiments with the help of A.C.W, S.M.K, R.A.M and Y.U under the direction of D.J.K. D.A.M and M.J.R performed bioinformatics analysis on the RNA sequencing data. A.K.E, A.C.W, S.M.K, and R.A.M analyzed all other data and interpreted results. All authors discussed the results and commented on the manuscript.

This PDF file includes:

Main Text

Main Figures 1-6

ABSTRACT

Tauopathies are a class of neurodegenerative diseases associated with pathological tau. Despite many advances in our understanding of these diseases, the direct mechanism through which tau contributes to neurodegeneration remains poorly understood. Previously, our lab implicated the histone demethylase LSD1 in tau-induced neurodegeneration by showing that LSD1 localizes to pathological tau aggregates in Alzheimer's disease cases, and that it is continuously required for the survival of hippocampal and cortical neurons in mice. Here, we utilize the P301S tauopathy mouse model to demonstrate that pathological tau can exclude LSD1 from the nucleus in neurons. In addition, we show that reducing LSD1 in these mice is sufficient to highly exacerbate tau-mediated neurodegeneration and tau-induced gene expression changes. Finally, we find that overexpressing LSD1 in the hippocampus of tauopathy mice, even after pathology has formed, is sufficient to significantly delay neurodegeneration and counteract tau-induced expression changes. These results suggest that inhibiting LSD1 via sequestration contributes to tau-mediated neurodegeneration. Thus, LSD1 is a promising therapeutic target for tauopathies such as Alzheimer's disease.

SIGNIFICANCE STATEMENT

We have made the novel discovery that pathological tau functions through the histone demethylase LSD1 in the Alzheimer's disease pathway. Thus, we have identified a mechanism that links tau to the downstream neuronal dysfunction pathways. This step can potentially be targeted therapeutically, after the onset of dementia symptoms, to block the progression of dementia in Alzheimer's disease patients.

INTRODUCTION

Tauopathies such as corticobasal degeneration, progressive supranuclear palsy, and frontotemporal lobar degeneration with tau inclusions are neurodegenerative diseases pathologically defined by different forms of tau positive intraneuronal deposits (*1-5*). In addition to these primary tauopathies, neuropathological observations of postmortem Alzheimer's disease (AD) brains show the presence neurofibrillary tangles (NFTs) of hyperphosphorylated tau protein, as well as plaques containing β-amyloid (Aβ) peptide (*6-9*). AD is the leading cause of age-related dementia, resulting from neuronal cell death in the frontal and temporal cortices, as well as the hippocampus (*7*). As dementia progresses, the spatial pattern of tau pathology highly correlates with the level of cognitive impairment (*10-13*). In addition, Aβ oligomers and/or plaques can enhance tau pathology in various mouse models (*14, 15*), and there is increasing evidence that accumulation of Aβ plaques can contribute to tau pathology (*3, 16, 17*). The most well-defined physiological role of tau is in stabilizing microtubules, particularly in neuronal axons (*2*). However, in the pathological state, tau becomes aberrantly phosphorylated (*2, 18, 19*), truncated (*1, 4*), and aggregates into oligomers and larger insoluble filaments (*20, 21*). This pathology is thought to trigger synaptic loss, dramatic genome-wide expression changes, increased inflammatory response, and neuronal cell death (*22-25*). These data suggest that pathological tau may be a downstream mediator of the neurotoxic effects leading to neuronal degeneration in AD.

Previously, our lab demonstrated that deleting the histone demethylase *Lsd1* in adult mice leads to significant neuronal cell death in the hippocampus and cortex with associated learning and memory defects (*26*). In this mouse model, loss of *Lsd1* induces genome-wide expression changes that significantly overlap with those observed in the brains of postmortem human AD

cases, but not other neurodegenerative diseases, such as Parkinson's disease or amyotrophic

lateral sclerosis (ALS) cases. Consistent with this overlap, we observed LSD1 protein

mislocalized to cytoplasmic NFTs, but not associated with Aβ plaques in AD cases or Lewy

bodies of α-synuclein in Parkinson's disease cases (26). In control cases, LSD1 remains strictly

confined to the nucleus (26), due to it's well-defined nuclear localization signal (27). These data

highlight the requirement for LSD1 in neuronal survival and suggest that the nuclear function of

the histone demethylase LSD1 could be disrupted by mislocalization to pathological tau.

   To investigate how LSD1 may contribute to tau-mediated neurodegeneration, we utilized

the PS19 P301S tauopathy mouse model (hereafter referred to as PS19 Tau). PS19 Tau mice

express a P301S mutated form of the human tau protein, originally identified in a frontotemporal

dementia with parkinsonism (FTDP-17) patient, driven by the prion promoter throughout the

nervous system (28). When expressed in mice, the P301S tau protein is prone to

hyperphosphorylation and somatodendritic aggregation, without the presence of Aβ plaques.

PS19 Tau mice develop a heavy pathological tau burden and have been well characterized for the

temporal progression of tau pathology and disease-related phenotypes (29, 30). However, the

mechanism of neuronal cell death caused by pathological tau is still unknown.

   Here, we provide functional data that the inhibition of LSD1 function contributes to tau

induced neurodegeneration. We demonstrate in PS19 Tau mice that pathological tau sequesters

LSD1 in the cytoplasm of neurons throughout the brain. This results in depletion of LSD1 from

the nucleus. Additionally, we provide genetic and molecular evidence that pathological tau

contributes to neurodegeneration by disrupting LSD1. Finally, we show that overexpressing

LSD1 in hippocampal neurons is sufficient to suppress neuronal cell death even after

pathological tau has formed. We propose that pathological tau contributes to neuronal cell death

by sequestering LSD1 in the cytoplasm, depleting the nuclear pool of LSD1 that is required for

neuronal survival.

RESULTS

Tau pathology depletes LSD1 from the nucleus in the PS19 Tau mouse model

Previously, we showed in human AD cases that LSD1 protein inappropriately colocalizes with

NFTs in the cell body of hippocampal and cortical neurons, while in unaffected controls LSD1

was properly localized exclusively to the nucleus (*26*). However, because neurons in AD cases

with intracellular NFTs presumably die and are cleared, it was difficult to determine whether tau

prevents LSD1 from localizing to the nucleus in a dying neuron. To address this possibility, we

performed LSD1 immunofluorescence on 12 month old PS19 Tau mice, which have significant

tau pathology (*28*). Because PS19 Tau mice undergo neurodegeneration over a shortened period,

there are more neurons undergoing neurodegeneration at any given time point. Thus, we

reasoned that it may be possible to observe LSD1 depletion from the nucleus. Similar to what we

observe in humans, LSD1 protein in 12 month old Wild Type mice was strictly localized to the

nucleus of neurons in the cerebral cortex (Fig. 1A-C) and the hippocampus (Fig. S1 A-C). This

was expected since LSD1 has a well-defined nuclear localization signal (*27*). However, in 12

month old PS19 Tau mice, LSD1 protein was sequestered in the cytoplasm and depleted from the

nucleus both in the cerebral cortex (Fig. 1D-F,M,N) and the hippocampus (Fig. S1 D-F). These

are both regions where we observe substantial cytoplasmic tau pathology (Fig. 1G-I; Fig. S1 G-

I). Similarly, in other brain regions that accumulate tau pathology, such as the thalamus and

amygdala, LSD1 was localized to the nucleus in 12 month old Wild Type control mice (Fig. S1

J-O), but abnormally localized to the cytoplasm in PS19 Tau mouse littermates (Fig. S1 P-U).

Overall, we observed sequestration of LSD1 in 6 out of 7 mice analyzed. In each of the 6 mice,

there were varying levels of sequestration ranging from LSD1 found in both the nucleus and

cytoplasm (Fig. S1 V-X, M,N), to depletion from the nucleus (Fig. S1 Y-DD, M,N). The mouse

that has little sequestration (denoted by the ^ in Fig. 1M) has extensive tau pathology, but this

pathology is largely extracellular rather than cytoplasmic, indicating that neurons with extensive

LSD1 sequestration may have died and been cleared.

Reduction of LSD1 increases the mouse tauopathy phenotype

If the presence of pathological tau in the cytoplasm is leading to neuronal cell death through the

sequestration and nuclear depletion of LSD1, we would expect that lowering the overall levels of

LSD1 would accelerate depletion and exacerbate the progression of disease. To test this, we

made PS19 Tau mice heterozygous for *Lsd1* (hereafter referred to as PS19;*Lsd1*$^{\Delta/+}$, Fig. S2 A).

Lsd1 was deleted using *Vasa-Cre* (*31*). The *Lsd1* deletion created when the floxed allele

recombines is a null allele that has previously been characterized (*26, 32, 33*). Once the *Lsd1*

deletion passes through the germline, *Lsd1* is heterozygous throughout the mouse. These *Lsd1*

heterozygotes (hereafter referred to as *Lsd1*$^{\Delta/+}$) have a functioning copy of *Lsd1* and don't have

any phenotypes associated with the heterozygous loss of LSD1 function (*33-35*). Thus, *Lsd1*

heterozygosity does not completely compromise LSD1 function. Instead mice that are

heterozygous for *Lsd1* are sensitized to mechanisms affecting LSD1 localization and function.

Consistent with this, we observed a 20% reduction in transcript levels (Fig. S2 B) and a 35%

reduction in protein levels (Fig. S2 C,D) in *Lsd1*$^{\Delta/+}$ mice compared to their *Lsd1*$^{+/+}$ littermates.

Surprisingly, PS19 Tau mice have a 20% increase in LSD1 protein levels compared to *Lsd1*$^{+/+}$

littermates. It is possible that this is due to some type of compensation because LSD1 is being

sequestered. Nevertheless, consistent with the reduction in LSD1 that we observe in $Lsd1^{\Delta/+}$

mice, PS19;$Lsd1^{\Delta/+}$ mice have a similar 14% reduction in transcript levels (Fig. S2 B) and a 31%

reduction in protein levels (Fig. S2 C,D) compared to PS19 Tau littermates. This enables us to

directly assess the effect of reducing LSD1 on tau-induced neurodegeneration. In addition, all

genotypes were born at normal Mendelian ratios with equal male/female ratios.

To determine if reducing LSD1 protein levels accelerates tau-induced depletion of LSD1

from the nucleus, we analyzed localization of LSD1 in PS19;$Lsd1^{\Delta/+}$ mice versus PS19 Tau

controls. When one copy of $Lsd1$ was removed from PS19 Tau mice, there was a significant

decrease in the nuclear localization of LSD1 compared to PS19 Tau mice (Fig. 1J-N),

accompanied by an increase in the number of cells that have LSD1 in both the nucleus and the

cytoplasm (Fig 1J-N). This suggests that reducing LSD1 enables tau to accelerate the depletion

of LSD1.

Since lowering LSD1 accelerates tau-induced depletion of LSD1 from the nucleus, we

determined if this acceleration exacerbates the progression of disease, by analyzing survival,

paralysis, neurodegeneration and genome-wide gene expression changes in PS19;$Lsd1^{\Delta/+}$ mice

versus PS19 Tau controls. As expected, $Lsd1^{\Delta/+}$ mice had normal survival (Fig. 2A). In contrast,

PS19 Tau mice had a reduced overall survival (Fig. 2A) (28). When one copy of $Lsd1$ was

removed from PS19 Tau mice, their reduced survival was significantly exacerbated (P-value =

0.0017, Fig. 2A). As expected, there was little effect on the onset of reduced viability. The initial

decline in the survival of PS19;$Lsd1^{\Delta/+}$ mice started only slightly earlier than PS19 Tau mice, but

after the appearance of hyperphosphorylated (AT8 positive) pathological tau in neurons, as

quantified by the average AT8 positive tau immunoreactivity per area in the hippocampus and

cortex (Fig. S6 K-M)(28). This suggests that pathological tau may have to be present in order for

Lsd1 heterozygosity to have deleterious effects. Subsequently, PS19;*Lsd1$^{Δ/+}$* mice had a 14%

reduction in median lifespan compared to PS19 Tau mice and reached median survival 44 days

earlier then PS19 Tau mice. In addition, there was a further exacerbation of reduced lifespan as

pathology became more severe. PS19;*Lsd1$^{Δ/+}$* mice reached the point when there was only 10%

of the population remaining 83 days earlier than PS19 Tau mice, and all but one of the last 25%

of PS19;*Lsd1$^{Δ/+}$* mice died between 11.5-13.5 months, compared to 13.5-19 months in PS19 Tau

mice. As a result, 28% of PS19 Tau mice were still alive after all but one of the PS19;*Lsd1$^{Δ/+}$*

mice had died (Fig. 2A).

PS19 Tau mice develop paralysis starting with hind limb clasping which progresses until

they are unable to feed (*28*). In our hands*,* PS19 Tau mice displayed intermittent hind limb

clasping starting at approximately 6 months of age. At 12 months, these mice had a severe clasp,

but were still mobile. This is delayed compared to what was originally reported by Yoshiyama

and colleagues (*28*). PS19;*Lsd1$^{Δ/+}$* mice also displayed intermittent hind limb clasping beginning

at approximately 6 months of age. However PS19;*Lsd1$^{Δ/+}$* mice became terminally paralyzed at a

faster rate compared to PS19 Tau mouse littermates. At 12 months, when PS19 Tau mice were

still mobile, PS19;*Lsd1$^{Δ/+}$* mice were severely paralyzed and typically terminal (Movie S1). To

quantitatively assess paralysis we performed rotarod and grid performance tests. In the rotarod,

we assessed the ability of the mice to stay on the rotating rod (latency to fall) (Fig. 2B), the speed

of the rod at which they fall off the rotarod (rotations per minute) (Fig. 2C), and the total distance

traveled (Fig. 2D). All genotypes performed the same at 6 months and 8 months (Fig. 2B-D).

However, at 10 months, when PS19 Tau mice still performed normally, PS19;*Lsd1$^{Δ/+}$* mice had a

significant deficit in mobility (P-value < 0.01, Fig. 2B,C). A deficit in PS19;*Lsd1$^{Δ/+}$* mice was

also observed at 10 months in the total distance traveled (Fig. 2D) and in grid performance

testing (Fig. S3 A), though neither were statistically significant.

To further investigate the exacerbation of paralysis we examined spinal cord motor

neurons. Healthy motor neurons from $Lsd1^{+/+}$ control mice express LSD1 (Fig. S3 B-D) and are

classically identified by circular nuclei at the center of a large cell body. In contrast to the

healthy motor neurons we observed in 12 month old PS19 Tau mice (Fig. S3 E), many of the

motor neurons in PS19;$Lsd1^{\Delta/+}$ mice at 12 months had abnormal morphology, with the nucleus

skewed to the edge of the cell body (Fig. S3 E vs. S3 F) and a ballooned cell body (Fig. S3 F).

Within the cell body we found aberrant hyperphosphorylated NFH (heavy chain neurofilament),

which is a sign of activated neuronal stress pathways (Fig S3 G vs. S3 H) (*36, 37*). This

abnormal morphology is highly reminiscent of a well-established process known as

chromatolysis, which is characterized by swelling of the neuronal cell body, disruption of Nissl

granules, and pyknotic or shrunken nuclei abnormally skewed to the edge of the cell body (*38,*

39). Chromatolysis, which is linked to neuronal stress and often leads to apoptosis (*39*), has been

observed in AD and other neurodegenerative diseases (*38, 40-42*).

Reduction of LSD1 exacerbates PS19 Tau neurodegeneration

In addition to accelerating the paralysis phenotype, reducing the level of *Lsd1* in PS19 Tau mice

exacerbated neuronal cell death in the brain. At 6 months and 8 months, we observed no

difference between genotypes in the overall morphology in the hippocampus (Fig. S4 A-H),

based on histological analysis. There was also no difference between $Lsd1^{+/+}$ and $Lsd1^{\Delta/+}$ mice at

10 months or 12 months (Fig. 3A,B; Fig. S4 I,J,M,N; Fig. S5 A,B,E,F,I,J,M,N). At 10 months,

PS19 Tau mice had very little cell loss in the hippocampus compared to $Lsd1^{+/+}$ and $Lsd1^{\Delta/+}$

control mice (Fig. S4 I-K, M-O). In contrast, at 10 months, PS19;$Lsd1^{\Delta/+}$ mice had dramatic cell

loss both in the CA1 region of the hippocampus and throughout the posterior hippocampus (Fig.

S4 L,P). At 12 months, the PS19 Tau mice had a slight decrease in CA1 and CA3 neurons

spanning the hippocampal pyramidal cell layer compared to $Lsd1^{+/+}$ control mice (17% and

19.4% respectively, Fig. 3A,B; Fig. S5 A-C, E-G). In comparison, PS19;$Lsd1^{\Delta/+}$ mice had a 52%

and 54% decrease in the CA1 and CA3 respectively (Fig. 3A,B; Fig. S5 D,H) compared to

$Lsd1^{+/+}$ control mice. This resulted in decreased overall brain size (Fig. 3C) and brain weight

(Fig. 3D) in PS19;$Lsd1^{\Delta/+}$ mice compared to PS19 Tau and $Lsd1^{\Delta/+}$ mouse littermates.

Additionally, at 12 months there were increased levels of cell loss in the Dentate Gyrus (Fig. S5

I-L), and throughout the posterior hippocampus (Fig. S5 M-P).

Along with the histology, we monitored the progression of neuronal cell death in the

same individual over time by performing magnetic resonance imaging (MRI) at 6 months and

again at 10 months (Movie S2). At 6 months, there was no sign of cell loss or ventricular

dilatation in $Lsd1^{\Delta/+}$, PS19 Tau, or PS19;$Lsd1^{\Delta/+}$ mice (Fig. 3E-G). However, at 10 months the

MRI showed that there was dramatic ventricular dilatation in PS19;$Lsd1^{\Delta/+}$ mice, as evidenced by

high-intensity areas in T2- weighted imaging, with substantial hippocampal and neocortical

atrophy (Fig. 3J vs. 3H, Movie S2). At this timepoint, PS19 Tau mice had some ventricular

dilatation and hippocampal atrophy throughout the hippocampus (Fig. 3I), but much less than

PS19;$Lsd1^{\Delta/+}$ mice (Fig. 3J, Movie S2. 0:51sec vs. 1:00sec).

Tau pathology is not affected by change in LSD1 levels

Since LSD1 is a chromatin regulator, it is possible that reducing LSD1 protein levels affects the

PS19 Tau transgene. However, we confirmed that there was no difference between PS19 Tau

mice and PS19;*Lsd1*$^{\Delta/+}$ mice in the endogenous mouse *Mapt* RNA expression, nor in the human

P301S MAPT transgene expression (Fig. S6 A). It is also possible that LSD1 affects tau

pathology. To test this, we performed immunohistochemistry staining for a hyperphosphorylated

form of tau (AT8). As expected, we did not observe any AT8 positive staining in *Lsd1*$^{\Delta/+}$ at 6, 8,

or 10 months (Fig. S6 B,E,H). Additionally, we observed very little AT8 positive

immunoreactivity at 6 months in both PS19 Tau or PS19;*Lsd1*$^{\Delta/+}$ mice (Fig. S6 C,D). At 8

months both PS19 Tau and PS19;*Lsd1*$^{\Delta/+}$ mice had low but consistent levels of AT8 positive

immunoreactivity (Fig. S6 F,G), and by 10 months both PS19 Tau and PS19;*Lsd1*$^{\Delta/+}$ mice

developed the same high level of AT8 positive tau immunoreactivity (Fig. S6 I,J). This was

consistent throughout both the CA1 and CA3 regions of the hippocampus and the cerebral cortex

(Fig. S6 K-M). We also did not observe any difference between PS19 Tau mice and

PS19;*Lsd1*$^{\Delta/+}$ mice when assaying PHF1 (an alternative phospho-tau antibody) immunoreactivity

in the CA1 region of the hippocampus at 8 months (Fig. S7 A-C, G) and 10 months (Fig. S7 D-F,

G), nor in the CA3 region of the hippocampus or the cerebral cortex at 8 and 10 months (Fig. S7

H,I).

The functional interaction between tau pathology and LSD1 inhibition is specific

To test the specificity of the functional interaction between tau pathology and LSD1, we

investigated the overlap in the effected molecular pathways associated with both pathological tau

and *Lsd1* heterozygosity. To address this, we performed RNA sequencing on the hippocampus of

9 month *Lsd1*$^{+/+}$, *Lsd1*$^{\Delta/+}$, PS19 Tau, and PS19;*Lsd1*$^{\Delta/+}$ littermates. As opposed to analyzing

transcriptional changes at the terminal stage of disease, this time point allows us to assess

molecular changes prior to the onset of neuronal cell death. This is also the time point that we

observed the earliest signs of exacerbation of paralysis in PS19;$Lsd1^{\Delta/+}$ mice. Because of this

early stage in the progression of the disease, we would not expect dramatic changes in

transcription overall. Nevertheless, if tau pathology is inhibiting LSD1 function, we would

expect that the genome-wide expression changes induced by tau might be exacerbated by a

reduction in LSD1. The RNA-seq analysis detected 112 significant gene expression changes in

PS19 Tau mice compared to $Lsd1^{+/+}$ (Fig S8 A,B), and 295 significant gene expression changes

in PS19;$Lsd1^{\Delta/+}$ mice compared to $Lsd1^{+/+}$ (Fig S8 C,D). Importantly, $Lsd1^{\Delta/+}$ mice had only 4

gene expression changes observed genome-wide (Fig. S8 E,F), indicating that the partially

reduced level of LSD1 expression had very little effect on transcription on its own. This is

consistent with the lack of phenotype in these animals.

We first examined the relationship between tau-induced expression changes and the

effects of *Lsd1* heterozygosity by comparing the transcriptional changes observed in PS19 Tau

mice with PS19;$Lsd1^{\Delta/+}$ mice. In PS19 Tau mice that do not yet have significant

neurodegeneration, we identified 112 genes (107 up and 5 down) that were differentially

expressed. All of these 112 genes were changed in the same direction in PS19;$Lsd1^{\Delta/+}$ mice. In

addition, amongst the 112 genes changed in the same direction, 104 (93%) had exacerbated

expression in PS19;$Lsd1^{\Delta/+}$ mice compared to PS19 Tau mice (Fig. 4A). Based on this overlap,

we further compared the expression changes between PS19 Tau mice and PS19;$Lsd1^{\Delta/+}$ mice

genome-wide. Amongst the transcripts that were changed in both PS19 Tau mice and

PS19;$Lsd1^{\Delta/+}$ mice compared to $Lsd1^{+/+}$ mice, 77% changed in the same direction (either up or

down). Consistent with this overlap in gene expression, Gene Set Enrichment Analysis

demonstrated that the pathways that are affected in both PS19 Tau mice and PS19;$Lsd1^{\Delta/+}$ mice

are very similar (Fig. S8 G-J). However, the genes affected in both sets of mice tended to be

further exacerbated in PS19;*Lsd1*$^{\Delta/+}$ mice compared to PS19 Tau mice. Amongst the 77% of

genes that changed in the same direction in both PS19 Tau mice and PS19;*Lsd1*$^{\Delta/+}$ mice, 75% of

these transcripts had a higher fold-change in PS19;*Lsd1*$^{\Delta/+}$ mice compared to PS19 Tau mice

(Fig. 4B). Importantly, the gene expression pathways that are induced by the PS19 transgene

overlap substantially with those observed in human AD case (*43*). This suggests that reduction of

Lsd1 exacerbates human AD pathways.

Overexpression of LSD1 rescues neurodegeneration in the hippocampus of PS19 Tau mice

Our data demonstrate that reduction of LSD1 protein exacerbates the tauopathy phenotype in

PS19 Tau mice. Based on this, we considered the possibility that overexpression of LSD1 might

counter the loss of LSD1 from the nucleus and protect against neurodegeneration in PS19 Tau

mice. To address this possibility, we injected PS19 Tau mice with a neuronal specific virus

(AAV-DJ driven by the synapsin promoter) expressing either the full length LSD1 protein with

an N-terminal HA tag (hereafter referred to as PS19- LSD1 inj) or a control virus expressing

only the HA tag (hereafter referred to as PS19- HA inj). Additionally, to control for the effects of

viral injection, we injected Wild Type littermates with the HA only expressing virus (hereafter

referred to as WT- HA inj). All injections were performed directly into the hippocampus at 8-8.5

months, when tau pathology is already present throughout the nervous system. Immunolabeling

for the HA tag demonstrated that the virus is specific to NeuN+ neurons (Fig. S9 A-D), with no

HA expression observed in IBA+ microglia (Fig. S9 E-H), or GFAP+ astrocytes (Fig. S9 I-L). It

also confirmed that virally expressed LSD1 is nuclear (Fig. S9 M) and confined to the

hippocampus (Fig. S9 N), even at 11-11.5 months, when LSD1 is normally becoming

sequestered to the cytoplasm (Fig. S9 O-R). After 3 months of overexpression, 11-11.5 month

old mice were euthanized, and the brains were analyzed. Injections resulted in a ~6-fold increase

in expression of LSD1 in the hippocampus compared to endogenous LSD1 in the PS19- HA inj

mice, but no increase in the cerebellum (Fig. S9 S,T). As expected, because the viral injections

were restricted to the hippocampus, the mice injected with LSD1 still developed paralysis

(Movie S3). This confirms that the tau transgene is expressed and functioning. Additionally, we

did not observe a difference in total levels of AT8 positive tau immunoreactivity (Fig. S9 U-X).

Therefore, any modulation to the phenotype was not due to changes in tau pathology.

Injected mice were evaluated for cell death by neuronal cell counts in the hippocampus.

Injection of LSD1 virus into the hippocampus of PS19 Tau mice rescued the neurodegeneration

phenotype. At 11 months, compared to WT-HA inj control mice, 70% of the PS19-HA inj mice

had hippocampal cell counts that were below the lowest WT-HA inj control, while none of the

Tau mice injected with the LSD1 virus were below this level. Overall, we observed significantly

more neurons (P-value <0.05) spanning the pyramidal cell layer (84% of WT- HA inj CA1

counts) compared to their PS19- HA inj littermates (59% of WT- HA inj CA1 counts), such that

overall the neuronal cell count in PS19- LSD1 inj mice was not statistically different from the

WT- HA inj (Fig. 5A-D). Additionally, in the histological analysis we observed a large number

of cells infiltrating the hippocampus in PS19- HA inj mice compared to WT- HA inj littermates

(Fig. 5B vs. 5C). Marker analysis demonstrated that this was due to a strong inflammatory

response, with a large increase in the number of GFAP+ astrocytes (Fig. S10 A-C vs. D-F) and

TRL2+ activated microglia (Fig. S10 J-L vs. M-O, S-V vs. W,Z, EE). Injection of PS19 Tau

mice with LSD1 virus rescued this inflammatory response. For example, all but one (9 out of 10

analyzed) of the PS19- LSD1 inj mice had a reduction in the number of GFAP+ astrocytes (Fig.

S10 G-I vs. D-F) and TRL2+ activated microglia (Fig. S10 P-R vs. M-O, AA-DD vs. W-Z, EE).

Of note, the one PS19- LSD1 inj mouse where we did observe increased glial cells, similar to

PS19- HA inj mice, had the lowest neuronal cell count (74% of WT- HA inj CA1 neurons). It is

possible that this mouse was already undergoing neurodegeneration prior to the injection of the

LSD1 overexpression virus.

Although the number of hippocampal neurons in PS19- LSD1 inj mice did not differ

from WT- HA inj controls, in 6 of the 10 PS19- LSD1 inj mice we observed cells with abnormal

blebbed nuclei at varying numbers throughout the hippocampus (Fig. 5E). These abnormal cells

are rare in PS19- HA inj mice, which have a reduced overall number of cells in the pyramidal

cell layer compared to WT- HA inj control mice. One possibility is that these abnormal cells

with blebbing nuclei represent an intermediate state between a healthy neuron and a dying

neuron that is prolonged by rescue via LSD1 overexpression. Interestingly, these abnormal cells

also differed in the localization of HA-tagged LSD1. The four mice with normal nuclei had HA-

tagged LSD1 protein localized uniformly throughout the nucleus (Fig. 5F). In contrast, the six

mice with abnormal nuclear blebbing had some HA-tagged LSD1 that was mislocalized to the

cytoplasm (Fig. 5G). This includes the one PS19- LSD1 inj mouse that had an elevated number

of astrocytes and TRL2 positive microglia. Thus, the blebbing state correlates with when the

viral produced LSD1 begins to be sequestered in the cytoplasm, similar to the endogenously

produced LSD1.

To test the specificity of the rescue that we observe when LSD1 is overexpressed in the

hippocampus of PS19 Tau mice, we performed RNA sequencing on the hippocampus of rescued

11-11.5 month old PS19- LSD1 inj mice versus PS19- HA inj mice. The RNA-seq analysis

detected 144 significant gene expression changes PS19- HA inj mice compared to WT- HA inj

control mice (Fig. S11 A,B), and 57 significant gene expression changes in PS19- LSD1 inj mice

compared to WT- HA inj control mice (Fig. S11 C,D). All of the 144 significant gene expression

changes induced by tau are ameliorated by overexpression of LSD1 (Fig. 6A). As a result, 116

out of the 144 significantly changed genes in the PS19 Tau- HA mice are no longer significantly

changed in the rescued PS19- LSD1 inj mice. Based on this specific rescue of the differentially

expressed genes induced by tau, we further compared the expression changes between PS19-

LSD1 inj mice and PS19- HA inj mice genome-wide (Fig. 6B). Amongst the genome-wide

expression changes in PS19- HA inj mice compared to WT- HA inj control mice, 76% were

changed in the same direction (either up or down), and of these 69% were ameliorated. It should

be noted, that this is the opposite of what we observe when we reduce LSD1 (Fig. 4). Thus, tau-

induced gene expression changes can be specifically modulated in both directions by increasing

or decreasing LSD1 levels. Importantly, the pathways identified by Gene Set Enrichment

Analysis are similar between PS19- LSD1 inj mice and PS19- HA inj mice, and we fail to detect

any new pathways induced by overexpression of LSD1 that could potentially account for a non-

specific rescue (Fig. S11 E-H).

DISCUSSION

In this study we investigate a potential downstream mediator of tau pathology in

neurodegenerative disease. We find that modulation of the chromatin modifying enzyme LSD1

can alter neurodegeneration in a tauopathy mouse model. Previously, we showed that LSD1

colocalizes with tau pathology in the cell body of neurons in AD cases (26). This suggested that

LSD1 might be disrupted in tauopathies such as AD, by being excluded from the nucleus. To

address this directly, we utilize the PS19 tauopathy mouse model. In these mice, we find that

LSD1 is sequestered in the cytoplasm, in some cases being completely depleted from the

nucleus. This provides the first cytological evidence that pathological tau can prevent LSD1 from

properly localizing to the nucleus in hippocampal and cortical neurons, where we have

previously shown it is continuously required (*44*).

Based on the ability of pathological tau to sequester LSD1, we hypothesized that

neuronal cell death may be due, at least partly, to LSD1 being sequestered in the cytoplasm and

depleted from the nucleus. In this case, reducing LSD1 levels should make it easier for tau to

deplete LSD1 from the nucleus, resulting in a faster progression of neurodegeneration and/or a

more severe neurodegenerative phenotype. We find that reducing LSD1 in the PS19 Tau

transgenic background accelerates the depletion of LSD1 from the nucleus compared to PS19

Tau mice alone. Surprisingly, in PS19;*Lsd1$^{\Delta/+}$* mice we also observe a slight decrease in number

of cells in the cortex with LSD1 localized completely to the cytoplasm. However, this is likely

because many of the cells in mice with decreased LSD1 have died and been cleared. Consistent

with this possibility, we observe less cells/area overall in PS19;*Lsd1$^{\Delta/+}$* mice compared to PS19

Tau mice.

LSD1 heterozygosity alone induces only 4 significant gene expression changes and does

not lead to any neurodegeneration. This suggests that any effects observed in PS19;*Lsd1$^{\Delta/+}$* mice

are not simply due to LSD1 haploinsufficiency. Normally, PS19 Tau mice develop paralysis and

neurodegeneration, along with reduced survival. When we reduce LSD1 in the PS19 Tau mice,

mice die significantly earlier, most likely due to the increased rate of paralysis. Additionally,

PS19;*Lsd1$^{\Delta/+}$* mice have increased neuronal cell death and clearance in the hippocampus. This

suggests that pathological tau functions through LSD1 to cause neurodegeneration *in vivo* in

mice.

PS19;*Lsd1*$^{\Delta/+}$ mice have a 31% reduction in LSD1 protein levels compared to PS19 Tau

mice from birth. This reduction should theoretically make mice sensitive to LSD1 depletion from

birth. However, tau pathology starts at 6-8 months in PS19 Tau mice. As a result, if *Lsd1*

heterozygosity is functioning by making it easier for pathological tau to deplete LSD1 from the

nucleus, we would not expect to see any exacerbation until after pathological tau is present. The

exacerbation of the PS19 Tau mouse neurodegenerative phenotype does not occur until after

pathological human tau was present. This suggests that the effect of *Lsd1* heterozygosity requires

the presence of pathological tau, placing LSD1 downstream of tau. Consistent with LSD1 being

downstream of pathological tau, we found no evidence that *Lsd1* heterozygosity affects the

expression of the tau transgene, or the buildup of pathological tau in PS19 Tau mice.

To test whether the functional interaction between pathological tau and reduced LSD1 is

specific, we used RNA-seq to determine whether the downstream molecular pathways altered in

PS19 Tau mice are exacerbated in PS19;*Lsd1*$^{\Delta/+}$ mice. This analysis was performed at the time of

earliest signs of neuronal distress, allowing us to assess molecular changes prior to cell death and

clearance. LSD1 heterozygosity induces only 4 significant expression changes. In addition, the

pathways that are affected in both PS19 Tau mice and PS19;*Lsd1*$^{\Delta/+}$ mice are very similar. This

suggests that reducing LSD1 did not induce any additional neurodegeneration pathways. In

contrast, when LSD1 is reduced in PS19 Tau mice, the genome-wide expression changes

induced by pathological tau are specifically exacerbated. This suggests that the functional

interaction that we observe between pathological tau and reduced LSD1 is occurring through the

tau pathway.

Our previous data implicated LSD1 in the tau-mediated neurodegeneration pathway.

Utilizing the PS19 Tau mouse model, we now show a functional interaction between

pathological tau and LSD1. Importantly, because the tau mutation utilized in PS19 Tau mice

comes from an FTD patient, the functional interaction that we detected may also be relevant to

FTD and other taupathies, such progressive supranuclear palsy, corticobasal degeneration and

chronic traumatic encephalopathy. However, because PS19 Tau mice do not have Aβ plaque

accumulation, this functional interaction would not be relevant to dysfunction induced directly

by pathological Aβ.

Based on the data we uncovered showing a specific functional interaction between

pathological tau and LSD1, we propose the following model (Movie S4): in healthy hippocampal

and cortical neurons, LSD1 is translated in the cytoplasm and transported into the nucleus where

it is continuously required to repress inappropriate transcription. In tauopathy, pathological tau

accumulates in the cytoplasm blocking LSD1 from nuclear import. This interferes with the

continuous requirement for LSD1, resulting in neuronal cell death. It is possible that pathological

tau inhibits LSD1 function indirectly, by interfering with some other process that subsequently

disrupts LSD1 function. However, based on our observations that LSD1 mislocalizes to

cytoplasmic pathological tau in PS19 Tau mice and human AD cases (*44*), we favor a model

where pathological tau directly masks LSD1 from interacting with the nuclear import complex.

Regardless of whether the interaction is direct or indirect, it should be noted that breakdown of

the nuclear pore, which has been observed in human AD cases (*45-47*), would potentially

exacerbate the model that we propose.

This model makes a direct prediction: if tau is predominantly functioning through LSD1,

then increasing the levels of LSD1 should rescue the tau-induced neurodegenerative phenotype.

To address this, we overexpressed LSD1 in the hippocampal neurons of PS19 Tau mice.

Overexpression of LSD1 specifically in hippocampal neurons rescues the neuronal cell death and

limits the inflammatory response. This rescue is neuronal specific, suggesting that the functional

interaction between LSD1 and tau is occurring in neurons. In addition, this rescue occurs despite

there being no effect on tau aggregation. This negates the possibility that the tau transgene is

simply not functioning when LSD1 is overexpressed. The ability of LSD1 overexpression to

overcome tau-mediated neurodegeneration in the presence of pathological tau aggregates,

provides further evidence that pathological tau is functioning through the inhibition of LSD1.

A priori, we would not expect the gene expression changes induced by overexpression of

LSD1 to be related to the gene expression changes induced by pathological tau. However, the

data presented here, along with our previous work (*44*), raise the possibility that pathological tau

may be causing neurodegeneration by sequestering and inhibiting tau. If this is the case, we

would expect that overexpression of LSD1, even after pathological tau is present, would

specifically counteract tau-induced expression changes. Strikingly, we find that overexpression

of LSD1 ameliorates tau-induced gene expression changes genome-wide. This suggests that the

inhibition of LSD1 may be the critical mediator of neurodegeneration caused by pathological tau.

Overexpressing LSD1 should not prevent it from being sequestered. Rather

overexpressing LSD1 should make it more difficult for pathological tau to sequester all of the

LSD1 protein, allowing some LSD1 to be transported to the nucleus. Thus, overexpressing LSD1

would be expected to temporarily rescue the ability of pathological tau to kill neurons.

Consistent with this, LSD1 overexpression delays the effect of pathological tau rather than

permanently rescuing. In 60% of the mice, the surviving neurons have abnormal morphology,

and the overexpressed version of LSD1 is also sequestered. The observation that neurons fail to

maintain their morphology when the overexpressed LSD1 begins to be sequestered in the

cytoplasm provides further support for the model that tau mediates neurodegeneration through

the sequestration of LSD1. Nevertheless, our data suggest that overexpression of LSD1 cannot

permanently overcome pathological tau. To permanently overcome pathological tau, it would

likely be necessary to permanently disrupt the interaction between pathological tau and LSD1.

This work is currently ongoing in the lab. Overall, our data establish LSD1 as a major

downstream effector of tau-induced neurodegeneration. Based on these data, we propose that the

LSD1 pathway is a potential late stage target for intervention in tauopathies, such as AD.

MATERIALS AND METHODS

The mouse strains are described in SI Materials and Methods. All relevant techniques are also

included there. These include mouse tissue fixation, histology and histological studies,

immunohistochemistry and immunofluorescence, protein quantification, quantitative analysis of

paralysis, MRI of brain atrophy, RNA sequencing, RNA sequencing analysis, stereotaxic surgery

and viral infusion, and statistical analysis.

ACKNOWLEDGMENTS

We thank M. Rosenfeld (U.C.S.D) for providing *Lsd1* deletion mice and D. Castrillon for

providing the *Vasa-Cre* mice; N. Seyfried, R. Betarbet, Mr. Gearing, from the Emory ADRC

(P50 AG025688), NINDS Emory Neuroscience Core Facilities (P30NS055077) for tissue

processing and providing PHF1 antibody, J. Alcudia of the Stanford Gene Vector and Virus Core

for the help with virus generation and production, the Georgia Genomics and Bioinformatics

Core, which provided the RNA library preparation and sequencing service, J. Schroeder from

Emory Rodent Behavioral Core for help with Rotatord and Grid Performance assays supported

by the Emory Neuroscience NINDS Core Facilities (P30NS055077) with further support

provided by the Georgia Clinical & Translational Science Alliance of the National Institutes of

Health under Award Number UL1TR002378; and J. Park from the Emory Center for Systems

Imagining for preforming MRI supported by the National Center for Advancing Translational

Sciences of the National Institutes of Health under Award Number UL1TR000454. We would

also like to thank M.J. Rowley and V. Corces for assistance on RNA sequencing analysis, K.

Porter-Stransky and D. Weinshenker for providing PS19 Tau mouse and teaching stereotaxic

surgery procedure, L.R. Lym and Dr. Lerit for assistance with confocal imagining; D.

Weinshenker, A. Levey, C. Bean and T. Caspary for comments on the manuscript and assistance

throughout. Additionally the authors would like to thank fellow Katz lab members for assistance

experimentally and intellectually. **Funding:** A.K.E was supported by the National Institute of

General Medicine training grant (T32 GM008367-26) and an NRSA from National Institute of

Neurological Disorders and Stroke (NINDS) (F31 NS098663-02). D.A.M was supported by a

research supplement to promote diversity in health-related research from the NINDS

(1R01NS087142). S.M.K was supported the NINDS Training in Translational Research in

Neurology grant (5T32 NS007480-17). M.J.R. is supported by the National Institutes of Health

(NIH) Pathway to Independence Award K99/R00 GM127671.The work was supported by a

grant to D.J.K from the National Institute of Neurological Disorders and Stroke

(1R01NS087142). **Competing Interests:** The authors declare that they have no competing

interests. **Data and materials availability:** FastQ files for the RNA sequencing experiments are

bring deposited in the GEO and will be available upon publication. Results from DESEQ2 are

available in Data S1. Correspondence and requests for materials should be addressed to D.J.K.

(djkatz@emory.edu).

REFERENCES

1. M. E. Orr, A. C. Sullivan, B. Frost, A Brief Overview of Tauopathy: Causes, Consequences, and Therapeutic Strategies. *Trends in pharmacological sciences* **38**, 637-648 (2017).
2. Y. Wang, E. Mandelkow, Tau in physiology and pathology. *Nat Rev Neurosci* **17**, 5-21 (2016).
3. D. M. Holtzman *et al.*, Tau: From research to clinical development. *Alzheimers Dement* **12**, 1033-1039 (2016).
4. C. M. Wischik *et al.*, Isolation of a fragment of tau derived from the core of the paired helical filament of Alzheimer disease. *Proc Natl Acad Sci U S A* **85**, 4506-4510 (1988).
5. M. Goedert, C. M. Wischik, R. A. Crowther, J. E. Walker, A. Klug, Cloning and sequencing of the cDNA encoding a core protein of the paired helical filament of Alzheimer disease: identification as the microtubule-associated protein tau. *Proc Natl Acad Sci U S A* **85**, 4051-4055 (1988).
6. J. Hardy, D. J. Selkoe, The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* **297**, 353-356 (2002).
7. S. H. Barage, K. D. Sonawane, Amyloid cascade hypothesis: Pathogenesis and therapeutic strategies in Alzheimer's disease. *Neuropeptides* **52**, 1-18.
8. G. S. Bloom, Amyloid-β and Tau: The Trigger and Bullet in Alzheimer Disease PathogenesisAmyloid-β and TauAmyloid-β and Tau. *JAMA Neurology* **71**, 505-508 (2014).
9. C. Duyckaerts, B. Delatour, M.-C. Potier, Classification and basic pathology of Alzheimer disease. *Acta Neuropathologica* **118**, 5-36 (2009).
10. P. V. Arriagada, J. H. Growdon, E. T. Hedley-Whyte, B. T. Hyman, Neurofibrillary tangles but not senile plaques parallel duration and severity of Alzheimer's disease. *Neurology* **42**, 631-639 (1992).
11. E. Karran, M. Mercken, B. De Strooper, The amyloid cascade hypothesis for Alzheimer's disease: an appraisal for the development of therapeutics. *Nat Rev Drug Discov* **10**, 698-712 (2011).
12. T. Gomez-Isla *et al.*, Neuronal loss correlates with but exceeds neurofibrillary tangles in Alzheimer's disease. *Ann Neurol* **41**, 17-24 (1997).
13. A. Bejanin *et al.*, Tau pathology and neurodegeneration contribute to cognitive impairment in Alzheimer's disease. *Brain* **140**, 3286-3300 (2017).
14. D. E. Hurtado *et al.*, A{beta} accelerates the spatiotemporal progression of tau pathology and augments tau amyloidosis in an Alzheimer mouse model. *Am J Pathol* **177**, 1977-1988 (2010).
15. Z. He *et al.*, Amyloid-beta plaques enhance Alzheimer's brain tau-seeded pathologies by facilitating neuritic plaque tau aggregation. *Nat Med* **24**, 29-38 (2018).
16. S. H. Choi *et al.*, A three-dimensional human neural cell culture model of Alzheimer/'s disease. *Nature* **515**, 274-278 (2014).
17. H. Braak, E. Braak, Frequency of Stages of Alzheimer-Related Lesions in Different Age Categories. *Neurobiology of Aging* **18**, 351-357 (1997).
18. D. P. Hanger, J. C. Betts, T. L. Loviny, W. P. Blackstock, B. H. Anderton, New phosphorylation sites identified in hyperphosphorylated tau (paired helical filament-tau) from Alzheimer's disease brain using nanoelectrospray mass spectrometry. *J Neurochem* **71**, 2465-2476 (1998).
19. M. Morishima-Kawashima *et al.*, Hyperphosphorylation of tau in PHF. *Neurobiol Aging* **16**, 365-371; discussion 371-380 (1995).
20. T. Arendt, J. T. Stieler, M. Holzer, Tau and tauopathies. *Brain Research Bulletin* **126**, 238-292 (2016).

21. R. J. Castellani, G. Perry, Tau Biology, Tauopathy, Traumatic Brain Injury, and
Diagnostic Challenges. *J Alzheimers Dis* **67**, 447-467 (2019).
22. E. Masliah *et al.*, Altered expression of synaptic proteins occurs early during progression
of Alzheimer's disease. *Neurology* **56**, 127-129 (2001).
23. M. Ingelsson *et al.*, Early Abeta accumulation and progressive synaptic loss, gliosis, and
tangle formation in AD brain. *Neurology* **62**, 925-931 (2004).
24. J. W. Kinney *et al.*, Inflammation as a central mechanism in Alzheimer's disease.
Alzheimer's & Dementia: Translational Research & Clinical Interventions **4**, 575-590
(2018).
25. J. M. Rubio-Perez, J. M. Morillas-Ruiz, A review: inflammatory process in Alzheimer's
disease, role of cytokines. *TheScientificWorldJournal* **2012**, 756357-756357 (2012).
26. M. A. Christopher *et al.*, LSD1 protects against hippocampal and cortical
neurodegeneration. *Nature Communications* **8**, 805 (2017).
27. Y. Jin *et al.*, Nuclear import of human histone lysine-specific demethylase LSD1. *J
Biochem* **156**, 305-313 (2014).
28. Y. Yoshiyama *et al.*, Synapse Loss and Microglial Activation Precede Tangles in a
P301S Tauopathy Mouse Model. *Neuron* **53**, 337-351.
29. A. Bellucci *et al.*, Induction of Inflammatory Mediators and Microglial Activation in Mice
Transgenic for Mutant Human P301S Tau Protein. *The American Journal of Pathology*
165, 1643-1652 (2004).
30. M. Iba *et al.*, Tau pathology spread in PS19 tau transgenic mice following locus
coeruleus (LC) injections of synthetic tau fibrils is determined by the LC's afferent and
efferent connections. *Acta Neuropathol* **130**, 349-362 (2015).
31. T. Gallardo, L. Shirley, G. B. John, D. H. Castrillon, Generation of a germ cell-specific
mouse transgenic Cre line, Vasa-Cre. *Genesis (New York, N.Y. : 2000)* **45**, 413-417
(2007).
32. J. Wang *et al.*, Opposing LSD1 complexes function in developmental gene activation
and repression programmes. *Nature* **446**, 882-887 (2007).
33. D. B. Lyons *et al.*, An epigenetic trap stabilizes singular olfactory receptor expression.
Cell **154**, 325-336 (2013).
34. C. T. Foster *et al.*, Lysine-specific demethylase 1 regulates the embryonic transcriptome
and CoREST stability. *Molecular and cellular biology* **30**, 4851-4863 (2010).
35. L. Jin *et al.*, Loss of LSD1 (lysine-specific demethylase 1) suppresses growth and alters
gene expression of human colon cancer cells in a p53- and DNMT1(DNA
methyltransferase 1)-independent manner. *The Biochemical journal* **449**, 459-468
(2013).
36. B. I. Giasson, W. E. Mushynski, Aberrant stress-induced phosphorylation of perikaryal
neurofilaments. *J Biol Chem* **271**, 30404-30409 (1996).
37. Y. L. Liu *et al.*, Alternation of neurofilaments in immune-mediated injury of spinal cord
motor neurons. *Spinal Cord* **47**, 166 (2008).
38. L. D. F. Moon, Chromatolysis: Do injured axons regenerate poorly when ribonucleases
attack rough endoplasmic reticulum, ribosomes and RNA? *Dev Neurobiol* **78**, 1011-1024
(2018).
39. B. G. Cragg, What is the signal for chromatolysis? *Brain Res* **23**, 1-21 (1970).
40. S. Sil, A. R. Goswami, G. Dutta, T. Ghosh, Effects of naproxen on immune responses in
a colchicine-induced rat model of Alzheimer's disease. *Neuroimmunomodulation* **21**,
304-321 (2014).
41. J. Riancho *et al.*, Compensatory Motor Neuron Response to Chromatolysis in the Murine
hSOD1(G93A) Model of Amyotrophic Lateral Sclerosis. *Front Cell Neurosci* **8**, 346
(2014).

42.   A. M. Cataldo, D. J. Hamilton, R. A. Nixon, Lysosomal abnormalities in degenerating
neurons link neuronal compromise to senile plaque development in Alzheimer disease.
Brain Res **640**, 68-80 (1994).

43.   Y.-W. Wan *et al.*, Meta-Analysis of the Alzheimer's Disease Human Brain Transcriptome
and Functional Dissection in Mouse Models. *Cell Reports* **32**, 107908 (2020).

44.   M. A. Christopher *et al.*, LSD1 protects against hippocampal and cortical
neurodegeneration. *Nat Commun* **8**, 805 (2017).

45.   B. Eftekharzadeh *et al.*, Tau Protein Disrupts Nucleocytoplasmic Transport in
Alzheimer's Disease. *Neuron* **99**, 925-940.e927 (2018).

46.   L. G. Sheffield, H. B. Miskiewicz, L. B. Tannenbaum, S. S. Mirra, Nuclear Pore Complex
Proteins in Alzheimer Disease. *Journal of Neuropathology & Experimental Neurology* **65**,
45-54 (2006).

47.   B. Frost, F. H. Bardai, M. B. Feany, Lamin Dysfunction Mediates Neurodegeneration in
Tauopathies. *Curr Biol* **26**, 129-136 (2016).

48.   E. Afgan *et al.*, The Galaxy platform for accessible, reproducible and collaborative
biomedical analyses: 2016 update. *Nucleic Acids Res* **44**, W3-w10 (2016).

49.   E. Afgan *et al.*, The Galaxy platform for accessible, reproducible and collaborative
biomedical analyses: 2018 update. *Nucleic Acids Research* **46**, W537-W544 (2018).

50.   B. B. Gregory R. Warens, Lodewijk Bonebakker, Robert Gentleman, Wolfgang Huber
Andy Liaw, Thomas Lumley, Martin Maechler, Arni Magnusson, Steffen Moeller, Marc
Schwartz, Bill Venables. (2019).

51.   K. Blighe, in *R. package version 1.2.0*. (2019).

52.   J. Wang, D. Duncan, Z. Shi, B. Zhang, WEB-based GEne SeT AnaLysis Toolkit
(WebGestalt): update 2013. *Nucleic Acids Res* **41**, W77-83 (2013).

53.   J. Wang, S. Vasaikar, Z. Shi, M. Greer, B. Zhang, WebGestalt 2017: a more
comprehensive, powerful, flexible and interactive gene set enrichment analysis toolkit.
Nucleic Acids Res **45**, W130-w137 (2017).

54.   Y. Liao, J. Wang, E. J. Jaehnig, Z. Shi, B. Zhang, WebGestalt 2019: gene set analysis
toolkit with revamped UIs and APIs. *Nucleic Acids Res* **47**, W199-w205 (2019).

55.   B. Zhang, S. Kirov, J. Snoddy, WebGestalt: an integrated system for exploring gene sets
in various biological contexts. *Nucleic Acids Res* **33**, W741-748 (2005).

Fig. 1: LSD1 sequestration and tau accumulation in the presence of pathological tau. A-C, Representative immunofluorescence of 12 month old control Wild Type mice showing DAPI (**A**), LSD1 (**B**), and merged (**C**) in the cerebral cortex where LSD1 is localized specifically to DAPI positive nuclei. **D-F**, Representative image of the cerebral cortex in 12 month old PS19 Tau mice. Staining for DAPI (**D**), LSD1 (**E**), and merged (**F**) shows that LSD1 is localized outside the nucleus, and depleted from the DAPI positive nucleus. Arrows denote cells where LSD1 is localized outside of the nucleus, and asterisks denote LSD1 localized specifically to the nucleus. **G-I,** Representative immunofluorescence of 12 month old PS19 Tau mouse with staining for DAPI (**G**), AT8 positive hyper-phosphorylated tau (**H**) and merge (**I**) where hyper-phosphorylated tau accumulates in the cytoplasm of the cell body. Arrowheads denote hyper-phosphorylated tau. **J-L,** Representative immunofluorescence of 12 month old control Tau mice that are heterozygous for *Lsd1* (PS19; *Lsd1$^{\Delta/+}$*) showing DAPI (**J**), LSD1 (**K**), and merged (**L**) in the cerebral cortex. Arrows denote cells where LSD1 is localized outside of the nucleus, asterisks denote LSD1 localized specifically inside the nucleus, and § denotes cells where LSD1 is both nuclear and cytoplasmic. Scale bars=25μm. **M,N**, Quantification of the LSD1 localization represented in **A-F, J-L**.(**M**) Each bar represents a single mouse. LSD1 was scored as being localized to the nucleus, the cytoplasm, or both areas. ^ denotes the one PS19 Tau mouse with low levels of observed cytoplasmic LSD1 localization. (**N**) Combined analysis from **M** in each cell region for Control (WT *n*=3), PS19 Tau (*n*=7), and PS19;*Lsd1$^{\Delta/+}$* (*n*=7) mice. Values are mean ± SEM, two-way analysis of variance (ANOVA) with Tukey's post hoc test *P<0.05, **P<0.01,****P<0.001.

Fig. 2: Reduction of *Lsd1* exacerbates the PS19 Tau mouse paralysis phenotype. A, Lifespan curve showing that no *Lsd1^{Δ/+}* mice died before 18 months (orange, *n*=20). PS19;*Lsd1^{Δ/+}* mice (purple, *n*=44) have a significant reduction in survival compared to PS19 Tau mice with wild-type levels of *Lsd1* (green, *n*=37)(Log-rank Mantle-Cox test ***P<0.005). **B-D**, Rotarod testing of latency to fall (in seconds) (**B**), rotations per minute (when the mouse fell) (**C**), and distance traveled (in centimeters) (**D**) for mice at age 6, 8 and 10 months. *Lsd1^{Δ/+}* (orange, *n*=10,11,14), PS19 Tau (green, *n*=11,22,9), and PS19;*Lsd1^{Δ/+}* (purple, *n*=8,17,11). Values are mean ± SEM (two-way analysis of variance (ANOVA) with Tukey's post hoc test *P<0.05 **P<0.01, ns=not significant).

672
673
Fig. 3: Reduction of *Lsd1* exacerbates neurodegeneration in PS19 Tau mice. A,B, Average nuclei per area in the
CA1 (**A**) and CA3 (**B**) regions of the hippocampus in 12 month old *Lsd1*$^{+/+}$, *Lsd1*$^{Δ/+}$, PS19 Tau, and PS19;*Lsd1*$^{Δ/+}$
mice. Quantification from histology represented in **Fig. S5 A-H**. Values are mean ± SD (**A**, *n*=13 & **B,** *n*=9). **C,**
Representative image of the brains of 12 month old *Lsd1*$^{Δ/+}$, PS19 Tau, and PS19;*Lsd1*$^{Δ/+}$ littermates. **D,** Total brain
weight of 12 month old littermates represented in **Fig. 3C.** Values are mean ± SD (*n*=5). For all graphs: one-way
analysis of variance (ANOVA) with Tukey's post hoc test (two-sided) *P<0.05,****P<0.001. **E-J,** Representative
image of T2- weighted RARE coronal MRI taken from 6 months (**E-G**) and 10 months (**H-jJ**) of age in *Lsd1*$^{Δ/+}$
(**E,H**), PS19 (**F,J**), and PS19;*Lsd1*$^{Δ/+}$ (**G,J**) mice (*n*=3). Arrow denotes region of hippocampal atrophy.

682
683
Fig. 4: Molecular overlap between loss of LSD1 function and tauopathy. A, Histogram ($\log_2$ fold change) of the
112 genes that have significant changes in gene expression in the PS19 Tau mice (green, *n*=4) and their
corresponding gene expression change in PS19;*Lsd1*$^{Δ/+}$ mice (purple, *n=4*). **B,** Scatter plot showing the correlation
between the genome-wide $\log_2$ fold change in gene expression between PS19 Tau and PS19;*Lsd1*$^{Δ/+}$. The most
significantly changed genes in PS19;*Lsd1*$^{Δ/+}$ mice are shown in red (upregulated) and green (downregulated). All
other genes are shown in grey. Dotted line represents 1:1 relationship between gene expression changes in PS19 Tau
vs. PS19;*Lsd1*$^{Δ/+}$. Exacerbated genes fall to the right of the dotted line in the positively correlated quadrant and to
the left of the dotted line in the negatively correlated quadrant. Genes with correlated expression changes are found
in the top right and bottom left quadrants, while genes that do not correlate are found in the opposite quadrants.

693
Fig. 5: LSD1 overexpression rescues the neurodegenerative phenotype in the hippocampus of 11 month old
PS19 Tau mice. A-C, Representative image of H&E stained CA1 region of the hippocampus of 11 month old Wild
Type mice injected with HA control virus (WT- HA inj) (**A**), PS19 Tau mice injected with HA control virus (PS19-
HA inj) (**B**), and PS19 Tau mice injected with *Lsd1* overexpressing virus (PS19- LSD1 inj) (**C**). Square brackets
denote thickness of pyramidal layer of the CA1 of the hippocampus and curvy brackets denote hippocampal region
with or without infiltrating cells. **D,** Quantification of the average number of nuclei in the pyramidal layer of the
hippocampus per area per mouse from histology represented in **Fig. 5A-C**. Values are mean ± SD (*n*=10, one-way
analysis of variance (ANOVA) with Tukey's post hoc test, **p<0.01, ns=not significant). **E,** Representative H&E of
PS19- LSD1 inj mouse with abnormal nuclei blebbing in the CA1 region of the hippocampus. **E'-E''',** High
magnified image of cells denoted by arrows in **Fig. 5E** of individual nuclei that are either abnormally blebbed (**E'**,
E'') or normal (**E'''**). **F,G,** Immunohistochemistry staining of HA(LSD1) in 11 month PS19- LSD1 inj mice. HA is
either localized specifically to the nucleus in all nuclei (**F**) or in only a few nuclei while it is partially sequestered in
the cytoplasm in others (**G**). **F'-F''',** High magnified image of cells denoted by arrows in **Fig. 5F** of nuclear HA
localization in individual nuclei. **G'-G''',** High magnified image of cells denoted by arrows in **Fig. 5G** of individual
nuclei with HA(LSD1) either sequestered to the cytoplasm (**G'**, **G''**) or confined to the nucleus (**G'''**). Scale
bars=50µm.

710
Fig. 6: Molecular overlap between loss of LSD1 function and tauopathy. A, Histogram ($\log_2$ fold change) of the
144 genes that have significant changes in expression in the PS19 Tau- HA inj mice (green, n=3) and their
corresponding gene expression changes in the PS19 Tau- LSD1 inj mice (blue, n=3). **B,** Scatter plot showing the
correlation between the genome-wide $\log_2$ fold change in gene expression between PS19 Tau- Ha inj and PS19
Tau- LSD1 inj. The most significantly changed genes in the PS19 Tau- HA inj mice are shown in red (upregulated)
and green (downregulated). All other genes are shown in grey. Dotted line represents 1:1 relationship between gene
expression changes in PS19 Tau- Ha inj and PS19 Tau- LSD1 inj. Rescued genes fall to the left of the dotted line in
the positively correlated quadrant and to the right of the dotted line in the negatively correlated quadrant. Genes with
correlated expression changes are found in the top right and bottom left quadrants, while genes that do not correlate
are found in the opposite quadrants.
721